Meson - nucleon vertex form factors at finite temperature.

A. Rakhimov^{*a*}, U. Yakhshiev^{*b*} and F.C. Khanna^{*c*} ^{*a*}Institute of Nuclear Physics, Academy of Sciences,

Uzbekistan (CIS)

^bTheoretical Physics Labaratory, Institute of Applied Physics,

Tashkent State University Tashkent-174, Uzbekistan (CIS)

^c Physics Department, University of Alberta,

Edmonton, Alberta T6G 2J1,

and

TRIUMF, 4004 Wesbrook Mall, Vancouver, BC V6T 2A3

Abstract

In this paper the dependence of meson-nucleon-nucleon vertex form factors is studied as a function of termperature. The results are obtained starting from a zero temperature Bonn potential. The temperature dependence of the vertex form factors and radii is studied in the thermofield dynamics, a realtime operator formalism of finite temperature field theory. It is anticipated that these results will have an impact on the study of relativistic heavy-ion collisions as the critical temperature for the phase transition from hadronic to quark-gluon system is approached.

Typeset using REVT_{EX}

I. INTRODUCTION

In-medium properties of hadrons and their interactions is a field of high current interest. For studying phase transitions and thermodynamical properties of the nuclear system under extreme conditions, it is essential to determine the temperature (T) and density (ρ) dependence of the nuclear force. Below a critical temperature T_c , meson exchange picture of nuclear physics provides the natural description for the nucleon - nucleon (NN) interaction. Quarks and gluons are certainly present in all NN interactions, but it is not always necessary to take them explicitly into account especially below a critical temperature ${\cal T}_c$, when the transition into a quark gluon phase takes place. At distances smaller than a fermi, the inner structure of hadrons is probed since it involves the short distance or the high momentum components of the wave function. This structure is typically taken into account by including vertex form factors. In hot and dense matter, the structure of hadrons undergoes changes which should lead to a modification of the meson - nucleon coupling constants or, in general, the form factors. The temperature [1] and density dependence [2] of some coupling constants have been investigated recently, while that of form factors is still fairly unknown and will be studied in the present paper for the first time. For this purpose we shall use the formalism of Thermo Field Dynamics (TFD) in order to obtain the required temperature and density dependence.

The Thermo Field Dynamics (TFD) which was first suggested by Takahashi and Umezawa [3] is a real time operator formalism of finite temperature quantum field theory. In this framework, all the operator formalism of quantum field theory at zero temperature can be extended directly to finite temperature and density. Therefore in TFD it is possible to continue using Wick's theorem and the Feynman diagrammatic approach as in the case of zero temperature theory. An auxiliary field is introduced which leads to a field doublet; the propagators become 2 * 2 matrices, the Feynman rules are now algebraic operational

rules in the space of 2 * 2 matrices. Recently, some attempts to study in medium properties of hadrons at finite temperature have been obtained using TFD [4, 5].

A study of relativistic heavy-ion collisions [6] is expected to lead us to a proper equation of state of hot and dense nuclear matter. The behaviour of nuclear matter as a function of temperature and density is relevant for investigation in nuclear physics, astrophysics, cosmology and particle physics. The phase transition from hadronic system to a quarkgluon plasma phase and the subsequent hadronisation of the quark-gluon plasma phase can provide information on the asymptotic freedom as well as confinement behaviour in Quantum Chromodynamics (QCD). Therefore it is important to study theoretically the behaviour of the hadronic system at high temperatures. The present report attempts to study such a behaviour. In sec. II, a brief review of the TFD is given. In sec. III, we calculate the renormalization of meson nucleon vertices using the Feynman propagators from TFD. The results of calculations and their discussions will be presented in sec. IV.

II. FINITE TEMPERATURE FORMALISM FOR MESON AND NUCLEON PROPAGATORS.

The Thermo Field Dynamics is a real time operator formalism of quantum field theory at finite temperature. The main feature of the TFD is that thermal average of operator \hat{A} is defined as the expectation value with respect to a temperature dependent vacuum, $|0(\beta)\rangle$, which is obtained from the regular vacuum by a Bogoliubov transformation. Therefore, we have

$$\langle \hat{A} \rangle \equiv \operatorname{Tr}(\hat{A}e^{-\beta(H-\mu)})/\operatorname{Tr}(e^{-\beta(H-\mu)}) = \langle 0(\beta)|\hat{A}|0(\beta)\rangle$$
 (1)

where $\beta \equiv 1/k_B T$ with k_B being the Boltzmann constant, H is the total Hamiltonian of the system, and μ is the chemical potential. Due to the doubling of all degrees of freedom, every field is represented in the thermal doublet form in TFD. For example, the thermal doublet for nucleon field is

$$\psi^{(a)}(x) \equiv \left\{ \begin{array}{c} \psi(x) \\ i \ {}^{t} \tilde{\psi}^{\dagger} \end{array} \right\}$$
(2)

where $\psi(x)$ is the ordinary nucleon field and $\tilde{\psi}$ is the doublet partner of $\psi(x)$. As a consequence the thermal propagator of a fermion field is a 2 * 2 matrix defined as

$$iS_F^{(a,b)}(x_1, x_2) = <0(\beta)|T[\psi^{(a)}(x_1)\bar{\psi}^{(b)}(x_2)]|0(\beta)>$$
(3)

The free thermal propagator S_F to be used in perturbation theory can be separated into two parts, i.e. the usual Feynman part S_0 , propagator at zero temperature and the temperature dependent part S_T such that $S_F^{(ab)} = S_0^{(ab)} + S_T^{(ab)}$. For a fermion with mass M, the propagators have the form

$$S_0^{(ab)} = (\not p + M)_{(ab)} \begin{pmatrix} G_0(p^2) & 0\\ 0 & G_0^*(p^2) \end{pmatrix}$$
(4)

$$S_T^{(ab)} = 2\pi i \delta(p^2 - M^2) (\not p + M)_{\alpha\beta} \begin{pmatrix} \sin^2 \theta_{p_0} & \frac{1}{2} \sin 2\theta_{p_0} \\ \frac{1}{2} \sin 2\theta_{p_0} & -\sin^2 \theta_{p_0} \end{pmatrix}$$
(5)

with

$$G_0(p^2) = \frac{1}{p^2 - M^2 + i\epsilon}$$

$$\cos \theta_{p_0} = \frac{\theta(p_0)}{(1 + e^{-x})^{1/2}} + \frac{\theta(-p_0)}{(1 + e^x)^{1/2}},$$
(6)

$$\sin\theta(p_0) = \frac{e^{-x/2}\theta(p_0)}{(1+e^{-x})^{1/2}} - \frac{e^{x/2}\theta(-p_0)}{(1+e^x)^{1/2}},$$

where a and b are Dirac indices and $x = \beta(p_0 - \mu)$. The propagator of the scalar particle with mass m_B is given by

$$\Delta^B(p) = \Delta^B_0(p) + \Delta^B_T(p), \tag{7}$$

$$\Delta_0^B = \begin{pmatrix} D_0(p^2) & 0\\ 0 & -D_0^*(p^2) \end{pmatrix}$$
(8)

$$D_T = -2\pi i \delta(p^2 - m_B^2) \begin{pmatrix} \sinh^2 \phi_{p_0} & \frac{1}{2} \sinh 2\phi_{p_0} \\ \frac{1}{2} \sinh 2\phi_{p_0} & \sinh^2 \phi_{p_0} \end{pmatrix}$$
(9)

with

$$D_0(p^2) = \frac{1}{p^2 - m_B^2 + i\epsilon}$$

$$\cosh \phi_{p_0} = \frac{1}{(1 - e^{-|y|})^{1/2}},$$
(10)

$$\sinh\phi_{p_0} = \frac{e^{-|y|/2}}{(1 - e^{-|y|})^{1/2}},$$

where $y = \beta p_0$. Note that the propagator of vector mesons would be similar to that of the scalar mesons except for an additional factor of $(-g_{\mu\nu} + k_{\mu}k_{\nu}/m^2)$.

III. VERTEX FORM FACTORS

The OBE model of the N-N interaction [7] includes exchange of several mesons. Here we consider the meson-nucleon interactions of π ., σ ., ω - and ρ -mesons:

$$\mathcal{L}_{\pi NN} = -g_{\pi NN} \bar{\psi} i \gamma_5 \vec{\tau} \psi \vec{\varphi}_{\pi} ; \qquad \mathcal{L}_{\sigma NN} = g_{\sigma NN} \bar{\psi} \psi \varphi_{\sigma} ;$$

$$\mathcal{L}_{\omega NN} = -g_{\omega NN} \bar{\psi} \gamma_{\mu} \psi \omega^{\mu} - \frac{f_{\omega NN}}{4M} \bar{\psi} \sigma_{\mu\nu} \psi [\partial^{\mu} \omega_{\nu} - \partial^{\nu} \omega_{\mu}] ; \qquad (11)$$

$$\mathcal{L}_{\rho NN} = \left[-g_{\rho NN} \bar{\psi} \gamma_{\mu} \vec{\tau} \psi \vec{\rho}^{\mu} - \frac{f_{\rho NN}}{4M} \bar{\psi} \sigma_{\mu\nu} \psi [\partial^{\mu} \vec{\rho}_{\nu} - \partial^{\nu} \vec{\rho}_{\mu} \right] .$$

Here g_{BNN} are the meson-nucleon coupling constants and $f_{\omega NN}$ and $f_{\rho NN}$ are the tensor coupling constants for ω - and ρ - mesons respectively.

The corresponding meson nucleon form factors are usually defined as:

$$\langle N(p')|\Gamma_{\pi}^{\alpha}|N(p)\rangle = G_{\pi NN}(t)\bar{u}(p')i\gamma_{5}\tau_{\alpha}u(p)$$

$$\langle N(p')|\Gamma_{\sigma}|N(p)\rangle = -G_{\sigma NN}(t)\bar{u}(p')u(p)$$

$$\langle N(p')|\Gamma_{\omega}^{\mu}|N(p)\rangle = \bar{u}(p')\left[\gamma^{\mu}G_{\omega NN}(t) + \frac{i}{2M}F_{\omega NN}(t)\sigma^{\mu\nu}q_{\nu}\right]u(p)$$

$$\langle N(p')|\Gamma_{\rho}^{\alpha,\mu}|N(p)\rangle = \bar{u}(p')\tau_{a}\left[\gamma^{\mu}G_{\rho NN}(t) + \frac{i}{2M}F_{\rho NN}(t)\sigma^{\mu\nu}q_{\nu}\right]u(p)$$
(12)

where $G_{\pi NN}(t)$, $G_{\sigma NN}(t)$, $G_{\omega NN}(t)$ ($F_{\omega NN}(t)$) and $G_{\rho NN}(t)$ ($F_{\rho NN}(t)$) are form factors of $\pi - , \sigma - , \omega - , \rho - NN$ vertices respectively; $t = q^2 = q_0^2 - \bar{q}^2 = (p - p')^2$ is the 4-momentum transfer and M is the nucleon mass.

In order to investigate the T dependence we calculate three line vertex correction, as is illustrated in Fig. 1 for e.g. $G_{\pi NN}(t,T)$. In accordance with Feynman rules this may be written as:

$$\Gamma_A(t,T) = \Gamma_A(t) + i \sum_B \Lambda_{AB}(t,T),$$
(13)
$$\Lambda_{AB}(t,T) = \int \frac{d^4k}{(2\pi)^4} \Gamma_B(k^2) S_F(a) \Gamma_A(t) S_F(b) \Gamma_B(k^2) \Delta^B(k^2)$$

where
$$A, B$$
 - denote $\pi, \sigma, \omega, \rho$ mesons, e.g. $\Gamma_{\pi} = G_{\pi NN}(t), a = p' - k, b = p - k$. Here $S_F(a)$ and $\Delta^B(k^2)$ are the in - medium nucleon and meson propagators respectively. In the framework of TFD the thermal propagator has a 2 * 2 matrix structure, but only the 11-component refers to the physical field. So, we may use the following representation:

$$S_{F}^{11}(a) \equiv S_{F}(a) = (\hat{a} + M)[G_{0}(a) + G_{T}(a)],$$

$$\Delta^{11} \equiv \Delta(k^{2}) = D_{0}(k^{2}) + D_{T}(k^{2}),$$

$$D_{0}(k^{2}) = \frac{1}{k^{2} - m^{2} + i\varepsilon}, \quad D_{T}(k^{2}) = -2\pi i n_{B}(k)\delta(k^{2} - m^{2}),$$

$$G_{0}(a^{2}) = \frac{1}{a^{2} - M^{2} + i\varepsilon}, \quad G_{T}(a^{2}) = 2\pi i \delta(a^{2} - M^{2})N_{F}(a),$$
(14)

where $N_F(a) = \theta(a_0)n_F(a) + \theta(-a_0)\bar{n}_F(a)$, $\theta(a_0)$ is the step function, *m* is the mass of corresponding meson and

$$n_F(a) = \frac{1}{e^{\beta(|a_0|-\mu)} + 1}, \qquad \bar{n}_F(a) = \frac{1}{e^{\beta(|a_0|+\mu)} + 1}, \qquad n_B(k) = \frac{1}{e^{\beta|k_0|} - 1}$$
(15)

are the fermion, antifermion and boson distribution functions respectively. It is clear that when one substitutes Eq. (14) into Eq. (13), $\Lambda_{AB}(t,T)$ will be separated into two parts: one refers to the naive zero temperature contribution, and the other one depends on T and ρ [4]. We shall concentrate on the latter part, which may be rewritten as follows:

$$\Lambda_{AB}(t,T) = \int \frac{dk^4}{(2\pi)^4} \Gamma_B(k^2) (\hat{a} + M) \Gamma_A(t) (\hat{b} + M) \Gamma_B(k^2) \times$$
(16)

$$\times [G_0(a)G_0(b)D_T^B(k^2) + 2G_0(a)G_T(b)D_0^B(k^2)]$$

Note that we have neglected terms that are quadratic in the temperature dependent distribution function. Now, using (16) and (14) in (13) we get the following expressions for the form factors:

$$G_{ANN}(t,T)/G_{ANN}(t,T=0) = 1 + i\sum_{B} \int \frac{dk^4}{(2\pi)^4} W_{AB}(t,k^2,T) \times [G_0(a)G_0(b)D_T^B(k^2) + 2G_0(a)G_T(b)D_0^B(k^2)]$$
(17)

where the explicit formulas for $W_{AB}(t, k^2, T)$ may be found in the Appendix. Here we write expressions for $W_{\omega\pi}(t, k^2, T)$ and $W_{\rho\pi}(t, k^2, T)$:

$$W_{\omega\pi}(t,k^{2},T) = \frac{3G_{\pi NN}^{2}(k^{2})}{4} \left\{ 2[2M^{2} - (ab)] - \frac{3F_{\omega NN}(t)t}{2G_{\omega NN}(t)} \right\} ;$$

$$W_{\rho\pi}(t,k^{2},T) = \frac{-G_{\pi NN}^{2}(k^{2})}{4} \left\{ 2[2M^{2} - (ab)] - \frac{3F_{\rho NN}(t)t}{2G_{\rho NN}(t)} \right\} ;$$
(18)

where $W_{\omega\pi}(t, k^2, T)$ and $W_{\rho\pi}(t, k^2, T)$ denote the contribution from the π - exchange diagram to the $G_{\omega NN}(t, T)$ and $G_{\rho NN}(t, T)$ respectively.

IV. RESULTS AND DISCUSSIONS

In order to start the calculations, a set of free space meson - nucleon form factors are chosen. We choose the OBE monopole form factors [8] (Bonn A): $G_{BNN}(t) = g_{BNN}(\Lambda_B^2 - t)$ $m_B^2)/(\Lambda_B^2 - t)$, where , for example, $g_{\pi NN}^2/4\pi = 14.09$, and $\Lambda_{\pi NN} = 1005 MeV$. Let us first discuss the T dependence of meson - nucleon coupling constants - $g_{BNN}(T) \equiv G_{BNN}(t =$ m_B^2, T). The variation of the coupling constants $g_{\pi NN}(T)/g_{\pi NN}(T=0), g_{\sigma NN}(T)/g_{\sigma NN}(T=0)$ 0), $g_{\omega NN}(T)/g_{\omega NN}(T=0)$ and $g_{\rho NN}(T)/g_{\rho NN}(T=0)$ with temperature are displayed in Fig. 2. It is clear that they are nearly independent of the temperature below T_c^B , then change rapidly for $T > T_c^B$. A similar behavior was also predicted by Zhang et al. [4] and by Dominguez et al. [1] for pion - nucleon coupling constant $g_{\pi NN}(T)/g_{\pi NN}(T=0)$. But here in Fig. 2, an unexpected result is that, the $\omega - N$ coupling constant, $g_{\omega NN}(T)$ increases while the coupling constants of all other mesons decrease! Let's consider a possible origin of this controversy, by comparing the medium modification of ωNN and ρNN coupling constants. In the present model the modifications arise from the triangle diagrams (Fig. 1). Actual calculations show that, for any $G_{BNN}(t,T)$ the triangle diagram with pion exchange gives a dominant contribution. It is clear from Eq. (18) that at small t the explicit expressions for $g_{\omega NN}(T)$ and $g_{\rho NN}(T)$ formally coincide. However, $W_{\rho\pi} = -\frac{W_{\omega\pi}}{3}$ and hence the two have an opposite sign. The factor (-1/3) arises from the isotopic spin structure. In fact, using Eq. (13) and Fig.1, $g_{\omega NN}(T)$ and $g_{\rho NN}(T)$ may be written in a schematic way:

$$g_{\omega NN}(T) \approx g_{\omega NN}(T=0) + g_{\pi NN}^2 \sum_{\beta} \tau_{\beta} g_{\omega NN} \tau_{\beta} + \dots$$
(19)

$$g_{\rho NN}(T)\tau_{\alpha} \approx g_{\rho NN}(T=0)\tau_{\alpha} + g_{\pi NN}^2 \sum_{\beta} \tau_{\beta}\tau_{\alpha}g_{\rho NN}\tau_{\beta} + \dots$$

where α and β are isospin indices. We omit the spin variables, since ω and ρ have the same spin structure. Now, using $\sum_{\beta} \tau_{\beta}^2 = 3$ and $\sum_{\beta} \tau_{\beta} \tau_{\alpha} \tau_{\beta} = -\tau_{\alpha}$, it is clear that the contribution of the leading triangle diagram with pion exchange to $g_{\omega NN}(T)$ and $g_{\rho NN}(T)$ has different signs. In other words, the term $\Lambda_{\rho\pi}$ in Eq. (13) is negative, while $\Lambda_{\omega\pi}$ is positive.

In general, we conclude from the results (Fig. 2) that at a certain critical temperature T_c^B , the couplings g_{BNN} change drastically. The thermal behavior of $g_{\pi NN}(T) = G_{\pi NN}(t,T)|_{t\to 0}$ has been investigated earlier [1] and we get here a similar behaviour of $g_{\pi NN}(T)$ as a function of temperature. The temperature T_c , where $g_{\pi NN}(T)$ changes dramatically, was interpreted in [1] as a signal for the quark - gluon deconfinement phase transition. Indeed, near T_c the associated mean square radius $\langle r_{BNN}^2 \rangle = 6 \left[\frac{\partial}{\partial t} \ln G_{BNN}(t,T) \right] |_{t \to 0}$ is a monotonically increasing function of T and in fact diverges at the critical temperature. A similar behavior of $\langle r_{BNN}^2 \rangle$ is also found in our calculations and is illustrated in Fig. 3. As the critical temperature, T_c^B , is approached the strength of coupling of π, σ and ρ mesons to nucleons is quenched, at the same time, the size of nucleons as probed by the appropriate meson gets bigger. However, it is to be stressed that T_c^B is not the same for all mesons: $T_c^{\pi} \approx 360 MeV$, $T_c^\sigma \approx 95 MeV, \ T_c^\omega \approx 175 MeV$ and $T_c^\rho \approx 200 MeV$. In OBE picture this means that , in the temperature region e.g. 200 MeV < T < 300 MeV, the ρ meson exchange is no longer important while the pion exchange is still important. On the other hand, the σ and π mesons are mainly responsible for the attraction between nucleons. So, the quenching of the σNN coupling constant at $T > T_c^{\sigma}$ leads to the vanishing of the bound state, as it was predicted earlier [4].

The density dependence of vertex form factors has been studied in greater detail [2, 8].

It was found that most of the vertex form factors are quenched at high densities and it was anticipated that the temperature dependence is likely to yield results that are qualitatively similar to those of density dependence. Now, let's consider their temperature dependence in some detail. The form factors as a function of momentum transfer at several temperatures are displayed in Fig. 4. It is seen that, the in-medium effects lead to the suppression of $G_{\pi NN}(t,T)$, $G_{\sigma NN}(t,T)$ and $G_{\rho NN}(t,T)$. The temperature dependence of $G_{\omega NN}(t,T)$, is opposite to that of $G_{\rho NN}$ due to it's isotopic structure as outlined above.

For practical calculations a parametrization of these $G_{BNN}(t, T, \rho)$ form factors is needed. At small momentum transfer we can parametrize them by a monopole form :

$$G_{BNN}(t,T,\rho) = g_B(T,\rho)(\Lambda_B^2(T,\rho) - m_B^2) / (\Lambda_B^2(T,\rho) + t)$$
(20)

where in general the effective mass of a meson, m_B , is also temperature dependent. Consideration of this dependence is beyond the scope of the present paper. It will be carried out in a self consistent way, taking into account both the self energy and the triangle diagrams, in a forthcoming paper. But here, for simplicity, one may consider this as just a parametrization and choose the parameters $g_B(T, \rho)$ and $\Lambda_B^2(T, \rho)$. Their temperature dependence is still unknown. Here the T dependence maybe represented, for $T \ll T_c^B$, in a polynomial form as:

$$\frac{g_B(T,\rho)}{g_B(T=0,\rho=0)} = 1 - \alpha_B^g(\rho)(T/T_c^B)^2 - \beta_B^g(\rho)(T/T_c^B)^4$$

$$\frac{\Lambda_B(T,\rho)}{\Lambda_B(T=0,\rho=0)} = 1 - \alpha_B^\lambda(\rho)(T/T_c^B)^2 - \beta_B^\lambda(\rho)(T/T_c^B)^4$$
(21)

The calculated form factors are fitted to this form and the paramters α and β are determined. The results are presented in tables I, II, and III for densities $\rho = 0$, $\rho = \rho_0$ and $\rho = 3\rho_0$, respectively (with $\rho_0 = 0.17 f m^{-3}$ - the density of normal nuclear matter). The results show that the density dependence of parameters, and hence, the form factors is rather weak. What is the reason for such a result? The answer may be found by considering expression in Eqs. (14) - (17). It is clear that the density, ρ , influences only through the chemical potential μ defined from the particle and antiparticle distribution functions as

$$\rho = \frac{4}{(2\pi)^3} \int d\vec{k} [n_F(k) - \bar{n}_F(k)], \qquad (22)$$

where μ is involved in Eq. (15) in the exponent. The argument of this exponent is $\beta(|a_0|\pm\mu)$. Since $|a_0| \approx M = 939 MeV$ is much larger at moderate densities than μ ($\mu \approx 40 MeV$ for $\rho = 3\rho_0$, T = 150 MeV), the result is weakly sensitive to μ .

In summary, we have considered the temperature dependence of meson - nucleon form factors and coupling constants in TFD formalism. It is shown that at a critical temperature, where the coupling constant changes drastically and the associated mean square radius diverges is different for different mesons. The temperature dependent vertex form factors are parametrized in a simple monopole form and the T-dependence of these parameters is clarified. These form factors may be used in the calculation of the in - medium NN cross sections [9] and in investigations of the properties of hot dense matter.

In the present calculations, the monopole form factors with the parameters (at T = 0) given by the Bonn group, [7] are used. But the next question which may arise is: are the results sensitive to the shape of the input form factors? To answer this question a different set of form factors obtained by Meissner et. al., [10] in the framework of a topological soliton model have been used. The thermal behaviour of such form factors is shown in Fig. 5. Now, comparing Fig. 4 and Fig. 5 it is clear that only the σNN form factor is affected the most. The reason is that even in free space and at T = 0 the σNN form factor in the topological soliton model [10] is much harder ($\Lambda_{\sigma} \approx 2700 MeV$) than those of the Bonn potential ($\Lambda_{\sigma} \approx 2000 MeV$). Nevertheless our main conclusions about thermal behavior of different meson nucleon vertices remain valid.

The variation of the meson-nucleon coupling constant with temperature (Fig. 1) clearly

indicates that the attractive NN force due to σ -exchange decreases quite rapidly while the repulsive N-N force due to ω -exchanges increases. This would lead to the fact that the nuclear matter is quite likely un-bound at high temperature. In fact it will look more like a hard sphere gas since the repulsive interaction at short distances will dominate. The attractive interactions in such a case play only a small perturbative role. The hot and dense nuclear matter could be approximated as a free gas with an excluded volume around each nucleon. Perhaps the extensive studies [11] along this line can be justified on the basis of the results obtained in this paper. The nuclear matter would be hard to compress.

It would be quite interesting to know, if the critical temperature, T_c^B , maybe uniquely considered as a point of transition from the hadron state to a quark gluon state, as it was suggested [1]. The usual way of estimating T_c is based on QCD lattice calculations [12] or temperature dependent quark - gluon potentials [13]. In such calculations it is usual to start from high temperatures and then decreases the temperature, looking for a phase transition from the quark gluon state to a hadron state i.e. hadronization of the quarks. Clearly this method gives a unique T_c , since at any stage there is either a hadron or a quark gluon plasma phase. Contrary to such studies the present calculations explore the transition from a hadron to a quark-gluon phase and find that the critical temperature for hadron \rightarrow quark-gluon transition depends on the kind of hadrons under consideration. A similar result, where hadrons and quark - gluon plasma coexist have been obtained in lattice calculations [12]. In conclusion it is important to emphasize that the dynamics of hadrons and of quark-gluon plama at finite temperature is poorly understood. The question of critical phenomenon, in particular critical temperature, in the hadronic systems is not well-known. Studies in nuclear matter are needed urgently to clarify the critical transition from a hadronic system to the quark gluon phase which will depend on both density and temperature. The experiments with relativistic heavy ion colliders would need such an understanding to clarify the production of quark-gluon plasma from hadrons and the eventual hadronisation of the quark-gluon system. It is possible that there will be clear and unambiguous signals for the formation and subsequent hadronisation of the quark-gluon plasma [14].

ACKNOWLEDGEMENTS.

We are indebted to Prof. M.M. Musakhanov, Prof. A. Nakamura and Prof. W.M. Alberico for their useful discussions. We thank Prof. C.P. Singh for a critical reading of the manuscript. U. Yakhshiev acknowledges partial support by the grant 11/97 of the State Committee for Science and Technology of Uzbekistan. The research of F.C. Khanna is supported in part by the National Sciences and Engineering research council of Canada.

REFERENCES

- [1] C.A. Dominguez, C. van Gend and M. Loewe Phys. Lett. B429, 64, (1998).
- [2] A.M. Rakhimov, F.C. Khanna, U.T. Yakhshiev and M.M. Musakhanov Nucl. Phys. A643, 383, (1998).
- [3] Y. Takahashi and H. Umezawa, Collective Phenomenon 2, 55, (1975) reprinted in Int.
 J. Mod. Physics B10, 1755, (1996); H. Umezawa et. al., Thermo Field Dynamics and condensed states (North- Holland, Amsterdam, 1982); H. Umezawa, Advanced Field Theory (American Institute of Physics, New York, 1993); P.A. Henning, Phys. Rep. 253, 235, (1995).
- [4] Yi Jun Zhang, Song Gao and Ru Keng Su Phys. Rev. C56, 3336, (1997) and references there in.
- [5] L. Alvarez Ruso, P.F. de Cordoba and E. Oset, Nucl. Phys. A606, 407, (1996); Y. Kim, H.K. Lee nucl-th/9609020.
- [6] J.W. Harris, Proc. Lake Louise Winter Institute, Ed. A. Astbury et. al., (World Scientific, 1999). This review contains numerous references to both experimental and theoretical articles.
- [7] R. Machleidt Adv. Nucl. Phys. 19, 189, (1989).
- [8] Ulf-G. Meissner, Nucl. Phys. A503, 801, (1989).
- [9] G.Q. Li and R. Machelidt Phys. Rev. C 48, 1702, (1993).
- [10] U. Meissner, A. Rakhimov, U. Yakhshiev nucl-th/9901067.
- [11] R. Hagedorn and J. Rafelski, Phys. Lett. B97, 136, (1980); R. Hagedorn, Z. Phys. C17, 265, (1985); C.P. Singh, B.K. Patra and K.K. Singh, Phys. Lett. B387, 680, (1996); J.

Sollfrank, J. Phys. G23, 1903, (1997).

- [12] H. Satz talk at Int. Conf. on Physics and Astrophysics of Quark-Gluon plasma, Ed.B.C. Sinha et. al., (Naros publications, 1998).
- [13] W.M. Alberico, M. Nardi and S. Quatrocolo Nucl. Phys. A589, 620, 1995.
- [14] C.P. Singh, Phys. Rep. 236, 147 (1993); S.A. Bass, M. Gyulassy, H. Stoker and W. Greiner, J. Phys. G25, R1 (1999) and the references therein.

APPENDIX

Here explicit expressions for $W_{AB}(t, k^2, T)$ introduced in eq. (7) are given in detail.

$$\begin{split} W_{\pi\pi} &= G_{\pi}(k^{2})[M^{2} - (ab)]; \quad W_{\pi\sigma} = W_{\pi\pi}|_{G_{\pi} \to G_{\sigma}}; \\ W_{\pi\omega} &= A_{\pi}^{\omega}[M^{2} - (ab)] + 3F_{1}^{\omega}(k^{2})F_{2}^{\omega}(k^{2})z; \quad W_{\pi\rho} = -W_{\pi\omega} \begin{vmatrix} F_{i}^{\omega} \to F_{i}^{\rho} \\ A_{\pi}^{\omega} \to A_{\pi}^{\rho} \end{vmatrix}; \\ A_{\pi}^{\omega} &= [F_{1}^{\omega}(k^{2})]^{2}[4 - \frac{k^{2}}{m_{\omega}^{2}}] + \frac{3k^{2}[F_{2}^{\omega}(k^{2})]^{2}}{4M^{2}}; \\ W_{\sigma\pi} &= -3g_{\pi}^{2}(k^{2})[M^{2} + (ab)], \quad W_{\sigma\sigma} = -\frac{1}{3}W_{\sigma\pi}|_{G_{\pi} \to G_{\sigma}}, \\ W_{\sigma\omega} &= [M^{2} + (ab)]A_{\sigma}^{\omega}; \quad W_{\sigma\rho} = 3W_{\sigma\omega}|_{A_{\sigma}^{\omega} \to A_{\sigma}^{\rho}}; \\ A_{\sigma}^{\omega} &= -[F_{1}^{\omega}(k^{2})]^{2}[4 - \frac{k^{2}}{m_{\omega}^{2}}] + \frac{3k^{2}[F_{2}^{\omega}(k^{2})]^{2}}{4M^{2}}; \\ W_{\omega\pi} &= \frac{3g_{\pi}^{2}(k^{2})}{4}\left\{2[2M^{2} - (ab)] - \frac{3F_{2}^{\omega}(t)t}{2F_{1}^{\omega}(t)}\right\}; \quad W_{\omega\sigma} = \frac{1}{3}W_{\omega\pi}|_{G_{\pi}} \to G_{\sigma}; \\ W_{\omega\omega} &= \frac{1}{4}\left\{2\Gamma_{1}^{\omega}[2M^{2} - (ab)] + \frac{\Gamma_{2}^{\omega}}{M^{2}}[k^{2}[M^{2} - (ab)] + 2(ak)(bk)] + \\ +3z\Gamma_{3}^{\omega} - \frac{F_{2}^{\omega}(t)}{2M^{2}F_{1}^{\omega}(t)}[3\Gamma_{1}^{\omega}t + \Gamma_{2}^{\omega}(k^{2}t - z^{2})] + \\ + \frac{\Gamma_{3}^{\omega}}{M^{2}}[z[(ab) + 3M^{2}] - 2(ak)(bq) - 2(aq)(bk)]\right\}; \end{split}$$

$$\begin{split} \Gamma_{1}^{\omega} &= [F_{1}^{\omega}(k^{2})]^{2} [2 - \frac{k^{2}}{m_{\omega}^{2}}] + \frac{k^{2} [F_{2}^{\omega}(k^{2})]^{2}}{4M^{2}}, \\ \Gamma_{2}^{\omega} &= \frac{2M^{2}}{m_{\omega}^{2}} [F_{1}^{\omega}(k^{2})]^{2} - [F_{2}^{\omega}]^{2}; \\ \Gamma_{3}^{\omega} &= F_{1}^{\omega}(k^{2}) F_{2}^{\omega}(k^{2}), \qquad W_{\omega\rho} = 3W_{\omega\omega}|_{\Gamma_{i}^{\omega} \to \Gamma_{i}^{\rho}}, \qquad W_{\rho\pi} = -\frac{1}{3} W_{\omega\pi}|_{\omega \to \pi}; \\ W_{\rho\sigma} &= -W_{\rho\pi}|_{G_{\pi} \to G_{\sigma}}; \qquad W_{\rho\omega} = W_{\omega\omega} \left|_{\Gamma_{i}^{\omega} \to \Gamma_{i}^{\omega}}; \qquad W_{\rho\rho} = -W_{\omega\omega}|_{\omega \to \rho}; \\ F_{i}^{\omega} \to F_{i}^{\rho} \end{split}$$

where $G_{\pi} \equiv G_{\pi NN}(k^2)$, $G_{\sigma} \equiv G_{\sigma NN}(k^2)$, $F_1^{\omega} \equiv G_{\omega NN}$, $F_2^{\omega} \equiv F_{\omega NN}$, z = (kq) and $q^2 = t$.

FIGURE CAPTIONS

- Fig. 1. Feynman diagrams for pion nucleon vertex, The solid line is for nucleon. Dashed, dotted, dot dashed and wavy lines for π , σ , ω and ρ mesons.
- Fig. 2. The ratio of meson-nucleon coupling constants at finite temperature T and at T = 0 as a function of temperature, T.
- Fig. 3. The ratio of mean square radius at finite temperature T and at T = 0 as a function of temperature, T.
- Fig. 4. Meson nucleon form factors at several temperatures as a function of q^2 using parametrisation of BONN group [7]. a) πNN vertex; b) σNN vertex; c) ωNN vertex and d) ρNN vertex.
- Fig. 5. Meson-nucleon form factors at several temperatures as a function of q^2 using parametrisation of Meissnar et al [10]. a) πNN vertex; b) σNN vertex; c) ωNN vertex and d) ρNN vertex.

TABLES

Meson	α_q	β_q	α_{λ}	eta_λ	T_c
π	-0.201	0.884	0.6237	-0.08419	≈ 360
σ	0.3252	0.8381	-0.3245	0.9184	≈ 95
ho	0.4384	0.03194	1.198	-0.5245	≈ 200
ω	-0.8951	-0.2235	0.7322	-0.5258	≈ 175

TABLE I. Parameters of vertex form factors in Eq.s (20)-(21) at zero density $\rho = 0$.

TABLE II. The same as in Table I , but for $\rho=\rho_0$

Meson	α_g	eta_{g}	$lpha_{\lambda}$	eta_λ	T_c
π	-0.1966	0.8899	0.6181	-0.0750	≈ 360
σ	0.3254	0.8378	-0.3301	0.9329	≈ 95
ho	0.4379	0.03347	1.2734	-0.6071	≈ 200
ω	-0.8966	-0.2168	0.7332	-0.5366	≈ 175

TABLE III. The same as in Table I , but for $\rho=3\rho_0$

Meson	$lpha_g$	$eta_{m{g}}$	$lpha_{\lambda}$	eta_λ	T_c
π	-0.1868	0.9012	0.6056	-0.05452	≈ 360
σ	0.3258	0.8366	-0.3429	0.9674	≈ 95
ho	0.4379	0.03347	1.1934	-0.5047	≈ 200
ω	-0.8994	-0.2026	0.7350	-0.5609	≈ 175



Fig. 1

Fig.2.



Fig.3.

















