

Determination of apparent activating energy of solubilizing MLV-DPPC and MLV-egg-yolk lecithin with surfactants by rectangular optic diffusimetry

ȘTEFAN HOBAI

Department of Biochemistry, University of Medicine and Pharmacy, Tg. Mureș, Romania

Correspondence address:

Ștefan Hobai, Department of Biochemistry, University of Medicine and Pharmacy, Târgu Mureș, Romania. Fax: 040-065-210407, E-mail: fazy@netsoft.ro

Abstract

It were determined the apparent activating energy of dipalmitoylphosphatidylcholine (DPPC) multilamellar liposome (MLV) solubilizing with sodium deoxycholate (DOCNa) by turbidimetric measurements and the apparent activating energy of egg-yolk lecithin (EYL)-MLV solubilizing with different surfactants (DOCNa, TX-100, CTMB) by rectangular optic diffusimetric measurements. The apparent activating energy of MLV-DPPC solubilizing with DOCNa is $E_a = 127.7\text{kJ/mol}$. The apparent activating energy of MLV-EYL solubilizing with cetyltrimethylammonium bromide (CTMB) is 22.3kJ/mol and with DOCNa is 47.3kJ/mol (average value) and with Triton X-100 is 99.3kJ/mol . On the basis of the above data it can be said that besides the utilisation of the liposomes in the modelling of cellular membranes, they can be used as experimental models, too, in order to investigate the solubilization capacity of surfactants.

Introduction

The interaction of various types of surfactants with lamellar phospholipid aggregates has been described in the literature [4,5,6]. Surfactants have been applied to study the size and structure of the formed mixed micelles [7,8,9] and the kinetics and the mechanism of membrane solubilization [2,3]. In some reviews is report the different mechanism of solubilization multilamellar and unilamellar liposomes [1,5].

Materials

Dipalmitoylphosphatidylcholine($\text{di-C}_{16}\text{PC}$) (DPPC) was purchased from Sigma Chemical Co.

Egg yolk lecithine was purchased from Sigma Chemical Co. type X-E (cat.no.P5394/1996) purified by us by neutral alumina column chromatography, verified by thin layer chromatography (silicagel). The mobile phase was a mixture $\text{CHCl}_3:\text{CH}_3\text{OH}:\text{H}_2\text{O}$ (65:30:4,vol). The identification of the phospholipid was made with iodine and shows a single spot. Sodium deoxycholate (DOCNa) was purchased from Merck, the Triton X-100 (TX-100) from Sigma Chemical Co. and Cetyltrimethylammoniumbromid (CTMB) from Fluka AG. Tris(hydroxymethyl)aminomethane-HCl (Tris-Cl) was purchased from Australanal.

All chemicals and inorganic solvents were of analytical and of spectroscopic grades, respectively.

Preparation of MLV-DPPC and MLV-egg-yolk lecithin (EYL) suspensions

2.5 mg DPPC (SIGMA) dissolved in 0.2 ml ethanol (redistilled on P₂O₅) or 0.2ml EYL (78% purity from SIGMA purified by us up to thin layer chromatographic single spot iodine) in chloroform:methanol (9:1,vol) solution which is in a glass test tube was evaporated under a nitrogen stream until it reached the consistence of a lipid film on the wall tube settled down. The film was purged with the gas for 30 minutes in order to eliminate the alcohol traces. The film hydration and bilayer suspension were made with sucrose (CHEMAPOL) solution of 16.6% at 0.6 mg/ml (lipid/solution) by vortexing 10 sec. above the chain-melting temperature ($T_m \sim 41,5^\circ\text{C}$).

Methodology

1. By using a spectrophotometer Specol and K-201 I titrated samples of 1.5ml solution of MLV-DPPC 0.6mg/ml (total 1.23 μmol phospholipid) with DOCNa 20mM at a 4 $\mu\text{l}/\text{sec}$ titration rate up to a total addition of 16 μmol surfactant. The titration was made at several temperatures between 16 and 22 °C.
2. By using a spectrophotometer Specol with FR optical diffusion system and photocells and voltage source type HQE 40 and UV lamp type OSRAM it was possible to determine the rectangular optic diffusion at $\lambda=437\text{nm}$ of some MLV-egg-yolk lecithin suspension (1.4ml, 0.7 μmol lipid/sample) in the course of the titration with the surfactant solution. The titration rate was 0.08 μmol surfactant/sec. The tables below include the experimental data concerning the total volume of titrant (expressed in μl), the total quantity of surfactant that was added (expressed in μmol) and modification of the percentage diffusion calculated by us being 1cm, as the deflexion unity of the recording pen ($\Delta\text{Dif}^\circ/\text{cm}$). $\Delta\text{Dif}^\circ/\mu\text{mol}$ surfactant is the variation of the unity of percentage diffusion, calculated for the 1 μmol surfactant addition.

Results and discussion

1. It is known that the titration curves in the transmission system of SUV-egg-yolk lecithin have a sigmoidal form, and in the case of MLV-DPPC they are straight.

Table 1 includes the experimental data obtained by turbidimetric titrations.

Table 1. Variation speed of optical transmission percentage (Tr%) during the titration of the MLV-DPPC solutions with DOCNa at different temperatures.


Nr. samples	t (°C)	T (K)	$10^3/T$	ΔTr %	$\ln(\Delta Tr \%)$
1	16	289	3.458	3.36	1.21
2	17.1	290.1	3.445	5.08	1.62
3	18.4	291.4	3.430	6.35	1.85
4	20.2	293.2	3.409	6.98	1.94
5	21	294	3.400	9.44	2.24
6	22	295	3.389	10.95	2.39

Table 2 presents the necessary data for the calculation of the regression line of dependence $\ln(\Delta Tr\%) \sim f(10^3/T)$ necessary to determine the activating energy of MLV-DPPC solubilization with DOCNa.

Table 2. Statistical data of titration with DOCNa 20mM with a rate 4 μ l/sec of 1.5ml MLV-DPPC (1.23 μ mol of phospholipid) at different temperatures. $X_n=10^3/T$; $Y_n= \ln(\Delta Tr\%)$. T-absolute temperature (K); $\Delta Tr\%$ - variation of optical transmission percentage. The horizontal bar indicates the mean values. Σ is the sum of the individual values.

n	X_n	Y_n	$X_n - \bar{X}$	$Y_n - \bar{Y}$	$(X_n - \bar{X})(Y_n - \bar{Y})$	$(X_n - \bar{X})^2$	$(Y_n - \bar{Y})^2$
1	3.389	2.39	-0.032	0.515	-0.01648	0.001024	0.265
2	3.400	2.24	-0.021	0.365	-0.007665	0.000441	0.1332
3	3.409	1.94	-0.012	0.065	-0.00078	0.000144	0.00422
4	3.430	1.85	0.009	-0.025	-0.000225	0.000081	0.000625
5	3.445	1.62	0.024	-0.255	-0.00612	0.000576	0.065
6	3.458	1.21	0.037	-0.665	-0.0246	0.001369	0.4422
Σ	20.531	11.25	-	-	-0.055875	0.003635	0.91027
mean	3.421	1.875					

Figure 1. Arrhenius logarithmic representation of change of speed of optical transmission percentage (logarithmic) of sample MLV-DPPC during the titration with DOCNa at different temperatures. The straight line in the middle of experimental points was obtained by linear regression. Statistical data appear in Table 2.



The correlation coefficient (r) was calculated based on data in Table 2.

$$r = \frac{\sum_{1}^{6} (X_n - \bar{X})(Y_n - \bar{Y})}{\sqrt{\sum_{1}^{6} (X_n - \bar{X})^2 \sum_{1}^{6} (Y_n - \bar{Y})^2}} = \frac{-0.055875}{\sqrt{0.003635 \cdot 0.91027}} = \frac{-0.055875}{0.05752} = -0.9714 \quad (1)$$

The correlation coefficient value indicates a good linear correlation between the two experimental data X_n and Y_n . The calculus of the straight line of linear regression produced the following data:

$$a = \frac{\sum_{1}^{6} (X_n - \bar{X})(Y_n - \bar{Y})}{\sum_{1}^{6} (X_n - \bar{X})^2} = \frac{-0.055875}{0.003635} = -15.37 \quad (2)$$

$$b = \bar{Y} - \bar{X} \cdot a \approx 1.875 - 3.421(-15.37) = 54.45 \quad (3)$$

The expression of straight line regression is:

$$Y = -15.37X + 54.45 \quad (4)$$

Straight line regressin slope is equal to E_a / R ratio, so the apparent activating energy of solubilization is $E_a = R \cdot 15.37 = 127.7 \text{ kJ/mol}$.

2. Determination of apparent activating energy of solubilizing MLV-EYL with surfactants (DOCNa, TX-100, CTMB) by rectangular optic diffusimetry was carried out by the same procedure as above.

Table 3. Titrant: DOCNa 20mM; $t = 17.5^\circ\text{C}$; $\Delta\text{Dif.}\% / \text{cm} = 5.5$

	$\mu\text{l DOCNa}$	$\mu\text{mol DOCNa}$	$\Delta\text{Dif.}\% / \mu\text{mol DOCNa}$
1	-	-	-
2	12	0.24	22.92
3	8	0.16	34.37
4	8	0.16	34.37
5	24	0.48	11.46
6	16	0.32	17.19
7	10.4	0.2	27.50
8	29.6	0.6	9.16
9	28	0.56	9.82
10	28	0.56	9.82
Mean $\Delta\text{Dif.}\% / \mu\text{mol DOCNa} =$			19.7
$t = 22.5^\circ\text{C}$; $\Delta\text{Dif.}\% / \text{cm} = 5.68$			
1	-	-	-
2	8	0.16	35.5
3	20	0.4	14.2
4	8	0.16	35.5
5	4	0.08	71
6	8	0.16	35.5
7	28	0.56	10.1
8	16	0.32	17.7
9	24	0.48	11.8
10	20	0.4	14.2
Mean $\Delta\text{Dif.}\% / \mu\text{mol DOCNa} =$			27.3

For the calculus of the apparent activating energy, E_a , of solubilizing MLV-egg-yolk we use the following formulae:

$$\frac{\ln(\Delta\text{Dif.}\% / \mu\text{mol})_2 - \ln(\Delta\text{Dif.}\% / \mu\text{mol})_1}{1000/T_2 - 1000/T_1} = \frac{3.307 - 2.981}{3.384 - 3.442} = 5.62$$

$$E_a = 5.62 \cdot R = 46.7 \text{ kJ/mol}$$

Table 4. Titrant: DOCNa 20mM; $t = 17.5^{\circ}\text{C}$; $\Delta\text{Dif.}\%/\text{cm} = 6.58$

	$\mu\text{l DOCNa}$	$\mu\text{mols DOCNa}$	$\Delta\text{Dif}\%/\mu\text{mol DOCNa}$
1	-	-	-
2	12	0.24	27.4
3	24	0.48	13.7
4	21.6	0.43	15.3
5	26.4	0.53	12.4
6	36	0.72	9.14
7	42	0.84	7.83
8	42	0.84	7.83
9	132	2.64	2.49
Mean $\Delta\text{Dif}\%/\mu\text{mol DOCNa} =$			12
$t = 21^{\circ}\text{C}$; $\Delta\text{Dif.}\%/\text{cm} = 5.26$			
1	-	-	-
2	6	0.12	43.8
3	27.6	0.55	9.56
4	20.4	0.41	12.8
5	15.6	0.31	16.97
6	32.4	0.65	8.1
7	24	0.48	10.9
8	60	1.2	4.38
Mean $\Delta\text{Dif}\%/\mu\text{mol DOCNa} =$			15.2

$$\frac{\ln(\Delta\text{Dif}\%/\mu\text{mol})_2 - \ln(\Delta\text{Dif}\%/\mu\text{mol})_1}{1000/T_2 - 1000/T_1} = \frac{2.722 - 2.485}{3.401 - 3.442} = 5.78$$

$$E_a = 5.78 \cdot R = 48 \text{ kJ/mol}$$

It can be observed that the apparent activating energy of solubilizing MLV-egg-yolk with DOCNa, determined in two different experiments (made on different days), had really close values (46.7 kJ/mol and 48kJ/mol, respectively) proving a good reproductibility of the determination.

Table 5. Titrant: Triton X-100 20mM; $t = 17.5^\circ\text{C}$; $\Delta\text{Dif.}\%/\text{cm} = 5.34$

	$\mu\text{l T X-100}$	$\mu\text{mols T X-100}$	$\Delta\text{Dif}\%/\mu\text{mol T X-100}$
1	-	-	-
2	12	0.24	22.2
3	12	0.24	22.2
4	12	0.24	22.2
5	12	0.24	22.2
6	12	0.24	22.2
7	96	0.19	28.1
8	12	0.24	22.2
9	26.4	0.53	10.1
10	36	0.72	7.42
Mean $\Delta\text{Dif}\%/\mu\text{mol TX-100} =$			19.87
$t = 21^\circ\text{C}; \Delta\text{Dif.}\%/\text{cm} = 6.06$			
1	-	-	-
2	6	0.12	50.5
3	8.4	0.17	35.6
4	7.2	0.14	43.3
5	12	0.24	25.25
6	4.8	0.1	60.6
7	19.2	0.38	15.9
8	8.4	0.17	35.6
9	18	0.36	16.8
10	36	0.72	8.4
Media $\Delta\text{Dif}\%/\mu\text{mol TX-100} =$			32.44


$$\frac{\ln(\Delta\text{Dif}\%/\mu\text{mol})_2 - \ln(\Delta\text{Dif}\%/\mu\text{mol})_1}{1000/T_2 - 1000/T_1} = \frac{3.479 - 2.989}{3.401 - 3.442} = 11.95$$

$$E_a = 11.95 \cdot R = 99.314 \text{ kJ/mol}$$

Table 6. $\Delta\text{Dif\%}/\mu\text{mol}$ is calculated on the basis of knowledge of the variation of percentage diffusion determined, by adding $8\mu\text{mol}$ s surfactant to the sample.

Nr	(°C)	1000/T (grad ⁻¹)	$\Delta\text{Dif\%}/\mu\text{mol}$	$\ln(\Delta\text{Dif\%}/\mu\text{mol})$ CTMB
1	18.4	3.431	8.89	2.185
2	20.5	3.407	9.517	2.253
3	22.5	3.384	10.08	2.311

Figure 2. The Arrhenius representation of the logarithm of the variation of optical diffusion percentage to $1\mu\text{mol}$ CTMB at different temperatures. The straight line was obtained by linear regression.



$$\frac{\Delta \ln(\Delta\text{Dif\%}/\mu\text{mol})}{1000/T_2 - 1000/T_1} = \frac{2.311 - 2.185}{3.384 - 3.431} = 2.68$$

$$E_a = 2.68 \cdot R = 22.3 \text{ kJ/mol}$$

In the Table 7 the apparent activating energies for the solubilizations with the three surfactants are presented.

Table 7. *The apparent activating energy of solubilizing MLV-EYL with surfactants.*

Surfactants	E _a (kJ/mol)
CTMB	22.3
DOCNa	48 and 46.7
Triton X-100	99.3

It is observed that the solubilization powers expressed in kinetic terms, in the condition of titrating at a rate of 0.08 μmols surfactant/sec, for the three surfactants, can be arranged in the order: CTMB < DOCNa < TX-100.

Glossary

CMC- critical micellar concentration

CTMB – cetyltrimethylammonium bromide

Dif%- percentage optical diffusion

DOCNa-sodium deoxycholate

DPPC-dipalmitoylphosphatidylcholine

EYL – egg-yolk lecithin

E_a – apparent activating energy

MLV- multilamellar vesicles

r- correlation coefficient

SUV- small unilamellar vesicles

Tr%- percentage optical transmission

TX-100- Triton X-100, polyoxyethylene-p-t-octilfenol

References

1. Almog, S., Kushnir, T., Nir, S., Lichtenberg, D., *Kinetic and structural aspects of reconstitution of phosphatidylcholine vesicles by dilution of phosphatidylcholine-sodium cholate mixed micelles*, *Biochemistry*, 1986, **25**, 2597-2605
2. Fazakas, Z., *Effect of surfactants on the thermodynamic and kinetic properties of liposome emulsions*, Ph.D. Thesis, University Babeş-Bolyai, Cluj-Napoca, 1999
3. Helenius, A., Simons, K., *Solubilization of membranes by detergents*, *Biochim. Biophys. Acta*, 1975, **415**, 29-79
4. Hobai, Şt., *Kinetics of solubilization with Triton X-100 of egg-yolk lecithin bilayers containing cholesterol*, unpublished
5. Lichtenberg, D., Robson, R.J., Dennis, E.A., *Solubilization of phospholipids by detergents: structural and kinetic aspects*, *Biochim. Biophys. Acta*, 1983, **737**, 285-304
6. Lichtenberg, D., *Characterization of the solubilization of lipid bilayers by surfactants*, *Biochim. Biophys. Acta*, 1985, **821**, 470-478
7. Mazer, N.A., Benedek, G.B., Carey, M.C., *Quasielastic light-scattering studies of aqueous biliary lipid systems*, *Biochemistry*, 1980, **19**, 601-615

8. Müller, K., Schuster, A., *Solubilization of multilamellar liposomes of egg yolk lecithin by the bile salt sodiumtaurodeoxycholate and the effect of cholesterol- a rapid-ultrafiltration study*, Chem.Phys.Lipids, 1990, **52**, 111-127
9. Schurtenberger, P., Mazer, N.A., Kanzig, W., *Micelle to vesicle transition in aqueous solutions of bile salt and lecithin*, J.Phys.Chem., 1985, **89**, 1042-1049