

# Comment on ‘Self-dressing and radiation reaction in classical electrodynamics’\*

V Hnizdo

*National Institute for Occupational Safety and Health, 1095 Willowdale Road, Morgantown, WV 26505, USA*

## Abstract

Using the canonical formalism, Compagno and Persico [*J. Phys. A: Math. Gen.* **35** (2002) 3629–45] have calculated the ‘radiation-reaction’ force on a uniform spherical charge moving rigidly, slowly and slightly from a point where the transverse electric field is assumed to vanish. This force is shown to result in the same time-averaged self-force as that which has been obtained by different means for the test charge of a Bohr–Rosenfeld field-measurement procedure and which Compagno and Persico claimed to be incorrect.

PACS numbers; 03.50.De, 03.70.+k, 12.20.-m

Typeset using REVTeX

---

\*This work was conducted independently of official duties in the Centers for Disease Control and Prevention (CDC), and the views do not necessarily represent those of the CDC.

In a recent paper [1], Compagno and Persico (CP) have calculated, by solving the coupled charge-field Hamilton equations of motion, the ‘radiation-reaction’ force on a spherically symmetric charge that moves rigidly, slowly and only a little from a point where the transverse electric field is assumed to vanish. For a charge  $q$  that is uniformly distributed within a sphere of radius  $a$ , they obtain a ‘radiation-reaction’ force

$$F_{\text{RR}}(t) = -\frac{2q^2}{a^3} \int_0^t dt' \dot{Q}(t') \left[ 1 - \frac{3(t-t')}{2a} + \frac{(t-t')^3}{4a^3} \right] \Theta[2a - (t-t')] \quad (1)$$

where  $\dot{Q}(t)$  is the time derivative of a one-dimensional trajectory of the charge and  $\Theta(x)$  is the Heaviside step function; here and henceforth, we use units such that the speed of light  $c = 1$  and put  $t = 0$  for the time at which the transverse electric field vanishes.

CP remark that the result (1) is relevant to the issues raised in recent papers [2–5] in connection with the Bohr–Rosenfeld (BR) analysis of the measurability of the electromagnetic field [6], as it should apply to a BR measurement procedure with only minor modifications. In the present note, we show that the force (1), which we prefer to call the electromagnetic self-force, results in the same time-averaged self-force as that which has been obtained by different means in [3,5] for the test charge of a BR measurement procedure and which CP have rejected in [4] as incorrectly calculated.

A condition on the one-dimensional trajectory  $Q(t)$  of the test charge in a BR measurement procedure occupying a time interval  $(0, T)$  is that  $Q(t) = 0$  for  $t \leq 0$ . While this means that the transverse electric field of the test charge vanishes at  $t = 0$ , the initial condition  $\dot{Q}(t)|_{t=0} = 0$  that is implied would result according to (1) in  $Q(t) = 0$  also for  $t > 0$  if there were no other force acting on the test charge in addition to the self-force. It is presumably for this reason that when CP touch on the applicability of (1) to the test charge of the BR measurement procedure, they invoke the ‘neutralization’ of the test charge at  $t = 0$  by the stationary neutralizing charge employed in the procedure. Beside the question whether such a neutralization alone indeed guarantees a vanishing transverse electric field at  $t = 0$ , the fact remains that then there is at least one other force acting on the test charge, namely the electrostatic force of attraction to the neutralizing charge. It thus appears inescapable that any meaningful use of formula (1) in an analysis of the BR field-measurement procedure requires the presence of external forces. We shall first write (1) in a different form before returning to this point.

Using integration by parts, we write (1) as

$$F_{\text{RR}}(t) = Q(t')u(t-t')\Big|_{t'=0}^t - \int_0^t dt' Q(t') \frac{du(t-t')}{dt'} \quad (2)$$

where

$$u(t-t') = -\frac{2q^2}{a^3} \left[ 1 - \frac{3(t-t')}{2a} + \frac{(t-t')^3}{4a^3} \right] \Theta[2a - (t-t')]. \quad (3)$$

Now,  $u(t-t')|_{t'=t} = -2q^2/a^3$  and

$$\frac{du(t-t')}{dt'} = -\frac{3q^2}{2a^4} \left[ 2 - \frac{(t-t')^2}{a^2} \right] \Theta[2a - (t-t')]. \quad (4)$$

Using this and an initial condition  $Q(t')|_{t'=0} = 0$ , we obtain the self-force (2) for  $t < T$  as

$$F_{\text{RR}}(t) = -\frac{2q^2}{a^3}Q(t) + \frac{3q^2}{2a^4} \int_0^T dt' Q(t') \left[ 2 - \frac{(t-t')^2}{a^2} \right] \Theta(t-t')\Theta[2a-(t-t')] \quad (5)$$

where the factor  $\Theta(t-t')$  is introduced in the integrand in order to be able to fix the integration range as  $(0, T)$ . Apart from differences in notation, this expression is the same as that for the self-force  $F_x(t_2)$  given in [5] by equations (26) and (28), with a factor of 2 instead of 3 in the delta-function term of equation (28) so that the electrostatic force due to a BR neutralizing charge is subtracted.

The self-force  $F_x(t_2)$  was obtained in [5] by using the electromagnetic self-field of a uniform spherical charge whose trajectory was *prescribed* to be a given trajectory  $Q(t)$  satisfying the conditions imposed on it by the BR field-measurement procedure apart from the stipulation that it is to have a step-like character with respect to the measurement period  $(0, T)$ . This fact invalidates the caution of CP that the force which they obtained ‘is unambiguously the radiation-reaction force only in the absence of other forces’. A prescribed trajectory can result to a given degree of accuracy only when there is a suitable external force acting on the charge in addition to the self-force. In the BR field-measurement procedure, the test charge is acted on by an external force that is the resultant of forces originating from several sources: the momentum-measurement system, the neutralizing charge, the spring that compensates the time-averaged effects of the self-force and the neutralizing charge, and, of course, the measured external field itself; the approximate attainment of the prescribed step-like trajectory is there facilitated by the fact that the test charge is allowed to have an arbitrarily great mass.

The test charge’s self-force averaged over the duration  $T$  of the field measurement,

$$\bar{F}_{\text{RR}} = \frac{1}{T} \int_0^T dt F_{\text{RR}}(t) \quad (6)$$

can now be written using (5) as

$$\bar{F}_{\text{RR}} = -\frac{2q^2}{a^3 T} \int_0^T dt Q(t) + \frac{q^2}{T} \int_0^T dt' Q(t') \int_0^T dt v(t-t') \quad (7)$$

where

$$v(t-t') = \frac{3}{2a^4} \left[ 2 - \frac{(t-t')^2}{a^2} \right] \Theta(t-t')\Theta[2a-(t-t')]. \quad (8)$$

The integration with respect to  $t$  in the second term of (7) is straightforward, yielding

$$\int_0^T dt v(t-t') = \frac{2}{a^3} + f(t') \quad (9)$$

where

$$f(t') = -\frac{1}{2a^3}(2-\chi)(2-2\chi-\chi^2)\Theta(2-\chi) \quad \chi = \frac{T-t'}{a}. \quad (10)$$

Using (9) in (7), the time-averaged self-force is obtained finally as

$$\bar{F}_{\text{RR}} = \frac{q^2}{T} \int_0^T dt' Q(t') f(t'). \quad (11)$$

Expressions (10) and (11) are, apart from differences in notation and the absence in (10) of the electrostatic term  $-1/a^3$  due to the neutralizing charge, the same as those given for the time-averaged self-force in equations (3) and (9) of [3] and equations (10) and (22) of [5]; these expressions have been claimed by CP to be incorrect [4].

It has been shown in [5] that the rejection [4] of CP of the criticism [3] of their re-analysis [2] of the BR field-measurement procedure is based on erroneous calculations. It has to be concluded that that rejection is invalidated also by the results of the latest work of CP themselves.

## REFERENCES

- [1] Compagno G and Persico F 2002 Self-dressing and radiation reaction in classical electrodynamics *J. Phys. A: Math. Gen.* **35** 3629–45
- [2] Compagno G and Persico F 1998 Limits on the measurability of the local quantum electromagnetic-field amplitude *Phys. Rev. A* **57** 1595–1603
- [3] Hnizdo V 1999 Comment on Limits of the measurability of the local quantum electromagnetic-field amplitude *Phys. Rev. A* **60** 4191–95
- [4] Compagno G and Persico F 1999 Reply to Comment on Limits of the measurability of the local quantum electromagnetic-field amplitude *Phys. Rev. A* **60** 4196–97
- [5] Hnizdo V 2000 The electromagnetic self-force on a charged spherical body slowly undergoing a small, temporary displacement from a position of rest *J. Phys. A: Math. Gen.* **33** 4095–103; lanl.arXiv e-print math-ph/0005014
- [6] Bohr N and Rosenfeld L 1933 Zur Frage der Messbarkeit der elektromagnetischen Feldgrößen *Mat. Fys. Medd. K. Dan. Vidensk. Selsk.* **12** no 8 (Engl. translation 1983 On the question of the measurability of electromagnetic field quantities *Quantum Theory and Measurement* ed J A Wheeler and W H Zurek (Princeton NJ: Princeton University Press) pp 479–522)