Response to Compagno and Persico^{*}

V Hnizdo

National Institute for Occupational Safety and Health, 1095 Willowdale Road, Morgantown, WV 26505, USA

Abstract

Some mathematically incorrect claims of Compagno and Persico in their reply (2002 J. Phys. A: Math. Gen. **35** 8965) to my comment on their recent paper on self-dressing and radiation reaction in classical electrodynamics are pointed out.

Compagno and Persico (CP) have replied [1] to my comment [2] on their paper on selfdressing and radiation reaction in classical electrodynamics [3]. CP acknowledge the main point of the comment, namely that the expression for the time-averaged electromagnetic selfforce obtained in [4] for the test charge of a Bohr–Rosenfeld field-measurement procedure and rejected in [5] as incorrect can be obtained also using a formula for the self-force which they derived by different means in [3]. In view of this fact, CP now endorse the expression in question as correct. However, some claims in their reply call for my response.

The expression for the time-averaged self-force on a spherical uniform charge q of radius a obtained in [4] reads

$$\bar{F} = \frac{q^2}{T} \int_0^T dt' \, Q(t') f(t') \tag{1}$$

where

$$f(t') = -\frac{1}{2a^3}(2-\chi)(2-2\chi-\chi^2)\Theta(2-\chi) \qquad \chi = \frac{T-t'}{a}.$$
 (2)

Here, the speed of light c = 1 and Q(t) is the charge's one-dimensional trajectory, which is subject to the conditions that Q(t) = 0 for $t \leq 0$ and $|Q(t)| \ll a$, $|dQ(t)/dt| \ll c$ for 0 < t < T; $\Theta(x)$ is the Heaviside step function. Instead of the simple closed-form expression (2) for the function f(t'), CP counter-proposed in [5] the expression (normalized here to conform with (2)):

$$f(t') = -\frac{2}{3V^2} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} [\delta^{(n+1)}(T-t') - \delta^{(n+1)}(-t')] \langle r^{n-1} \rangle \qquad V = \frac{4}{3}\pi a^3 \tag{3}$$

where

$$\langle r^{n-1} \rangle = \int_{|\boldsymbol{r}_1| < a} \mathrm{d}\boldsymbol{r}_1 \int_{|\boldsymbol{r}_2| < a} \mathrm{d}\boldsymbol{r}_2 \, |\boldsymbol{r}_2 - \boldsymbol{r}_1|^{n-1} = \frac{72V^2(2a)^{n-1}}{(n+5)(n+3)(n+2)}.$$
 (4)

^{*}This article is written by V Hnizdo in his private capacity. No official support or endorsement by the Centers for Disease Control and Prevention is intended or should be inferred.

(The above closed-form expression for $\langle r^{n-1} \rangle$ was given subsequently in [6].) My first point here has to be unfortunately of a rather trivial nature. The sign of expression (3) is correct, if, as in the standard notation of elementary calculus, $u^{(n)}[v(x)] \equiv d^n u(y)/dy^n|_{y=v(x)}$. Thus, e.g., $\delta^{(n)}(T-t) \equiv d^n \delta(x)/dx^n|_{x=T-t}$. With this notation, there is no sign misprint in the function f(t') in [5] that CP now want to correct—but if $\delta^{(n)}(T-t)$ meant instead $d^n \delta(T-t)/dt^n = (-1)^n d^n \delta(x)/dx^n|_{x=T-t}$, as CP suggest, then not only the sign but also the factor $(-1)^n$ would be there in error. In either case, the function f(t') is now given incorrectly by the expression (3) of [1]—it either has the wrong overall sign, or the factor $(-1)^n$ there must be omitted.

Contrary to an assertion of CP, I have never claimed that the expression (3) for the function f(t') is incorrect. I have rather pointed out in [6] that, in order to obtain the time-averaged self-force (1), this expression was used incorrectly by CP in the requisite integration. Since this is an integration with finite limits involving high-order derivatives of the delta function, it cannot be performed as simplistically as CP have done, obtaining [5]

$$\bar{F}_{\rm CP} = -\frac{2\rho^2}{3T} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \left[Q^{(n+1)}(T) - Q^{(n+1)}(0) \right] \langle r^{n-1} \rangle \qquad \rho = \frac{q}{V}.$$
(5)

This is the 'exact' expression for the time-averaged self-force \overline{F} that CP still claim in [1] to be correct, despite the fact that it is inconsistent with their newly adopted approval of the expression (2) for the function f(t'). This can be shown by evaluating the time-averaging integral (1) for \overline{F} using the expression (2) for f(t') and the Taylor expansion of the trajectory Q(t) about the point t = T:

$$\bar{F} = \frac{q^2}{T} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} Q^{(n)}(T) \int_0^T dt' (T-t')^n f(t')$$

= $\frac{q^2}{T} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} Q^{(n)}(T) \left\{ \alpha_n - \left[a^{n-2} \left(\frac{2\kappa^{n+1}}{n+1} - \frac{3\kappa^{n+2}}{n+2} + \frac{1}{2}\kappa^{n+4} \right) + \alpha_n \right] \Theta(2-\kappa) \right\}$ (6)

where

$$\alpha_n = \frac{48(2a)^{n-2}n}{(n+1)(n+2)(n+4)} \qquad \qquad \kappa = \frac{T}{a}.$$
(7)

This is the correct expression for the time-averaged self-force in terms of the derivatives $Q^{(n)}(T) \equiv \lim_{t \to T^-} d^n Q(t)/dt^n$. (Note for completeness that the correct expression for \bar{F} in terms of the derivatives $Q^{(n)}(0) \equiv \lim_{t \to 0^+} d^n Q(t)/dt^n$ is given by equation (25) of [6].) As $\alpha_n = \frac{2}{3}n\langle r^{n-2}\rangle/V^2$, using (6) we can write \bar{F} for $T \geq 2a$ as

$$\bar{F} = \frac{q^2}{T} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} Q^{(n)}(T) \alpha_n = -\frac{2\rho^2}{3T} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} Q^{(n+1)}(T) \langle r^{n-1} \rangle \qquad T \ge 2a \qquad (8)$$

which happens to be equal to the first of the two terms in the expression (5) of CP for \overline{F} . This demonstrates that expression (5) is incorrect; for $T \ge 2a$, (5) can be corrected by dropping its second term, but for T < 2a, no such simple correction is possible.

Far from being 'convenient' for an 'exact' evaluation of the time-averaged self-force \overline{F} , formula (3) is a purely formal expression that has no practical application in an integration

with finite limits. Its use by CP has led to the erroneous expression (5) for \overline{F} ; when CP use it in [1] to prove the equivalence of expressions (2) and (3), they revert the Taylor expansions $\sum_{n}(-1)^{n}\delta^{(n+1)}(t)r^{n}/n!$ to $\delta'(t-r)$ before performing the finite-limit integration.

So far, CP have responded to my criticism [4] of their re-analysis [7] of the Bohr–Rosenfeld field-measurement procedure only by making mathematically incorrect claims. All these involved rather simple mathematical points about which there should have been no need of explicating since my paper [6] of 2000 (where I derived expression (2) using elementary calculus of the delta function), if not already since my comment [4] of 1999 (where I used Fourier transform methods). It is regrettable that such points have deflected from the interesting issues of physics relating to the famous Bohr–Rosenfeld analysis.

- Compagno G and Persico F 2002 Reply to Comment on Self-dressing and radiation reaction in classical electrodynamics J. Phys. A: Math. Gen. 35 8965–67
- [2] Hnizdo V 2002 Comment on Self-dressing and radiation reaction in classical electrodynamics J. Phys. A: Math. Gen. 35 8961–63 (E-print physics/0205022)
- [3] Compagno G and Persico F 2002 Self-dressing and radiation reaction in classical electrodynamics J. Phys. A: Math. Gen. 35 3629–45
- [4] Hnizdo V 1999 Comment on Limits of the measurability of the local quantum electromagneticfield amplitude Phys. Rev. A 60 4191–95 (E-print quant-ph/0210074)
- [5] Compagno G and Persico F 1999 Reply to Comment on Limits of the measurability of the local quantum electromagnetic-field amplitude *Phys. Rev.* A **60** 4196–97
- [6] Hnizdo V 2000 The electromagnetic self-force on a charged spherical body slowly undergoing a small, temporary displacement from a position of rest J. Phys. A: Math. Gen. 33 4095–103 (E-print math-ph/0005014)
- [7] Compagno G and Persico F 1998 Limits on the measurability of the local quantum electromagnetic-field amplitude *Phys. Rev.* A **57** 1595–603