Relativistic Brownian Motion

O. Oron and L.P. Horwitz*

School of Physics and Astronomy Raymond and Beverly Sackler Faculty of Exact Sciences Tel Aviv University, Ramat Aviv 69978, Israel

Abstract:

We solve the problem of formulating Brownian motion in a relativistically covariant framework in 1 + 1 and 3 + 1 dimensions. We obtain covariant Fokker-Planck equations with (for the isotropic case) a differential operator of invariant d'Alembert form. Treating the spacelike and timelike fluctuations separately, we show that it is essential to take into account the analytic continuation of "unphysical" fluctuations in order to achieve these results.

^{*} Also at Department of Physics, Bar Ilan University, Ramat Gan 529000, Israel

Introduction

Nelson¹, in 1966, constructed the Schrödinger equation from an analysis of Brownian motion by identifying the forward and backward average velocities of a Brownian particle with the real and imaginary parts of a wave function. He pointed out that the basic process involved is defined nonrelativistically, and can be used if relativistic effects can be safely neglected. The development of a relativistically covariant formulation of Brownian motion could therefore provide some insight into the structure of a relativistic quantum theory.

In recent years, Brownian motion has been applied as a mechanism for the description of irreversible evolution of quantum mechnical states, for example, collapse of wave functions as result of measurement. One finds that the wave function with a stochastic term added to the Schrödinger equation (represented in the Hilbert space of states² or in their projective representation³) evolves to a mixed state of alternative outcomes of a measurement with the same probability distribution as given by the calculation of *a priori* (Born) probabilities in the framework of the standard quantum theory ^{2,3}. One of the motivations for the work of Adler and Horwitz³ was based on the existence of a statistical mechanics, developed by Adler⁴ and Adler and Millard⁵ for the description of the equilibrium state of a general class of quantum fields; in this theory, the thermal average of commutator expressions take on the values implied by the familiar complex canonical quantum theory and quantum field theory. The Brownian motion corrections to the standard theories may be thought of as arising from the fluctuations around the equilibrium state. The development of a relativistically covariant theory of Brownian motion could make an extension of these ideas to relativistic quantum theory and quantum field theory more accessible.

The program of stochastic quantization of Parisi and Wu^6 assumes the existence of relativistic wave equations, and applies a statistical approach closely related to path integrals. The Wiener distribution of nonrelativistic Brownian motion, as is well known, is associated with (imaginary time) path integral formulations, and one would expect that covariant Brownian motion would similarly be associated with Parisi-Wu stochastic quantization.

Nelson¹ has pointed out that the formulation of his stochastic mechanics in the context of general relativity is an important open question. The Riemannian metric spaces one can achieve, in principle, which arise due to nontrivial correlations between fluctuations in spacetime directions, could, in the framework of a covariant theory of Brownian motion, lead to spacetime pseudo-Riemannian metrics in the structure of diffusion and Schrödinger equations.

In this paper we shall study the structure a covariant theory of Brownian motion.

We first point out some of the obvious difficulties in reaching a covariant theory of Brownian motion, and indicate the directions we have chosen to solve these problems.

Brownian motion, thought of as a series of "jumps" of a particle along its path, necessarily involves an ordered sequence. In the nonrelativistic theory, this ordering is naturally provided by the Newtonian time parameter. In a relativistic framework, the Einstein time t does not provide a suitable parameter. If we contemplate jumps in spacetime, to accomodate a covariant formulation, a possible spacelike interval between two jumps may appear in two orderings in different Lorentz frames. We therefore adopt the invariant parameter τ introduced by Stueckelberg⁷ in his construction of a relativistically covariant classical and quantum dynamics. For the many body theory, Piron and Horwitz⁸ postulated that this parameter is universal, as the Newtonian time in the nonrelativistic theory, and drives the classical particle trajectories $x_i^{\mu}(\tau)$ (worldlines labelled i = 1, 2, 3, ..., N) through equations of motion, and the evolution of the wave function in the Hilbert space $L^2(R^{4N}), \psi_{\tau}(\{x_i^{\mu}\})$ through the Stueckelberg-Schrödinger equation (the differential form of the action of a one-parameter unitary group with parameter τ).

A second fundamental difficulty in formulating a covariant theory of Brownian motion lies in the form of the correlation function of the random variables of spacetime. The straightforward generalization of the usual Brownian correlation property to special relativity, i.e.,

$$< dw_{\mu}(\tau)dw_{\nu}(\tau') >= \begin{cases} 0 & \tau \neq \tau' \\ 2\alpha\eta_{\mu\nu}d\tau & \tau = \tau', \end{cases}$$

contains the serious problem that $\langle dw_0(\tau) dw_0(\tau) \rangle < 0$, which is impossible. Brownian motion in spacetime, however, should be a generalization of the nonrelativistic problem, constructed by observing the nonrelativistic process from a moving frame according to the transformation laws of special relativity. Hence, as a first step, the process taking place in space in the nonrelativistic theory should be replaced by a spacetime process in which the Brownian jumps are spacelike. The pure time (negative) self-correlation does therefore not occur. In order to meet this requirement, we shall use a coordinatization in terms of generalized polar coordinates which assure that all jumps are spacelike. In this case, one would expect a distribution function of the form $e^{-\frac{\mu^2}{ad\tau}}$, where μ is the invariant spacelike interval of the jump, and a is some constant. As we shall see, a Brownian motion based on purely spacelike jumps does not yield the correct form for an invariant diffusion process. We must therefore consider the possibility as well that, in the framework of relativistic dynamics, there are timelike jumps. In a frame in which the timelike jumps are pure time, the construction of the Gaussian distribution from the central limit theorem can again be applied. The distribution would be expected to be of the form $e^{-\frac{\sigma^2}{bd\tau}}$, where σ is the timelike interval of these jumps, and b is some constant. By suitably weighting the occurrence of the spacelike process (which we take for our main discussion to be "physical", since its nonrelativistic limit coincides with the usual Brownian motion) and an analytic continuation of the timelike process, we show that one indeed obtains a Lorentz invariant Fokker-Planck equation in which the d'Alembert operator appears in place of the Laplace operator of the 3D Fokker-Planck equation. One may, alternatively, consider the timelike process as "physical" and analytically continue the spacelike ("unphysical") process to achieve a d'Alembert operator with opposite sign.

2. Brownian motion in 1+1 dimensions

We consider a Brownian path in 1 + 1 dimensions of the form

$$dx^{\mu}(\tau) = K^{\mu}(x(\tau))d\tau + dw^{\mu}(\tau).$$
(2.1)

We start by considering the second order term in the series expansion of a function of position of the particle on the world line, $f(x^{\mu}(\tau) + \Delta x^{\mu})$, involving the operator

$$\mathcal{O} = \Delta x^{\mu} \Delta x^{\nu} \frac{\partial}{\partial x^{\mu}} \frac{\partial}{\partial x^{\nu}}.$$
(2.2)

We have remarked that one of the difficulties in describing Brownian motion in spacetime is the possible occurrence of a negative value for the second moment of some component of the Lorentz four vector random variable. If the Brownian jump is timelike, or spacelike, however, the components of the four vector are not independent, but must satisfy the timelike or spacelike constraint. Such constraints can be realized by using parametrizations for the jumps in which they are restricted geometrically to be timelike or spacelike. We now separate the jumps into spacelike jumps and timelike jumps accordingly, i.e., for the spacelike jumps,

$$\Delta x = \mu \cosh \alpha \quad \Delta t = \mu \sinh \alpha \tag{2.3}$$

and for the timelike jumps,

$$\Delta x = \sigma \sinh \alpha \quad \Delta t = \sigma \cosh \alpha \tag{2.3'}$$

Here we assumed that the two sectors have the same distribution on the hyperbolic variable. We first look for the effects of a particle experiencing spacelike jumps only. In that case the operator \mathcal{O} takes the following form:

$$\mathcal{O} = \mu^2 \left[\cosh^2 \alpha \frac{\partial^2}{\partial x^2} + 2\sinh\alpha \cosh\alpha \frac{\partial^2}{\partial x \partial t} + \sinh^2 \alpha \frac{\partial^2}{\partial t^2}\right]$$
(2.4)

If the particle going under timelike jumps only we find the operator \mathcal{O} takes the following form:

$$\mathcal{O} = \sigma^2 [\sinh^2 \alpha \frac{\partial^2}{\partial x^2} + 2 \sinh \alpha \cosh \alpha \frac{\partial^2}{\partial x \partial t} + \cosh^2 \alpha \frac{\partial^2}{\partial t^2}]$$
(2.5)

In order to obtain the relativistically invariant d'Alembert diffusion operator, the expression obtained in the timelike region must be *subtracted* from the expression for the spacelike region, and furthermore, the amplitudes must be identified. In the physical timelike region, the coefficient σ^2 is, of course, positive, and using the law of large numbers on the random distribution, one obtains a Gaussian distribution analogous to that of the spacelike case.

We see, however, that we can obtain the d'Alembert operator only by considering the analytic continuation of the timelike process to the spacelike domain. This procedure is analogous to the effect, well-known in relativistic quantum scattering theory, of a physical process in the crossed (t) channel on the observed process in the direct (s) channel. Although we are dealing with an apparently classical process, as Nelson has shown, the Brownian motion problem gives rise to a Schrödinger equation, and therefore contains properties of the quantum theory. We thus see the remarkable fact that one must take into account the physical effect of the analytic continuation of processes occurring in a non-physical, in this case timelike, domain, on the total observed behavior of the system.

In the non-stochastic case, Einstein's relativity identifies $\Delta x/\Delta t$ with p/E, where p and E are the components of the energy-momentum four-vector of the particle. If we make an analogous identification, assigning these variables as properties of the fluctuations, then

$$\sigma^{2} = (\Delta t)^{2} - (\Delta x)^{2} \propto (E^{2} - p^{2}), \qquad (2.7)$$

defining a stochastic mass squared associated with the Brownian particle. If the relation between E and p becomes spacelike, the notion of stochastic mass can be retained undere the transformation to an imaginary representation $E \rightarrow iE'$ and $p \rightarrow ip'$, for E', p' reals, so the relation p'/E' > 1 remains, but $p'^2 - E'^2 > 0$. We assume sufficient symmetry in the spacelike and timelike distributions so that the absolute values of $\langle \sigma^2 \rangle$ and $\langle \mu^2 \rangle$ are equal. The preservation of the mean magnitude of the interval reflects the conservation of a mass-like property which remains, as an intrinsic property of the particle, for both spacelike and timelike jumps.

With these assumptions, the cross-term in hyperbolic functions cancels in the sum, which now takes the form

$$\langle \mathcal{O} \rangle = \langle \mu^2 \rangle \left[\frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial t^2} \right]$$
 (2.8)

Taking into account the drift term in (2.1), one then finds the relativistic Fokker-Planck equation

$$\frac{\partial D(x,\tau)}{\partial \tau} = \left\{ -\frac{\partial}{\partial x^{\mu}} K^{\mu} + \langle \mu^2 \rangle \frac{\partial^2}{\partial x^{\mu} \partial x_{\mu}} \right\} D(x,\tau),$$
(2.9)

where $\partial/\partial x^{\mu}$ operates on both K^{μ} and D.

We see that the procedure we have followed permits us to construct the Lorentz invariant d'Alembertian operator, as required for obtaining a relativistically covariant diffusion equation. Furthermore, since the expectation of $\sinh^2 \alpha$, $\cosh^2 \alpha$ could be infinite (e.g., for a uniform distribution on α), the result we obtain in this way constitutes an effective regularization.

3. Brownian motion in 3 + 1 dimensions

In the 3 + 1 case, we again separate the jumps into timelike and spacelike types. The spacelike jumps may be parametrized, in a given frame, by

$$\Delta t = \mu \sinh \alpha$$

$$\Delta x = \mu \cosh \alpha \cos \phi \sin \vartheta$$

$$\Delta y = \mu \cosh \alpha \sin \phi \sin \vartheta$$

$$\Delta z = \mu \cosh \alpha \cos \vartheta$$

(3.1)

We assume the four variables μ , α , ϑ , ϕ are independent random variables. In addition we demand in this frame that ϑ and ϕ are uniformly distributed in their ranges $(0, \pi)$ and $(0, 2\pi)$, respectively. In this case, we may average over the trigonometric angles, i.e., ϑ and ϕ and find that:

$$<\Delta x^{2} >_{\phi,\vartheta} = <\Delta y^{2} >_{\phi,\vartheta} = <\Delta z^{2} >_{\phi,\vartheta} = \frac{\mu^{2}}{3} \cosh^{2}\alpha$$

$$<\Delta t^{2} >_{\phi,\vartheta} = \mu^{2} \sinh^{2}\alpha$$
(3.2)

[§] This transformation is similar to the analytic continuation $p \to ip'$ in nonrelativistic tunneling, for which the particle appears as an instanton.

We may obtain the averages over the trigonometric angles of the timelike jumps by replacing everywhere in Eq.(3.2)

$$\cosh^2 \alpha \leftrightarrow \sinh^2 \alpha$$
$$\mu^2 \to \sigma^2$$

to obtain

$$<\Delta x^{2} >_{\phi,\vartheta} = <\Delta y^{2} >_{\phi,\vartheta} = <\Delta z^{2} >_{\phi,\vartheta} = \frac{\sigma^{2}}{3} \sinh^{2}\alpha$$

$$<\Delta t^{2} >_{\phi,\vartheta} = \sigma^{2} \cosh^{2}\alpha,$$
(3.3)

where σ is a real random variable, the invariant timelike interval. Assuming, as in the 1+1 case, that the likelihood of the jumps being in either the spacelike or (virtual) timelike phases are equal, and making an analytic continuation for which $\sigma^2 \rightarrow -\lambda^2$, the total average of the operator \mathcal{O} , including the contributions of the remaining degrees of freedom μ, λ and α is

$$<\mathcal{O}>= \left(<\mu^2><\sinh^2\alpha>-<\lambda^2><\cosh^2\alpha>\right)\frac{\partial^2}{\partial t^2}+ \frac{1}{3}\left(<\mu^2><\cosh^2\alpha>-<\lambda^2><\sinh^2\alpha>\right)\triangle$$
(3.4)

If we now insist that the operator $\langle \mathcal{O} \rangle$ be invariant under Lorentz transformations (i.e. the d'Alembertian) we impose the condition

$$<\mu^{2} > <\sinh^{2}\alpha > - <\lambda^{2} > <\cosh^{2}\alpha > = -\frac{1}{3} (<\mu^{2} > <\cosh^{2}\alpha > - <\lambda^{2} > <\sinh^{2}\alpha >)$$
(3.5)

Using the fact that $\langle \cosh^2 \alpha \rangle - \langle \sinh^2 \alpha \rangle = 1$, and defining $\gamma \equiv \langle \sinh^2 \alpha \rangle$, we find that

$$<\lambda^2>=\frac{1+4\gamma}{3+4\gamma}<\mu^2> \tag{3.6}$$

The Fokker-Planck equation then takes on the same form as in the 1 + 1 case, i.e., the form (2.9). We remark that for the 1 + 1 case, one finds in the corresponding expression that the 3 in the denominator is replaced by unity, and the coefficients 4 are replaced by 2; in this case the requirement reduces to $\langle \mu^2 \rangle = \langle \lambda^2 \rangle$ and there is no γ dependence.

We see that in the limit of a uniform distribution in α , for which $\gamma \to \infty$, $\langle \lambda^2 \rangle \to \langle \mu^2 \rangle$. In this case, the relativistic generalization of a nonrelativistic Gaussian distribution of the form $e^{-\frac{r^2}{dt}}$ becomes of the form $e^{-\frac{\mu^2}{d\tau}}$, which is Lorentz invariant. As in the 1 + 1 case, the result (3.6) corresponds to a regularization.

4. Conclusions and Discussion

We have constructed a relativistic generalization of Brownian motion, using an invariant world-time to order the Brownian fluctuations, and separated consideration of spacelike and timelike jumps to avoid the problems of negative second moments which might otherwise follow from the Minkowski signature. Associating the Brownian fluctuations with an underlying dynamical process, one may think of γ in the 3 + 1 case as an order parameter, where the distribution function (over α), associated with the velocities, is determined by the temperature of the underlying dynamical system (as we have remarked, the result for the 1 + 1 case is independent of the distribution on the hyperbolic variable).

At equilbrium, where $\partial D/\partial \tau = 0$, the resulting diffusion equation turns into a classical wave equation which, in the absence of a drift term K^{μ} , is the wave equation for a massless field. An exponentially decreasing distribution in τ of the form $\exp -\kappa \tau$ would correspond to a Klein-Gordon equation for a particle in a tachyonic state (mass squared $-\kappa$). We have considered the spacelike jumps as "physical" since they result in the usual Brownian motion in the nonrelativistic limit. If the timelike jumps were considered as "physical", one would analytically continue the "unphysical" spacelike process. The resulting diffusion equation would have the opposite sign for the d'Alembert operator, and an exponentially decreasing distribution would then result in a Klein-Gordon equation in a timelike particle state.

Nelson¹ has shown that non-relativistic Brownian motion can be associated with a Schrödinger equation. Equipped with the procedures we presented here, constructing relativistic Brownian motion, Nelson's methods can be generalized. One then can construct relativistic equations of Schrödinger (Schrödinger-Stueckelberg) type. The eigenvalue equations for these relativistic forms are also Klein-Gordon type equations. Moreover one can also generalize the case where the fluctuations are not correlated in different directions into the case where correlations exist, as discussed by Nelson¹ for three dimensional Riemannian spaces. In this case the resulting equation is a quantum equation in a curved Riemannian spacetime; as pointed out in ref.10, the eikonal approximation to the solutions of such an equation contains the geodesic motion of classical general relativity. The medium supporting the Brownian motion may be identified with an "ether" ¹¹ (Nelson¹ has remarked that the self-interaction of charged particles might provide a mechanism for the Brownian motion) for which the problem of local Lorentz symmetry is solved. This generalization of Nelson's method will be discussed elsewhere.

Acknowledgements

We would like to thank Linda Reichl, W.C. Schieve and Sungyan Kim at the Ilya Prigogine Center for Statistical Mechanics and Complex Systems at the University of Texas at Austin for helpful and stimulating discussions.

References

- 1. Edward Nelson, Dynamical Theories of Brownian Motion, Princeton University Press, Princeton (1967); Edward Nelson, Quantum Fluctuations, Princeton University Press Princeton (1985). See also, Ph. Blanchard, Ph. Combe and W. Zheng, Mathematical and Physical Aspects of Stochastic Mechanics, Springer-Verlag, Heidelberg (1987), for further helpful discussion.
- 2. For example, G.C. Ghirardi, P. Pearle and A. Rimini, Phys. Rev. A42, 78 (1990).
- L.P. Hughston, Proc. Roy. Soc. London A452, 953 (1996), and references cited there;
 S.L. Adler and L.P. Horwitz, Jour. Math. Phys. 41, 2485 (2000).

- S.L. Adler, Nuc. Phys. B415, 195 (1994); S.L. Adler, Quaternionic Quantum Mechanics and Quantum Fields, Oxford Univ. Press, N.Y. (1995); S.L. Adler, hep-th/0206120.
- 5. S.L. Adler and A.C. Millard, Nuc. Phys. B473, 199 (1996).
- G. Parisi, Y. Wu, Sci.Sin. 24 483; Mikio Namiki Stochastic Quantization, (Springer-Verlag, Heidelberg, 1992).
- 7. E.C.G. Stueckelberg, Helv. Phys. Acta 14, 322 (1941); 14, 588 (1941).
- 8. L.P. Horwitz and C. Piron, Helv. Phys. Acta 46, 316 (1973).
- 9. See, for example, S.S. Schweber, An Introduction to Relativistic Quantum Field Theory, Harper and Row, N.Y. (1961).
- 10. L.P.Horwitz , O.Oron, hep-ph/0205018
- 11. S. Liberati, S. Sonego and M. Visser, Ann. Phys. 167 298 (2002).