

# Teleparallel structures and background metric in monocism cosmology

Vladimir Trifonov  
American Mathematical Society,  
trifonov@e-math.ams.org

## Abstract

The monocism model shows that Boolean observer will always see his spacetime as a certain four-dimensional pseudo-Riemannian manifold of signature 2. In this note we study the geometry of this manifold in some detail. In particular, we compute its natural teleparallel connection, torsion and background metric induced by teleparallel communications.

PACS: 02.19.-v, 04.50.+h, 98.80.Jk

(1) *Introduction.* The monocism model [1] links operational logic of rational observer with the dimensionality, possible types of metric and topology of the spacetime he perceives. Here is the basic technical setup. Let  $\mathbf{F}$  be a partially ordered field. An  $\mathbf{F}$ -observer is the category  $\mathcal{A}[\mathbf{F}]$  of linear algebras over  $\mathbf{F}$ . *Paradigms* of an  $\mathbf{F}$ -observer are  $\mathcal{A}[\mathbf{F}]$ -objects, his *effectors* are  $\mathcal{A}[\mathbf{F}]$ -arrows. For a paradigm  $\mathcal{A}$ : (a) *states of knowledge* are elements of the algebra  $\mathcal{A}$ ; (b) a *sensory metric* is the algebra- $\mathcal{A}$  structure constant tensor  $\mathbf{C}(\tilde{\omega}; a, b)$  together with a one-form  $\tilde{t}$  on the additive linear space  $\mathbf{S}_{\mathcal{A}}$  of the algebra  $\mathcal{A}$ , provided  $\mathbf{C}(\tilde{t}; a, b)$  is symmetric in vector arguments  $a$  and  $b$ ;  $\tilde{t}$  is then the *sensory time*; (c) the *sensory space* is  $\mathbf{S}_{\mathcal{A}}$  together with all the sensory metrics of the paradigm  $\mathcal{A}$ ; elements of  $\mathbf{S}_{\mathcal{A}}$  are *reflexors*; (d) the *motor space* is the multiplicative subgroupoid  $\mathbf{M}_{\mathcal{A}}$  of the algebra  $\mathcal{A}$ , generated by the set of all non-zero elements of  $\mathcal{A}$ . A paradigm  $\mathcal{A}$  is (*ir*)rational if  $\mathbf{M}_{\mathcal{A}}$  is (not) a monoid. If  $\mathcal{A}$  is a rational paradigm and the topos of  $\mathbf{M}_{\mathcal{A}}$  realizations,  $\mathbf{M}_{\mathcal{A}}\text{-Set}$ , is (not) Boolean, the paradigm  $\mathcal{A}$  is (*non*-)classical, the topos  $\mathbf{M}_{\mathcal{A}}\text{-Set}$  is the *proper world* of  $\mathcal{A}$ . A classical paradigm of maximal finite dimensionality, if it exists, is a *classic* paradigm.

Intuitively, we are to understand reflexors as elementary experiences (observations) of an individual, and, as emphasized in [1], the former are the basic construction material for everything that follows. In the absence of restrictions on logic, this axiomatic setup constitutes a rather weak system in the sense that there is a lot of room for pathological behavior (presence of *xenomorphs*), which justifies the somewhat extravagant terminology of [1]. In *our case*, however, i. e. for individuals with  $\mathbf{R}$  type *psychology* ( $\mathbf{F} \cong \mathbf{R}$ , where  $\mathbf{R}$  is the naturally ordered field of reals) and Boolean logic, a dramatic reduction of the specter of

possible structures takes place. For *decoherent* paradigms (i. e. rational finite-dimensional paradigms of  $\mathbf{R}$ -observer), it becomes possible to define *spacetime*, as the (multiplicative) Lie group of all invertible elements of  $\mathcal{A}$ . The main result of [1] is that in  $\mathbf{R}$  case the classic paradigm exists, is unique and isomorphic to the quaternion algebra  $\mathcal{H}$ , spacetime  $\Omega \cong \mathbf{R}^+ \otimes SU(2)$ , where  $\mathbf{R}^+$  is the multiplicative group of positive reals, and there is a unique Lorentz sensory metric on  $\mathbf{S}_{\mathcal{H}}$ , generated by a unique (up to real factor) sensory time  $\tilde{t}$  with components  $(1, 0, 0, 0)$  in the basis dual to the basis of the unit quaternions  $\mathbf{1}, \mathbf{i}, \mathbf{j}, \mathbf{k}$ . Thus in the monocism model the effective four-dimensionality of, and existence of a signature 2 metric are not axioms, as in the standard approach, but can be deduced from a specification of the operational logic of the observer.

There are two natural orthonormal frame fields  $e_n^*(x)$  and  $*e_n(x)$  on  $\Omega$  induced by translations of the basis  $\mathbf{1}, \mathbf{i}, \mathbf{j}, \mathbf{k}$ , in the tangent space  $\mathbf{S}_{\mathcal{H}}$  at  $\mathbf{1} \in \Omega$ , along the left and right-invariant vector fields, respectively, and by its subsequent normalization, in the sense of quaternion norm. Since the computations are similar, in this paper we deal only with the left-invariant fields. The frame field  $e_n^*(x)$  generates a natural teleparallel [2] connection on, and provides continuations,  $\tilde{t}(x)$  and  $\mathbf{C}(x)$ , of the sensory time and structure constant tensor to  $\Omega$ . Specifically,  $\tilde{t}(x)$  and  $\mathbf{C}(x)$  are declared constant in this frame field. Since the sensory metric is contraction of  $\tilde{t}$  and  $\mathbf{C}$ , this guaranties that the metric is canonical in  $e_n^*(x)$ .

(2) *Left-invariant vector fields and their integral curves in  $\Omega$ .* The quaternionic equation of the integral curves of the left-invariant vector fields over  $\Omega$  is

$$\frac{dx(\mu)}{d\mu} = p_0 * x(\mu), \quad (1)$$

where  $\mu$  is real parameter,  $*$  is quaternion multiplication,  $x(\mu) \in \Omega$ ,  $p_0 \in \mathbf{S}_{\mathcal{H}}$  is the initial tangent vector to the curve (at  $x(0)$ ). The basis  $\mathbf{1}, \mathbf{i}, \mathbf{j}, \mathbf{k}$  generates a global system of normal coordinates  $(x^n)$  in  $\Omega$ , in which a quaternion  $a\mathbf{1} + b\mathbf{i} + c\mathbf{j} + d\mathbf{k}$  has coordinates  $x^0 = a$ ,  $x^1 = b$ ,  $x^2 = c$ ,  $x^3 = d$ . Note that  $x^n$  are not all zero, since  $\mathbf{0} \notin \Omega$ . Letting the components of  $p_0$  be  $p_0^n$  in the coordinate basis  $\frac{\partial}{\partial x^n}$  at  $\mathbf{1}$ , we can present  $p_0$  in the left matrix [3] form

$$p_0^{left} = \begin{pmatrix} p_0^0 & -p_0^1 & -p_0^2 & -p_0^3 \\ p_0^1 & p_0^0 & -p_0^3 & p_0^2 \\ p_0^2 & p_0^3 & p_0^0 & -p_0^1 \\ p_0^3 & -p_0^2 & p_0^1 & p_0^0 \end{pmatrix}.$$

Then (1) becomes a fourth-order linear system of ordinary differential equations:

$$\begin{aligned} \frac{dx^0(\mu)}{d\mu} &= p_0^0 x^0(\mu) - p_0^1 x^1(\mu) - p_0^2 x^2(\mu) - p_0^3 x^3(\mu), \\ \frac{dx^1(\mu)}{d\mu} &= p_0^1 x^0(\mu) + p_0^0 x^1(\mu) - p_0^3 x^2(\mu) + p_0^2 x^3(\mu), \\ \frac{dx^2(\mu)}{d\mu} &= p_0^2 x^0(\mu) + p_0^3 x^1(\mu) + p_0^0 x^2(\mu) - p_0^1 x^3(\mu), \end{aligned}$$

$$\frac{dx^3(\mu)}{d\mu} = p_0^3 x^0(\mu) - p_0^2 x^1(\mu) + p_0^1 x^2(\mu) + p_0^0 x^3(\mu),$$

with the initial condition  $x^n(0) = x_0^n$ . The solution (integral curves of the left-invariant vector fields) is

$$\begin{aligned} x^0(\mu) &= \exp(p_0^0 \mu) \left( \frac{-p_0^1 x_0^1 - p_0^2 x_0^2 - p_0^3 x_0^3}{P_0} \sin(P_0 \mu) + x_0^0 \cos(P_0 \mu) \right), \\ x^1(\mu) &= \exp(p_0^0 \mu) \left( \frac{p_0^1 x_0^0 + p_0^2 x_0^3 - p_0^3 x_0^2}{P_0} \sin(P_0 \mu) + x_0^1 \cos(P_0 \mu) \right), \\ x^2(\mu) &= \exp(p_0^0 \mu) \left( \frac{p_0^2 x_0^0 + p_0^3 x_0^1 - p_0^1 x_0^3}{P_0} \sin(P_0 \mu) + x_0^2 \cos(P_0 \mu) \right), \\ x^3(\mu) &= \exp(p_0^0 \mu) \left( \frac{p_0^3 x_0^0 + p_0^1 x_0^2 - p_0^2 x_0^1}{P_0} \sin(P_0 \mu) + x_0^3 \cos(P_0 \mu) \right), \end{aligned}$$

where  $P_0 \equiv \sqrt{(p_0^1)^2 + (p_0^2)^2 + (p_0^3)^2}$ . Each curve is in the exponential four-funnel of radius  $\exp(p_0^0 \mu) r_0$ , where  $r_0 \equiv \sqrt{(x_0^0)^2 + (x_0^1)^2 + (x_0^2)^2 + (x_0^3)^2}$ . This induces a system of natural coordinates  $\eta, \chi, \theta, \phi$  in  $\Omega$ :

$$\begin{aligned} x^0 &= \exp(\eta) \cos(\chi), \\ x^1 &= \exp(\eta) \sin(\chi) \sin(\theta) \cos(\phi), \\ x^2 &= \exp(\eta) \sin(\chi) \sin(\theta) \sin(\phi), \\ x^3 &= \exp(\eta) \sin(\chi) \cos(\theta). \end{aligned} \tag{2}$$

(3) *Orthonormal frame field on  $\Omega$ .* The basis vector  $\mathbf{1} \in \mathbf{S}_H$  generates a left-invariant vector field over  $\Omega$  whose vector at a point  $x$  with coordinates  $x^n$  has components  $x^n$  in the coordinate basis  $\frac{\partial}{\partial x^n}$ , and the normalization gives  $\frac{x^n}{r}$ , where  $r \equiv \sqrt{(x^0)^2 + (x^1)^2 + (x^2)^2 + (x^3)^2}$ . Similarly for  $\mathbf{i}, \mathbf{j}, \mathbf{k}$ , so, at the point  $x$  we have four vectors, that are, as easily checked, linear independent, and constitute a (noncoordinate) basis  $e_m^*(x)$ :

$$\begin{aligned} e_0^*(x) &= \frac{x^0}{r} \frac{\partial}{\partial x^0} + \frac{x^1}{r} \frac{\partial}{\partial x^1} + \frac{x^2}{r} \frac{\partial}{\partial x^2} + \frac{x^3}{r} \frac{\partial}{\partial x^3}, \\ e_1^*(x) &= -\frac{x^1}{r} \frac{\partial}{\partial x^0} + \frac{x^0}{r} \frac{\partial}{\partial x^1} - \frac{x^3}{r} \frac{\partial}{\partial x^2} + \frac{x^2}{r} \frac{\partial}{\partial x^3}, \\ e_2^*(x) &= -\frac{x^2}{r} \frac{\partial}{\partial x^0} + \frac{x^3}{r} \frac{\partial}{\partial x^1} + \frac{x^0}{r} \frac{\partial}{\partial x^2} - \frac{x^1}{r} \frac{\partial}{\partial x^3}, \\ e_3^*(x) &= -\frac{x^3}{r} \frac{\partial}{\partial x^0} - \frac{x^2}{r} \frac{\partial}{\partial x^1} + \frac{x^1}{r} \frac{\partial}{\partial x^2} + \frac{x^0}{r} \frac{\partial}{\partial x^3}. \end{aligned}$$

The corresponding integral curves are  $e_0^*(x)$ :

$$\begin{aligned} x^0(\mu) &= \frac{x_0^0 \mu}{r_0} + x_0^0, x^1(\mu) = \frac{x_0^1 \mu}{r_0} + x_0^1, \\ x^2(\mu) &= \frac{x_0^2 \mu}{r_0} + x_0^2, x^3(\mu) = \frac{x_0^3 \mu}{r_0} + x_0^3; \end{aligned}$$

$e_1^*(x)$ :

$$\begin{aligned}x^0(\mu) &= -\sqrt{(x_0^0)^2 + (x_0^1)^2} \sin\left(\frac{\mu}{r_0} + \arctan\left(\frac{x_0^3}{x_0^2}\right)\right), \\x^1(\mu) &= \sqrt{(x_0^0)^2 + (x_0^1)^2} \cos\left(\frac{\mu}{r_0} + \arctan\left(\frac{x_0^3}{x_0^2}\right)\right), \\x^2(\mu) &= \sqrt{(x_0^2)^2 + (x_0^3)^2} \cos\left(\frac{\mu}{r_0} + \arctan\left(\frac{x_0^3}{x_0^2}\right)\right), \\x^3(\mu) &= \sqrt{(x_0^2)^2 + (x_0^3)^2} \sin\left(\frac{\mu}{r_0} + \arctan\left(\frac{x_0^3}{x_0^2}\right)\right);\end{aligned}$$

$e_2^*(x)$ :

$$\begin{aligned}x^0(\mu) &= \sqrt{(x_0^0)^2 + (x_0^2)^2} \sin\left(\frac{\mu}{r_0} + \arctan\left(\frac{x_0^1}{x_0^3}\right)\right), \\x^1(\mu) &= \sqrt{(x_0^3)^2 + (x_0^1)^2} \cos\left(\frac{\mu}{r_0} + \arctan\left(\frac{x_0^1}{x_0^3}\right)\right), \\x^2(\mu) &= -\sqrt{(x_0^0)^2 + (x_0^2)^2} \cos\left(\frac{\mu}{r_0} + \arctan\left(\frac{x_0^1}{x_0^3}\right)\right), \\x^3(\mu) &= \sqrt{(x_0^3)^2 + (x_0^1)^2} \sin\left(\frac{\mu}{r_0} + \arctan\left(\frac{x_0^1}{x_0^3}\right)\right);\end{aligned}$$

$e_3^*(x)$ :

$$\begin{aligned}x^0(\mu) &= \sqrt{(x_0^0)^2 + (x_0^3)^2} \cos\left(\frac{\mu}{r_0} + \arctan\left(\frac{x_0^3}{x_0^0}\right)\right), \\x^1(\mu) &= \sqrt{(x_0^1)^2 + (x_0^3)^2} \sin\left(\frac{\mu}{r_0} + \arctan\left(\frac{x_0^3}{x_0^0}\right)\right), \\x^2(\mu) &= -\sqrt{(x_0^1)^2 + (x_0^3)^2} \cos\left(\frac{\mu}{r_0} + \arctan\left(\frac{x_0^3}{x_0^0}\right)\right), \\x^3(\mu) &= \sqrt{(x_0^0)^2 + (x_0^3)^2} \sin\left(\frac{\mu}{r_0} + \arctan\left(\frac{x_0^3}{x_0^0}\right)\right).\end{aligned}$$

(4) *Telegravity in  $\Omega$ .* To compute the Christoffel symbols, torsion tensor and the background metric explicitly, we make the standard teleparallel transport declaration: a vector at  $x \in \Omega$  is parallel (equal) to a vector at  $y \in \Omega$  if their components are proportional (equal) in the orthonormal frame  $e_n^*$  at those points. In the coordinate basis  $\frac{\partial}{\partial x^n}$  the nonzero Christoffels are

$$\begin{aligned}\Gamma_{11}^0 &= \Gamma_{22}^0 = \Gamma_{33}^0 = -\Gamma_{01}^1 = -\Gamma_{23}^1 = \Gamma_{32}^1 = \\&= -\Gamma_{02}^2 = \Gamma_{13}^2 = -\Gamma_{31}^2 = -\Gamma_{03}^3 = -\Gamma_{12}^3 = \Gamma_{21}^3 = \frac{x^0}{r^2},\end{aligned}$$

$$\begin{aligned} -\Gamma_{10}^0 &= \Gamma_{23}^0 = -\Gamma_{32}^0 = \Gamma_{00}^1 = \Gamma_{22}^1 = \Gamma_{33}^1 = \\ -\Gamma_{03}^2 &= -\Gamma_{12}^2 = \Gamma_{31}^2 = \Gamma_{02}^3 = -\Gamma_{20}^3 = -\Gamma_{13}^3 = \frac{x^1}{r^2}, \end{aligned}$$

$$\begin{aligned} -\Gamma_{13}^0 &= -\Gamma_{20}^0 = \Gamma_{31}^0 = \Gamma_{03}^1 = -\Gamma_{21}^1 = -\Gamma_{30}^1 = \\ \Gamma_{00}^2 &= \Gamma_{11}^2 = \Gamma_{33}^2 = -\Gamma_{01}^3 = \Gamma_{10}^3 = -\Gamma_{23}^3 = \frac{x^2}{r^2}, \end{aligned}$$

$$\begin{aligned} \Gamma_{12}^0 &= -\Gamma_{21}^0 = -\Gamma_{30}^0 = \Gamma_{02}^1 = \Gamma_{20}^1 = -\Gamma_{31}^1 = \\ \Gamma_{01}^2 &= -\Gamma_{10}^2 = -\Gamma_{32}^2 = \Gamma_{00}^3 = \Gamma_{11}^3 = \Gamma_{22}^3 = \frac{x^3}{r^2}. \end{aligned}$$

The nonzero components of torsion tensor in the basis  $\frac{\partial}{\partial\eta}, \frac{\partial}{\partial\chi}, \frac{\partial}{\partial\theta}, \frac{\partial}{\partial\phi}$  of natural coordinates (2) are

$$T_{10}^1 = -T_{01}^1 = T_{20}^2 = -T_{02}^2 = T_{30}^3 = -T_{03}^3 = 1, T_{21}^3 = -T_{12}^3 = \frac{2}{\sin(\theta)},$$

$$T_{13}^2 = -T_{31}^2 = 2\sin(\theta), T_{32}^1 = -T_{23}^1 = 2\sin(\chi)^2\sin(\theta).$$

In the basis  $\frac{\partial}{\partial\eta}, \frac{\partial}{\partial\chi}, \frac{\partial}{\partial\theta}, \frac{\partial}{\partial\phi}$  the background metric takes form

$$\mathbf{g} = \begin{pmatrix} \exp(2\eta) & 0 & 0 & 0 \\ 0 & -\exp(2\eta) & 0 & 0 \\ 0 & 0 & -\exp(2\eta)\sin(\chi)^2 & 0 \\ 0 & 0 & 0 & -\exp(2\eta)\sin(\chi)^2\sin(\theta)^2 \end{pmatrix}.$$

(5) *Cosmological time and structure field on  $\Omega$ .* In the basis  $e_n^*(x)$  the continuation of the sensory time (*cosmological time* of the paradigm  $\mathcal{H}$ )  $\tilde{t}(x)$  has components  $t_0(x) = 1, t_1(x) = t_2(x) = t_3(x) = 0$ ; in the basis  $\frac{\partial}{\partial x^n}$  we have  $t_m(x) = \frac{x^m}{r}$ , and in the basis  $\frac{\partial}{\partial\eta}, \frac{\partial}{\partial\chi}, \frac{\partial}{\partial\theta}, \frac{\partial}{\partial\phi}$  we have  $t_0(x) = \exp(\eta), t_1(x) = t_2(x) = t_3(x) = 0$ . Note that cosmological time is a tensor.

In the basis  $e_n^*(x)$  the components of  $\mathbf{C}(x)$  (the *structure field* of  $\Omega$ ) are constant and equal to the quaternion structure constants. This means, in particular, that  $\mathbf{C}(x)$  induces the quaternion multiplication at each point  $x \in \Omega$ , i.e. each fiber of the tangent bundle of  $\Omega$  is an isomorphic copy of the quaternion algebra. The nonzero components  $C_{nm}^q(x)$  in the basis  $\frac{\partial}{\partial\eta}, \frac{\partial}{\partial\chi}, \frac{\partial}{\partial\theta}, \frac{\partial}{\partial\phi}$  are

$$\begin{aligned} C_{00}^0 &= -C_{11}^0 = C_{01}^1 = C_{10}^1 = C_{02}^2 = C_{20}^2 = C_{03}^3 = C_{30}^3 = \exp(\eta), \\ C_{33}^0 &= -\exp(\eta)\sin(\chi)^2\sin(\theta)^2, C_{23}^1 = -C_{32}^1 = \exp(\eta)\sin(\chi)^2\sin(\theta), \end{aligned}$$

$$C_{31}^2 = -C_{13}^2 = \exp(\eta) \sin(\theta), C_{12}^3 = -C_{21}^3 = \frac{\exp(\eta)}{\sin(\theta)}.$$

Thus obtained information about natural geometry of  $\Omega$  is actually quite remarkable: all three basic features of this simple structure connected in a unique and unexpected way to Boolean logic - the effective dimensionality, signature and global properties of the metric - conform to the corresponding properties of the world we currently observe.

#### *References*

- [1] V. Trifonov, *Europhys. Lett.*, **32 (8)** (1995) pp. 621-626, (physics/0301044).
- [2] K. Hayashi, T. Shirafuji, *Phys. Rev.*, **D19** (1979) pp. 3524-3553.
- [3] Pertti Lounesto, *Clifford Algebras and Spinors*, London Mathematical Society Lecture Notes, Vol. 239 (Cambridge University Press), 1997, p. 72.