

# The effective inertial acceleration due to oscillations of the gravitational potential: footprints in the solar system

D.L. Khokhlov

*Sumy State University, R.-Korsakov St. 2,*

*Sumy 40007, Ukraine*

*E-mail: khokhlov@cafe.sumy.ua*

## Abstract

The conjecture is considered that every body induces the wave field which imposes oscillations on the gravitational potential of a body. The function for oscillations is chosen to prevent the gravitational collapse of the matter at the nucleus energy density. Oscillations of the gravitational potential of a body produce effective inertial outward acceleration for a particle orbiting around the body. Footprints of the effective inertial acceleration due to oscillations of the gravitational potentials of the Sun and Earth are investigated. The conjecture allows to explain the anomalous shift of the perihelion of Mercury and Icarus, the anomalous shift of the perigee of LAGEOS II, the anomalous acceleration acting on Pioneer 10, 11, the anomalous increase in the lunar semi-major axis. The advance of the Keplerian orbit for Earth, Jupiter, Neptune, Uranus caused by the effective inertial acceleration due to oscillations of the gravitational potential of the Sun is in agreement with the observational bounds from the planetary ephemeris.

The theory of gravity [1] faces the problem of singularities. In particular, singularities may arise as a result of the gravitational collapse of massive stars. A body contracted to the nucleus energy density resembles the neutron star [2]. There is a maximum mass for the neutron star of order of the mass of the Sun,  $m_{max} \sim m_{\odot}$ . For the neutron star with the mass less than the maximum mass, the pressure due to the degenerated neutron Fermi gas balances the gravity of the star. For the neutron star with the mass more than the maximum mass, the gravity of the star overcomes the pressure due to the degenerated neutron Fermi gas, and the star goes to the singularity.

The radius of the first Bohr's orbit of the Hydrogen atom  $a_0$  and the classical radius of electron  $r_e$  are related as

$$\frac{r_e}{a_0} = \alpha^2 \quad (1)$$

where  $\alpha$  is the fine structure constant. The velocity of the electron at the first Bohr's orbit is given by

$$v_e = \alpha c \quad (2)$$

where  $c$  is the velocity of light. The radius of the nucleus of the Hydrogen atom  $r_n$  is of order of the classical radius of electron

$$r_n \sim r_e. \quad (3)$$

If to contract the Hydrogen atom to the radius of the nucleus,  $r_H \rightarrow r_n$ , the fine structure constant tends to unity,  $\alpha \rightarrow 1$ , and the velocity of the electron tends to the velocity of light,  $v_e \rightarrow c$ . Since the velocity of light is a limiting one, it is reasonable to think that the

radius of the nucleus is a limiting one, and hence the nucleus energy density is a limiting one.

It is reasonable to assume that there is some mechanism which allows to switch off the gravity at a limiting scale thus preventing appearance of singularities in the gravitational collapse of the matter. Below we shall consider such a mechanism, with the nucleus energy density being chosen as a limiting scale.

Assume that every body induces the wave field which carries scalar perturbation of the gravitational potential. Thus the wave field imposes oscillations on the gravitational potential of a body

$$\Phi = -\frac{Gm}{r} + \Delta\Phi \sin \omega_{gr}t \quad (4)$$

where  $G$  is the Newtonian constant,  $m$  is the mass of the body,  $\omega_{gr}$  is the frequency of oscillations. On the average oscillations do not change the gravitational potential of a body. When a particle orbits around a body, oscillations of the gravitational potential of the body produce radial oscillations around the Keplerian orbit of a particle.

Assume that the value of oscillations of the gravitational potential is a function of the matter density such that the value of oscillations of the gravitational potential is equal to the Newtonian potential at the radius of neutron star

$$\Delta\Phi = \frac{Gmr_{NS}^2}{r_0^3} \quad (5)$$

where  $r_0$  is the radius of a body,  $r_{NS}$  is its radius of neutron star. Suppose that the frequency of oscillations is given by

$$\omega_{gr} = \left( \frac{Gm}{4r_{NS}^3} \right)^{1/2}. \quad (6)$$

The frequency of oscillations is large,  $\omega_{gr} \gg 1/T = (Gm/r^3)^{1/2}$ . Then the radial displacement around the Keplerian orbit of a particle is small [3]

$$\delta r = \frac{\Delta\Phi}{\omega_{gr}^2 r}. \quad (7)$$

Kinetic energy due to oscillation motion is given by

$$E_{osc} = \frac{\Delta\Phi^2}{4\omega_{gr}^2 r^2} = \frac{Gmr_{NS}^7}{r_0^6 r^2}. \quad (8)$$

Thus we arrive at the effective Newtonian gravity

$$\Phi = \Phi_N + E_{osc} = \Phi_N \left( 1 - \frac{r_{NS}^7}{r_0^6 r} \right). \quad (9)$$

The value of oscillations of the gravitational potential grows under the contraction of a body. When the radius of a body reaches the radius of neutron star,  $r_0 \rightarrow r_{NS}$ , oscillations of the gravitational potential tends to the value,  $\Delta\Phi \rightarrow Gm/r_{NS}$ . The radial displacement due to oscillations of the gravitational potential tends to the radius of neutron star,  $\delta r \rightarrow r_{NS}$ , and the kinetic energy due to oscillations of the gravitational potential tends

to the value,  $E_{osc} \rightarrow Gm/r_{NS}$ . Then oscillations of the gravitational potential switches off the gravity of a body contracted to the radius of neutron star. Thus the radius of neutron star is a limiting one. Oscillations of the gravitational potential prevent the gravitational collapse of the matter to singularity. Thus the conjecture proposed may resolve the problem of singularities in the theory of gravity.

Let a particle orbit around a body. Oscillations of the gravitational potential of the body produce effective oscillations of the circular velocity of the particle

$$\Delta v = (\Delta\Phi \sin \omega_{gr} t)^{1/2}. \quad (10)$$

As a result effective inertial acceleration directed outward the body arises

$$w_{eff} = \frac{\Delta\Phi}{r}. \quad (11)$$

The effective inertial acceleration does not modify the Newtonian gravity. Therefore displacement of the radius of the Keplerian orbit of a particle due to the effective inertial acceleration cannot be detected in the measurement of the orbit's radius.

The effective inertial acceleration due to oscillations of the gravitational potential of a body should lead to the precession of the Keplerian orbit of a particle under its motion in the gravitational field of a body. A small addition  $\delta\Phi$  to the potential of a body causes the shift of the perihelion of particle's orbit per revolution by the value [3]

$$\delta\varphi = \frac{\partial}{\partial M} \frac{2m'^2}{M} \int_0^\pi r^2 \delta\Phi d\varphi \quad (12)$$

where  $m'$  is the mass of the particle,  $M$  is the angular momentum of the particle. The non-perturbed orbit is given by

$$r = \frac{p}{1 + e \cos \varphi} \quad (13)$$

with

$$p = \frac{M^2}{Gm'^2m} \quad p = a(1 - e^2) \quad (14)$$

where  $p$  is the orbit's latus rectum,  $e$  is the eccentricity,  $a$  is the semi-major axis. Let the effective potential due to oscillations of the gravitational potential of a body give a small addition to the potential of a body,  $\delta\Phi = \Delta\Phi$ . Then integration of eq. (12) with the use of eqs. (13,14) gives the shift (advance) of the perihelion of a particle due to oscillations of the gravitational potential of a body per revolution

$$\delta\varphi \approx \frac{6\pi a(1 - e^2)\Delta\Phi}{Gm}. \quad (15)$$

Here we omit the terms containing  $\cos^2 \varphi$  since an addition  $\delta\Phi$  leads to the deviation from the Keplerian orbit in this order.

Investigate footprints of the effective inertial acceleration due to oscillations of the gravitational potential in the solar system. Suppose that the Sun induces the wave field which imposes oscillations on the gravitational potential of the Sun. Estimate the value of oscillations of the gravitational potential of the Sun adopting the density of neutron

star,  $\rho_{NS} \sim 10^{14}$  g/cm<sup>3</sup>. Then the radius of neutron star for the Sun is equal to  $r_{NS} = (3m_{\odot}/4\pi\rho_{NS})^{1/3} = 1.7 \times 10^6$  cm. The value of oscillations of the gravitational potential of the Sun is equal to  $\Delta\Phi_{\odot} = Gm_{\odot}r_{NS}^2/r_{\odot}^3 = 1.1 \times 10^6$  cm<sup>2</sup>/s<sup>2</sup>.

Observations [4] give the value, 43 arcseconds per century, for the anomalous shift (advance) of the perihelion of Mercury. This anomalous shift is now explained within the framework of the Einstein general relativity [4]. Suppose that the value, 43 arcseconds, is brought about by the effective inertial acceleration due to oscillations of the gravitational potential of the Sun. Determine the value of oscillations of the gravitational potential of the Sun from the data on the anomalous shift of the perihelion of Mercury with the use of the Mercury data, the semi-major axis  $a = 0.387$  AU, the eccentricity  $e = 0.206$ , the period of revolution  $T = 0.241$  yr. Then from eq. (15) we obtain the value of oscillations of the gravitational potential of the Sun,  $\Delta\Phi_{\odot} = 6.4 \times 10^5$  cm<sup>2</sup>/s<sup>2</sup>.

Determine the shift of the perihelion of the asteroid Icarus due to oscillations of the gravitational potential of the Sun with the use of the Icarus data, the semi-major axis  $a = 1.076$  AU, the eccentricity  $e = 0.827$ , the period of revolution  $T = 1.116$  yr. In accordance with eq. (15) the value of oscillation of the gravitational potential of the Sun,  $\Delta\Phi_{\odot} = 6.4 \times 10^5$  cm<sup>2</sup>/s<sup>2</sup>, gives the shift (advance) of the perihelion of Icarus, 8.5 arcseconds per century. The observed anomalous shift (advance) is  $9.8 \pm 0.8$  arcseconds per century [4]. Thus the calculated value lies below the uncertainty margin by around 5% of the observed effect.

Oscillations of the gravitational potential of the Sun yield the effective outward acceleration of the Earth. This effective outward acceleration may be seen in ranging of distant spacecraft. The effective outward acceleration due to oscillations of the gravitational potential of the Sun should give the blue shift into the frequency of light at the Earth

$$\frac{\Delta\omega}{\omega} = \frac{\Delta\Phi_{\odot}t}{cr_{SE}} \quad (16)$$

where  $r_{SE}$  is the distance between the Earth and Sun,  $t$  is the time of two-leg light travel. In ranging of distant spacecraft, the acceleration of the Earth outward the Sun looks like the acceleration of the spacecraft inward the Sun. Then one may interpret the shift of the reference frequency given by eq. (16) as a shift of the observed re-transmitted frequency due to the acceleration of the spacecraft inward the Sun. The acceleration of the Earth gives contribution into the shift of the reference frequency during the time of two-leg light travel, while the acceleration of the spacecraft gives contribution into the shift of the observed re-transmitted frequency during the time of one-leg light travel. Then, if to interpret the observed anomalous shift of frequency as the anomalous acceleration of the spacecraft inward the Sun, its value is doubled in comparison with the effective outward acceleration of the Earth. Thus the effective outward acceleration of the Earth due to oscillations of the gravitational potential of the Sun,  $\Delta\Phi_{\odot} = 6.4 \times 10^5$  cm<sup>2</sup>/s<sup>2</sup>, may be interpreted as the inward acceleration of the spacecraft,  $w = 2\Delta\Phi_{\odot}/r_{SE} = 8.5 \times 10^{-8}$  cm/s<sup>2</sup>. Analyses of radio Doppler and ranging data from distant spacecraft in the solar system indicated that an apparent anomalous acceleration is acting on Pioneer 10 and 11, with a magnitude  $w_P = (8.74 \pm 1.25) \times 10^{-8}$  cm/s<sup>2</sup>, directed towards the Sun [5]. Thus the effective outward acceleration of the Earth due to oscillations of the gravitational potential of the Sun mimics the anomalous acceleration acting on Pioneer 10 and 11. Thus the value of oscillations of the gravitational potential of the Sun explaining

the anomalous shift of the perihelion of Mercury may explain the anomalous acceleration acting on Pioneer 10 and 11.

Oscillations of the gravitational potential of the Sun yield the radial displacement around the Keplerian orbit of a planet. In view of eq. (7), the radial displacement around the Keplerian orbit of the Earth is of order  $\delta r \sim 10^{-15}$  cm. This value is too small to be seen in observations. The Viking ranging data [6] determine the Earth-Mars radius to around a 7.5 m accuracy.

The effective inertial acceleration due to oscillations of the gravitational potential of the Sun yields the advance of the Keplerian orbit,  $\delta\varphi \sim 6\pi\Delta\Phi_\odot/\Phi_N$ , where  $\Phi_N$  is the Newtonian potential of the Sun at the radius  $r$ . The dark matter restricted within the radius  $r$  yields the advance of the Keplerian orbit,  $\delta\varphi \sim 2\pi\Phi_{DM}/\Phi_N$ , where  $\Phi_{DM}$  is the potential of the dark matter restricted within the radius  $r$ . Hence the effective inertial acceleration due to oscillations of the gravitational potential of the Sun mimics the effective mass restricted within the radius  $r$

$$m_{eff} = \frac{\Delta\Phi_\odot r}{3G}. \quad (17)$$

Compare the effective mass due to oscillations of the gravitational potential of the Sun,  $\Delta\Phi_\odot = 6.4 \times 10^5 \text{ cm}^2/\text{s}^2$ , with the observational bounds for the dark matter from the planetary ephemeris. Within the Earth's orbit the effective mass is  $m_{eff} = 2.4 \times 10^{-8}m_\odot$  while the observational bound is  $m_{DM} \leq 7 \times 10^{-8}m_\odot$  [7]. Within Jupiter's orbit the effective mass is  $m_{eff} = 1.2 \times 10^{-7}m_\odot$  while the observational bound is  $m_{DM} = (0.12 \pm 0.027) \times 10^{-6}m_\odot$  [8]. Within Uranus's orbit the effective mass is  $m_{eff} = 4.7 \times 10^{-7}m_\odot$  while the observational bound is  $m_{DM} \leq 0.5 \times 10^{-6}m_\odot$  [8]. Within Neptune's orbit the effective mass is  $m_{eff} = 7.3 \times 10^{-7}m_\odot$  while the observational bound is  $m_{DM} \leq 3 \times 10^{-6}m_\odot$  [8]. Thus the effective mass due to oscillations of the gravitational potential of the Sun is in agreement with the observational bounds for the Earth's, Jupiter's, Neptune's, and Uranus's orbits.

Oscillations of the gravitational potential of the Sun yield the effective inertial acceleration of the Moon with respect to the Earth,  $w \approx \Delta\Phi_\odot r_{EM} \sin \Omega t / r_{SE}^2$ , where  $r_{EM}$  is the distance between the Earth and Moon,  $\Omega$  is the angular velocity of the Moon. This leads to the polarization of the Moon's orbit in the direction of the Sun, with the Earth-Moon distance decreasing. As a result the Moon's circular velocity increases by the value,  $\Delta v = (\sqrt{2}\Delta\Phi_\odot r_{EM} / 2r_{SE})^{1/2}$ , average for the period of revolution of the Moon. Then inertial acceleration of the Moon outwards the Earth increases by the value,  $w = \sqrt{2}\Delta\Phi_\odot / 2r_{SE}$ , average for the period of revolution of the Moon. Oscillations of the gravitational potential of the Sun,  $\Delta\Phi_\odot = 6.4 \times 10^5 \text{ cm}^2/\text{s}^2$ , yield the effective inertial acceleration of the Moon outwards the Earth,  $w = 3.0 \times 10^{-8} \text{ cm/s}^2$ .

As follows from telescopic observations of the Earth's rotation collected for the last 300 years, e.g. [10], the rate of deceleration of the Earth's rotation is  $\dot{T}_E = 1.4 \times 10^{-5} \text{ s/yr}$ . As a result the lunar semi-major axis increases with a rate,  $\dot{a}_{tel} = 2.53 \text{ cm/yr}$ . Measurement of the Earth-Moon distance in lunar laser ranging [11] gives the value,  $\dot{a}_{LLR} = 3.82 \text{ cm/yr}$ , hence points to the anomalous increase in the lunar semi-major axis with an excessive rate,  $\dot{a}_{LLR} - \dot{a}_{tel} = 1.29 \pm 0.2 \text{ cm/yr}$ . This corresponds to the anomalous outward acceleration of the Moon,  $w = (\dot{a}_{LLR} - \dot{a}_{tel})c/r_{EM} = (3.2 \pm 0.5) \times 10^{-8} \text{ cm/s}^2$ . Thus the effective inertial acceleration of the Moon due to oscillations of the gravitational potential of the

Sun explains the main part (around 94%) of the anomalous outward acceleration of the Moon.

Suppose that the Earth induces the wave field which imposes oscillations on the gravitational potential of the Earth. Estimate the value of oscillations of the gravitational potential of the Earth. The radius of neutron star for the Earth is equal to  $r_{NS} = (3m_{\oplus}/4\pi\rho_{NS})^{1/3} = 2.4 \times 10^4$  cm. The value of oscillations of the gravitational potential of the Earth is equal to  $\Delta\Phi_{\oplus} = Gm_{\oplus}r_{NS}^2/r_{\oplus}^3 = 9 \times 10^2$  cm<sup>2</sup>/s<sup>2</sup>.

Observations of the Earth's satellite LAGEOS II [9] give the value, 3.4 arcseconds per year, for the anomalous shift (advance) of the perigee of LAGEOS II. This anomalous shift is now explained within the framework of the Einstein general relativity [9]. Suppose that the value, 3.4 arcseconds, is brought about by the effective inertial acceleration due to oscillations of the gravitational potential of the Earth. Determine the value of oscillations of the gravitational potential of the Earth from the data on the anomalous shift of the perigee of LAGEOS II with the use of the LAGEOS II data, the semi-major axis  $a = 1.2163 \times 10^9$  cm, the eccentricity  $e = 0.014$ , the period of revolution  $T = 3.758$  hr, the inclination  $I = 52.65^\circ$ . Then from eq. (15) we obtain the value of oscillations of the gravitational potential of the Earth,  $\Delta\Phi_{\oplus} = 1.2 \times 10^2$  cm<sup>2</sup>/s<sup>2</sup>. Estimate the effect of oscillations of the gravitational potential of the Sun on the motion of LAGEOS II,  $\Delta\Phi = \sqrt{2}\Delta\Phi_{\odot}a \cos I/2r_{SE} = 0.3 \times 10^2$  cm<sup>2</sup>/s<sup>2</sup>. In view of this effect, we obtain the value of oscillations of the gravitational potential of the Earth,  $\Delta\Phi_{\oplus} = 0.9 \times 10^2$  cm<sup>2</sup>/s<sup>2</sup>.

Oscillations of the gravitational potential of the Earth,  $\Delta\Phi_{\oplus} = 0.9 \times 10^2$  cm<sup>2</sup>/s<sup>2</sup>, yield the effective inertial outward acceleration of the Moon,  $w = \Delta\Phi_{\oplus}/r_{EM} = 0.2 \times 10^{-8}$  cm/s<sup>2</sup>. This acceleration explains around 6% of the anomalous outward acceleration of the Moon. The overall effective inertial acceleration of the Moon due to oscillations of the gravitational potential of the Sun and Earth is  $w = 3.0 + 0.2 = 3.2 \times 10^{-8}$  cm/s<sup>2</sup> and thus completely explains the anomalous outward acceleration of the Moon.

We have considered the conjecture that every body induces the wave field which imposes oscillations on the gravitational potential of a body. The function for oscillations is chosen to prevent the gravitational collapse of the matter at the nucleus energy density. The radial displacement around the Keplerian orbit of a planet due to oscillations of the gravitational potential of the Sun is too small to be seen in observations. Oscillations of the gravitational potential produce effective inertial outward acceleration for a particle orbiting around the body. Footprints of the effective inertial acceleration due to oscillations of the gravitational potential of the Sun may be revealed as an anomalous shift of the perihelion of the Keplerian orbit of a planet or as an anomalous shift of the frequency of light seen in ranging. Although we have rough estimation of the value of oscillations of the gravitational potential of the Sun it is possible to perform the accurate testing of the conjecture by comparing the results of different observations. The value of oscillations of the gravitational potential of the Sun obtained from the anomalous shift of the perihelion of Mercury is consistent with that obtained from the anomalous shift of the frequency of light at the Earth now interpreted as the anomalous acceleration acting on Pioneer 10, 11. Also the value of oscillations of the gravitational potential of the Sun approximately explains the anomalous shift (advance) of the perihelion of Icarus, with the calculated value lies below the uncertainty margin by around 5% of the observed effect. Also the value of oscillations of the gravitational potential of the Sun explains the main part (around 94%) of the anomalous outward acceleration of the Moon. The value of oscillations of the

gravitational potential of the Earth obtained from the anomalous shift of the perigee of LAGEOS II explains the retaining part (around 6%) of the anomalous outward acceleration of the Moon. The advance of the Keplerian orbit for Earth, Jupiter, Neptune, Uranus caused by the effective inertial force due to oscillations of the gravitational potential of the Sun is in agreement with the observational bounds from the planetary ephemeris.

The conjecture considered provides an explanation of the anomalous precession (advance) of the Keplerian orbit alternative to the general relativity. So from the point of view of the conjecture considered the general relativity is questioned. Note that excluded the anomalous precession (advance) of the Keplerian orbit we may describe the planetary motion in the solar system within the Newtonian mechanics. At the same time the relativistic effects for the electromagnetic waves in the field of the gravitational potential are confirmed by the experiments. If to adopt the conjecture considered we should reinterpret the theory of relativity in the way that relativistic effects pertain only to the electromagnetism while the gravitation remains non-relativistic. That is the motion of the electromagnetic waves follows the relativistic dynamics while the motion of the massive bodies follows the Newtonian dynamics.

## References

- [1] C.W. Misner, K.S. Thorne, J.A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973).
- [2] S.L. Shapiro and S.A. Teucolsky, *Black Holes, White Dwarfs, and Neutron Stars* (Jhon Wiley and Sons, New York, 1983).
- [3] L. Landau and Ye. Lifshitz, *Mechanics* (Pergamon Press, Oxford, 1960).
- [4] S. Weinberg, *Gravitation and Cosmology* (Jhon Wiley and Sons, New York, 1972).
- [5] J.D. Anderson et al., Phys. Rev. **D65** (2002) 082004.
- [6] R.D. Reasenberget al., Astrophys. J. **234** (1979) L219.
- [7] D.R. Mikkelsen and M.J. Newman, Phys. Rev. **D16** (1977) 919.
- [8] J.D. Anderson et al., Astrophys. J. **448** (1995) 885.
- [9] L. Iorio, J. Geodetic Soc. Japan **48**(1) (2002) 13.
- [10] B.P. Pertsev, Izvestiya: Phys. Solid Earth **36** (2000) 218.
- [11] J.O. Dickey et al., Science **265** (1994) 482.