THE SUBGRID PROBLEM OF THE THERMAL CONVECTION IN THE EARTH'S LIQUID CORE

M. Reshetnyak¹², B. Steffen³

Abstract

The problem of the turbulent thermal convection in the Earth's liquid core is considered. Following assumptions on decreasing of the spatial scales due to the rapid rotation, we propose the subgrid model of the eddy diffusivity, which is used in the large-scale model. This approach makes it possible to model realistic regimes with small Ekman and Rossby numbers ($E \sim 10^{-14}$, $R_o \sim 10^{-8}$) and a sufficiently large Rayleigh number $R_a \sim 10^{12}$. The obtained estimate of the averaged kinetic energy is comparable with observations. The model includes rotation of the solid core due to the viscous torque.

Keywords: Rotating turbulence, geodynamo, control-volume method.

1 Introduction

Convection in the liquid core of the Earth, caused by the radiactive heating and compositional processes [1], is the subject of numerous researches, usually concerned with the geomagnetic field generation, also. The last few decades saw a fascinating development in this area [2]. Based on the MHD large-scale equations, numerical simulations can reproduce different geomagnetic and geophysical phenomena: various properties of the geomagnetic field (e.g., its reversals and spectrum), eastward rotation of the inner core of the Earth as well as the realistic ratio of the kinetic and magnetic energies [3].

However, the wide range of spatial and temporal scales make the direct numerical simulations (DNS) very cumbersome. The difficulty is caused by the small values of the transport coefficients: for instance the kinematic viscosity of the liquid core is $\nu^M = 10^{-6} \text{m}^2 \text{s}^{-1}$, and the thermal diffusivity: $\kappa^M = 10^{-5} \text{ m}^2 \text{s}^{-1}$ (here the subscript M corresponds to the molecular values). This gives estimates

⁰¹Institute of the Physics of the Earth, Russian Acad. Sci, 123995 Moscow, Russian Federation; e-mail: maxim@uipe-ras.scgis.ru

²Research Computing Center of Moscow State University, 119899, Moscow, Russian Federation; e-mail: rm@uipe.srcc.msu.su

³Central Institute for Applied Mathematics (ZAM) of Forshungszentrum Jülich, D-52425, Jülich, Germany; e-mail: b.steffen@fz-juelich.de

of the molecular Reynolds and Peclet numbers of $Re^M = \frac{V_{wd}L}{\nu^M} \sim 10^9$ and $Pe^M = \frac{V_{wd}L}{\kappa^M} \sim 10^8$, where $V_{wd} = 0.2^{\circ} \text{ year}^{-1}$ is the west drift velocity and $L = 3 \cdot 10^6$ m is the scale of the liquid core [4], which corresponds to the regime of the highly developed turbulence. In the case of the Kolmogorov's turbulence 3D DNS, simulations require $\sim Re^{9/4} = 10^{20}$ grid nodes [5]. Attempts to use the exact values of these parameters on the coarse grid lead to the numerical instabilities. The first intuitive models in geodynamo theory which suppressed instabilities at the small scales, e.g., the model of hyperdiffusivity [3], gave rise to new questions concerned with interpretation of the results obtained [6]. The more consequential way is an application of the semiempirical models of turbulence [7]. Usually, these models are based on assumptions on the cascade transfer similar to Kolmogorov's, which give descriptions of the average effect of the small-scale field fluctuations onto the large-scale flow in terms of the eddy diffusivity. The recent studies of the subgrid [8] and complex models [9, 10] of the thermal convection and dynamo problems in the rotating sphere revealed the principal possibility of describing the small-scale fluctuations in the turbulence with the desired Reynolds and Peclet numbers much like Kolmogorov's model.

These models work up to the regime of moderate rotation speed. Further increase of the Coriolis force can reduce the total kinetic energy and even suppress convection at all. From linear analysis it follows that the critical Rayleigh number depends on the Ekman number like $R_a^{cr} \sim E^{-1/3}$ [11]. Even though the molecular estimate of the Rayleigh number gives huge numbers $R_a^M \sim 10^{14}$ [12, 13], this value is only $5 \cdot 10^2$ times larger then the critical value R_a^{cr} [2]. Due to the rapid rotation of the Earth, the situation in the liquid core is more complicated and assumptions on similarity of the spectral characteristics of the fields must be checked very carefully. We show that the direct applications of the traditional models of the turbulence, based on the mix-length assumptions, lead to results that differ from the observations by orders of magnitude. The cause of such disagreement lies in the daily rapid rotation of the Earth, which gives rise to new characteristic spatial scales in the core [14]. As a result, the energy distribution in the spectrum changes, which makes the application of the Prandtl-Kolmogorov's approach to the eddy diffusivity estimate difficult. Convection at these new scales plays the crucial role in the energy balance of the whole system and changes the estimate of the total energy by orders of magnitude. Even a simple account of these effects leads to essential change of the rate of the energy dissipation and thus to a better agreement of LSS models with the observations.

In the section 2 we introduce the large-scale equations of the thermal convection and consider the Prandtl-Kolmogorov's assumptions on the eddy diffusion. In the section 3 we recall the basics of convection in a rapidly rotating body and estimate the subgrid diffusion. Afterwards, this estimate is used in the large-scale model, section 4. The discussion of results is in section 5.

2 The large-scale equations

The problem of the thermal convection in the Earth's core can be reduced to the problem in the spherical shell. Let the surface of the sphere, radius r_0 (in the spherical system of coordinates (r, θ, φ)), rotating with angular velocity Ω around the z-axis. This sphere contains a concentric solid inner sphere, radius r_i , and the outer spherical layer $(r_i < r < r_0)$ is filled with an incompressible liquid ($\mathbf{V} = 0$). The inner sphere is allowed to rotate freely around the z-axis due to viscous torque. Convection in the Boussinesq approximation in the outer sphere is described by the Navier-Stokes equation and by the heat flux equation. Choosing $L = r_0$ as the unit of length, velocity \mathbf{V} , time t and pressure p can be measured in units of κ^M/L , L^2/κ^M and $2\Omega\rho\kappa^M$, respectively. Then, the governing equations can be written in the form

$$R_o^M \left[\frac{\partial \mathbf{V}}{\partial t} + (\mathbf{V} \cdot \nabla) \mathbf{V} \right] = -\nabla p + -\mathbf{1}_{\mathbf{z}} \times \mathbf{V} + R_a^M Tr \mathbf{1}_{\mathbf{r}} + E^M \nabla \cdot \overset{\leftrightarrow}{\mathbf{S}},\tag{1}$$

$$\frac{\partial T}{\partial t} + (\mathbf{V} \cdot \nabla) \left(T + T_0 \right) = \nabla \cdot (\nabla T), \tag{2}$$

where $\mathbf{1}_{\mathbf{z}}$ is the unit vector in z-direction, $\mathbf{\ddot{S}}$ is the rate of the strain tensor and T the temperature fluctuations relatively to the imposed profile $T_0 = \frac{r_i/r-1}{1-r_i}$. The molecular Rossby R_o^M , Ekman E^M and Rayleigh R_a^M numbers appear in the equations

$$R_o^M = \frac{\kappa^M}{2\Omega L^2}, \qquad E^M = \frac{\nu^M}{2\Omega L^2}, \qquad R_a^M = \frac{\alpha g_0 \delta T L}{2\Omega \kappa^M}, \qquad (3)$$

where α is the coefficient of thermal expansion, g the gravity acceleration and δT is a temperature unit ($\delta T \sim 10^{-4}$ K, see [2]). It should be mentioned that the Rayleigh number for non-rotating bodies is usually given in the form $\tilde{R}_a^M = \alpha g \delta T L^3 / \kappa^M \nu^M$ and that $R_a^M = E^M \tilde{R}_a^M$.

The solid inner sphere is allowed to rotate freely around the z-axis due to viscous torque. The dimensionless momentum equation for the angular velocity ω of the inner sphere ($0 < r < r_i$) has the form

$$R_o^M I \frac{\partial \omega}{\partial t} = r_i E^M \oint_{\mathcal{S}} S_{\varphi r}|_{r=r_i} \sin \vartheta \, d\mathcal{S},\tag{4}$$

where I is the moment of inertia of the inner sphere S and $S_{\varphi r}$ is a component of the strain tensor in the spherical system of coordinates [15]. Equations (1–2, 4) are accompanied by the non-penetrating and no-slip boundary condition for velocity **V** and zero temperature fluctuations at the shell boundaries.

The system (1-2,4) was successfully studied in the regimes of the laminar convection using different numerical approaches [16, 2]. However, these regimes are still very far from the desired estimates for the Earth's liquid core $R_o^M = 10^{-8}$, $E^M = 10^{-14}$ and $R_a^M = 10^{14}$ [2]. Attempts to approach to these

parameters using DNS caused numerical instabilities and required application of turbulencs models [8]. However, even the direct usage of the known models of turbulence is not trivial.

To support this point we offer a simple estimate of the eddy diffusivity, based on the most popular mix-length model of the turbulence. Following the Prandtl-Kolmogorov hypotheses, the eddy diffusion at the scale l can be estimated as $\nu^T = (\varepsilon l^4)^{1/3}$, where $\varepsilon = v^3/l$ is the rate of energy dissipation and v is a velocity at the scale l. Even the largest estimate, based on the main scale l = L and the west drift velocity $V = 3 \cdot 10^{-3} \text{m s}^{-1}$, gives $\nu^T = 2 \cdot 10^3 \text{m}^2 \text{s}^{-1}$, giving an Ekman number of order $E^T = 2 \cdot 10^{-5}$. The more realistic estimate with $V_l = \delta V \sim V(\frac{l}{L})^{1/3}$ and the usual grid scale of $l \sim 3 \cdot 10^{-2}L$ gives $\nu^T = 15$ and $E^T = 10^{-7}$. On the other hand, this estimate of E^T would require resolution of about $N_{\varphi} \sim 2 \cdot 10^2$ columns [11], which need use of the most powerful modern computers. All this means that this estimate of ν^T will not provide the smooth field behaviour of fields assumed in the Kolmogorov's turbulence, when $E^T \geq 1$. Thus, the traditional methods underestimate the eddy diffusion ν^T .

Such situation corresponds to the case, where the classical ideas on the direct cascade of energy from the main scale L to the dissipative scale are violated and additional information on ε at the dissipative scale is needed. As we see below in the section 3, it appears that in the case of the rapid rotating body the energy in the spectrum is shifted to the small scales, those which DNS cannot resolve even at the onset of convection. This is the reason, why any attempts to estimate ν^T in the turbulent regime at scales compared with the grid resolution, lead to the non-selfconsistent behaviour of the turbulent model.

The way out of such difficulties is make proper assumptions on the spectral properties of the solution in the range of the high wave numbers.

3 The model of columns

The origin of the problem can be seen from the analysis of the linerized system (1-2) at the onset of convection in the limit of small Rossby and Ekman numbers. As it was shown in [11] already (see also recent paper [17]), at the onset of convection the structure of the flow tends to develop columns along z-direction, such that $\partial/\partial \varphi \sim \mathcal{O}(E^{-1/3})$, $\partial/\partial s \sim \mathcal{O}(E^{-1/6})$, $\partial/\partial z \sim \mathcal{O}(1)$, when $E = R_o \to 0$. Linerization of the system (1-2) leads to the balance of the Archemedean and viscous terms in the Navier-Stokes equation: $R_a T \sim E^{-1/3}V$. The balance of the convective and viscous terms in the heat-flux equation gives $V \sim E^{-2/3}T$, from which follows the estimate of the critical Rayleigh number $R_a^{\rm cr} \sim E^{-1/3}$. (For convenience we omitted index ^M.) Such, at the onset of convection for system (1-2), the flow is anisotropical with the smallest scale $l_E \sim E^{1/3}L$, defined by the balance of the Coriolis and viscous forces. Note that the scale $l_E \sim 10^{-5}$ is beyond the level of DNS. If this asymptotic is correct,

the critical Rayleigh number in the Earth's core is $R_a^{cr} \sim 10^5$ [2]. As we show below, the predicted column-like form of the flow is very important for estimates of the subgrid dissipation in the liquid core.

The main assumption is, that even in the turbulent regime believed to be in the Earth's liquid core, the flow tends to elongated structures with the smallest scale l_E , predicted by the linear analysis. It is from this scale ideas of the direct cascade of energy are applicable. To simplify the problem, we estimate the isotropical eddy diffusion, based on the scale (l_E) . In particular, instead of the estimate of velocity gradient at the subgrid scale l: $V' \sim \delta V/l$, we use $V' \sim E^{-1/3}\delta V$, where $\delta V \sim 0.3V$ is the average variation of velocity at the scale l. In this case the estimate of the eddy diffusion gives $\nu^T \sim l^2 V' \approx 5 \cdot 10^4 \text{m}^2 \text{s}^{-1}$ and $E^T \sim 5 \cdot 10^{-4}$. This estimate of the turbulent Ekman number corresponds to $N_{\varphi} \sim 10$ columns which can be resolved in the large-scale models with the desired accuracy. To demonstrate these arguments, we propose a simulations of the system (1-2, 4) with the given eddy diffusion ν^T estimated as above.

4 Turbulent model. Results of calculations.

Equations (1–2, 4) are solved using the control-volume method (Simple algorithm) [18] on the staggered grid $(n_{\rm r}, n_{\theta}, n_{\varphi}) = (45, 45, 64)$. This method is based on the finite-difference approximation and demonstrates very high numerical stability for the regimes with strong convection¹. For ease of calculation, we renormalize equations (1–2, 4) using turbulent diffusion units, so that instead of κ^M the $\hat{\kappa} = 1\text{m}^2 \text{s}^{-1}$ was used. Then, the dimensionless parameters are: $R_o^T = \frac{\hat{\kappa}}{2\Omega L^2} = 4 \cdot 10^{-2}$, $E^T = \frac{\nu^T}{2\Omega L^2} = 10^{-3}$. We consider three regimes with turbulent Rayleigh numbers $R_a^T = \frac{\alpha g_0 \delta T L}{2\Omega \hat{\kappa}} = 10^6$, 10^7 , 10^8 (see the time evolution of the kinetic energy E_K in Fig. 1). The corresponding Reynolds numbers averaged over the shell volume $Re^M = \frac{\hat{\kappa}}{\nu^M}\sqrt{2E_K}$ are $3 \cdot 10^9$, $6 \cdot 10^9$ and $2 \cdot 10^{10}$, c. f. with the molecular Reynolds number for the Earth's core based on the west drift velocity $R_e^{Earth} \sim 10^9$.

Characteristic snapshots of the large-scale velocity r, θ , φ -components are presented in Fig. 2. The observed curls in r, φ -projections corresponds to the columns parallel to z-axis. These columns may drift in the φ -direction. In its turn, the non-zero viscous gradient $\nu \frac{\partial}{\partial r} \left(\frac{V_{\varphi}}{r}\right)_{r=r_i}$ causes rotation of the inner core, (see evolution of the angular velocity of the inner core ω in Fig.1). Here the positive value of ω corresponds to the eastward direction, known to occur in the Earth [20]. We emphasize that these maps are a product of averaging of the small-scale ($l_E \sim \mathcal{O}(E^{M 1/3}) = 10^{-5}$) structures. So far, the micro-scale Reynolds number r_e at the scale l_E is still larger then unity, and the inertial spectrum for the scales smaller then l_E exists. An estimate of $r_e = \frac{vl}{\nu^M}$ with $l = E^{M 1/3}$ and v = 0.1V gives

¹See also some special questions of the control-volume method for the full dynamo problem in the sphere in [19].

 $r_e \sim 10^3$. This spectrum has two parts with the transition point defined by the balance of the inertial and Coriolis terms: $l_{\Omega} \sim R_o v$. The turbulence in the range of $l_E \leq l_{\Omega}$ is influenced by rotation and the kinetic energy spectrum is $E_l \sim l^2$ [21]. For the scales smaller then l_{Ω} up to the dissipative scale $l_d = R_e^{-3/4}$ the Kolmogorov's spectrum $E_l \sim l^{5/3}$ reappears.

Summarizing the obtained results we conclude, that based on the realistic values of the Rossby and Rayleigh numbers and on assumptions on the spectrum of the flow in the liquid core we obtained a value of the kinetic energy E_K comparable with the observations. Having in mind that the velocity field and the eddy diffusion are connected in our model, we consider this agreement to be worth to note.

5 Conclusions

We propose the scenario of the turbulent thermal convection in the rapid rotating body, when the Coriolis force shifts the system to the origin of the small scales already at the onset of convection, and show that further increasing of the intensity of the heat sources leads to a turbulent regime, which is still far from the Kolmogorov's case. It appears that predictions of the linear analysis at the onset of convection are applicable to the eddy diffusion estimate in the regime of the fully developed turbulence. Though the original problem is highly anisotropical, the "isotropical" estimate of the eddy diffusion gives a kinetic energy of the system comparable with the observations. Note that introduction of the magnetic field will not change the problem in principal, because at the scales l_E considered the corresponding micro-scale magnetic Reynolds number is already $r_m \ll 1$ and the magnetic field decays due to the Ohmic dissipation process. On the other hand, it is not yet clear how the west drift velocity relates to the flow at the scales l_E and different interpretations of observations can exist. This question requires the solution of the full dynamo problem.

Acknowledgements

RM is grateful to Central Institute for Applied Mathematics (ZAM) of Forshungszentrum in Jüulich for hospitality. This work was supported by the Russian Foundation of Basic Research (grant 03-05-64074).

References

- S. I. Braginsky, P. H. Roberts, "Equations governing convection in Earth's core and the geodynamo", *Geophys. Astrophys. Fluid Dynamics*, **79**: 1–95, 1995.
- [2] C. A. Jones, "Convection-driven geodynamo models", Phil. Trans. R. Soc. London, A 358: 873– 897, 2000.
- [3] G. A. Glatzmaier, and P. H. Roberts, "A three-dimension convective dynamo solution with rotating and finitely conducting inner core and mantle", *Phys. Earth Planet. Inter.*, **91**: 63–75, 1995.
- [4] D. Gubbins, P. H. Roberts, "Magnetohydrodynamics of the Earth's core". In *Geomagnetism* (ed. Jacobs), Academic. 2: 1–184, 1988.
- [5] U. Frisch, Turbulence: the Legacy of A.N. Kolmogorov. Cambridge University Press, Cambridge, 1995.
- [6] K. Zhang, C. A. Jones, "The effect of hyperviscosity on geodynamo models." *Geophys. Res. Lett.*, 24: 2869-2872, 1997.
- [7] W. Kollman, Prediction Methods for Turbulent Flows. Hemisphere Publishing Corporation, 1980.
- [8] B. A. Buffett, "A comparison of subgrid-scale models for large-scale models for large-eddy simulations of convection in the Earth's core", *Geophys. J. Int.*, 153: 753–765, 2003.
- [9] P. Frick, M. Reshetnyak, D. Sokoloff, "Combined grid-shell approach for convection in a rotating spherical layer", *Europhys. Lett.* 59: 212–217, 2002.
- [10] P. G. Frick, M. Yu. Reshetnyak, D. D. Sokoloff, Cascade models of turbulence for the Earth's liquid core. *Doklady Earth Sciences*, 387: 988-991, 2002.
- [11] P. H. Roberts, "On the thermal instability of a rotating-fluid sphere containing heat sources", *Phil. Trans. R. Soc.*, A 263: 93–117, 1968.
- M. Kono, P. H. Roberts, "Definition of the Rayleigh number for geodynamo simulation", *Phys. Earth Planet. Int.*, **128**: 13–24, 2001.
- [13] D. Gubbins, "The Rayleigh number for convection in the Earth's core", *Phys. Earth Planet. Int.*, 128: 2–12, 2001.
- [14] S. I. Braginsky, V. P. Meytlis, "Local turbulence in the Earth's core", Geophys. Astrophys. Fluid Dynam., A 55: 71–87, 1991.

- [15] L. D. Landau, and E. M. Lifshits, Gidrodinamika (Hydrodynamics). Moscow: Nauka, 1988.
- [16] U. R. Christensen, J. Aubert, P. Cardin, E. Dormy, S. Gibbons, G. A. Glatzmaier, E. Grote, Y. Honkura, C. Jones, M. Kono, M. Matsushima, A. Sakuraba, F. Takahashi, A. Tilgner, J. Wicht, K. Zhang, "A numerical dynamo benchmark", *Phys. Earth Planet. Inter.*, **128**: 25–34, 2001.
- [17] C. A. Jones, A. M. Soward, A. Mussa, "The onset of thermal convection in a rapidly rotating sphere", J. Fluid. Mech., 405: 157–179, 2000.
- [18] S. V. Patankar, Numerical Heat Transfer And Fluid Flow, Taylor & Francis, 1980.
- [19] P. Hejda, M. Reshetnyak, "Control volume method for the dynamo problem in the sphere with the free rotating inner core", *Studia geoph. et. geod*, **47**: 147–159, 2003.
- [20] X. Song, P. G. Richards, "Observational evidence for differential rotation of the Earth's Inner Core", Nature, 382: 221-224, 1996.
- [21] Y. Zhou, "A phenomenological treatment of rotating turbulence", Phys. Fluids, 7: 8, 2092–2094, 1995.



Figure 1: Evolution of the angular velocity of the liquid core ω and averaged over the volume kinetic energy E_k for $R_o = 4 \cdot 10^{-7}$, $E = 10^{-x}$; $R_a^T = 10^6$ – thick line, (2) – $R_a^T = 10^7$ – thin line, (3) – $R_a^T = 10^8$ – dashed line.



Figure 2: The snapshots of the velocity field components $(v_r, v_\vartheta, v_\varphi)$ (from left to right) for the equatorial sections (the left half of the plane): (-700, 1200), (-4700, 4100), (-1200, 1800) and meridional sections for axi-symmetrical parts of the fields (the right half): (-2400, 1800), (-3800, 3100), (-400, 1000). Numbers in round brackets indicate ranges.