QuestionsConcerningSchwarzschild'sSolutionofEinstein'sEquat ions.

J.Dunning-Davies, DepartmentofPhysics, UniversityofHull, HullHU67RX; England, j.dunning-davies@hull.ac.uk

Abstract.

NowthatanEnglishtranslationofSchwarzschild'soriginalworkexists,thatworkhasbecomeaccessibletomorepeople.HerehisoriginalsolutiontotheEinsteinfieldequations is examined and it is noted that it does not contain the mathematicalsingularity normally associated with the existence of a black hole. Einstein's ownviews on this subject are considered also and it is seen that, atthe very least, gravequestionsexistoverthepossibleexistenceofthesesomewhatesotericstellarobjects.

Introduction.

InmanyofthestandardtextbooksontheGeneralTheoryofRelativity [1],timeis devoted to discussing Schwarzschild's solution of the Einstein field equa tions. Normally,thissolutionisstatedasbeingeither

$$ds^{2} = \left\{1 - \frac{2Gm}{rc^{2}}\right\}c^{2}dt^{2} - \left\{1 - \frac{2Gm}{rc^{2}}\right\}^{-1}dr^{2} - r^{2}\left(d\theta^{2} + \sin^{2}\theta d\phi^{2}\right)(1)$$

ormoreusually

$$ds^{2} = \left\{1 - \frac{2m}{r}\right\} dt^{2} - \left\{1 - \frac{2m}{r}\right\}^{-1} dr^{2} - r^{2} \left(d\theta^{2} + \sin^{2}\theta d\phi^{2}\right) (2)$$

where the universal constant of gravitation, G, and the speed of light, c, have both been put equal to unity. Here r, θ , and ϕ appear to be taken to be normal polar coordinates.

Intheaboveexpressions, amathematical singularity is seen to oc curwhen *r*=0.as mightbeexpected for polar co-ordinates. However, due to the form of the coefficient of dr^2 , it follows that a second mathematical singularity occurs when, i $n(1), rc^2 =$ 2Gmor, in(2), r=2 m. The first singularity is regularly dismissed as being mere lya property of polar co-ordinates and, therefore, of no physical significance .The second singularity, however, tends to have a physical interpretation attribute dtoit-namely that it is said to indicate the existence of a blackhole. Somewhat tironically, as will be seenlater.thisisreferredtoasaSchwarzschildblackhole.I fthisinterpretationwere valid, it would imply that, for an object of mass *m* andradius *r*tobeablackhole, it wouldneedtosatisfytheinequality

$$m$$
 $/r \ge c^2/2G=6.7$ $\times 10^{26}$ kg/m(3)Incidentally, it has always seemed fascinating to realise tha
ratioofmasstoradiusisthesameasthatderived, using purelyt this expression for the
Newtonianmechanics,
by Michell in 1784 for a material body having an escape speed equal to, or
greater than, the speed of light [2].

As stated above, many modern texts quote one of equations (1) or (2) as t he Schwarzschild solution of the Einstein field equations, but is this so? Recently, an English translation of Schwarzschild's article of 1916[3], has appear ed and this has made the original work accessible to many more people. For this the scientific community owes the translators, S. Antoci and A. Loinger, a tremendous de bt of gratitude.Italsoenablestheabovequestiontoberaisedbymorepeople.

TheSchwarzschildSolution.

An excellent discussion of the Schwarzschild solution and its derivati on is provided in chapter eighteen of the little book on the General Theory of R elativity by Dirac [4]. Here it is presented in the form (2) above and r, θ , and ϕ are quite clearly stated to be the usual polar co-ordinates. It is pointed out that the ca se being considered is that of a static, spherically symmetric field pr oduced by a spherically symmetric body at rest. After the completion of the derivation, it is noted that the said solution holds only outside the surface of the body producing the field, where there is no matter and, hence, it holds fairly accurately outside the surface of fastar. The following chapter is then devoted to the topic of black holes. It is noted that theSchwarzschildsolution(2)becomes singular when r=2 mandsoitmightappear thatthatvaluefor rindicatedaminimumradiusforabodyofmass mbutitisclaimed that a closer investigation reveals that this is not so. In the dis cussionwhichfollows, thecontinuationoftheSchwarzschildsolutionforvaluesof *r* <2 *m*isinvestigated.To achieve this, it is found necessary to use a non-static system of co-ordinates so that components of the metric tensor may vary with the time co-ordinate. This is accomplished by retaining θ and ϕ as co-ordinates but, instead of tand r, using τ and ρ definedby

 $\tau = t + f(r)$ and $\rho = t + g(r), (4)$ where the functions f and g are at the disposal of the investigator.

Ittranspires that,fortheregion r < 2 m,theSchwarzschildsolutionis foundto adopt theform

$$ds^{2} = d\tau^{2} - \frac{2m}{\mu(\rho - \tau)^{2/3}} d\rho^{2} - \mu^{2} (\rho - \tau)^{4/3} (d\theta^{2} + \sin^{2}\theta d\phi^{2}), (5)$$

where $\mu = \left(\frac{3}{2}\sqrt{2m}\right)^{2/3}$. From the actual derivation, it follows that the cr it is a large real value r = 2m corresponds to ρ - $\tau = 4 m/3$ and there is no singularity at this point in this smetric.

From this point onwards, Dirac's argument bec omes extremely interesting. He notesthatthemetric given by (5) satisfies Einste in'sequationsforemptyspaceinthe region r > 2 m because it may be transformed into the Schwarzschi ld solution by a simple change of co-ordinates. By analytic continua tion, it is seen to satisfy the equations for $r \leq 2 m$ also, because there is now no singularity at r = 2 m. The singularitynowappears, viaequations(4), in the connectionbetweenoldandnewcoordinates. Diracthencomments that, once the new c o-ordinate system is established, theoldonemaybeignored and then the singularity appearsnolonger.

He comments further that the region of space for which r > 2 m may not communicate with that for which *r* <2 *m*.Also,anysignal,evenalightsignal,would take an infinite time to cross the boundary at r=2 m. Thus, there can be no direct observational knowledge of the region for which r < 2 m. If this argument were true, *r* <2 *m*wouldlieoutsideouruniverse;wouldnotreallyb surelytheregionforwhich e apartofit?Diraccallstheregionforwhich r < 2 mablackholebutisthisanobjectin our physical three-dimensional space or one in an a bstract, four-dimensional, mathematicalspace-time?

Finally, Dirac asks whether such a region exist s and notes that the only definite statement which may be made is that the Einstein eq uations allow it. This is a question which will be considered further shortly b utsufficeittosayatthisjuncture d physically [5]. It is noted that a that Einstein himself did not accept that it existe small radius where the forces of massive stellar object may collapse to an extremely gravity might become so strong that no known physic alforcescouldwithstandthem and prevent further collapse. Such a situation woul d herald the collapse to a black hole but, as measured by our clocks, the final stat e would be achieved only after an infinitetime. This argument would appear to stem f romtheideasofOppenheimerand Snyder [6]. They predicted that, when all sources o f thermonuclear energy were exhausted, a large enough star would collapse and t he contraction would continue indefinitely unless the star was able to reduce its mass sufficiently by some means. Theyalsomadethepointthatthetotaltimeforsu chacollapsewouldbefiniteforan observer co-moving with the stellar matter, althoug h it would appear to take an infinite time for a distant observer. This was take ntoindicate that the startended to 'close itself off from any communication with a dis tant observer'; only its gravitational field persisting. Accepting this argu ment as valid for the moment, it might be asked, if such an object existed, would it ever be detectable by an external observer? On the other hand, if its gravitational f ield persists, and presumably the effects of that gravitational field on the surround ings, then, in a sense, the star is retainingsomecontact, albeitindirect, withadis tantobserver. Also, forverymanyyears, it has been noted thatthetransformation

$$t + u + \log(r - 2m)$$

applied to the Schwarzschild solution in the form (singularity. This was taken to indicate that the si physical. This conclusion agrees with that of Einst 1939[5], concluded that the result of the investig 'clear understanding as to why the "Schwarzschild s physical reality'. He went on to point out that, hi clusters whose particles moved along circular paths feel that more general cases would have analogous r categoricallythat 'the"Schwarzschildsingularity" matter cannot be concentrated arbitrarily'. This se notionofblackholesbytheverymanoftenheralde of his book is an indication of his view, then its with this interpretation also. This point concernin of a mathematical singularity has been raised previ publishedanumberofarticlesonarXivinwhichth beenclaimed.However,whatofSchwarzschildhimsel equationswhichisreallyattheheartofthismatt er?

 $\begin{pmatrix} r-2 & m \end{pmatrix}$ m $\begin{pmatrix} 2 \end{pmatrix}$ would remove the offending

ngularity was mathematical, *not* ein himself who, in an article of ation contained in that paper was a ld s ingularities" do not exist in s investigation dealt only with buthe feltit not unreasonable to us r esults. He then stated quite does not appear for the reason that ems a very definite rejection of the das their father. If the general to ne eemst obe the case that Dirac agreed g a possible physical interpretation vi ously by Loinger [7], who has enon-existence of blackholes has sel f; it's his solution of Einstein's er?

Schwarzschild'sOriginalSolution.

Asnotedearlier.thetranslationofSchwarzs child'spaperof1916[3]intoEnglish has made his work accessible to many more people. I n his article, everything is written initially in terms of variables denoted by x_1, x_2, x_3, x_4 and the point is made that the field equations 'have the fundamental prop erty that they preserve their form under the substitution of other arbitrary variables as long as the determinant of the substitution equals one'. The first three of the ab ove co-ordinates are then taken to stand for rectangular co-ordinates, and the fourth is taken to be time. If these are denoted by x, y, z, and tthemost general acceptable line elementis then s tated.butit is noted immediately that, when one goes over to po larco-ordinates according to the usual rules, the determinant of the transformation is not one. Hence, the field equations would not remain unaltered. Schwarzschild then employs the trick of putting

$$x_1 = r^3 / 3, x_2 = -\cos\theta, x_3 = \phi,$$

where r, θ , ϕ are the normal polar co-ordinates. These new varia bles are then polar coordinates but with a determinant of the transformat ion equal to one. Schwarzschild then proceeds to derive his solution and presents in the form

$$ds^{2} = (1 - \alpha / R)dt^{2} - (1 - \alpha / R)^{-1}dR^{2} - R^{2}(d\theta^{2} + \sin^{2}\theta d\phi^{2}),$$
$$R = (r^{3} + \alpha^{3})^{1/3}.$$

Hence, Schwarzschild'sactual solution does co ntainasingularitywhen $R = \alpha$, but Risnotthepolarco-ordinate. It is clearly seen f romabove that, when $R = \alpha$, r = 0: that is, the singularity actually occurs at the ori gin of polar co-ordinates, as is usual. Therefore, according to Schwarzschild'sownwriting thereissimplynosingularityat or general relativity r = 2 m, to use the modern notation, and so the argument f predictingtheexistenceofblackholescannotbej ustifiedbyreferencetotheso-called Schwarzschild solution and it seems not a little ir onic that non-rotating, uncharged blackholesshouldbecalledSchwarzschildblackho les

Conclusions.

where

Thesedays, claims for the identification of bl ackholes appear fairly regularly in the scientific literature. Quite often, the supposed ex so-called massive black holes - is invoked to expla in some otherwise puzzling phenomenon. However, so far, on no occasion has the requirement mentioned earlier that, for ablackhol radius - or more specifically in general relativist horizon-must besubject to the restriction expla in some otherwise puzzling is consistent to expla in some otherwise puzzling postulated object satisfied the e, the radius of its event horizon-must besubject to the restriction explanation of the supposed explanation of the s

$$m/r \ge 6.7 \times 10^{-26} \text{kg/m}$$

ty at the centre of the discussion [8]. Now it emerges that the mathematical singulari simply did not appear in Schwarzschild's original solution of Einstein's equations. Obviously mathematics was used by Schwarzschild to find this solution, but it was used meticulously. It was noted carefully that, if a transformation of coordinates for which the determinant of the transformation does no t equal unity, is used, then the field equations themselves would not remain in an u naltered form. Hence, Schwarzschild adopted a transformation for which th e value of the said determinant wasoneandwentontoderiveanexact,-notappro ximate,-solutiontotheequations. Also, Einsteinhimselfproved that the singularity appearinginthepopularformofthe Schwarzschild solution has no physical significance . In all that Schwarzschild and Einsteindidonthistopic, the mathematics was at ooltohelpthemachievewhatthey wanted. At no point was physical reality modified t o fit a mathematical conclusion. This is the way things should be and provides an ob ject lesson to many; - the mathematicsisatooland,assuch,mustbesubserv ienttothephysics.

Wherethendoesthatleavethemodernnotion of as those above, undoubtedly raise major questions a work. The idea of a body being so dense that it's e speed of light remains not unreasonable though but, quantity-proportional, for example, to the square assuggested by Thornhill [9], Moffatt [10] and, mo [11]-many new and interesting questions arise.

ablackhole?Considerationssuch bout the basis of much modern scape speed is greater than the if the speed of light is avariable root of the background temperature, rerecently,Albrecht and Magueijo

References.

[1]Adler, R., Bazin, M. & Schiffer, M., 1965, IntroductiontoGeneralRelativity, (McGraw-Hill,NewYork) [2]Michell, J., 1784, Philos. Trans. R. Soc., 74.35 [3]Schwarzschild,K.,1916,Sitzungsberichteder KöniglichPreussischenAkademie derWis senschaftenzuBerlin, Phys.-Math.Klasse, 189 Antoci&A.Loinger,arXiv:physics/9905030) (translationbyS. [4]Dirac,P.A.M.,1996, GeneralTheoryofRelativity , (PrincetonUniversityPress,Princeto n,NewJersey) [5]Einstein, A., 1939, Annalsof Mathematics, **40**,922 [6]Oppenheimer, J.R. & Snyder, H., 1939, Phys. Re v. **56**,455 [7]Loinger, A., arXiv: physics/0402088 [8]Dunning-Davies, J., 2004, Science, **305**,1238 Tech., 8,263 [9]Thornhill,C.K.,1985,SpeculationsinSci.& [10]Moffatt,J.,1993,Int.J.Mod.Phys.D., **2**,351;1993,Found.Phys., **23**,411 [11]Albrecht, A.&Magueijo, J., 1999, Phys. Rev. D., **59**,043516