## "Somecommentsonextremevaluestatistics"

by

J.Dunning-Davies, DepartmentofPhysics, UniversityofHull, HullHU67RX.

J.Dunning-Davies@hull.ac.uk

## Abstract.

The question of whether extreme value statistics should be introducephysics' students has been broached recently. Here it is argued tha betaught but the many and varied physical applications should be stresshouldnotbeconfinedpurelytoabstracttheory.

dintocoursesfor t the topic should sed; emphasis It is claimed by many that the oldest problem concerned wit hextreme values is that arisingfromfloods.Infact,naturaldisasters,suchasfloodsanddroug htsareextreme valueswhichhavebeenrecognisedinonewayoranothersincethebeg inningoftime. Both have such important and dire consequences for mankind. However, the systematic examination of such problems and, indeed, of so-called ex treme value statisticsthemselvesisarelativelyrecentoccurrence.Ne vertheless.itisanareawhich hasattracted the attention of people working in a wide variety of very differe ntfields. very welcome and In a fairly recent article, Zetie and James [1] provided a timely mention of this subject of extreme value statistics. M yonlyqueryrelatingto thearticlewasthatitconcentratedonthenotionthattheconc eptofriskwasentering physics courses in a variety of guises; nowhere was mention ma de of the range of areas of physics where the theory of extreme values becomes important.Certainlyin universities, many physics students constantly question the need to s tudy topics in d? Physics students mathematics. How often is the protest, 'But that's maths.!', hear r relevance to a seem happier with mathematical concepts when they can see thei particulararea, or even problem, in physics quite clearly.

Althoughthestudyofextremevaluesaroseinitiallybecauseof theoccurrenceof natural disasters, such as floods or droughts, it was also seen rel evant to problems associated with the breaking strength of materials. For this pr oblem, it proved unsatisfactory simply to put a material under stress and determine its breakingpointin order to be in a position to predict its lifespan. Doing this takes no a ccount, for example, of flaws in a sample which may weaken it. To gain a sa tisfactory answer, the statistical nature of the problem needs to be considered. The fl aws are then assumed distributed at random and a distribution is assigned. Then, of cour se. the forcerequiredtorupturethematerialwillbedifferentindifferentplace s.Thisleadsto the idea of a chain being no stronger than its weakest link but, possibl y more importantly, also provides the motive for studying the distribution in th e smallest valueandforapplyingstatisticaltechniquesingeneral.

The fact that the extreme value has to be treated as a random value has the treated as a random value hasriablebecauseof fluctuations in its environmental conditions means it must be describedintermsofa probability distribution. The big difficulty in developing a statistic al theory of extremes lies in the determining of the probability distribution f rom which the sampleshavebeendrawn.Inphysicsitself,untilveryrecently,t hestatisticaltheories of extremes have remained virtually unknown except in a few special fields. The concept does enter physics, however, especially in the general fiel d of statistical thermodynamics and even here, if it wasn't for the idea of entrop v. the whole area would reduce to a simple exercise in mathematical statistics .Entropy,whichmightbe thought to bridge the gap between the microscopic world of atoms and the macroscopic worldof heatengines, changes all that. It was Bol tzmannwhooriginally linkedincrease of entropy with 'disorder' and, later, thermodynamic equilibriumwas identified as the state of maximum disorder and this corresponds to t he state of maximum entropy. The average values of the variables of state coi ncide with the most probable values so that the size of fluctuations decreases as the sample size increases. All this is guaranteed essentially by the law of large numbers and the centrallimittheorem.

In the early days, observations which lay outside a general or expected pattern wereviewed with suspicion and often regarded as being due to errors in measurement . Indeed, methods for the elimination of such 'extreme values' weres ort and this led to the connection between the method of leasts quares and Gauss' law of error. Althoug h

it seemed at one point that looking seriously at extremely rare events was going against the grain of probability theory, the appreciation that probabil itytheorycould be applied fruitfully to events with small associated probabilities was developing. It was, in fact, at the end of the nineteenth century that Poisson's law, or the law of small numbers, found acceptance. It was at this time also that P areto's work was accomplished and published but that work, which has been known by economistsfor roughlyacentury, has come only recently to the attention of physici sts. However.it waslaterinthetwentiethcenturythatthetopicofextreme valuestatisticsgrewapace with the work of people such as Fréchet, Lévy, and others. (Detailed r eferences to theseandothercontributorsmaybefoundinthebookbyLavenda[2]). Nowadays, however, the subject is no longer simply a branch of mat hematics. Extremevaluestatisticshavebeenappliedinaverywidevarietyof fields.Inthebook referred to above [2], attention is focussed more on the applications t han the basic techniques. These statistics are shown to be important when consideri ngsuchdiverse problems as the model of a rubberband, problems associated with spectr allineshapes, the thermostatistics of polymer chains, a whole variety of probl ems associated with cosmologyand(whatLavendaterms)thermogravity,problemsofthe thermostatistics of materials, and, more recently, he has looked at problems assoc iated with earthquakes [3]. The applications to physics of this field are im mense and are so varied and topical that justifying the study of this seemingl y abstruse branch of mathematicstoscepticalphysicsstudentsismaderelativelyeasy.

## References

[1]Zetie,K.P.andJames,J.E.M.,PhysicsEd. **37(5)**,(2002),381-383

[2]Lavenda,B.H., *ThermodynamicsofExtremes*, AlbionPublishing,Chichester,1995

[3]Lavenda,B.H.&Cipollone,E.,AnnalidiGeofisica **43**,(2000),469-484 and967-982