

The total cross-sections for the photoeffect for K-shell bound electrons and pair production with the created electron in the ground state for photon energies above 1 MeV

A. Costescu^{a)}, S. Spanulescu^{a), b)}, C. Stoica^{a)}

^{a)} Department of Physics, University of Bucharest, MG11, Bucharest-Magurele 76900, Romania

^{b)} Department of Physics, Hyperion University of Bucharest, Postal code 030629, Bucharest, Romania

PACS number: 32.80.Cy

Abstract

Considering the contributions of the main term of the relativistic Coulombian Green function given by Hostler [J. Math. Phys **5**, 591 (1964)] to the second order of S matrix element and taking into account only the large components of Dirac spinor of the ground state we are able to obtain the imaginary part of the Rayleigh amplitudes in terms of elementary functions. Thereby a simple and high accurate total cross-sections for photoeffect and pair production with the electron created in the K-shell are obtained *via* the optical theorem.

Comparing the predictions given by our formulae with the full relativistic numerical calculations of Kissel *et al* [Phys. Rev. A **22**, 1970 (1980)] and Scofield [LLRL, Internal Report, 1973], a very good agreement within 7% is found for photon energies above the pair production threshold up to 5 MeV for any Z elements.

We present our numerical results for the total photoeffect and pair production cross-sections, for various photon energies for Ag and Pb.

I. INTRODUCTION

The S matrix element for Rayleigh scattering of an initial photon with momentum $\vec{k}_1 = \omega \vec{v}_1$ and polarization vector \vec{s}_1 and a final photon with momentum $\vec{k}_2 = \omega \vec{v}_2$ and polarization vector \vec{s}_2 , by K-shell electrons, if we take into account only the large components of Dirac spinor of the ground state and only the main term of the relativistic coulombian Green function [1], is [4]:

$$\mathcal{M}_R = M(\omega, \theta)(\vec{s}_1 \vec{s}_2) + N(\omega, \theta)(\vec{s}_1 \vec{v}_2)(\vec{s}_2 \vec{v}_1), \quad (1.1)$$

where θ is the photon scattering angle and

$$M(\omega, \theta) = [\mathcal{G} - P(\Omega_1, \theta) - P(\Omega_2, \theta)], \quad (1.2)$$

$$N(\omega, \theta) = -[Q(\Omega_1, \theta) + Q(\Omega_2, \theta)], \quad (1.3)$$

$$P(\Omega, \theta) = \frac{128\lambda^5 X^3}{d^4(\Omega)} \frac{F_1(2-\tau; 2, 2; 3-\tau; x_1, x_2)}{2-\tau}, \quad (1.4)$$

$$Q(\Omega, \theta) = \frac{2048\lambda^5 X^5 \omega^2}{d^6(\Omega)} \frac{F_1(3-\tau; 3, 3; 4-\tau; x_1, x_2)}{3-\tau}, \quad (1.5)$$

where $\lambda = \alpha Z m$,

$$\Omega_1 = \gamma m + \omega, \quad \Omega_2 = \gamma m - \omega, \quad \gamma = (1 - \alpha^2 Z^2)^{1/2} \quad (1.6)$$

$$X_j^2 = m^2 - \Omega_j^2, \quad \tau_j = \frac{\alpha Z \Omega}{X_j}, \quad \text{Re}[X_j] > 0, \quad j=1,2 \quad (1.7)$$

The function \mathcal{G} is the relativistic atomic form factor, while $F_1(a; b_1, b_2; c; x_1, x_2)$ is the Appell hypergeometric function of four parameters and two complex variables given by the relationships:

$$x_1 x_2 = p = \left[\frac{d^*(\Omega)}{d(\Omega)} \right]^2 \equiv \xi^2, \quad (1.8)$$

$$x_1 + x_2 = s = 2\xi - \frac{16X^2 \omega^2 \sin^2 \frac{\theta}{2}}{d^2(\Omega)}. \quad (1.9)$$

with

$$d(\Omega) = 2m(\gamma\omega - \alpha^2 Z^2 m + \alpha Z X), \quad d^*(\Omega) = 2m(\gamma\omega - \alpha^2 Z^2 m - \alpha Z X). \quad (1.10)$$

Obviously, in the case of the forward scattering, the amplitude $N(\omega, \theta)$ is no longer present in the Rayleigh amplitude. Also, the imaginary part of the amplitude \mathcal{M}_R is given only by the terms $P(\Omega_1, 0)$ and $P(\Omega_2, 0)$ which are present in the expressions of the photoeffect and the pair production cross-sections respectively.

II. The total cross-section of the photoelectric effect and pair production

According to the optical theorem the imaginary part of the Rayleigh amplitude for forward scattering allows to get the total photoeffect cross-section :

$$\sigma_{ph} = \frac{4\pi}{\alpha} \frac{m}{\omega} r_0^2 |\text{Im} P(\Omega_1)|_{\theta=0} = \frac{32\pi^2}{3} r_0^2 m^2 \alpha^5 Z^6 \frac{\Omega_1 |X_1^2|}{\omega^5} (1 + |\tau_1|^2) \frac{(-\xi_1)^{\tau_1}}{e^{\pi|\tau_1|} - e^{-\pi|\tau_1|}} \frac{\omega + \omega_{pp}}{2m} \quad (2.1)$$

where

In a similar way we get:

$$\sigma_{pp} = \frac{4\pi}{\alpha} \frac{m}{\omega} r_0^2 |\text{Im} P(\Omega_2)|_{\theta=0} = \frac{32\pi^2}{3} r_0^2 m^2 \alpha^5 Z^6 \frac{\Omega_2 |X_2^2|}{\omega^5} (1 + |\tau_2|^2) \frac{(-\xi_2)^{\tau_2}}{e^{\pi|\tau_2|} - e^{-\pi|\tau_2|}} \frac{\omega - \omega_{pp}}{2m} \quad (2.2)$$

We point out that

$$(-\xi_1)^{\tau_1} = e^{-|\tau_1| \chi_1} \quad \text{with} \quad \chi_1 = \pi - 2 \arctan \left(\frac{\lambda |X_1|}{\gamma \omega m - \lambda^2} \right); \quad \omega \geq \frac{\lambda^2}{\gamma \omega m} \quad (2.3)$$

and

$$(-\xi_2)^{\tau_2} = e^{-|\tau_2| \chi_2} \quad \text{with} \quad \chi_2 = \pi - 2 \arctan \left(\frac{\lambda |X_2|}{\gamma \omega m + \lambda^2} \right); \quad \omega \geq \omega_{pp} \quad (2.4)$$

III. Numerical results and conclusions

Using our analytical formulae for the cross sections for K-shell electrons we get the numerical numerical results in table 1 and 2 and figure 1:

Table 1. Photoeffect and pair production cross sections for Ag K-shell .

Energy (keV)	Pair prod cross section ($10^{-26}\text{cm}^2/\text{sr}$)	Photoeffect cross section ($10^{-26}\text{cm}^2/\text{sr}$)	Cross sections ratio
1000.	9.14527×10^{-8}	49.3836	5.39991×10^8
1100.	0.00875335	40.1376	4585.39
1249.	0.0851678	30.6868	60.309
1500.	0.311888	21.1705	67.8785
2000.	0.716782	12.2626	17.1079
2754.	0.996235	7.02819	7.05475
3000.	1.03218	6.10716	5.91674
3500.	1.06112	4.78107	4.50568
4000.	1.05516	3.899	3.69516
4807.	1.01208	2.97828	2.94272
5000.	0.998901	2.81578	2.81887
5500.	0.962702	2.46344	2.55888
6000.	0.925496	2.18602	2.362
6500.	0.888765	1.96248	2.2081
7000.	0.853296	1.77887	2.0847
7500.	0.81948	1.62559	1.98369

Table 2. Photoeffect and pair production cross sections for Pb K-shell .

Energy (keV)	Pair prod cross section ($10^{-26}\text{cm}^2/\text{sr}$)	Photoeffect cross section ($10^{-26}\text{cm}^2/\text{sr}$)	Cross sections ratio
1000.	0.0113701	509.907	44846.5
1100.	0.236329	415.189	1756.82
1249.	1.22336	318.111	260.03
1500.	3.71725	220.044	59.1953
2000.	7.94739	127.886	16.0916
2754.	10.7831	73.5099	6.81712
3000.	11.1338	63.919	5.74101
3500.	11.3924	50.0947	4.39719
4000.	11.2949	40.8876	3.61999
4807.	10.8012	31.265	2.89459
5000.	10.6549	29.5652	2.7748
5500.	10.2571	25.8781	2.52295
6000.	9.85179	22.9731	2.33187
6500.	9.45392	20.6311	2.18228
7000.	9.07119	18.7065	2.06219
7500.	8.70731	17.0993	1.96378




Figure1. Photoeffect and pair production cross sections for Pb K-shell

Comparing the predictions given by our formulae with the full relativistic numerical calculations of Kissel *et al* [Phys. Rev. A **22**, 1970 (1980)] and Scofield [LLRL, Internal Report, 1973], a very good agreement within 7% is found for photon energies above the pair production threshold up to 5 MeV for any Z elements.

The good agreement of our calculation with the full relativistic results shows that, for the presented energies regime, the main relativistic kinematics terms are cancelled by retardation and multipoles terms, and the spin effects are small.

ACKNOWLEDGEMENTS

This work was partially supported by the Romanian National Research Authority (ANCS) under Grant CEEEX PC-D11-PT00-582/2005.

-
- [1] L. Hostler, J. Math. Phys **5**, 591 (1964);
 - [2] Lynn Kissel, R. H. Pratt, and S. C. Roy, Phys. Rev. A **22**, 1970 (1980);
 - [3] J.H. Scofield, Lawrence Radiation Laboratory Report No. CRL 51326, Livermore, CA, 1973 (unpublished);
 - [4] A. Costescu, S. Spanulescu, C. Stoica (to be published).