

The total cross-sections for the photoeffect for 2S-subshell bound electrons and pair production with the created electron in the 2S subshell for photon energies above 1 MeV

A. Costescu^{a)}, S. Spanulescu^{a), b)}, C. Stoica^{a)}

^{a)} Department of Physics, University of Bucharest, MG11, Bucharest-Magurele 76900, Romania

^{b)} Department of Physics, Hyperion University of Bucharest, Postal code 030629, Bucharest, Romania

PACS number: 32.80.Cy

Abstract

Considering the contributions of the main term of the relativistic Coulombian Green function given by Hostler to the second order of S matrix element and taking into account only the large components of Dirac spinor of the 2s subshell we are able to obtain the imaginary part of the Rayleigh amplitudes in terms of elementary functions. Thereby a simple and high accurate total cross-sections for photoeffect and pair production with the electron created in the 2S-shell are obtained *via* the optical theorem.

Comparing the predictions given by our formulae with the full relativistic numerical calculations of Kissel *et al* [Phys. Rev. A **22**, 1970 (1980)] and Scofield [LLRL, Internal Report, 1973], an excellent agreement is found for photon energies above the pair production.

We present our numerical results for the total photoeffect and pair production cross-sections, for various photon energies for lead.

I. INTRODUCTION

The S matrix element for Rayleigh scattering by 2s bound electrons of an initial photon with momentum $\vec{k}_1 = \omega \vec{v}_1$ and polarization vector $\vec{s}_1$ and a final photon with momentum $\vec{k}_2 = \omega \vec{v}_2$ and polarization vector $\vec{s}_2$ is [1]:

$$\mathcal{M}_{RET}^{NR} = M_{RET}(\omega, \theta)(\vec{s}_1 \vec{s}_2) + N_{RET}(\omega, \theta)(\vec{s}_1 \vec{v}_2)(\vec{s}_2 \vec{v}_1), \quad (1.1)$$

where θ is the photon scattering angle.

According to the optical theorem the imaginary part of the Rayleigh amplitude for forward scattering allows to get the total photoeffect cross-section :

$$\sigma_{ph} = \frac{4\pi}{\alpha} \frac{m}{\omega} r_0^2 |\text{Im} P(\Omega_1)|_{\theta=0} \quad (1.2)$$

$$\sigma_{pp} = \frac{4\pi}{\alpha} \frac{m}{\omega} r_0^2 \left| \text{Im} P(\Omega_2) \right|_{\theta=0} \quad (1.3)$$

where,

$$\Omega_1 = \varepsilon m + \omega, \quad \Omega_2 = \varepsilon m - \omega, \quad \varepsilon = \left(\frac{1+\gamma}{2} \right)^{1/2} \quad \gamma = (1 - \alpha^2 Z^2)^{1/2} \quad (1.4)$$

II. The total cross-section of the photoelectric effect and pair production in the case of 2s sub-shell.

Using our analytical formulae for the imaginary part of the elastic scattering of photons by 2s sub-shell electrons [5] we get the total cross-section for the photoelectric effect as follows:

$$\begin{aligned} \sigma_{ph} = & 2 \frac{\omega + \omega_{pp}}{2m} \frac{\lambda^5 X_1^2 (1 + |\tau_1|^2) \pi |\tau_1| \|X_1\|}{6m^4 \omega^4} \left\{ \frac{\eta^2 X_1^2}{m^2 \omega^2} \left(1 + \frac{|\tau_1|^2}{4} \right) \left(1 - \frac{4\eta^2}{5m^2} \right) \right. \\ & \left. - \left(1 - \frac{\eta^2}{m^2} \right) \left[1 + 2 \frac{\eta^2 X_1^2}{m^2 \omega^2} - \frac{2\eta |\tau_1| \|X_1\|}{m^2 \omega^2} (\eta^2 - \varepsilon m \omega) \right] \right\} \frac{e^{-\chi |\tau_1|}}{e^{\pi |\tau_1|} - e^{-\pi |\tau_1|}} \end{aligned} \quad (2.1)$$

where

$$\chi = -\pi + 2 \arctan \frac{\eta \|X_2\|}{\varepsilon m + \eta^2}, \quad X_j^2 = m^2 - \Omega_j^2, \quad \text{Re}(X_j) > 0, \quad j = 1, 2, \quad \tau_j = \frac{\alpha Z \Omega_j}{X_j}, \quad \eta = \frac{\alpha Z m}{2\varepsilon} \quad (2.2)$$

In a similar way we get for the total cross-section pair production:

$$\begin{aligned} \sigma_{pp} = & 2 \frac{\omega - \omega_{pp}}{2m} \frac{\lambda^5 X_2^2 (1 + |\tau_2|^2) \pi |\tau_2| \|X_2\|}{6m^4 \omega^4} \left\{ \frac{\eta^2 X_2^2}{m^2 \omega^2} \left(1 + \frac{|\tau_2|^2}{4} \right) \left(1 - \frac{4\eta^2}{5m^2} \right) \right. \\ & \left. - \left(1 - \frac{\eta^2}{m^2} \right) \left[1 + 2 \frac{\eta^2 X_2^2}{m^2 \omega^2} - \frac{2\eta |\tau_2| \|X_2\|}{m^2 \omega^2} (\eta^2 - \varepsilon m \omega) \right] \right\} \frac{e^{-\chi |\tau_2|}}{e^{\pi |\tau_2|} - e^{-\pi |\tau_2|}} \end{aligned} \quad (2.3)$$

III. Numerical results and conclusions

Using our analytical formulae for the cross sections for 2s sub-shell electrons we get the numerical numerical results in table 1 and 2 and figure 1:

Table 1. Photoeffect and pair production cross sections for Ag 2S subshell .

Energy (keV)	Pair prod cross section	Photoeffect cross section	Cross sections ratio
1022.	7.18629×10^{-6}	0.305755	42547.
1100.	0.000217646	0.260446	1196.65
1200.	0.000742358	0.216344	291.428
1500.	0.00277867	0.137254	49.3956
2000.	0.00526318	0.0796745	15.1381
2754.	0.00669276	0.0458612	6.85237
3000.	0.00683286	0.0399039	5.84001
3500.	0.006878	0.0313161	4.55308
4000.	0.00674389	0.0255934	3.79504
4807.	0.00637291	0.0196062	3.0765
5000.	0.00627312	0.0185476	2.95668
5500.	0.00601054	0.0162499	2.70357
6000.	0.0057513	0.014438	2.51038
6500.	0.00550198	0.0129759	2.35841
7000.	0.00526563	0.0117735	2.23592
7500.	0.00504337	0.0107686	2.1352

Table 2. Photoeffect and pair production cross sections for Pb 2S subshell .

Energy (keV)	Pair prod cross section	Photoeffect cross section	Cross sections ratio
1022.	0.00156954	4.74653	3024.16
1100.	0.00927769	4.10317	442.262
1200.	0.0226335	3.46417	153.055
1500.	0.0630386	2.27928	36.1569
2000.	0.102649	1.37392	13.3846
2754.	0.119336	0.816785	6.84441
3000.	0.119748	0.715838	5.97786
3500.	0.117396	0.568407	4.84179
4000.	0.112942	0.468712	4.15001
4807.	0.104455	0.362929	3.47451
5000.	0.102405	0.344045	3.35966
5500.	0.0972347	0.302852	3.11465
6000.	0.0923508	0.270154	2.9253
6500.	0.0877981	0.243621	2.77478
7000.	0.0835822	0.221692	2.65239
7500.	0.0796897	0.203287	2.55098

Cross sections (cm²)




Figure1. Photoeffect and pair production cross sections for Pb 2S subshell

Comparing the predictions given by our formulae with the full relativistic numerical calculations of Kissel *et al* [Phys. Rev. A **22**, 1970 (1980)] and Scofield [LLRL, Internal Report, 1973], an excellent agreement is found for photon energies above the pair production threshold up to 5 MeV for any Z elements.

ACKNOWLEDGEMENTS

This work was partially supported by the Romanian National Research Authority (ANCS) under Grant CEE X PC-D11-PT00-582/2005.

-
- [1] L. Hostler, J. Math. Phys **5**, 591 (1964)
 - [2] Lynn Kissel, R. H. Pratt, and S. C. Roy, Phys. Rev. A **22**, 1970 (1980).
 - [3] L. Hostler, R.H. Pratt, Phys. Rev. Lett. **10**, 469 (1963)
 - [4] J.H. Scofield, Lawrence Radiation Laboratory Report No. CRL 51326, Livermore, CA, 1973 (unpublished).
 - [5] A. Costescu, S. Spanulescu, C. Stoica (to be published).
 - [6] S. Hultberg, B. Nagel, P. Olsson, Ark. Fys. **38**, 1 (1968)