

Model of photon emission from luminescent center

Eugeniusz Chimczak

Poznań University of Technology, Faculty of Technical Physics, ul. Nieszawska 13 a, 60-965 Poznań, Poland*

(Dated: August 12, 2018)

Luminescence is the phenomenon investigated and applied in many disciplines of science and technique. Spectral and kinetic measurements of luminescence provide much information concerning the mechanism of luminescent devices. Better understanding of the mechanism allows to improve the devices. It has been known for a long time that the spectral and kinetic properties of luminescence depend on the temperature [1]. It is assumed that fluorescence does not depend on temperature but practically lifetime of fluorescence changes insignificantly with temperature [1, 2]. Until now, to our knowledge, nobody describes mathematically the dependence. Here is reported the model of photon emission permitting a description of the dependence. It is found that lifetime of fluorescence decreases with increasing the square root of temperature. In the model presented, the new application of Maxwell distribution is shown.

Temperature influences the luminescence properties of materials. It is shown, in many papers, that temperature changes the shape of luminescence spectra [3, 4, 5, 6, 7, 8, 9, 10, 11]. With increasing temperature, some bands of luminescence spectrum can increase or decrease in intensity. Also, in some cases, some bands or lines can be shifted. Temperature effects also on kinetic properties of luminescence. Measurements of the kinetic properties are highly important since they provide much information concerning the luminescent centers and the mechanism of luminescence in general. Therefore, numerous papers have been devoted to the lifetime of luminescence because the parameter determines properties of luminescence centers [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20]. In several papers there is mathematical description of the lifetime. There is mathematical description of dependence of the fluorescence lifetime on crystal thickness, average absorption coefficient and intrinsic fluorescence decay time but not on temperature [11]. In order to explain the temperature dependence of the lifetime of fluorescence, the model of photon emission from luminescent centers is proposed.

Let us assume that: (i) Luminescent center is some kind of trap for photons. (ii) Trapped photon and electron create new particle like radioactive atom. (iii) Motion of the particle is periodic. The period t_c of the motion is the time required to complete one round trip of the motion, that is, one complete cycle. (iv) Speed of the particle in the motion is governed by Maxwell distribution. (v) Photon can only be emitted from luminescent center at the same point of the cycle path. (vi) During time dt , the particle has speeds between v and $v + dv$; $dt = dn$ seconds, where dn plays the same role as number of molecules in classical Maxwell distribution.

Under such assumptions the average speed of the particle is

$$\bar{v} = \frac{a}{t_c}, \quad (1)$$

where a is constant. The average speed obtained from

Maxwell distribution is

$$\bar{v} = b\sqrt{kT}, \quad (2)$$

where k is Boltzmann's constant, T is the absolute temperature and b is constant. From (1) and (2) we obtain

$$t_c = \frac{c}{\sqrt{T}}, \quad (3)$$

where c is constant. Because lifetime of luminescence τ and t_c are constant, we can write

$$\tau = \frac{d}{\sqrt{T}}, \quad (4)$$

where d is constant. The temperature dependence is considerably weaker than that observed in phosphorescence. The model proposed explains why, contrary to — for example — Cherenkov's radiation, luminescence appears after some time longer than about 10^{-10} s.

Conclusions resulted from the model are following: (i) If the luminescent centers are monomolecular and if the rectangular exciting pulse is very long then, during the time $t = t_c$ after the end of the exciting pulse, the luminescence intensity is constant and equal to luminescence intensity at the end of the pulse, when direct impact excitation is the only process. Time dependence of the luminescence after the end of the pulse is staircase curve. (ii) Trapped photon can only be emitted from luminescent center at time $t = nt_c$, where n is natural number.

This work was supported by Poznań University of Technology Research Program 62-208/07 - BW.

* Electronic address: chimczak@phys.put.poznan.pl

[1] D. Curie, *Luminescence Cristalline* (Dunod, Paris, 1960).

[2] E. Chimczak, W. S. Gordon, and M. Bertrandt-Żytkowiak, *Phys. Stat. Sol.(a)* **82**, 527 (1984).

[3] W. Busse, H. E. Gumlich, B. Meissner, and D. Theis, *J. Lumin.* **12-13**, 693 (1976).

- [4] S.-Y. Seo, M.-J. Kim, and J. H. Shin, *Appl. Phys. Lett.* **83**, 2778 (2003).
- [5] H. Weman, C. J. Harris, J. P. Bergman, M. S. Miller, J. C. Yi, and J. L. Merz, *Superlatt. and Microstruct.* **17**, 61 (1995).
- [6] N. Harukawa, S. Murakami, S. Tamon, S. Ijuin, A. Ohmori, K. Abe, and T. Shigenari, *J. Lumin.* **87-89**, 1231 (2000).
- [7] M. F. Joubert, B. Jacquier, C. Linares, J. P. Chaminade, and B. M. Wanklyn, *J. Lumin.* **37**, 239 (1987).
- [8] M. C. de Lucas, F. Rodrigez, and M. Moreno, *Phys. Stat. Sol. (b)* **184**, 247 (1994).
- [9] A. Wolfert, E. W. J. L. Oomen, and G. Blasse, *J. Sol. Stat. Chem* **59**, 280 (1985).
- [10] P. A. M. Berdowski, J. van Herk, and G. Blasse, *J. Lumin.* **34**, 9 (1985).
- [11] A. B. Suchocki, G. D. Gilliland, C. Powell, M. Bowen, and J. C. Walling, *J. Lumin.* **37**, 29 (1987).
- [12] K. Thilsing-Hansen, M. T. Neves-Petersen, S. B. Petersen, R. Neuendorf, K. Al-Shamery, and H. G. Rubahn, *Phys. Rev. B* **72**, 115213 (2005).
- [13] Y. Zhang, M. D. Sturge, K. Kash, B. P. van der Gaag, A. S. Gozdz, L. T. Florez, and J. P. Harbison, *Superlatt. and Microstruct.* **17**, 201 (1995).
- [14] M. Kapoor, V. A. Singh, and G. K. Johri, *Phys. Rev. B* **61**, 1941 (2000).
- [15] T. C. Leslie and J. W. Allen, *Phys. Stat. Sol. (a)* **65**, 545 (1981).
- [16] E. Chimczak and J. W. Allen, *J. Phys. D: Appl. Phys.* **18**, 951 (1985).
- [17] P. Bergman and B. Monemar, *J. Lumin.* **38**, 87 (1987).
- [18] M. O. Ramirez, D. Jaque, L. E. Bausá, I. R. Martín, F. Lahoz, E. C. A. Speghini, and M. Bettinelli, *J. Appl. Phys.* **97**, 093510 (2005).
- [19] H. Aizawa, M. Sekiguchi, T. Katsumata, S. Komuro, and T. Morikawa, *Rev. Sci. Instr.* **77**, 044902 (2006).
- [20] Y. Zhang, M. D. Sturge, K. Kash, B. P. van der Gaag, A. S. Gozdz, L. T. Florez, and J. P. Harbison, *Phys. Rev. B* **51**, 13303 (1995).