Comment on "Röntgen Quantum Phase Shift: A Semiclassical Local Electrodynamical Effect?"

Tomislav Ivezić

Ruđer Bošković Institute, P.O.B. 180, 10002 Zagreb, Croatia ivezic@irb.hr

Recently Horsley and Babiker [1] reported semiclassical calculations of the Röntgen and the Aharonov-Casher phase shifts. Their derivation shed light on a quantum phenomenon without using full apparatus of quantum theory. They argued that the Röntgen phase shift is due to the action of the nonvanishing three-dimensional (3D) force $\mathbf{F} = \nabla(\mathbf{d} \cdot \mathbf{E})$, where $\mathbf{E} = \mathbf{v} \times \mathbf{B}$. Here, the main conclusion is as in [1], i.e., that the mentioned phase shifts can be derived using a semiclassical calculation. However, instead of the 3D force \mathbf{F} from [1] the Lorentz invariant expression for the 4D force K^a is introduced.

First, in [1], **F** is written in the dipole rest frame S', since the electric field (it is **E**') is obtained by means of the usual transformations (UT) of **B** (e.g., [2], Eq. (11.149)) from the laboratory frame S to S', and **d** from [1] is also in S', i.e., it is **d**'. For comparison with experiments the force has to be in S and not in S'. Furthermore, it is proved in [3] that the UT of **E**, **B** are not the Lorentz transformations (LT) and that they have to be replaced by the LT of the corresponding 4D geometric quantities.

Here, as in [4], we deal with 4D geometric quantities that are defined without reference frames. They will be called the absolute quantities (AQs). The notation is the same as in [4]. Different covariant expressions for a charged particle with a dipole moment are recently presented in [5]. (Here we consider an uncharged particle.) They include, e.g., the Lagrangian and the equations of motion for 4-position. However the usual covariant formulation deals with components, which are implicitly taken in some basis, mainly the standard basis $\{e_{\mu}; 0, 1, 2, 3\}$ of orthonormal 4-vectors with e_0 in the forward light cone. In [4] we have already generalized the interaction term from the Lagrangian [5] and expressed it in terms of AQs as $-(1/2)F_{ab}D^{ab}$. Also, in [4], D^{ab} and F^{ab} are decomposed and written in terms of AQs, the electric and magnetic dipole moments d^a and m^a and 4-vectors E^a and B^a . Here the equation of motion for the 4-position, [5] Eq. (11), is similarly generalized as $mx^c = K^c = (1/2)D^{ab}\partial^c F_{ab}$ (the dot means the derivation with respect to the proper time s). This 4D force K^a replaces the forces written with the 3D d, m and E, B from [1] and many others.

For comparison with the usual formulation with the 3-vectors one needs to write the representation of the AQ K^a in the $\{e_{\mu}\}$ basis as $K^a = K^{\alpha}e_{\alpha} = ((1/2)D^{\mu\nu}\partial^{\alpha}F_{\mu\nu})e_{\alpha}$. Then, [4], we also need to choose the frame of "fiducial" observers in which the observers who measure E^a , B^a are at rest. That frame with the $\{e_{\mu}\}$ basis will be called the e_0 -frame. In the e_0 -frame the velocity of "fiducial" observers $v^a = ce_0$ and $E^0 = B^0 = 0$. In that frame K^{α} is the sum of two terms, one with the electric dipole $(1/c^2)[cd_0(u_{\nu}\partial^{\alpha}E^{\nu}) - c^2(d_{\nu}\partial^{\alpha}E^{\nu}) + c^2\varepsilon^{0ijk}u_jd_k\partial^{\alpha}B_i]$ (responsible for the Röntgen effect) and another one (responsible for the Aharonov-Casher effect) in which d^{μ} is replaced by m^{μ} , while E^{μ} and B^{μ} are interchanged. The last term in the above square braces describes the direct action of the magnetic field on the electric dipole moment. This is very important difference relative to all previous treatments, e.g., [1].

For comparison with experiments the laboratory frame has to be taken as the e_0 -frame. Then, for the Röntgen effect from [1], $m^{\mu} = 0$, $E^{\mu} = 0$, $B^{\mu} = (0, B^1, B^2, 0)$ and $u^{\mu} = (u^0, u^1, u^2, 0)$, since in $S' u'^{\mu} = (c, 0, 0, 0) (u^a = dx^a/ds)$ is the 4-velocity of the particle). K^{α} becomes $K^{\alpha} = \varepsilon^{0ijk} u_j d_k \partial^{\alpha} B_i$, or explicitly, $K^0 = K^3 = 0$, $K^{1,2} = d_3 [u_2 \partial^{1,2} B_1 - u_1 \partial^{1,2} B_2]$. This force is in S and not in S' as in [1]. Also K^{α} is not zero even in the case when the electric dipole moment is aligned parallel to the magnetic line charge. Only in the case when S' is chosen to be the e_0 -frame then it will be $K^{\alpha} = 0$, but that case is not physically realizable. The second term in K^{α} gives the analogous result for the Aharonov-Casher effect. Hence, as in [1], both phase shifts could be calculated using the concept of force, but not the 3D force than the 4D force K^a . Then the phase shifts can be calculated, e.g., as in [6].

References

- [1] S.A.R. Horsley and M. Babiker, Phys. Rev. Lett. 95, 010405 (2005).
- [2] J.D. Jackson, *Classical Electrodynamics* 3rd ed. (Wiley, New York, 1998).
- [3] T. Ivezić, Found. Phys. 33, 1339 (2003); Found. Phys. Lett. 18, 301 (2005); Found. Phys. 35, 1585 (2005).
- [4] T. Ivezić, Phys. Rev. Lett. **98**, 108901 (2007).
- [5] A. Peletminskii, S. Peletminskii, Eur. Phys. J. C 42, 505 (2005).
- [6] J. Anandan, Int. J. Theor. Phys. 19, 537 (1980).