The Supersymmetry of Relativistic Pöschl-Teller Systems

Ion I. Cotăescu

The West University of Timişoara, V. Pârvan Ave. 4, RO-1900 Timişoara, Romania

September 25, 2018

Abstract

We analyze the supersymmetry and the shape invariance of the potentials of the (1+1) relativistic oscillators we have recently proposed [1, 2].

In the general relativity, the geometric models play the role of kinematics, helping us to understand the characteristics of the classical or quantum free motion on a given background. One of the simplest (1+1) geometric models is that of the quantum relativistic harmonic oscillator represented by a free massive scalar field on the anti-de Sitter static background [3, 4]. Recently, we have generalized this model to a family of quantum models the metrics of which depend on a real parameter [1]. We have shown [2] that this family contains a set of models which can be considered as the relativistic correspondents of the usual nonrelativistic Pöschl-Teller (PT) systems [5, 6]. These models have been referred as relativistic PT systems [2]. They have countable discrete energy spectra and the same square integrable energy eigenfunctions as those known from the nonrelativistic quantum mechanics. However, the significance of the parameters as well as the formula of the energy levels are different. An important property of all these models is that they lead to the usual harmonic oscillator in the nonrelativistic limit.

These analytically solvable relativistic problems may have similar properties of supersymmetry and shape invariance as those known from the nonrelativistic theory [7]. Here, our aim is to analyze these properties, pointing out the changes requested by the specific form of the Klein-Gordon operator and by the relativistic expression of the energy levels.

Let us start with a short review of the main results concerning the relativistic PT problem (in natural units, $\hbar = c = 1$) [2]. There exists an holonomic frame, (t, x), in which the metrics of our family of models is given by the line element

$$ds^{2} = \left(1 + \frac{1}{\epsilon^{2}} \tan^{2} \hat{\omega}x\right) (dt^{2} - dx^{2}).$$

$$\tag{1}$$

where we have denoted $\hat{\omega} = \epsilon \omega$ understanding that ω is the frequency of the harmonic oscillator obtained as the nonrelativistic limit of these models. Here ϵ is a new parameter which determines the domain of the quantum free motion, $D = (-\pi/2\hat{\omega}, \pi/2\hat{\omega})$, for a fixed ω . Note that for $\epsilon = 1$ we obtain the exact anti-de Sitter metric. Particularly, in the frame (t, x), the scalar product of the square integrable functions defined on D is just the usual one. The Klein-Gordon equation of a spinless particle of the mass m has the form

$$\left(-\frac{d^2}{dx^2} + \frac{m^2}{\epsilon^2}\tan^2\hat{\omega}x\right)U(x) = (E_n{}^2 - m^2)U(x).$$
 (2)

It has a countable discrete energy spectrum. The corresponding squareintegrable energy eigenfunctions are

$$U_{n_s,s}(x) = N_{n_s,s} \cos^k \hat{\omega} x \sin^s \hat{\omega} x F(-n_s, k+s+n_s, s+\frac{1}{2}, \sin^2 \hat{\omega} x), \qquad (3)$$

where $N_{n_s,s}$ is the normalization factor and

$$k = \frac{1}{2} \left[1 + \sqrt{1 + 4\frac{m^2}{\epsilon^2 \hat{\omega}^2}} \right] > 1$$
 (4)

is the positive solution of the equation

$$k(k-1) = \frac{m^2}{\epsilon^2 \hat{\omega}^2}.$$
(5)

The quantum numbers, $n_s = 0, 1, 2...$ and s = 0, 1, can be embedded into the main quantum number $n = 2n_s + s$. This will take even values if s = 0and odd values for s = 1. Hence, the functions $U_{n_s,s}(x)$ are completely determined by n only. They are real polynomials of the degree n in $\sin \hat{\omega}x$, with the factor $\cos^k \hat{\omega}x$, which vanishes at $x = \pm \pi/2\hat{\omega}$. The energy levels also depend on the main quantum number. These can be obtained from the quantization condition

$$E_n^2 - m^2 \left(1 - \frac{1}{\epsilon^2} \right) = \hat{\omega}^2 (n+k)^2, \quad n = 0, 1, 2....$$
(6)

We specify that, for the pure anti-de Sitter oscillator, the resulting energy spectrum is equidistant [4] since $\epsilon = 1$.

We have seen that the relativistic PT systems depend only on three parameters, the mass m, the frequency ω and the new parameter ϵ . For our future needs it is convenient to change this parametrization by using the parameter k instead of m. Then, from (5) we have

$$m^2 = k(k-1)\epsilon^2 \hat{\omega}^2,\tag{7}$$

with the help of which the quantization condition (6) gives

$$E_n^2 = \hat{\omega}^2 [(n+k)^2 + (\epsilon^2 - 1)k(k-1)].$$
(8)

Moreover, the second term of the left-hand side of Eq.(2), which has been called the relativistic PT potential [2], can be written now as

$$V_{PT}(k,x) = k(k-1)\hat{\omega}^2 \tan^2 \hat{\omega} x.$$
(9)

In the following we shall consider that the parameters ω and ϵ remain fixed, while k is variable. For this reason, and by taking into account that the eigenfunctions (3) are defined, in fact, only by the main quantum number, we shall denote $U_{n_{s,s}} \equiv U_{k,n}$.

In order to construct the supersymmetric quantum mechanics [7] of the relativistic PT systems we have to adjust the Klein-Gordon operator (2) so that its spectrum is

$$(E_n^2 - E_0^2)/\hat{\omega}^2 = n(n+2k).$$
(10)

Thus, we obtain the new operator $\Delta[V]$ defined by

$$\{\Delta[V]U\}(x) = \left(-\frac{1}{\hat{\omega}^2}\frac{d^2}{dx^2} + V(x)\right)U(x).$$
 (11)

which will play here the same role as the Hamiltonian of the nonrelativistic theory. With its help we can rewrite the Klein-Gordon equation in the form

$$\Delta[V_{-}(k)]U_{k,n} = n(n+2k)U_{k,n}$$
(12)

where we have denoted

$$V_{-}(k,x) = \frac{1}{\hat{\omega}^2} [V_{PT}(k,x) + m^2 - E_0^2] = k(k-1) \tan^2 \hat{\omega} x - k.$$
(13)

Now, from the (normalized [8]) ground-state eigenfunction

$$U_{k,0}(x) = \left(\frac{\hat{\omega}^2}{\pi}\right)^{\frac{1}{4}} \left[\frac{\Gamma(k+1)}{\Gamma(k+\frac{1}{2})}\right]^{\frac{1}{2}} \cos^k \hat{\omega} x.$$
(14)

we obtain the superpotential

$$W(k,x) = -\frac{1}{\hat{\omega}} \frac{1}{U_{k,0}(x)} \frac{dU_{k,0}(x)}{dx} = k \tan \hat{\omega} x$$
(15)

and the supersymmetric partner of V_{-}

$$V_{+}(k,x) = -V_{-}(k,x) + 2W(k,x)^{2} = k(k+1)\tan^{2}\hat{\omega}x + k.$$
(16)

Furthermore, following the standard procedure [7], we can define the pair of the lowering and raising operators, A_k and A_k^+ , having the action

$$(A_k U)(x) = \left(\frac{1}{\hat{\omega}}\frac{d}{dx} + W(k,x)\right)U(x), \qquad (17)$$

$$(A_k^+U)(x) = \left(-\frac{1}{\hat{\omega}}\frac{d}{dx} + W(k,x)\right)U(x).$$
(18)

and the commutation rule

$$[A_k, A_k^+] = 2k + \frac{1}{2k}(A_k + A_k^+)^2.$$
(19)

Then, the operators Δ of the systems with superpartner potentials, V_{-} and V_{+} , can be written as

$$\Delta[V_{-}(k)] = A_{k}^{+}A_{k}, \qquad \Delta[V_{+}(k)] = A_{k}A_{k}^{+}.$$
(20)

Let us observe now that these potentials are shape invariant since

$$V_{+}(k,x) = V_{-}(k+1,x) + 2k + 1.$$
(21)

Consequently, we can verify that

$$\Delta[V_{+}(k)]U_{k+1,n-1} = n(n+2k)U_{k+1,n-1}.$$
(22)

From this equation combined with (12) it results that the normalized energy eigenfunctions satisfy

$$A_k U_{k,n} = \sqrt{n(n+2k)} U_{k+1,n-1}, \qquad A_k^+ U_{k+1,n-1} = \sqrt{n(n+2k)} U_{k,n}.$$
 (23)

Hence, we have obtained the relation between a pair of models with superpartner potentials. These can correspond to any pair of values k and k + 1 of our variable parameter. Therefore, if we consider the sequence of models with $k = k_0, k_0 + 1, k_0 + 2, ..., (k_0 > 1)$, then the of successive models are supersymmetric partners. What is interesting here is that the masses of these models appear as being quantized because of Eq. (7) where k play the role of the quantum number. On the other hand, we observe that the action (23) of the operators A_k and A_k^+ leaves invariant the value of n + k. This indicates that the operator H, defined by $HU_{n,k} = (n+k)U_{n,k}$, will commute with these operators. Finally we note that the very simple form of the shape invariance here allows one to write any normalized energy eigenfunction as

$$U_{k,n} = \frac{1}{\sqrt{n!}} \left[\frac{\Gamma(n+2k)}{\Gamma(2n+2k)} \right]^{\frac{1}{2}} A_k^+ A_{k+1}^+ \dots A_{k+n-1}^+ U_{k+n,0}.$$
 (24)

where $U_{k+n,0}$ is the normalized ground-state eigenfunction given by (14).

As a conclusion, we can say that our example of relativistic supersymmetric quantum mechanics of a massive spinless particle follows the same general rules as in the nonrelativistic case. The unique major difference is that the Hamiltonian of the nonrelativistic theory is replaced here by the operator Δ which is linearly dependent on the squared Hamiltonian. For this reason, the commutation relations of the Hamiltonian with the raising and lowering operators will be different too. In our opinion this could lead to new algebraic properties of the relativistic PT systems.

References

- I. I. Cotăescu, preprint physics/9704009, to appear in Int. J. Mod. Phys. A.
- [2] I. I. Cotăescu, preprint physics/9704007
- C. Dullemond, E. van Beveren, Phys. Rev. D 28, 1028 (1983); V. Aldaya,
 J. Bisquert, J. Navarro-Salas, Phys. Lett. A156, 315 (1991); V. Aldaya,
 J. Bisquert, R. Loll, J. Navarro-Salas, J. Math. Phys. 33, 3087 (1992)
- [4] D. J. Navarro, J. Navarro-Salas, preprint hep-th/9406001
- [5] G. Pöschl, E. Teller, Z. Phys. 83, 149 (1933)
- [6] M. M. Nieto, L. M. Simmons Jr., Phys. Rev. Lett. 41, 207 (1978); Phys. Rev. D 20, 1332 (1979)
- [7] R. Dutt, A. Khare, U. P. Sukhatme, Am. J. Phys. 56, 163 (1987)
- [8] M. Abramowitz, I. A. Stegun, Handbook of Mathematical Functions, Dover, New York (1964)