

On the codon assignment of chain termination signals and the minimization of the effects of frameshift mutations

Jean-Luc Jestin^{*}

Centre for Protein Engineering and Laboratory of Molecular Biology
Medical Research Council, Cambridge CB2 2QH, UK

Achim Kempf[†]

Corpus Christi College in the University of Cambridge
Cambridge CB2 1RH, UK

Abstract

It has been suggested that the minimization of the probability for lethal mutations is a major constraint shaping the genetic code [1]. Indeed, the genetic code has been found highly protective against transitions [2]. Here, we show that data on polymerase-induced frameshifts provide a rationale for the codon assignment of chain termination signals (CTS).

We work on the assumption that the mutational spectra of in vitro polymerization [3]-[11] for DNA-polymerases belonging to families found in at least two of the three living kingdoms [12] are relevant with respect to primordial polymerases. We will here not take into account DNA polymerases beta, which are family X DNA polymerases exclusively found among eukaryotes so far [13], and HIV-reverse transcriptases, which are thought to be active as dimers [14] and which emerged very ‘late’ in evolution [15].

As it is believed that RNA preceded DNA in evolution, data for RNA-replicases would be more adapted but are not available. Recent evidence shows however that DNA- and RNA-replicases are very closely related [16]: a single tyrosine to phenylalanine substitution changes DNA-replicases into RNA-replicases [17]; E.coli DNA-polymerase I is also an accurate RNA-dependent DNA-polymerase [18]. A single mutation confers to the MMLV-reverse transcriptase the ability to replicate RNA [19].

Polymerase-induced mutations are mainly substitutions and frameshifts [3]-[9]. The frameshifts’ error-rate is about half the substitutions’ error-rate for the Klenow-polymerase domain [3], which has no nuclease domain as can be assumed for a primordial polymerase. Frameshifts result mostly from the addition or deletion of one

^{*}email: jljestin@mrc-lmb.cam.ac.uk

[†]email: a.kempf@amtp.cam.ac.uk

base [3]-[7]. Frameshifts are highly deleterious as they prevent translation in the correct reading frame of the codons downstream the mutation. They happen in directly repeated and palindromic sequences [10] (where the assumption of polymerase-error tolerance can be shown to be consistent) and in non-reiterated runs, where single-base deletions occur more frequently than single-base additions [3, 4],[6],[9]. Additions will therefore be neglected here. For polymerases with or without nuclease domains we noticed no significant differences in the consensus sequence for single-base deletion sites in non-reiterated runs. It has been defined as YR [3], TTR [11], YTG [8], TR [10], and refined here from the current data [3]-[11] as YTRV (V= C, A or G [3]).

If the genetic code has been optimized for frameshift tolerance, then it should allow to code for most amino acid sequences without using YTRV sequences, or the TRV, YTR and NYT potential deletion site codons (PDSC) nor using their reverse-complementary (rc) sequences which are also expected to yield deletions during replication (Fig. 1):

If the base T is the first base of a codon and in case the previous codon has a pyrimidine as third codon base, then the amino acid should be encoded without using the six codons TRV; if the base T is the second base of a codon and in case the first base of the following codon is C, A or G, then the amino acid should be encoded without using the codons YTR; if the base T is the third base of a codon and in case the following amino acid has a RVN-type codon, then the amino acid should be encoded without using the eight codons NYT.

The most deleterious codons are TAA and TAG and their rc-sequences TTA and CTA that are both PDSC and rc-PDSC. Deletions at codons encoding amino acids are likely to yield non-functional proteins, as all downstream codons are not translated. However, deletions at codons encoding CTS should result in addition of peptides to the proteins' carboxy-termini, thereby likely providing functional proteins. *The least deleterious effects are therefore obtained by assigning the most deleterious codons to CTS and not to amino acids.* Frameshift tolerance may therefore have been the major constraint in the codon assignment of CTS. The codons TTA and CTA encode leucine having the highest, six-fold degeneracy. Frameshift tolerance may therefore be one of the constraints imposing a high degeneracy to these amino acids.

We point out, however, that substitution tolerance and frameshift mutation tolerance are to be considered as competing constraints on the selection of an optimal genetic code: substitution tolerance favors a code in which an amino acid is encoded by triplets differing only by single-base mutations. On the other hand, given that single-base deletions in non-reiterated runs occur mostly on a specific template sequence, (YTRV), tolerance of these frameshift mutations favors a code in which amino acids are encoded by triplets differing strongly from another, so that amino acid sequences are more likely to be able to be coded for without using the YTRV sequences. Our argument is based on the consideration that, for the consensus sequences, the ratio between single-base deletions and base substitutions is much greater than in other sequences, so that, for the assignment of specific codons, frameshift mutation tolerance should be a stronger constraint than substitution tolerance.

In conclusion, these results provide an insight into the constraints yielding the genetic code's fixation and suggest that the codon assignment of CTS may be contemporary with the emergence of polymerases being enzymes rather than ribozymes. Polymerase-error tolerance arguments similar to the one presented here may be useful in the investigation of alternative terrestrial or exobiological genetic codes and possibly also for the engineering of new genetic codes.

Tables available from authors via email and mail.

References

- [1] Sonneborn T.M., in *Evolving genes and proteins* (ed. Bryson V. & Vogel H.J.) 377-397, (Academic Press, New York, 1965).
- [2] Goldberg A.L., Wittes R.E., *Science*, 153, 420-424, (1966).
- [3] Bebenek K., Joyce C.M., Fitzgerald M.P., Kunkel T.A., *J. Biol. Chem.*, 265, 13878-13887, (1990).
- [4] Bell J.B., Eckert K.A., Joyce C.M., Kunkel T.A., *J. Biol. Chem.*, 272, 7345- 7351, (1997).
- [5] Minnick D.T., Astatke M., Joyce C.M., Kunkel T.A., *J. Biol. Chem.*, 271, 24954-24961, (1996).
- [6] Kunkel T.A., Patel S.S., Johnson K.A., *Proc. Natl. Acad. Sci. USA*, 91, 6830- 6834, (1994).
- [7] Kunkel T.A., *J. Biol. Chem.*, 260, 12866-12874, (1985).
- [8] Papanicolaou C., Ripley L.S., *J. Mol. Biol.*, 207, 335-353, (1989).
- [9] Cai H., Yu H., McEntee K., Kunkel T.A., Goodman M.F., *J. Biol. Chem.*, 270, 15327-15335, (1995).
- [10] Wang F.J., Ripley L.S., *Genetics*, 136, 709-719, (1994).
- [11] De Boer J.G., Ripley L.S., *Genetics*, 118, 181-191, (1988).
- [12] Braithwaite D.K., Ito J., *Nucl. Acid. Res.*, 21, 787-802, (1993).
- [13] Bork P., Ouzounis C., Sander C., Scharf M., Schneider R., Sonnhammer E., *Prot. Sci.*, 1, 1677-1690, (1992).
- [14] Restle T., M=FCller B., Goody R.S., *J. Biol. Chem.*, 265, 8986-8988, (1990).
- [15] Eigen M., Nieselt-Struwe K., *AIDS*, 4, S85-S93, (1990).
- [16] Joyce C.M., *Proc. Natl. Acad. Sci. USA*, 94, 1619-1622, (1997).

- [17] Sousa R., Padilla R., EMBO J., 14, 4609-4621, (1995).
- [18] Ricchetti M., Buc H., EMBO J., 12, 387-396, (1993).
- [19] Gao G., Orlova M., Georgiadis M.M., Hendrickson W.A., Goff S.P., Proc. Natl. Acad. Sci. USA, 94, 407-411, (1997).