NARROWING OF BENNETT HOLE IN COLLISIONAL PLASMA

E.V.Podivilov, D.A.Shapiro, M.G.Stepanov Institute of Automation & Electrometry, Novosibirsk, 630090, Russia

The profile of a Bennett hole induced by laser field in ionic distribution in collisional plasma is calculated. Influence of Chandrasekhar's dependence of coefficients of velocity space transport on the profile is included into the calculation for the first time. It is found that the hole narrows down as the field detuning frequency increases. Physical cause of the effect is the falling dependence of Coulomb collision frequency on the ionic velocity. Estimations show that the effect is quite observable under conditions of high-current gas-discharge plasma.

52.20Hv, 42.62Fi, 52.80.-s

The nonlinear spectroscopic technique as a plasma diagnostic tool is a subject of considerable recent interest because of its novelty and potential in basic and applied plasma physics. Of particular interest are the spectroscopic effects of Coulomb scattering [1], since it differs radically in appearance from binary collisions in gases [2,3]. In linear absorption spectra Doppler width usually exceeds collisional broadening, so it masks effects of the scattering. However, the velocity distribution of particles interacting with a monochromatic field near the resonance is changed significantly by the scattering. If the frequency ω of the field is near-resonant with Bohr's frequency $\omega_{mn} = (E_m - E_n)/\hbar$ between levels m and n of the atoms with velocity \vec{v} that satisfies the condition $\vec{k}\vec{v} = \omega - \omega_{mn} \equiv \Omega$, then the induced transitions $m \leftrightarrow n$ occur (\vec{k} denotes the wavevector). As a result, the narrow peak or dip (Bennett hole [4]) arises in the velocity distribution of level populations. The steady-state profiles of holes are controlled by both the homogeneous width of transition $m \leftrightarrow n$ and the scattering processes with a change in velocity. Measuring the width, shift and asymmetry of nonequilibrium structures in the distribution with the help of probe-field technique one can extract information about the scattering.

Previous calculations of Coulomb broadening [1,5] were based on the assumption of the constancy of the diffusion coefficient in the velocity space. This assumption is applicable when the detuning frequency Ω is less than the Doppler width kv_T , where $v_T = \sqrt{2T_i/m}$ is the thermal velocity of ions and m is the ion mass. The constant diffusion model was enough to interpret Lamb dip measurement in argon laser plasma or another with near- resonance pumping. In contrast, farther away from the resonance, at $\Omega > kv_T$, one must take Chandrasekhar's velocity dependence of the diffusion tensor that complicates the evaluation. One way around this problem was shown in linear theory of Dicke narrowing in the ionic spectrum, which had been built up for low intensity of the incident wave [6].

In the present Letter we study a first-order nonlinear problem, the influence of collisions in plasma on the profile of the Bennett hole. The calculation presented below for nonlinear structure in the velocity distribution is performed in the basis of density matrix formalism [3,7]. First we derive the formula for the distribution, then analyze it under some simple but realistic conditions, and at the end estimate whether the effect obtained be observable in experiment. In particular, we show that the width of hole decreases with the detuning of the electromagnetic field from the resonance. By observing the effect, one could directly measure the diffusion tensor. It would be helpful to confirm our understanding and offer scope for new diagnostic methods.

The diffusion coefficient as a function of the velocity was measured for Q-machine plasma at low density by Bowles et al. [8]. The effect of transient spread of a Bennett hole in the distribution of tagged ion population in a metastable state was exploited. However, that effect cannot be used for denser plasma, because the diffusion time becomes smaller and the measurement requires a faster registration technique. In this paper we shall examine the stationary hole shape. Analysis of its width is useful for diagnostics at high as well as at low density.

To develop the theory of nonequilibrium structures in the velocity distribution, let us describe the two-level subsystem of a probe ion in an ideal nondegenerate plasma by the spectroscopic density matrix $\rho_{ij}(\vec{r}, \vec{v}, t)$. This matrix over internal states i, j is at the same time the Wigner function of position \vec{r} and velocity \vec{v} . It has been shown [5,6] that elements of the matrix satisfy the quantum kinetic equation with the classical Landau collision term. This equation makes it possible to analyze resonant transitions between levels in the field of the traveling wave $\vec{E} = \vec{E}_0 \exp(-i\omega t + i\vec{k}\vec{r}) +$ c.c.

Denoting the relaxation constants of level populations as Γ_j and that of the coherence as Γ , write steady - state equations for off-diagonal $\rho_{mn}(\vec{r}, \vec{v}, t) = \rho(\vec{v}) \exp(-i\Omega t + i\vec{k}\vec{r})$ and diagonal $\rho_{jj} \equiv \rho_j$ elements of the density matrix

$$\left(\Gamma - i\Omega + i\vec{k}\vec{v}\right)\rho - \nu\hat{V}\rho = -iG\Delta N, \quad \Gamma_j\rho_j - \nu\hat{V}\rho_j - \lambda_j(\vec{v}) = \mp 2\Re(iG^*\rho), \tag{1}$$

where the upper/lower sign corresponds to upper/lower level. Here $\Delta N \equiv \rho_m - \rho_n$, $G = \vec{E}_0 \vec{d}_{mn}/2\hbar$, \vec{d}_{mn} is the matrix element of the dipole moment,

$$\nu = \frac{16\sqrt{\pi}LN_i(Z_a e^2)^2}{3m^2 v_T^3} \tag{2}$$

is the Coulomb collision frequency, $Z_a e$ is the charge of a probe ion, L is the Coulomb logarithm. We assume that there are only singly charged ions in plasm; N_i is their concentration. The excitation of ionic levels j = m, n usually occurs from the ground or metastable state, so we are justified in assuming the shape of the excitation function $\lambda_j(\vec{v})$ to be Maxwellian

$$\lambda_j(\vec{v}) = Q_j W(\vec{v}), \quad W(\vec{v}) = \frac{1}{v_T^3 \pi^{3/2}} \exp\left(-\frac{v^2}{v_T^2}\right),$$
(3)

where Q_j is the excitation rate of level j.

For plasma in a state close to the thermodynamic equilibrium with ionic temperature T_i , the distribution of buffer particles has the Maxwellian shape. Then it is possible to write an explicit expression for collision operator \hat{V} involving the dynamic friction and the diffusion in the velocity space

$$\hat{V} = \frac{1}{2} \frac{\partial}{\partial \xi_{\alpha}} \Phi_{\alpha\beta} \left(\frac{\partial}{\partial \xi_{\beta}} + 2\xi_{\beta} \right), \tag{4}$$

where $\xi_{\alpha} = v_{\alpha}/v_T$ is the α - component of the dimensionless velocity, the differential operator $\partial/\partial\xi_{\alpha}$ acts on the right. Functions $\Phi_l(\xi)$ and $\Phi_{tr}(\xi)$, occurring in the expression for the diffusion tensor

$$\Phi_{\alpha\beta} = \Phi_l(\xi) \frac{\xi_\alpha \xi_\beta}{\xi^2} + \Phi_{tr}(\xi) \left(\delta_{\alpha\beta} - \frac{\xi_\alpha \xi_\beta}{\xi^2} \right), \tag{5}$$

can be written as single integrals

$$\Phi_l(\xi) = 3 \int_0^1 \lambda^2 e^{-\lambda^2 \xi^2} d\lambda, \quad \Phi_{tr}(\xi) = \frac{3}{2} \int_0^1 (1 - \lambda^2) e^{-\lambda^2 \xi^2} d\lambda.$$
(6)

The velocity distribution of populations ρ_j of both levels j = m, n needs to be found. To simplify analysis, we assume the electromagnetic field to be weak $|G|^2 \ll \Gamma \Gamma_j$ and will neglect the effects of strong saturation. The right sides of Eqs. (1) can therefore be regarded as perturbation. The formal solution of (1) in terms of operator exponents is

$$\rho_{j}^{(0)} = \frac{Q_{j}}{\Gamma_{j}} W(\vec{v}), \quad \rho^{(0)} = 0; \quad \Delta N(\vec{v}) = \rho_{m}^{0}(\vec{v}) - \rho_{n}^{(0)}(\vec{v}) = N_{mn} W(\vec{v});$$

$$\rho^{(1)}(\vec{v}) = -iG \int_{0}^{\infty} dt \exp(-\hat{A}t + \nu\hat{V}t) \Delta N(\vec{v}), \quad \hat{A} = \Gamma - i\Omega + i\vec{k}\vec{v}, \tag{7}$$

$$\rho_j^{(1)}(\vec{v}) = \mp \int_0^\infty dt \exp(-\Gamma_j t + \nu \hat{V} t) 2\Re \left(i G^* \rho^{(1)}(\vec{v}) \right).$$
(8)

It remains to write out the explicit solution and analyze its limiting cases. For this purpose we must calculate the commutators of operators \hat{A} and \hat{V} and get, in the general case, the infinite series. Fortunately, the relation $[[[\hat{V}, \hat{A}], \hat{A}], \hat{A}] = 0$ allows us to break the series within the first order in ν :

$$\rho_{j}^{(1)}(\vec{v}) = \mp 2|G|^{2}\Delta N(\vec{v})\Re \int_{0}^{\infty} dt \int_{0}^{\infty} d\tau \exp\left(-\Gamma_{j}\tau - (\Gamma - i\Omega + ikv_{z})t\right) \times \\ \times \exp\left[\frac{\nu}{2}\left(-\Phi_{zz}(\vec{\xi})k^{2}v_{T}^{2}(t^{2}\tau + t^{3}/3) + 2ikv_{z}\Phi_{l}(\xi)(2t\tau + t^{2})\right) + \mathcal{O}(\nu^{2})\right],$$
(9)

where the axis z is chosen along the wavevector \vec{k} . Function

$$\Phi_{zz}(\vec{\xi}) = (\Phi_l(\xi) - \Phi_{tr}(\xi)) \ \frac{\xi_z^2}{\xi^2} + \Phi_{tr}(\xi)$$
(10)

is the zz - component of the diffusion tensor. This function depends on two variables $\xi \equiv |\vec{\xi}|$ and ξ_z . The integrand in (9), obtained by the expansion, is inappropriate at long times t, τ . Consequently, for application of this expansion it is necessary to have the main contribution to integral (9) in the domain where the quadratic terms in ν remain small. Since $|\partial \Phi_{\alpha\beta}/\partial \xi_{\gamma}| < |\Phi_{\alpha\beta}|$, estimation of those terms gives the following applicability conditions of the expansion (9)

$$\alpha = \frac{\nu}{\Gamma_j} \ll 1, \quad \beta = \frac{\Gamma}{kv_T} \ll 1. \tag{11}$$

The latter inequality means the Doppler limit; the former one coincides with the condition of smallness of the diffusion broadening compared to the Doppler width. The collision frequency in high-current gas-discharge plasma is about $\nu \sim 10^5 - 10^7 \text{ s}^{-1}$ (see [1]). It is by a few orders of magnitude less than the Doppler width $kv_T \sim 10^{10}s^{-1}$ and, as a rule, less than the relaxation rates $\Gamma \sim 10^8 - 10^9 s^{-1}$, $\Gamma_j \sim 10^6 - 10^9 s^{-1}$. That is why we may use small parameters α , β and it is enough to hold only terms linear in ν .

Taking an integral over τ we obtain the correction to the distribution function as a single integral over t

$$\rho_j^{(1)}(\vec{v}) = \mp 2|G|^2 \Delta N(\vec{v}) \Re \int_0^\infty \frac{\exp\left[-\hat{A}t - \Phi_{zz}(\vec{\xi})\nu k^2 v_T^2 t^3/6 + i\nu k v_z \Phi_l(\xi) t^2\right]}{\Gamma_j + \Phi_{zz}(\vec{\xi})\nu k^2 v_T^2 t^2/2 - 2i\nu k v_z \Phi_l(\xi) t} dt.$$
(12)

The integrand in (12) regarded as a function of t at $\Omega = kv_z$ is maximal at t = 0 and decreases with t. The shape of the distribution depends on a few parameters. Considering conditions (11) and the inequality $\Gamma_j \leq \Gamma$ we can set apart two limiting cases of parameters:

$$\nu k^2 v_T^2 \ll \Gamma^2 \Gamma_j \quad (\alpha \ll \beta^2), \tag{13}$$

$$\Gamma^2 \Gamma_j \ll \nu k^2 v_T^2 \quad (\alpha \gg \beta^2). \tag{14}$$

For the domain (13) integral (12) is accumulated at $t \leq T_1 = 1/\Gamma$. The diffusion and friction cause the Bennett hole to somewhat change its width and position. For the limit (14) integral (12) is gathered at $t \leq T_2 = (\alpha/2)^{-1/2}/kv_T$. Diffusion and friction effects control the width and shift of the hole. The width is defined by the maximum among the homogeneous width $\Delta v_H = \Gamma/k$ and diffusion width $\Delta v_D = v_T (\nu/\Gamma_j)^{1/2}$. Thus, the scattering has stronger impact on shape of the hole, while the relaxation constants of selected levels satisfy inequality (14).

Let us find the width as a function of detuning in this limit $\epsilon(\vec{\xi}) = 2\beta^2/\alpha \Phi_{zz}(\vec{\xi}) \ll 1$. It is convenient to write the expression for the nonlinear correction to the Maxwellian distribution as a convolution of two functions

$$\rho_j^{(1)}(\vec{v}) = W(\vec{v}) \int_{-\infty}^{\infty} f(x - x') g_j(x') \, dx', \quad x = k v_z - \Omega, \tag{15}$$

where

$$g_j(x) = \mp \frac{\pi |G|^2 N_{mn}}{\Gamma \Gamma_j} \frac{\epsilon}{\sqrt{\mu}} \exp\left(-\frac{|x|\sqrt{\mu} + x\sigma}{\Gamma}\right), \tag{16}$$

$$\sigma(\xi) = \frac{1}{kv_T \Phi_{zz}(\vec{\xi})}, \quad \mu(\xi) = \epsilon(\xi) + \sigma^2 \simeq \epsilon(\xi),$$

$$f(x) = \Re \int_0^\infty \frac{dt}{\pi} \exp\left[-(\Gamma + ix)t - \Phi_{zz}(\vec{\xi})\nu k^2 v_T^2 t^3/6 + i\nu k v_z \Phi_l(\xi) t^2\right]. \tag{17}$$

In the limit $\epsilon \ll 1$, the width of function f(x) is much less than that of $g_j(x)$. Accordingly, the central part of the distribution is given by expression (16) $\rho_j^{(1)} \simeq g_j(x)$ at $|x| < \Gamma/\sqrt{\epsilon}$. Meanwhile function $g_j(x)$ drops exponentially with |x|, so the asymptotic of the distribution $\rho_j^{(1)}(v_z)$ at $|x| \gg \Gamma/\sqrt{\epsilon}$ is given by a narrow but slowly falling function f(x) and has the Lorentzian shape.

The most interesting feature of the distribution obtained is the narrowing of its central part as the detuning Ω increases, due to the falling velocity dependence of the collision frequency. One can estimate the half-width by substituting the extreme value of the longitudinal velocity $v_z = \Omega/k$ into all components of the diffusion tensor:

$$\Delta x \simeq \frac{\Gamma}{\sqrt{\mu}} \simeq k v_T \sqrt{\frac{\nu \Phi_{zz}(\xi, \Omega/k v_T)}{\Gamma_j}}.$$

This quantity is proportional to the square root of the zz- component of the diffusion tensor. It falls off with detuning Ω . The effect is absent in the model of the constant collision frequency [9,10].

To describe this effect quantitatively we calculate the Ω - dependence of x^2 averaged over the distribution (15). Keeping only the terms linear in parameters α and β we obtain

$$F_{j}(\Omega) = \int d^{3}v \, x^{2} \rho_{j}^{(1)}(\vec{v}) \simeq \mp \frac{2|G|^{2} N_{mn}}{\Gamma_{j}} \left\{ \Gamma + \frac{\nu k^{2} v_{T}^{2}}{\Gamma_{j}} \Re \int_{0}^{\infty} dt \, \overline{\Phi}_{zz}(t) \exp\left(-(\Gamma - i\Omega)t\right) \right\},\tag{18}$$

$$\overline{\Phi}_{zz}(t) = \int d^3 v W(\vec{v}) \Phi_{zz}(\vec{\xi}) \exp(-i\xi_z t) = \int_{0}^{1/\sqrt{2}} 6y^2 dy \exp\left[-(1-y^2)\left(\frac{tkv_T}{2}\right)^2\right].$$

The first term on the right-hand side of (18) is independent of Ω and corresponds to the contribution of the Lorentzian wings of the distribution. The second term describes the collisional broadening and decreases with $|\Omega|$.

Under the approximations the line shape is Voigt's contour. The dependence of the variance $D = \langle (x - \langle x \rangle)^2 \rangle$ on detuning Ω consists of two terms: the contribution of the wings, which increases with detuning, and the decreasing contribution of the diffusion broadening. At small detuning $|\Omega| \ll kv_T$ we have the explicit expression

$$D \simeq \frac{\Gamma k v_T}{\sqrt{\pi}} \left(1 + \left(\frac{\Omega}{k v_T}\right)^2 \right) + \frac{\nu (k v_T)^2}{2 \Gamma_j} \left(\frac{3\pi}{2} - 3 - \left(\frac{\Omega}{k v_T}\right)^2 \left(15 - \frac{9\pi}{2}\right) \right).$$

Summarizing the results, we conclude that the Bennett hole narrows down as detuning increases. The physical reason of the effect is that the width of the hole is not affected by the homogeneous width Δv_H , but depends on the change in velocity due to diffusion $\Delta v_D = v_T \sqrt{\nu \tau_j}$, while $\Delta v_D \gg \Delta v_H$. The excited state lifetime $\tau_j = \Gamma_j^{-1}$ determines the stationary width of the hole. While the detuning Ω of the field remains inside the Doppler contour, the field interacts only with a group of slow ions $|v_z| \ll v_T$. When we turn the field off the resonance, the light interacts with faster excited ions. Collision frequency of those ions with ions in the ground state ν is smaller, therefore the width decreases. It would be more proper to call this effect decrease of the broadening instead of the narrowing.

To observe the narrowing experimentally, the probe-field spectroscopy can be used. One should measure the nearresonant absorption or gain of a weak probe wave as a function of its detuning on the same or adjacent transition. Another possibility is recording the spectrum of spontaneous emission in the presence of a strong resonant continuous field. Let us estimate two alternative effects independent of the detuning, namely, Stark broadening and change of velocity due to the interaction with electrons. The Stark effect is quadratic in the ArII spectrum. Its value [1, p.139] is about 100 MHz, which is small compared with both the homogeneous width Γ and the diffusion one $kv_T \sqrt{\nu/\Gamma_j}$. The velocity change owing to scattering on electrons is as small as the ratio of the ion and electron thermal velocities $v_{Ti}/v_{Te} \sim 10^{-2}$ [1, p.155]. The ion-ion collision frequency decreases no more than by a factor of 2–3 at the detuning about kv_T ; therefore, neither alternative broadening mechanism can compensate the effect in question. The measured diffusion broadening in low-temperature argon plasma at zero detuning is by a factor of 3–4 (see [1, p.199]). At the detuning $\Omega \sim kv_T$ the narrowing factor expected is more than 10%, so this effect seems observable, see Fig. 1.

We thank S.A.Babin and S.G.Rautian for stimulating discussions, E.G.Shapiro for excellent assistance in computation, and A.V. Shafarenko for valuable comments. The research described in this publication was made possible in part by Grant No. RCN000 from the International Science Foundation.

^[1] S.A. Babin, D.A. Shapiro, Phys. Rep. **241**, 119 (1994).

- [2] P.R. Berman, Adv. At. Mol. Phys. 13, 57 (1977).
- [3] S.G. Rautian, A.M. Shalagin, Kinetic Problems of Non-Linear Spectroscopy (Elsevier, Amsterdam, 1991).
- [4] W.R. Bennett, Jr., Phys. Rev. 126, 580 (1962); Appl. Opt., Supplement No 1, 24 (1962).
- [5] G.I. Smirnov, D.A. Shapiro, Zh. Eksp. Teor. Fiz. 76, 2084 (1979) [Sov. Phys. JETP 49, 1054 (1979)].
- [6] E.V. Podivilov, A.I. Chernykh, D.A. Shapiro, Zh. Eksp. Teor. Fiz. 105, 1214 (1994) [Sov. Phys. JETP 78, 653(1994)].
- [7] S. Stenholm, Phys. Rep. 43, 151 (1978).
- [8] J. Bowles, R. McWilliams, N. Rynn, Phys. Rev. Lett. 68, 1144 (1992).
- [9] S.G. Rautian, Zh. Eksp. Teor. Fiz. 51, 1176 (1966) [Sov. Phys. JETP 24, 788 (1967)].
- [10] L.A. Gel'medova, D.A. Shapiro, J. Mod. Opt. 38, 573 (1991).

LIST OF CAPTIONS

Fig.1 (a) Distribution $\rho_j^{(1)}(v_z)$ of *j*-level population in ion velocity v_z within the first order of the perturbation theory in the field intensity. Different curves correspond to distinct values of detuning Ω (from left to right $\Omega/kv_T = 0; 0.5; 1; 1.5; 2; 2.5$). Parameters are assumed to be $\Gamma = 10^{-2}kv_T$, $\Gamma_j = 10^{-3}kv_T$, $\nu = 10^{-5}kv_T$.

(b) Half-width on half a maximum of contour $\rho_j^{(1)}(v_z)$ as a function of detuning Ω/kv_T normalized to unit magnitude.



