

REPRESENTATIONS OF GENERALIZED OSCILLATOR ALGEBRA

PIOTR KOSIŃSKI*, MICHAŁ MAJEWSKI*, PAWEŁ MAŚLANKA**

ABSTRACT. The representations of the oscillator algebra introduced by Brzeziński et al. (Phys. Lett. **B 311** (1993), 202) are classified.

I. INTRODUCTION

In this paper we present an exhaustive discussion of representations of the oscillator algebra introduced in Ref. [1]. This algebra was obtained as a q -deformation of the modified oscillator algebra underlying the structure of Calogero model [2]. In the paper [1] the Fock representations of the algebra under consideration were found. However, it is known [3] that q -deformed oscillator algebra possesses some exotic representations which disappear in the limit $q \rightarrow 1$. Therefore, it is expected that some phenomenon occurs for the algebra considered by Brzeziński et al. Following the methods used in Ref. [3] we show that this is indeed the case and provide the complete classification of its representations.

II. CONSTRUCTION OF REPRESENTATIONS

The algebra under consideration reads :

$$\begin{aligned} aa^+ - qa^+a &= q^{-N}(1 + 2\alpha K), \\ [N, a] &= -a, \quad [N, a^+] = a^+, \\ \{K, a\} &= 0, \quad \{K, a^+\} = 0, \\ [N, K] &= 0, \\ N^+ &= N, \quad K^+ = K, \end{aligned} \tag{1}$$

where $q \in \mathbb{R}_+$, $\alpha \in \mathbb{R} - \{0\}$. This algebra possesses the following Casimir operators

$$C_1 = K^2, \quad C_2 = Ke^{i\pi N}, \quad C_3 = e^{2i\pi N}. \tag{2}$$

Obviously, they are not independent

$$C_1 C_3 = C_2^2. \tag{3}$$

*supported by KBN grant no 2P30221706p02

**supported by University of Łódź grant no 505/444

We will be looking for irreducible representations. Let γ be an eigenvalue of C_2 corresponding to a given representation; then

$$K = \gamma e^{-i\pi N}. \quad (4)$$

Let Ψ_0 be a common eigenvector of N and K :

$$\begin{aligned} N\Psi_0 &= \nu_0\Psi_0, \\ K\Psi_0 &= \gamma e^{-i\pi\nu_0}\Psi_0. \end{aligned} \quad (5)$$

Due to the commutativity of a^+a and aa^+ with N and K we may assume that

$$\begin{aligned} a^+a\Psi_0 &= \lambda_0\Psi_0, \\ aa^+\Psi_0 &= \mu_0\Psi_0 \end{aligned} \quad (6)$$

and $(\Psi_0, \Psi_0) = 1$. It is easy to see that the vectors Φ_n defined by

$$\Phi_n = \begin{cases} (a^+)^n \Psi_0 & \text{for } n \geq 0, \\ a^{-n} \Psi_0 & \text{for } n < 0 \end{cases}$$

are eigenvectors of a^+a and aa^+ :

$$\begin{aligned} a^+a\Phi_n &= \lambda_n\Phi_n, \\ aa^+\Phi_n &= \mu_n\Phi_n. \end{aligned} \quad (7)$$

Now, let us define the following vectors :

$$\Psi_n = \begin{cases} \frac{1}{\sqrt{\prod_{k=1}^n \lambda_k}} (a^+)^n \Psi_0 & \text{for } n \geq 0, \\ \frac{1}{\sqrt{\prod_{k=1}^{-n} \lambda_{n+k}}} a^{-n} \Psi_0 & \text{for } n < 0. \end{cases} \quad (8)$$

They are orthogonal (as eigenvectors of N corresponding to different eigenvalues) and normalized. The action of basic operators is given by

$$\begin{aligned} a^+\Psi_n &= \sqrt{\lambda_{n+1}}\Psi_{n+1}, \\ a\Psi_n &= \sqrt{\lambda_n}\Psi_{n-1}, \\ N\Psi_n &= (\nu_0 + n)\Psi_n, \\ K\Psi &= \frac{(-1)^n}{2\alpha} B\Psi_n, \end{aligned} \quad (9)$$

where, for later convenience, we have defined

$$B = 2\alpha\gamma e^{-i\pi\nu_0} \in \mathbb{R}. \quad (10)$$

The only additional condition we have to take into account is that λ_n and μ_n , being eigenvalues of nonnegative operators, should be nonnegative. Using the basic commutation rules applied to Ψ_n we obtain

$$\mu_n - q\lambda_n = -q^{-(n+\nu_0)} \left(1 + 2\alpha\gamma e^{-i\pi(n+\nu_0)}\right). \quad (11)$$

But $a(a^+a)\Psi_n = (aa^+)a\Psi_n$ which gives

$$\lambda_n = \mu_{n-1}. \quad (12)$$

Equations (11) and (12) imply the following recurrence relation

$$\lambda_{n+1} = q\lambda_n + q^{-\nu_0-n} (1 + (-1)^n B), \quad (14)$$

which can be explicitly solved to yield :

$$\lambda_n = \lambda_0 q^n + q^{-\nu_0} \left(\frac{q^n - q^{-n}}{q - q^{-1}} + B \frac{q^n - (-1)^n q^{-n}}{q + q^{-1}} \right). \quad (15)$$

Nonnegativity of λ_n implies

$$\lambda_0 q^{\nu_0} + \frac{1}{q - q^{-1}} + \frac{B}{q + q^{-1}} \geq q^{-4k} \left(\frac{1}{q - q^{-1}} + \frac{B}{q + q^{-1}} \right), \quad (16a)$$

$$\lambda_0 q^{\nu_0} + \frac{1}{q - q^{-1}} + \frac{B}{q + q^{-1}} \geq q^{-(4k+2)} \left(\frac{1}{q - q^{-1}} - \frac{B}{q + q^{-1}} \right). \quad (16b)$$

We have now to distinguish several cases.

- (i) Assume $q > 1$. Then at least one of the numbers $\frac{1}{q - q^{-1}} \pm \frac{B}{q + q^{-1}}$ is positive. Therefore, there exists n_0 such that for even and/or odd $n < n_0$, $\lambda_n < 0$, which implies $a\Psi_n = 0$ for some $n \leq n_0$. After possible renumbering we may assume

$$a\Psi_0 = 0, \quad \lambda_0 = 0 \quad (17)$$

Therefore, the representation is spanned by the vectors Ψ_n , $n \geq 0$, and λ_n are given by

$$\lambda_n = q^{-\nu_0+n} \left(\frac{1 - q^{-2n}}{q - q^{-1}} + B \frac{1 - (-1)^n q^{-2n}}{q + q^{-1}} \right) \quad (18)$$

The condition $\lambda \geq 0$ gives the following restriction on the possible values of B :

$$B \geq -1 \quad (19)$$

However, $B = -1$ must be considered separately. In this case $\lambda_1 = 0$, i.e. $\mu_0 = 0$ which, together with $\lambda_0 = 0$ and irreducibility implies

$$a = a^+ = 0, \quad N = \nu_0, \quad K = -\frac{1}{2\alpha}. \quad (20)$$

This representation is onedimensional. For $B > -1$ the representation is spanned by the vectors $\{\Psi_n\}_{n=0}^\infty$. We shall call it the Fock representation for obvious reasons. It is given by equations (9) and (18) with $n \geq 0$.

- (ii) $q < 1$ and one (and only one) of the values $\frac{1}{q - q^{-1}} \pm \frac{B}{q + q^{-1}}$ is positive. In this case there exists n_0 such that for $n > n_0$ λ_n is negative for even or odd n . This implies $a^+ \Psi_n = 0$ for some $n \geq n_0$. After possible renumbering we get

$$a^+ \Psi_0 = 0. \quad (21)$$

In order to find the restrictions on possible values of B we note that the condition (21) implies μ_0 , i.e. $\lambda_1 = 0$ or

$$\lambda_0 = -q^{-\nu_0-1}(1+B), \quad (22)$$

which gives $B \leq -1$. For $B = -1$ we obtain onedimensional representation (20). If $B < -1$ we have to consider the restrictions on B following from the formula

$$\lambda_n = q^{n-\nu_0} \left(-q^{-1}(1+B) + \frac{1-q^{-2n}}{q-q^{-1}} + B \frac{1-(-1)^n q^{-2n}}{q+q^{-1}} \right). \quad (23)$$

The condition $\lambda_n \geq 0$ implies $B \leq \frac{q+q^{-1}}{q-q^{-1}}$. For

$$B < \frac{q+q^{-1}}{q-q^{-1}} \quad (24)$$

we have $\lambda_n > 0$ and the representation is given by equations (9) and (23) with $n \leq 0$. We call this representation anti-Fock one. For

$$B = \frac{q+q^{-1}}{q-q^{-1}} \quad (25)$$

all λ_n with odd n are zero. Therefore the representation is twodimensional and given by

$$\begin{aligned} a \Psi_0 &= \sqrt{\frac{2q^{-\nu_0}}{q^{-1}-q}} \Psi_{-1}, & a^+ \Psi_0 &= 0, \\ a^+ \Psi_{-1} &= \sqrt{\frac{2q^{-\nu_0}}{q^{-1}-q}} \Psi_0, & a \Psi_{-1} &= 0, \\ N \Psi_0 &= \nu_0 \Psi_0, & N \Psi_{-1} &= (\nu_0 - 1) \Psi_{-1}, \\ K \Psi_0 &= \frac{q+q^{-1}}{2\alpha(q-q^{-1})}, & K \Psi_{-1} &= \frac{q+q^{-1}}{2\alpha(q-q^{-1})} \Psi_{-1}. \end{aligned} \quad (26)$$

- (iii) $q < 1$ and both values $\frac{1}{q-q^{-1}} \pm \frac{B}{q+q^{-1}}$ are nonpositive (at least one must be strictly negative). There are now the following possibilities :

(a)

$$\lambda_0 q^{\nu_0} + \frac{1}{q - q^{-1}} + \frac{B}{q + q^{-1}} < 0 \quad (27)$$

There exists then n_0 such that $\lambda_n < 0$ for $n < n_0$, n even or odd. Therefore the representation is given by formulae (9), (18); it is a Fock one. To provide $\lambda_n \geq 0$ for $n \geq 0$ we have to restrict B to lie in the interval

$$-1 \leq B < -\frac{q + q^{-1}}{q - q^{-1}}. \quad (28)$$

For $B = -1$ we get again onedimensional representation (20).

(b)

$$\lambda_0 q^{\nu_0} + \frac{1}{q - q^{-1}} + \frac{B}{q + q^{-1}} > 0 \quad (29)$$

Equation (29) implies $\lambda_n > 0$ for all $n \in \mathbb{Z}$. The representation is given by equations (9), (15) with $n \in \mathbb{Z}$.

(c)

$$\lambda_0 q^{\nu_0} + \frac{1}{q - q^{-1}} + \frac{B}{q + q^{-1}} = 0 \quad (30)$$

If $|B| < -\frac{q + q^{-1}}{q - q^{-1}}$ all $\lambda_n > 0$ and representation has the same form as in point (b). For $B = -\frac{q + q^{-1}}{q - q^{-1}}$ all λ_n with n even are vanishing; therefore, the representation is twodimensional and given by formulae :

$$\begin{aligned} a^+ \Psi_0 &= \sqrt{\frac{2q^{-\nu_0-1}}{q^{-1} - q}} \Psi_1, & a \Psi_0 &= 0, \\ a^+ \Psi_1 &= 0, & a \Psi_1 &= \sqrt{\frac{2q^{-\nu_0-1}}{q^{-1} - q}} \Psi_0, \\ N \Psi_0 &= \nu_0 \Psi_0, & N \Psi_1 &= (\nu_0 + 1) \Psi_1, \\ K \Psi_0 &= -\frac{q + q^{-1}}{2\alpha(q - q^{-1})} \Psi_0, & K \Psi_1 &= \frac{q + q^{-1}}{2\alpha(q - q^{-1})} \Psi_1. \end{aligned} \quad (31)$$

For $B = \frac{q + q^{-1}}{q - q^{-1}}$ all λ_n with n odd vanish. The representation is twodimensional and is given by formulae (26).

III. DISCUSSION

Let us summarize the results obtained in section II. For $q > 1$ there are two possibilities :

- (a) if $B > -1$ the spectrum of N is bounded from below – we have Fock representation which is irreducible and determined by the choice of ν_0 and B ; different choices correspond to inequivalent representations,
- (b) if $B = -1$ we get onedimensional irreducible representation labeled by the values of ν_0 ; again different values of ν_0 correspond to inequivalent representations.

The case $q < 1$ is more involved. The following possibilities have to be distinguished :

- (a) for $B < \frac{q + q^{-1}}{q - q^{-1}}$ we get irreducible anti-Fock representation; the representations are labelled by pairs (ν_0, B) and different choices correspond to inequivalent representations,
- (b) for $B = \frac{q + q^{-1}}{q - q^{-1}}$ one obtains twodimensional representations parametrized by ν_0 ; different values of ν_0 correspond to inequivalent representations,
- (c) for $B = -1$ we obtain again onedimensional representation parametrized by ν_0 ; it has the same form as for $q > 1$,
- (d) for $-1 < B < -\frac{q + q^{-1}}{q - q^{-1}}$ the representations are the Fock ones parametrized by ν_0 and B ; for different values of these parameters we obtain inequivalent representations,
- (e) for $B = -\frac{q + q^{-1}}{q - q^{-1}}$ the representations, parametrized by ν_0 , are twodimensional and mutually inequivalent,
- (f) finally, there exists a set of infinitedimensional representations for which the spectrum of N extends infinitely in both directions. They correspond to $|B| < -\frac{q + q^{-1}}{q - q^{-1}}$ and $\lambda_0 q^{\nu_0} + \frac{1}{q - q^{-1}} + \frac{B}{q + q^{-1}} \geq 0$ or $|B| = -\frac{q + q^{-1}}{q - q^{-1}}$ and $\lambda_0 q^{\nu_0} + \frac{1}{q - q^{-1}} + \frac{B}{q + q^{-1}} > 0$. It is easy to see that two such representations, labeled by (ν_0, B, λ_0) and (ν'_0, B', λ'_0) , are equivalent iff $\nu'_0 = \nu_0 + n$, $B' = (-1)^n B$, $\lambda'_0 = \lambda_0 q^n + q^{-\nu_0} \left(\frac{q^n - q^{-n}}{q - q^{-1}} + B \frac{q^n - (-1)^n q^{-n}}{q + q^{-1}} \right)$ for some integer n .

Let us now consider the limits $q \rightarrow 1$ or $B \rightarrow 0$. It is easy to see that only the onedimensional and Fock representations survive the limit $q \rightarrow 1$. On the other hand the $B \rightarrow 0$ limit coincides with the results obtained in Ref. [3]. Finally, the limit $q \rightarrow 1$, $B \rightarrow 0$ leaves only Fock representation as it should be. The results obtained are summarized in table 1.

TABLE 1.

Type of representation	q	Restrictions on B	Restrictions on λ_0 and ν_0	$q \rightarrow 1$ limit	$\alpha \rightarrow 0$ limit
onedimensional	arbitrary	$B = -1$	$\lambda_0 = 0$ ν_0 -arbitrary	exists	does not exist
twodimensional	$q < 1$	$B = \frac{q+q^{-1}}{q-q^{-1}}$	$\lambda_0 = \frac{2q^{-\nu_0}}{q^{-1}-q}$ ν_0 -arbitrary	does not exist	does not exist
		$B = -\frac{q+q^{-1}}{q-q^{-1}}$	$\lambda_0 = 0$ ν_0 -arbitrary	does not exist	does not exist
	$q > 1$	$B > -1$	$\lambda_0 = 0$ ν_0 -arbitrary	exists	exists
Fock	$q < 1$	$-\frac{q+q^{-1}}{q-q^{-1}} > B > -1$	$\lambda_0 = 0$ ν_0 -arbitrary	exists	exists
anti-Fock	$q < 1$	$B < \frac{q+q^{-1}}{q-q^{-1}}$	$\lambda_0 = -q^{\nu_0-1}(1+B)$ ν_0 -arbitrary	does not exist	does not exist
unbounded in both directions	$q < 1$	$ B < -\frac{q+q^{-1}}{q-q^{-1}}$	$\lambda_0 q^{\nu_0} + \frac{1}{q-q^{-1}} + \frac{B}{q+q^{-1}} \geq 0$	does not exist	exists
		$ B = -\frac{q+q^{-1}}{q-q^{-1}}$	$\lambda_0 q^{\nu_0} + \frac{1}{q-q^{-1}} + \frac{B}{q+q^{-1}} > 0$	does not exist	does not exist

REMARK to the case of representations unbounded in both directions : they are equivalent

$$(\nu'_0, \lambda'_0, B') \sim (\nu_0, \lambda_0, B)$$

iff

$$\begin{aligned} \nu'_0 &= \nu_0 + n, \quad B' = (-1)^n B, \\ \lambda'_0 &= \lambda_0 q^n + q^{-\nu_0} \left(\frac{q^n - q^{-n}}{q - q^{-1}} + B \frac{q^n - (-1)^n q^{-n}}{q + q^{-1}} \right), \\ n &\in \mathbb{Z}. \end{aligned}$$

REFERENCES

- [1] T. Brzeziński, J. L. Egusquiza, A. J. Macfarlane, Phys. Lett. **B 311** (1993), 202.
- [2] L. Brink, T. H. Hansson, M. A. Vasiliev, Phys. Lett. **B 286** (1992), 109.
- [3] M. A. Vasiliev, Int. J. Mod. Phys. **A 6** (1991), 1115.
- [4] G. Rideau, Lett. Math. Phys. **24** (1977), 147.

PIOTR KOSIŃSKI, MICHAŁ MAJEWSKI
DEPARTMENT OF THEORETICAL PHYSICS
UNIVERSITY OF ŁÓDŹ
POMORSKA 149/153
90-236 ŁÓDŹ, POLAND

PAWEŁ MAŚLANKA
INSTITUTE OF MATHEMATICS
UNIVERSITY OF ŁÓDŹ
BANACHA 22
90-238 ŁÓDŹ, POLAND