

Quadratic Poisson brackets and Drinfel'd theory for associative algebras

A.A. Balinsky*, Yu.M. Burman†

Department of Mathematics
Technion — Israel Institute of Technology,
32000, Haifa, Israel

Abstract

Quadratic Poisson brackets on associative algebras are studied. Such a bracket compatible with the multiplication is related to a differentiation in tensor square of the underlying algebra. Jacobi identity means that this differentiation satisfies a classical Yang–Baxter equation. Corresponding Lie groups are canonically equipped with a Poisson Lie structure. A way to quantize such structures is suggested.

1 Introduction

The main purpose of the present article is to construct an analog of Drinfel'd theory (see [1]) for associative algebras. Poisson Lie brackets involved in the original version of the theory are replaced by *quadratic* Poisson brackets compatible with the algebra structure. Restriction of such brackets to the group of invertible elements of the algebra leads to a Poisson Lie structure (see Theorem 4 below).

As usual, Poisson bracket $\{\cdot, \cdot\}$ on a smooth manifold M is understood as a Lie algebra structure on the space of smooth functions $C^\infty(M)$ satisfying the Leibnitz identity $\{fg, h\} = f\{g, h\} + \{f, h\}g$. If Jacobi identity is not

*E-mail : balin@leor.technion.ac.il

†E-mail : mat9304@technion.technion.ac.il

required one speaks about pre-Poisson brackets. A mapping $F : M_1 \rightarrow M_2$ of two manifolds equipped with Poisson brackets $\{\cdot, \cdot\}_1$ and $\{\cdot, \cdot\}_2$, respectively, is called *Poisson* if a canonical mapping $F^* : C^\infty(M_2) \rightarrow C^\infty(M_1)$ is a Lie algebra homomorphism, i.e.

$$\{f \circ F, g \circ F\}_1 = \{f, g\}_2 \circ F.$$

Suppose now that M is equipped with a multiplication, i.e. a mapping $*$: $M \times M \rightarrow M$. This immediately gives rise to a co-multiplication (diagonal) $\Delta : C^\infty(M) \rightarrow C^\infty(M) \otimes C^\infty(M)$ acting as

$$\Delta(f)(x, y) = f(x * y),$$

here a usual identification of $C^\infty(M \times M)$ with $C^\infty(M) \otimes C^\infty(M)$ is assumed.

A Poisson bracket is said to be *compatible* with the multiplication if $*$: $M \times M \rightarrow M$ is a Poisson mapping where $M \times M$ is equipped with a product Poisson bracket. In other words, the following identity should be satisfied:

$$\Delta(\{f, g\}) = \{\Delta(f), \Delta(g)\} \tag{1}$$

where

$$\{p \otimes q, r \otimes s\} \stackrel{\text{def}}{=} \{p, r\} \otimes qs + pr \otimes \{q, s\}.$$

In particular, a bracket on a vector space compatible with addition structure is exactly a Berezin–Lie one, i.e. a linear bracket (speaking about linear, quadratic, etc., brackets we will always mean that a bracket of two *linear* functions is linear, quadratic, etc., respectively).

The next-simplest case is quadratic Poisson brackets (see [2], [3], [4]). Such a bracket may be compatible only with a bilinear operation on the vector space, i.e. with an algebra structure. One of the most important cases, that of a full matrix algebra $\text{Mat}(n, K)$ where $K = \mathbb{R}$ or \mathbb{C} , was investigated in detail in a series of works by B. Kupershmidt [7, 8, 9].

A natural question arises, whether it is possible to obtain a quantization of these brackets. This can be done using quadratic algebras (see [5]) and will be a subject of the forthcoming paper. Quantization of the corresponding Poisson Lie groups may also be simpler than in general case (see e.g. [6]) because such Poisson Lie groups bear a preferred coordinate system (linear functions on the original algebra).

Authors are grateful to J. Donin, B. Kupershmidt, A. Stolin, and N. Zobin for many stimulating discussions.

2 Quadratic brackets compatible with algebra structure

Let A be an associative algebra, i.e. vector space with a bilinear associative multiplication $*$. The symbol A_L will denote an *adjacent* Lie algebra, i.e. Lie algebra on the same space with a commutator $[a, b] = ab - ba$. The dual space A^* is naturally embedded into functions algebra $C^\infty(A)$, as well as all its symmetric powers $\text{Sym}((A^*)^{\otimes n})$ are. Again, Poisson bracket is understood as a Lie algebra structure on $C^\infty(A)$, i.e. mapping $B: C^\infty(A) \wedge C^\infty(A) \rightarrow C^\infty(A)$ satisfying Jacobi identity.

Definition A Poisson bracket B is called *quadratic* if

$$B(A^* \wedge A^*) \subset \text{Sym}((A^*)^{\otimes 2}).$$

If $\{e_i\}$ is a basis in A and x^i are coordinate functions, then quadratic Poisson bracket is given by

$$\{x^i, x^j\} = c_{kl}^{ij} x^k x^l, \quad (2)$$

a summation over repeated indices will be always assumed.

With A being an algebra, its tensor square $A \otimes A$ can also be given an algebra structure by a componentwise multiplication. Then the set of symmetric tensors $\text{Sym}(A \otimes A) \subset A \otimes A$ is its subalgebra, while the set of skew-symmetric tensors $A \wedge A \subset A \otimes A$ is a $\text{Sym}(A \otimes A)$ -bimodule. A linear mapping $D: \mathfrak{A} \rightarrow V$ from algebra \mathfrak{A} to a \mathfrak{A} -bimodule V is called a *differentiation* if it obeys the condition

$$D(p \cdot q) = pD(q) + D(p)q \quad (3)$$

for all $p, q \in \mathfrak{A}$.

Theorem 1 *Quadratic Poisson bracket $\delta^*: A^* \wedge A^* \rightarrow \text{Sym}((A^*)^{\otimes 2})$ is compatible with the algebra structure in A if and only if its dual mapping*

$$\delta: \text{Sym}(A^{\otimes 2}) \rightarrow A \wedge A$$

is a differentiation.

See [3] for proof. It does not make use of Jacobi identity, and therefore applies for general pre-Poisson brackets as well.

Example 1 [10] Consider a Poisson bracket

$$\{x^i, x^j\} = x^i x^j \quad \text{for } i < j.$$

It is compatible with the multiplication of the “first column algebra” (algebra of $n \times n$ -matrices having nonzero elements only in the first column).

For a linear operator $P: V \otimes V \rightarrow V \otimes V$ symbol P^{12} will mean operator $V \otimes V \otimes V \rightarrow V \otimes V \otimes V$ acting as P on the first and second tensor component and as identity operator on the third one; notations like P^{13} have a similar meaning. A *Schouten bracket* $[[P, P]]$ is then defined as

$$[P^{12}, P^{13}] + [P^{12}, P^{23}] + [P^{13}, P^{23}].$$

with $[\cdot, \cdot]$ meaning an ordinary commutator.

Let δ^* be a quadratic pre-Poisson bracket, and $\tilde{\delta}: A \otimes A \rightarrow A \otimes A$ be an arbitrary linear extension of the dual operator δ .

Theorem 2 *Bracket δ^* is Poisson (i.e. satisfies Jacobi identity) if and only if*

$$[[\tilde{\delta}, \tilde{\delta}]](X) = 0 \tag{4}$$

for any fully symmetric tensor $X \in A \otimes A \otimes A$.

Remark In particular, it is asserted in Theorem that $[[\tilde{\delta}, \tilde{\delta}]](X)$ for fully symmetric X depends on δ only, and not on its specific extension $\tilde{\delta}$.

The proof is made by a straightforward (and rather tedious) computation.

Note also that Theorem 2 does not require bracket δ^* to be compatible with algebra.

3 Quadratic brackets and Poisson Lie groups

Lie bialgebra structures (cocommutators) on a Lie algebra L are in one-to-one correspondence with the *linear* Poisson brackets compatible with a commutator in L (see [1]). Quadratic Poisson brackets on associative unital algebras are also directly connected with Lie bialgebras:

Lemma 1 *Let A be an associative algebra with a unit u , and $\delta^*: A^* \wedge A^* \rightarrow \text{Sym}((A^*)^{\otimes 2})$ be a quadratic Poisson bracket compatible with it. Then the mapping (comultiplication)*

$$\Delta(x) = \delta(x \otimes u + u \otimes x) \quad (5)$$

equips A_L with a Lie bialgebra structure.

Proof: computation.

So, a differentiation $\delta: \text{Sym}(A^{\otimes 2}) \rightarrow A \wedge A$ satisfying condition (4) defines two brackets compatible with A_L , quadratic δ^* and linear Δ^* with Δ given by (5).

Theorem 3 *These two brackets are compatible with one another, i.e. their arbitrary linear combination $\alpha\delta^* + \beta\Delta^*$ is also a Poisson bracket compatible with A_L .*

Proof Let $u = u^i e_i$ be a unit of A (as usual, $\{e_i\} \subset A$ is an additive basis). Consider a change of coordinates $f_t(x) = x - tu$ ($t \in \mathbb{R}$ is a parameter). In new coordinates the bracket δ^* becomes $\delta^* + t\Delta^*$. This obviously means a compatibility. ■

The following Theorem directly relates quadratic brackets to Poisson Lie structures. Drinfel'd in [1] shows that there is a one-to-one correspondence between Poisson Lie groups and Lie bialgebra structures.

Theorem 4 *Let A be an associative algebra with a unit u , and $\delta^*: A^* \wedge A^* \rightarrow \text{Sym}((A^*)^{\otimes 2})$ be a quadratic Poisson bracket compatible with it. Then the group of invertible elements of A is a Poisson Lie group with respect to a restriction of the bracket δ^* to it. The corresponding bialgebra is exactly as in Lemma 1.*

Proof The group is an open submanifold of A , so, it is apparently a Poisson submanifold. Since the group inherits its multiplication from the algebra, it is a Poisson Lie group. To obtain bialgebra structure, one has to take a bracket of two coordinate function in the vicinity of the unit u , and to take its differential at the point u . Again, take a change of variables $f_t(x) = x - tu$, and the bracket of coordinate functions will be $\delta^* + t\Delta^*$; the differential at the point u is exactly Δ . ■

Corollary 1 *Let G be a Poisson Lie group whose Lie algebra \mathfrak{g} is adjacent to an unital associative algebra structure. Then G has a coordinate system in which its Lie Poisson bracket is quadratic.*

4 Coboundary brackets

Restrict now our consideration to a *coboundary* case when differentiation δ is internal i.e. given by a formula

$$\delta(x) = [r, x] \quad (6)$$

with some $r \in A \wedge A$ (mapping (6) is obviously a differentiation).

If $r = \sum_i \alpha_i \otimes \beta_i \in A \otimes A$ then r^{12} is, by definition, $\sum_i \alpha_i \otimes \beta_i \otimes 1 \in A \otimes A \otimes A$. Here 1 is a formal unit (a unit of an universal enveloping algebra of A_L). Then on Schouten bracket $[[r, r]] \in A \otimes A \otimes A$ is defined as

$$[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}]. \quad (7)$$

Theorem 5 *Differentiation δ satisfies condition (4) (and therefore defines a Poisson bracket) if and only if $[[r, r]]$ commutes with any fully symmetric tensor $X \in A \otimes A \otimes A$.*

Remark If A has a unit 1 then hypothesis of Theorem 5 is equivalent to the following property: $[[r, r]]$ commutes with all the tensors of the type $1 \otimes 1 \otimes x + 1 \otimes x \otimes 1 + x \otimes 1 \otimes 1$, for $x \in A$. Or, in other words, $[[r, r]]$ should be Ad-invariant with respect to the Lie algebra A_L .

Theorem 5 is just Theorem 2 for a coboundary bracket. Note also that for coboundary quadratic Poisson brackets one can introduce concepts of triangular, quasi-triangular, factorizable, etc., brackets, like those introduced in the theory of coboundary Lie bialgebras (see [6]).

Corollary 2 *Let A be associative algebra, and \mathfrak{g} is a Lie subalgebra of A_L . Then any element $r \in \mathfrak{g} \wedge \mathfrak{g}$ satisfying classical Yang–Baxter equation (see [6]) defines a quadratic Poisson bracket compatible with A .*

5 Examples

Besides those described here, several important examples can be found in the classical article [11].

Example 2 Let A be an associative algebra with a solvable A_L . By Ado theorem any representation of A_L has an eigenvector. Applying this to a regular representation, one obtains that A_L contains two elements, a and b , with $[a, b] = sb$ for some constant s . The element $r = a \otimes b - b \otimes a$ satisfies classical Yang–Baxter equation and thus, according to Corollary 2, defines a quadratic Poisson bracket on A .

Example 3 Consider a body \mathbf{H} of quaternions as a four-dimensional \mathbb{R} -algebra. Then any element $r \in \mathbf{H} \wedge \mathbf{H}$ of the type

$$r = a \mathbf{i} \wedge \mathbf{j} + b \mathbf{i} \wedge \mathbf{k} + c \mathbf{j} \wedge \mathbf{k} \quad (8)$$

satisfies conditions of Theorem 5. Corresponding Poisson bracket on \mathbf{H} is:

$$\{x^1, x^2\} = x^2(bx^3 - ax^4) + c((x^3)^2 + (x^4)^2) \quad (9)$$

$$\{x^1, x^3\} = -x^3(cx^2 + ax^4) - b((x^2)^2 + (x^4)^2) \quad (10)$$

$$\{x^1, x^4\} = x^4(-cx^2 + bx^3) + a((x^2)^2 + (x^3)^2) \quad (11)$$

$$\{x^2, x^3\} = x^1(-bx^2 + cx^3) \quad (12)$$

$$\{x^2, x^4\} = -x^1(ax^2 + cx^4) \quad (13)$$

$$\{x^3, x^4\} = x^1(ax^3 - bx^4) \quad (14)$$

where x^1, x^2, x^3, x^4 is a basis in \mathbf{H}^* dual to $1, \mathbf{i}, \mathbf{j}, \mathbf{k} \in \mathbf{H}$.

A Lie group corresponding to \mathbf{H} is a multiplicative group of all nonzero quaternions. It contains a subgroup $\{z \in \mathbf{H}, \|z\| = 1\}$ isomorphic to $\text{SU}(2)$. It can be checked (see [3]) that this subgroup is Poisson with respect to all the brackets (9)–(14). Thus, $\text{SU}(2)$ bears a three-parameter family of Poisson Lie structures.

Example 4 Consider a Lie algebra \mathfrak{G} satisfying an identity $[[\mathfrak{G}, \mathfrak{G}], \mathfrak{G}] = 0$ (i.e. $[[a, b], c] = 0$ for all $a, b, c \in \mathfrak{G}$) (a particular case here is Heisenberg algebra, a three-dimensional Lie algebra with generators p, q , and z , and relations $[p, q] = \hbar z$, and $[z, \cdot] = 0$). Then $a * b = \frac{1}{2}[a, b]$ is an associative operation with $[a, b] = a * b - b * a$. An arbitrary element $r \in \mathfrak{G} \wedge \mathfrak{G}$ commutes

with all the symmetric tensors from $\mathfrak{G} \otimes \mathfrak{G}$ and therefore defines a zero Poisson bracket on \mathfrak{G} .

Associative algebra \mathfrak{G} has no unit. Having it added, consider a space $A = \langle 1 \rangle \oplus \mathfrak{G}$. Then the Lie group G corresponding to \mathfrak{G} is an affine subspace $1 + \mathfrak{G} \subset A$ (it is the usual implementation of G via exponents in the universal enveloping algebra of \mathfrak{G} ; the thing is that $a^2 = 0$ for any $a \in \mathfrak{G}$). Fix a basis $\{e_i\} \in \mathfrak{G}$; it defines a natural coordinate system in the group G . The above construction applied to $r = r^{ij}e_i \otimes e_j \in \mathfrak{G} \wedge \mathfrak{G} \subset A \wedge A$ gives now the following Poisson Lie bracket on G :

$$\{x^p, x^q\} = 2(r^{pl}c_{li}^q + r^{lq}c_{li}^p)x^i \quad (15)$$

where $\{x^i\} \in \mathfrak{G}^*$ is a basis dual to $\{e_i\}$, and c_{ij}^k are structure constants of the associative operation $*$ on \mathfrak{G} . Note that the bracket is nonzero (unlike bracket on \mathfrak{G} itself) and linear which suggests possibility to obtain its quantization.

References

- [1] V.G. Drinfeld. Hamiltonian structures on Lie groups, Lie bialgebras and geometric meaning of classical Yang–Baxter equations. *Dokl. Akad. Nauk SSSR*, 268:285–287, 1983.
- [2] E.K. Sklyanin. Some algebraic structure connected with the Yang–Baxter equation. *Funct. Anal. Appl.*, 17:263–270, 1982.
- [3] A.A. Balinsky and Yu.M. Burman. Quadratic poisson brackets compatible with an algebra structure. *J. Phys. A: Math. Gen.*, 27:L693–L696, 1994.
- [4] B.A. Kupershmidt. Conformally multiplicative poisson structures in linear algebra and algebraic geometry. In J. Harnad and J.E. Marsden, editors, *Hamiltonian Systems, Transformation Groups and Spectral Transform Methods*, pages 177–188, 1990.
- [5] Yu. Manin. Quantum groups and non-commutative geometry. Montreal preprint, 1988.
- [6] V.G. Drinfeld. Quantum groups. In *Proc. ICM86*, volume 1, pages 798–820, Berkeley, 1987.

- [7] B.A. Kupershmidt. A q -analogue of the dual space of the Lie algebras $gl(2)$ and $sl(2)$. *J. Phys. A: Math. Gen.*, 26:L1–L4, 1993.
- [8] B.A. Kupershmidt. All quantum group structures on the supergroup $GL(1|1)$. *J. Phys. A: Math. Gen.*, 26:L251–L256, 1993.
- [9] B.A. Kupershmidt. Poisson relation between minors and their consequences. Preprint, 1994.
- [10] N.Yu. Reshetikhin L.D. Faddeev and L.A. Takhtajan. Quantization of Lie groups and Lie algebras. *Leningrad Math. J.*, 1:193–225, 1990.
- [11] M.A. Semenov-Tyan-Shanskii. What is classical R -matrix ? *Funct. Anal. Appl.*, 17(17):259–272, 1983.