GEOMETRIC INTERPRETATION OF THE POISSON STRUCTURE IN AFFINE TODA FIELD THEORIES

BENJAMIN ENRIQUEZ AND EDWARD FRENKEL

June 1996

ABSTRACT. We express the Poisson brackets of local fields of the affine Toda field theories in terms of the Drinfeld-Sokolov dressing operator. For this, we introduce a larger space of fields, containing "half screening charges" and "half integrals of motions". In addition to local terms, the Poisson brackets contain nonlocal terms related to trigonometric r-matrices.

INTRODUCTION

Since the work of Zakharov and Shabat [23], the dressing techniques have played an important role in the theory of classical integrable systems. These techniques have been developed by Drinfeld and Sokolov in [7] in the framework of affine Toda field theories. Later, Feigin and one of us proposed ([13], [14]) another approach to these theories; this approach was shown ([8], [10]) to be equivalent to that of [7]. In those works, the space of local fields of the Toda theory (equivalently, the mKdV hierarchy) associated to an affine Lie algebra \mathfrak{g} is described as the ring of functions on the coset space N_+/A_+ of a unipotent subgroup of the Kac-Moody group G corresponding to \mathfrak{g} . The mKdV flows are then identified with the right action of the principal commutative Lie algebra \mathfrak{a} normalizing A_+ , N_+ being viewed as an open subset of the flag manifold of G. This leads to a system of variables, in which the flows become linear and hence can be integrated.

In the works on quantization of the Toda theories, an important role is played by the vertex operator algebra structure on the space of local fields. At the classical level, this gives rise to what we call here a vertex Poisson algebra (VPA) structure on the space of local fields of a Toda theory. The notion of the VPA structure coincides with the notion of "coisson algebra" (on the disc) introduced by Beilinson and Drinfeld in [5]. The goal of this work is to define this and related structures on the space of fields of a Toda theory in the Lie group terms using the identification described above.

The research of the second author was partially supported by grants from the Packard Foundation, NSF and the Sloan Foundation

To this end, we make use of an idea introduced earlier by Feigin and one of us in [9], where a similar problem was solved in the setting of the classical lattice Toda theory associated to $\widehat{\mathfrak{sl}}_2$. In that work, the space of local fields was extended by the "half screening charges" and "half integrals of motions". The screening charges and integrals of motions (IM's) are sums of the lattice translates of certain expressions, and their "halfs" are just the sums of positive translates of the same expressions. The full space was then identified with the quotient G/H, where H is the Cartan subgroup of G. The smaller spaces of fields, without half screening charges (resp. without half IM's), can be obtained by taking the quotient of G/H from the left by a Borel subgroup B_- (resp. from the right by the positive part of the loop group of the Cartan subgroup). In this interpretation, the Poisson structure on the full space is given by the difference $\ell(R) - r(R^{\infty})$, where $\ell(R)$ stands for the left action of the trigonometric r-matrix $R \in \mathfrak{g}^{\otimes 2}$ and $r(R^{\infty})$ stands for the right action of its "infinitely twisted" version $R^{\infty} \in \mathfrak{g}^{\otimes 2}$. This leads to a description of the Poisson structures of the quotients.

In this work, we enlarge the space of local fields of the continuous affine Toda theories in a similar way, by adding continuous analogues of the half screening charges and IM's. Using results of [9, 10], we identify the full space $\bar{\pi}_0$ obtained this way with the space of functions on $B_- \times N_+$. We then study the structure of nonlocal VPA on $\bar{\pi}_0$. The axioms for this structure are given in Section 1. The main feature is that the the Poisson bracket $\{u(x), v(y)\}$, where u, v are elements of the algebra, can be expressed as a linear combination of local terms of the form $u_k(x)\partial_x^k\delta(x-y)$ with $k \ge 0$, and of nonlocal terms of the form $a(x)\partial_x^{-1}\delta(x-y)b(y)$.

A similar formalism was introduced by Radul [22] in the framework of formal variational calculus (see also [21]). Natural examples of nonlocal VPA's are given by the higher Adler-Gelfand-Dickey (AGD) structures (i.e. structures obtained from the pair of the first two AGD structures by application of the Magri recursion procedure) introduced in [11, 1].

In Sections 2 and 3 we give a geometric description of the nonlocal VPA structure of $\bar{\pi}_0 = \mathbb{C}[B_- \times N_+]$. A generic element $g = (b_-, n_+)$ of $B_- \times N_+$ can be considered as the product of expansions of the scattering matrix of the Lax operator from $-\infty$ to x for a small spectral parameter λ , and from x to $+\infty$ for a large λ .

The Poisson bracket on N_+ is obtained by a straightforward extension of the local VPA structure on N_+/A_+ . On the other hand, according to [4] and in the spirit of [12], [3], we define the Poisson bracket on B_- via the trigonometric *r*-matrix. We show that the trigonometric Poisson brackets on B_- are compatible with the Poisson brackets of local fields (see Lemma 2.2). The Poisson brackets of N_+ and between N_+ and B_- also have nonlocal parts which we determine in Lemmas 3.3 and 3.4. To derive the complete expression for the Poisson brackets (Thm. 3.1), we use the

evolution equation

(1)
$$\partial_x g(x) = g(x)p_{-1}$$

where p_{-1} is a degree -1 element of \mathfrak{a} .

After that we obtain another realization of the VPA structures on N_+ and N_+/A_+ purely in terms of the unipotent group elements (i.e. the Drinfeld-Sokolov dressing operators) – see Cor. 3.1 and formula (47). The latter formula could in principle be obtained directly in the framework of N_+ , but the simple form (1) of the action of ∂_x on a generic element g of the whole Kac-Moody group G makes the derivation easier on G.

Let us now say a few words about possible applications and extensions of this work. First, one may think of the following program of quantization of our results: to formulate quantum axioms corresponding to nonlocal VPA in the spirit of $[5]^1$; to quantize the geometric formulas (41), (47), (50) for nonlocal VPA structures; to realize these formulas in terms of the algebra of local fields of quantum Toda theories. The combination of small and large limits for the spectral parameter which we use is reminiscent of the work [4]. Second, it would be interesting to carry out the present work in the case of higher AGD structures; this would lead to a family of compatible nonlocal VPA structures on $B_- \times N_+$ (we construct such a family in sect. 4, but its connection with the AGD structures is not clear to us). Next, it would be interesting to obtain similar results for other soliton equations such as the nonlinear Schrödinger (NLS) equation; in that case \mathfrak{a} should be replaced by the loop algebra with values in the Cartan subalgebra \mathfrak{h} . The geometric interpretation of the NLS variables analogous to the one used here, was obtained by Feigin and one of us [15] (the case of \mathfrak{sl}_2 was also treated in [2]). Finally, the fact that the VPA structure given in Thm. 3.1 is left G-invariant, leads us to conjecture the existence of affine Weyl group symmetries of the mKdV hierarchies, mixing local and nonlocal terms (see remark 3.5). These symmetries are probably connected with the Darboux transformations.

The first author would like to thank B. Feigin for his collaboration in [9] where the ideas of extension of the space of local fields were used, and A. Orlov and V. Rubtsov for their collaboration in [11] on the subject of nonlocal Poisson structures, and for discussions about this work. The second author thanks B. Feigin for useful discussions. He is also grateful to P. Schapira for his hospitality at Université Paris VI, where this work was completed.

The first author would like to dedicate this paper to A. Guichardet, his senior colleague who is about to retire from Ecole Polytechnique.

¹In the course of writing this paper, we became aware of several works dealing with vertex operator structures containing logarithmic terms, see [19, 16] and references therein

1. Nonlocal vertex Poisson Algebras.

Let (R, ∂) be a differential ring, and let \mathcal{A} be the associated ring of formal pseudodifferential operators. The ring \mathcal{A} has generators $\tilde{\partial}, \tilde{\partial}^{-1}$, and i(r) for $r \in R$, and relations $\tilde{\partial}\tilde{\partial}^{-1} = \tilde{\partial}^{-1}\tilde{\partial} = 1$, $[\tilde{\partial}, i(r)] = i(\partial r)$ for $r \in R$, and $i : R \to \mathcal{A}$ is an algebra morphism. In what follows, we will denote by ∂, ∂^{-1} and $r, \tilde{\partial}, \tilde{\partial}^{-1}$ and i(r), respectively.

1.1. The modules \mathcal{R}_n . Let $n \geq 1$ be an integer. We will use the following notation in the algebra $\mathcal{A}^{\otimes n}$: $\partial_{x_i}, \partial_{x_i}^{-1}$ will stand for $\otimes_{j=1}^{i-1} 1 \otimes r \otimes_{j=i+1}^n 1$, $\otimes_{j=1}^{i-1} 1 \otimes \partial \otimes_{j=i+1}^n 1$, $\otimes_{j=1}^{i-1} 1 \otimes \partial^{-1} \otimes_{j=i+1}^n 1$, respectively. For $T \in \mathcal{A}$, we also denote $\otimes_{j=1}^{i-1} 1 \otimes T \otimes_{j=i+1}^n 1$ by $T(x_i)$.

We define \mathcal{R}_n to be the quotient of $\mathcal{A}^{\otimes n}$ by the left ideal I_n , generated by $\sum_{i=1}^n \partial_i$ and $r(z_i) - r(z_j)$ for $r \in R$ and $i, j = 1, \ldots, n$. We will sometimes denote \mathcal{R}_n by $\mathcal{R}_n(R)$. Let us consider \mathcal{R}_n as a left $\mathcal{A}^{\otimes n}$ -module and denote by δ_{x_1,\ldots,x_n} its generator $1 + I_n$. We then have the relations

(2)
$$(r(x_i) - r(x_j))\delta_{x_1,\dots,x_n} = 0, \quad (\sum_{i=1}^n \partial_{x_i})\delta_{x_1,\dots,x_n} = 0,$$

so that δ_{x_1,\ldots,x_n} plays the role of the distribution $\prod_{i=2}^n \delta(x_1 - x_i)$.

The module \mathcal{R}_2 admits the following simple description. Let $T \mapsto T^*$ be the anti-automorphism of \mathcal{A} , defined by $r^* = r$ and $\partial^* = -\partial$. Let us endow \mathcal{A} with the $\mathcal{A}^{\otimes 2}$ -module structure, defined by $(a \otimes b)c = acb^*$, for $a, b, c \in \mathcal{A}$. Then the linear map $op : \mathcal{R}_2 \to \mathcal{A}$, defined by $op((a \otimes b)\delta_{x_1x_2}) = ab^*$, is an isomorphism of $\mathcal{A}^{\otimes 2}$ -modules. The inverse map to op is given by $op^{-1}(a) = (a \otimes 1)\delta_{x_1x_2} = (1 \otimes a^*)\delta_{x_1x_2}$.

1.2. Definition of the nonlocal VPA structure. A nonlocal VPA structure on (R, ∂) is a linear map $P : R \otimes R \to \mathcal{R}_2$, satisfying the following conditions:

(3)
$$P(ab\otimes c) = a(x_1)P(b\otimes c) + b(x_1)P(a\otimes c),$$

(4)
$$P(\partial a \otimes b) = \partial_{x_1} P(a \otimes b),$$

(5)
$$P(b \otimes a) = -\sigma(P(a \otimes b)),$$

for $a, b, c \in R$, where σ is the involutive automorphism of \mathcal{R}_2 defined by $\sigma((a \otimes b)\delta_{x_1x_2}) = (b \otimes a)\delta_{x_1x_2}$, for $a, b \in \mathcal{A}$, and the Jacobi identity that we formulate below.

Let us define a map $P_{x,yz} : R \otimes \mathcal{R}_2 \to \mathcal{R}_3$ by the following rules (we attach indices y, z to \mathcal{R}_2 and x, y, z to \mathcal{R}_3):

$$P_{x,yz}(a \otimes \delta_{y,z}) = 0, \qquad P_{x,yz}(a \otimes \partial_y m) = \partial_y P_{x,yz}(a \otimes m),$$

 $P_{x,yz}(a \otimes b(y)T(z)\delta_{yz}) = (op \circ P)(a \otimes b)(x)T(z)\delta_{xyz} + b(y)P_{x,yz}(a \otimes T(z)\delta_{yz})$ for $a, b \in R, T \in \mathcal{A}, m \in \mathcal{R}_2$. One can check easily that $P_{x,yz}$ is well-defined by these conditions. This follows from the identity $P_{x,yz}(a \otimes \partial_y b(y)m) - P_{x,yz}(a \otimes b(y)\partial_y m) = P_{x,yz}(a \otimes (\partial b)(y)m)$, which can be checked by puting m in the form $(1 \otimes T)\delta_{yz}$, $T \in \mathcal{A}$.

The Jacobi identity is then expressed as

(6)
$$P_{x,yz}(a \otimes P(b \otimes c)) = \sigma_{xy}[P_{x,yz}(b \otimes P(a \otimes c))] + \sigma_{xz}[P_{x,yz}(c \otimes P(b \otimes a))]$$

for any $a, b, c \in \mathbb{R}$, where σ_{xy}, σ_{xz} are the automorphisms of \mathcal{R}_3 , defined by

$$\sigma_{xy}(T(x)U(y)V(z)\delta_{xyz}) = U(x)T(y)V(z)\delta_{xyz},$$

and

$$\sigma_{xz}(T(x)U(y)V(z)\delta_{xyz}) = V(x)U(y)T(z)\delta_{xyz}$$

for $T, U, V \in \mathcal{A}$.

Remark 1.1 We may consider elements of \mathcal{R}_2 as kernels in x_1 and x_2 , expressed as linear combinations of $r(x_1)\partial_{x_1}^k\delta(x_1-x_2)$, and think of $P(a \otimes b)$ as $\{a(x_1), b(x_2)\}$. The terms with $k \geq 0$ are called local and the terms with k < 0 are called nonlocal.

The expression $P_{x,yz}(a \otimes m)$ should then be thought of as expressing the Poisson bracket of the form $\{a(x), m(y, z)\}$, where m is some kernel.

Note that $P_{x,yz}$ also has the properties

$$P_{x,yz}(a \otimes \partial_z m) = \partial_z P_{x,yz}(a \otimes m),$$

 $P_{x,yz}(a \otimes T(y)b(z)\delta_{yz}) = (op \circ P)(a \otimes b)(x)T(z)\delta_{xyz} + b(z)P_{x,yz}(a \otimes T(y)\delta_{yz}).$

For $a, b, c \in R$, $\{a(x), \{b(y), c(z)\}\}$ is expressed as $P_{x,yz}(a \otimes P(b \otimes c))$. On the other hand, $\{b(y), \{a(x), c(z)\}\}$ is expressed as $\sigma_{xy}[P_{x,yz}(b \otimes P(a \otimes c))]$, and $\{c(z), \{b(y), a(x)\}\}$ as $\sigma_{xz}[P_{x,yz}(c \otimes P(b \otimes a))]$. This explains the connection between the standard Jacobi identity for the Poisson brackets and formula (6).

1.3. Connection with the Beilinson-Drinfeld formalism. Let \mathcal{A}_+ be the subalgebra of \mathcal{A} , generated by R and ∂ (the algebra of differential operators). Let us set $\mathcal{R}_2^+(R) = op^{-1}(\mathcal{A}_+)$. We will say that P defines a local VPA, if P takes values in $\mathcal{R}_2^+(R)$.

In this case, the notion described here coincides with that of "coisson algebra" (on the disc) of Beilinson and Drinfeld ([5]). Indeed, let $X = \operatorname{Spec} \mathbb{C}[t]$, and D_X be the ring of differential operators on X. Let A be the algebra R[t], considered as a D_X -module by the rule that $\frac{d}{dt}$ acts on R as ∂ . We extend P to an operation

 $\{,\} \in \operatorname{Hom}_{D_{X \times X}}(A \boxtimes A, \Delta_* A)$

 $(\Delta: X \to X \times X \text{ denotes the diagonal embedding})$ as in [5], (0.1.7), in the following way. Let us call $t_1 = t \otimes 1$ and $t_2 = 1 \otimes t$ the coordinates of $X \times X$; $A \boxtimes A$ is identified with $R \otimes R[t_1, t_2]$, $\frac{\partial}{\partial t_1}$ and $\frac{\partial}{\partial t_2}$ acting on $R \otimes R$ as $\partial \otimes 1$ and $1 \otimes \partial$; on the other hand, $\Delta_* A$ is identified with $\mathcal{A}_+[t]$, with $t_{1,2}$ acting as t and $\frac{\partial}{\partial t_{1,2}}$ as $\frac{\partial}{\partial t}$.

We then set $\{at_1^n, bt_2^m\} = t^{n+m}P(a \otimes b)$, for $a, b \in R$.

Remark 1.2 As explained in [5], some local VPAs can be obtained as a classical limit of a family of chiral algebras (or vertex operator algebras [6, 17]), in the same way as one obtains Poisson algebras as a classical limit of a family of associative algebras. In particular, the local VPA π_0 described below is the classical limit of the vertex operator algebra of a Heisenberg algebra (see Remark 2 in [14]).

It would be interesting to generalize the notion of chiral algebra to allow for nonlocality. П

1.4. A class of nonlocal VPA's. In this section we give a construction of a class of nonlocal VPA's. The results of this section will only be used in the proof of Prp. 3.3. However, the construction presented here might be of general interest.

Proposition 1.1. Let $E_k, k \ge 0$, be the subspace of $Der(R)^{\otimes 2}$, consisting of all tensors $\sum_{\alpha} u_{\alpha} \otimes v_{\alpha}$, such that

$$\sum_{\alpha} (u_{\alpha}a)(\partial^{i}v_{\alpha}b) = 0, \quad \forall a, b \in R, \quad i = 0, \dots, k$$

Suppose we are given elements $\sum_{\alpha} x_i^{(\alpha)} \otimes y_i^{(\alpha)}$ of $\operatorname{Der}(R)^{\otimes 2}$ for all $i \geq -1$.

Assume that

$$\sum_{\alpha} x_i^{(\alpha)} \otimes y_i^{(\alpha)} = (-1)^{i+1} \sum_{\alpha} y_i^{(\alpha)} \otimes x_i^{(\alpha)},$$

and that

$$\sum_{\alpha} [x_i^{(\alpha)}, \partial] \otimes y_i^{(\alpha)} \in \sum_{\alpha} x_{i-1}^{(\alpha)} \otimes y_{i-1}^{(\alpha)} + E_i$$

(here we set for $i \leq -2$, $\sum_{\alpha} x_i^{(\alpha)} \otimes y_i^{(\alpha)} = 0$; so that $\sum_{\alpha} x_{-1}^{(\alpha)} \otimes y_{-1}^{(\alpha)} \in (\operatorname{Der}(R)^{\partial})^{\otimes 2}$). Then the formula

(7)
$$P(a \otimes b) = \sum_{i \geq -1} \sum_{\alpha} (x_i^{(\alpha)} a)(x) (y_i^{(\alpha)} b)(y) \partial_x^i \delta_{xy},$$

for $a, b \in R$ defines a nonlocal VPA structure on R.

Proof. The first condition ensures the antisymmetry of P, the second condition is equivalent to the ∂ -linearity condition (4). The first condition being satisfied, the Jacobi identity for P is automatically satisfied: for example, the term $P_{xy,z}(P(a \otimes$ $b \otimes c$ is equal to

$$\sum_{i,j,\alpha,\beta} (x_j^{(\beta)} x_i^{(\alpha)} a)(x) (y_i^{(\alpha)} b)(y) (y_j^{(\beta)} c)(z) (-1)^{i+j} \partial_y^i \partial_z^j \delta_{xyz} + \sum_{i,j,\alpha,\beta} (x_i^{(\alpha)} a)(x) (x_j^{(\beta)} y_i^{(\alpha)} b)(y) (y_j^{(\beta)} c)(z) (-1)^j \partial_x^i \partial_z^j \delta_{xyz}$$

whose second term is cancelled by

$$\sum_{i,j,\alpha,\beta} (x_j^{(\beta)} x_i^{(\alpha)} b)(y)(y_i^{(\alpha)} c)(z)(y_j^{(\beta)} a)(x)(-1)^{i+j} \partial_z^i \partial_x^j \delta_{xyz}$$

which is a cyclic permutation of the first one.

Remark 1.3 Let us denote by Der(R) the Lie algebra of derivations of R and assume that we have $\varpi \in (Der(R)^{\partial})^{\otimes 2}$, and $P_+ : R \otimes R \to \mathcal{A}_+$, such that writing $\varpi = \sum_i \varpi_i \otimes \varpi'_i$ ($\varpi_i, \, \varpi'_i$ derivations of R commuting with ∂),

(8)
$$op \circ P(a \otimes b) = P_+(a \otimes b) + \varpi_i(a)\partial^{-1} \varpi'_i(b).$$

In view of the form of ϖ , the nonlocal part of axioms (5), (6) is satisfied. The l.h.s. of the Jacobi identity contains terms of the form

$$a(x)b(y)c(z)\partial_{x_i}^{-1}\partial_{x_j}^{-1}, a(x)b(y)c(z)\partial_{x_i}^{-1}\partial_{x_j}^k, a(x)b(y)c(z)\partial_{x_i}^k\partial_{x_j}^\ell,$$

 $a, b, c \in \mathbb{R}, k, \ell \geq 0, x_i \neq x_j$ run through x, y, z. The sum of the terms of the first type then cancels automatically (see Lemma 2.3).

All nonlocal VPA structures that we study in this work will be of the type described here.

Note that in particular, if P_+ is 0, then P given by (8) always defines a VPA structure.

We will denote by \mathcal{A}_{-1} the span of \mathcal{A}_{+} and the $a\partial^{-1}b$, $a, b \in \mathbb{R}$, and by $\mathcal{R}_{2}^{-1}(\mathbb{R})$ the space $op^{-1}(\mathcal{A}_{-1})$. \square

1.5. Hamiltonian vector fields. We wish to show briefly here how the notions introduced above can be related to the Gelfand-Dickey-Dorfman theory of formal variational calculus (see [18], and the introduction of [5]). Let us assume that P takes values in \mathcal{A}_+ . Define $\mathcal{V}_f \in \text{End}(R)$ by

(9)
$$\mathcal{V}_f(a) = -(op \circ P(a \otimes f)) \cdot 1,$$

the result of the action of the differential operator $P(a \otimes f)$ on $1 \in R$. Then \mathcal{V}_f is a derivation of R, commuting with ∂ . \mathcal{V}_f actually depends only on the class of f on $R/\partial R$, and is called the Hamiltonian vector field corresponding to the density f. We will also use the notation

$$\mathcal{V}_f(a) = \{\int_{-\infty}^{\infty} f, a\} = -\{a, \int_{-\infty}^{\infty} f\},\$$

where $\int_{-\infty}^{\infty} f$ denotes the class of f in $R/\partial R$.

It can be used to define a Lie algebra structure on $R/\partial R$, by the rule,

$$\{\int_{-\infty}^{\infty} f, \int_{-\infty}^{\infty} g\} = \int_{-\infty}^{\infty} \mathcal{V}_f(g), \qquad \forall f, g \in R.$$

A derivation D of R which commutes with ∂ defines an operation D_2 on \mathcal{R}_2 , in the following way. We set $D_2\delta_{xy} = 0$, and extend D_2 to the whole \mathcal{R}_2 by the condition that it commutes with ∂_x and ∂_y , and the formulae

$$D_2(a(x)m(x,y)) = a(x)D_2m(x,y) + (Da)(x)m(x,y),$$

BENJAMIN ENRIQUEZ AND EDWARD FRENKEL

$$D_2(b(y)m(x,y)) = b(y)D_2m(x,y) + (Db)(y)m(x,y)$$

for $m \in \mathcal{R}_2$, $a, b \in \mathcal{A}$; this definition makes sense because D commutes with ∂ .

We then say that D is an infinitesimal automorphism of R, if $P(Da \otimes b) + P(a \otimes Db) = D_2 P(a \otimes b)$ for $a, b \in R$.

Proposition 1.2. Under the conditions above, \mathcal{V}_f is an infinitesimal automorphism of R. Moreover, $f \mapsto \mathcal{V}_f$ is a Lie algebra homomorphism from $R/\partial R$ to $\text{Der}(R)^{\partial}$.

Proof. The first part is straightforward and the second is contained in [18].

2. Nonlocal extensions of the VPA of local fields π_0 .

2.1. Notation and definition of π_0 . Let $\tilde{\mathfrak{g}}$ be an affine Lie algebra, with generators $e_i, f_i, \alpha_i^{\vee}, i = 0, \ldots, l$, and d, subject to the relations of [20]. Let a_i, a_i^{\vee} be the labels of $\tilde{\mathfrak{g}}$, h be its Coxeter number; then $K = \sum_i a_i^{\vee} \alpha_i^{\vee}$ is a generator of the center of $\tilde{\mathfrak{g}}$. Let $\hat{\mathfrak{g}}$ be the subalgebra of $\tilde{\mathfrak{g}}$ with the same generators except d, \mathfrak{g} be the quotient of $\hat{\mathfrak{g}}$ by $\mathbb{C}K$, and let us denote the same way elements of $\tilde{\mathfrak{g}}$ and their images in \mathfrak{g} . Let \mathfrak{h} be the Cartan subalgebra of \mathfrak{g} , generated by the α_i^{\vee} 's, \mathfrak{n}_+ and \mathfrak{n}_- be the pronilpotent subalgebras generated by the e_i 's and the f_i 's respectively, and $\mathfrak{b}_- = \mathfrak{h} \oplus \mathfrak{n}_-$. Let α_i , $i = 0, \ldots, l$ be the simple roots of $\tilde{\mathfrak{g}}$, positive with respect to this decomposition. For each $x \in \mathfrak{g}$ we can write $x = x_+ + x_-$, where $x_+ \in \mathfrak{n}_+, x_- \in \mathfrak{b}_-$.

There is an invariant inner product \langle , \rangle on $\tilde{\mathfrak{g}}$; let us denote in the same way its restriction to $\hat{\mathfrak{g}}$. Let σ be any section of \mathfrak{g} to $\hat{\mathfrak{g}}$; the restriction of \langle , \rangle to $\sigma(\mathfrak{g})$ defines an inner product on \mathfrak{g} , independent of σ and again denoted by \langle , \rangle .

Let \mathfrak{h} be the Cartan subalgebra of $\tilde{\mathfrak{g}}$ spanned by $\alpha_i^{\vee}, i = 0, \ldots, l$ and d. The restriction of the inner product \langle, \rangle to $\tilde{\mathfrak{h}}$ is non-degenerate and hence defines an isomorphism $\tilde{\mathfrak{h}} \simeq \tilde{\mathfrak{h}}^*$. Let $\omega_i^{\vee} \in \tilde{\mathfrak{h}}^*$ be the *i*th fundamental coweight, i.e. it satisfies $\langle \omega_i^{\vee}, \alpha_j \rangle = \delta_{i,j}, (\omega_i^{\vee}, d) = 0$. Denote by h_i, h_i^{\vee} the elements of $\tilde{\mathfrak{h}}$, which are the images of $\alpha_i, \omega_i^{\vee} \in \tilde{\mathfrak{h}}^*$ under the isomorphism $\tilde{\mathfrak{h}} \simeq \tilde{\mathfrak{h}}^*$ (note that this is not a standard notation).

Let

$$p_{-1} = \sum_{i=0}^{l} \frac{(\alpha_i, \alpha_i)}{2} f_i,$$

and let \mathfrak{a} be the centralizer of p_{-1} in \mathfrak{g} . It is a commutative subalgebra of \mathfrak{g} , called the principal abelian subalgebra. We have $\mathfrak{a} = \mathfrak{a}_+ \oplus \mathfrak{a}_-$, where $\mathfrak{a}_{\pm} = \mathfrak{a} \cap \mathfrak{n}_{\pm}$. Let Ibe the set (with multiplicities) of integers, congruent to the exponents of \mathfrak{g} modulo h. Then \mathfrak{a}_{\pm} is generated by elements $p_n, n \in \pm I$.

We normalize the $p_n, n \in \pm I$ is such a way that

$$\langle p_n, p_{-n} \rangle = \frac{1}{h}, \quad n \in I,$$

where h is the Coxeter number of $\tilde{\mathfrak{g}}$. Let π_0 be the free differential ring generated by $u_i, i = 1, \ldots, l$; we have $\pi_0 = \mathbb{C}[u_i, \partial u_i, \ldots]$.

Proposition 2.1 ([5]). The space π_0 has a VPA structure, defined by the formula

(10)
$$P(A \otimes B) = \sum_{1 \le i, j \le l} (\alpha_i, \alpha_j) \sum_{n \ge 0} \sum_{m \ge 0} \left(\frac{\partial A}{\partial u_i^{(m)}} \partial^{n+1} \otimes \frac{\partial B}{\partial u_j^{(m)}} \partial^m \right) \delta_{xy}.$$

Remark 2.1 This structure is uniquely determined by the following formulas:

$$P(A \otimes 1) = 0, \qquad \forall A \in \pi_0,$$

and

(11)
$$P(u_i \otimes u_j) = (\alpha_i, \alpha_j) \partial_x \delta_{xy}.$$

The space π_0 also has the structures of \mathfrak{n}_+ - and \mathfrak{a}_- -modules, defined in [13]. Below we will extend these structures in three ways.

2.2. Extension by half IM's. Let B_{-} be the ind-algebraic group corresponding to \mathfrak{b}_{-} , and N_{+} be the pro-algebraic Lie group corresponding to \mathfrak{n}_{+} , A_{+} its subgroup corresponding to \mathfrak{a}_{+} . Let G be the group corresponding to \mathfrak{g} , containing B_{-} and N_{+} as subgroups.

Recall from [13], [14] the identification of π_0 and $\mathbb{C}[N_+/A_+]$ as rings and \mathfrak{n}_+ modules. Moreover, the Lie algebra \mathfrak{a}_- acts on $\mathbb{C}[N_+/A_+]$ from the right, since we can identify N_+ with an open subspace of $B_-\backslash G$. Let ∂_n be the derivation of π_0 corresponding to the right action of p_{-n} on N_+/A_+ . In particular, $\partial_1 \equiv \partial$.

Consider for $n \in I$, the hamiltonians $H_n \in \pi_0$ from [10]. They satisfy

$$\epsilon_{-\alpha_i} e_i \cdot H_n = \partial A_n^{(i)},$$

for certain $A_n^{(i)} \in \pi_{-\alpha_i}$ (in the notation of [10]). We have an isomorphism $\pi_0 \simeq \mathbb{C}[N_+/A_+]$ (see [13], [14]).

$$\partial_n H_m = \partial_m H_n = \partial H_{n,m}$$

for certain $H_{n,m} \in \pi_0$. Following [10], Sect. 4, define $\pi_0^+ = \pi_0 \otimes \mathbb{C}[F_n]_{n \in I}$, and extend the action of ∂_n to it by the formula

(12)
$$\partial_n F_m = H_{n,m}$$

(so that F_n can be viewed as a "half integrals of motion" $\int_{-\infty}^x H_n$). In particular, $\partial F_m = H_m$.

According to Thm. 5 and Prop. 9 of [10], the ring π_0^+ is isomorphic to $\mathbb{C}[N_+]$, and the action of ∂_n on π_0^+ defined this way corresponds to the right action of p_{-n} on N_+ .

Finally, the action of the generators of \mathfrak{n}_+ are defined by

(13)
$$e_i \cdot F_n = \epsilon_{-\alpha_i}^{-1} A_n^{(i)} = \phi_n(e_i)$$

(in the notation of [10]).

Now we extend the VPA structure on π_0 to a nonlocal VPA structure on π_0^+ . For $a \in \pi_0, n \in I$, we have

(14)
$$\{\int_{-\infty}^{\infty} H_n, a\} = n\partial_n a,$$

([7], prop. 4.5) so that

(15)
$$P(H_n \otimes a) \in n\partial_n a(y)\delta_{xy} + \partial_x \mathcal{R}_2^+(\pi_0)$$

Let $i_n(a)$ be the element of $\mathcal{R}_2^+(\pi_0)$, such that $P(H_n \otimes a) = n\partial_n a(y)\delta_{xy} + \partial_x(i_n(a))$ (this defines $i_n(a)$ uniquely, since $\xi \in \mathcal{A}, \ \partial \cdot \xi = 0$ implies $\xi = 0$). We then define

(16)
$$P(F_n \otimes a) = n\partial_n a(y)\partial_x^{-1}\delta_{xy} + i_n(a).$$

Next, we have:

$$P(H_n \otimes H_m) = n\partial_n H_m(x)\delta_{xy} + \partial_x(i_n(H_m)).$$

Integrating this expression w.r.t. the second variable, we obtain

$$n\partial_n H_m(x) + \partial_x (\int_{-\infty}^{\infty} i_n(H_m)(x,y)dy).$$

where

$$\int_{-\infty}^{\infty} i_n(H_m)(x,y)dy$$

is defined as $a_0(x)$, where $i_n(H_m) = \sum_{i \ge 0} a_i(x) \partial_x^i \delta_{xy}$.

On the other hand, we obtain from formula (14) that

$$\{H_n(x), \int_{-\infty}^{\infty} H_m\} = -m\partial_m H_n(x)$$

Comparing the last two formulas we find: $\int_{-\infty}^{\infty} i_n(H_m)(x,y)dy = -(n+m)H_{n,m}(x)$ (for degree reasons it cannot contain constant terms) and so $i_n(H_m) = -(n+m)H_{n,m}(x)\delta_{xy} + \partial_y Q_{n,m}$, with $Q_{n,m} \in \mathcal{A}_+$. Thus, we obtain:

$$P(H_n \otimes H_m) = n(\partial_n H_m)(x)\delta_{xy} - (n+m)H_{n,m}(y)\partial_x\delta_{xy} + (\partial Q_{n,m})(x)\partial_y\delta_{xy}$$

Using formula $H_i = \partial F_i$ we finally obtain:

(17)
$$P(F_n \otimes F_m) = (mH_{n,m}(x) + nH_{n,m}(y))\partial_x^{-1}\delta_{xy} + Q_{n,m}(x)\delta_{xy}$$

Proposition 2.2. The formulae (11), (16), (17) define a nonlocal VPA structure on π_0^+ .

Proof. To establish the antisymmetry condition, we should check that

$$P(F_n \otimes F_m) = -\sigma(P(F_m \otimes F_n));$$

this identity is $\partial_x^{-1} \partial_y^{-1}$ applied to the same identity, with H_n and H_m in place of F_n and F_m , which is true. The same argument works with the bracket $P(F_n \otimes a)$, $a \in \pi_0$, with only one application of ∂^{-1} .

Let us pass to the Jacobi identity. We should check it for the tensors $F_n \otimes F_m \otimes F_k$, $F_n \otimes F_m \otimes a$, $F_n \otimes a \otimes b$, $a, b \in \pi_0$. These identities are $\partial_x^{-1} \partial_y^{-1} \partial_z^{-1}$, resp. $\partial_x^{-1} \partial_y^{-1}$, ∂_x^{-1} applied to the same identities with H's replacing the F's, which are true.

Proposition 2.3. The action of ∂_n on π_0^+ is an infinitesimal automorphism of the nonlocal VPA structure on π_0^+ .

Moreover, we have for all $a \in \pi_0^+$,

(18)
$$P(H_n \otimes a) \in n\partial_n a(x)\delta_{xy} + \partial_x \rho_n(a),$$

with $\rho_n(a) \in \mathcal{R}_2^{-1}(\pi_0^+)$, such that

$$\rho_n(a) \in \sum_{m \in I} H_{n,m}(x)\rho_{n,m}(a)(y)\partial_x^{-1}\delta_{xy} + \mathcal{R}_2^+(\pi_0^+),$$

where $\rho_{n,m}$'s are linear endomorphisms of π_0^+ .

Proof. Since $n\partial_n$ coincides with \mathcal{V}_{H_n} , it satisfies the infinitesimal automorphism identity on $\pi_0^{\otimes 2}$. To see that this identity is also satisfied for the tensors $F_n \otimes a$ $(a \in \pi_0)$ and $F_n \otimes F_m$, we remark that they are ∂_x^{-1} , resp. $\partial_x^{-1}\partial_y^{-1}$ applied to the similar identities for $H_n \otimes a$, resp. $H_n \otimes H_m$.

According to formula (15), equation 18) is satisfied for $a \in \pi_0$, with $\rho_n(a)$ actually lying in $\mathcal{R}_2^+(\pi_0)$. Therefore (18) holds for $a = F_m$, as can be seen by applying ∂_x to (17). By Leibnitz rule, if (18) is true for $a, b \in \pi_0^+$, then it is true for ab. Moreover, we find that

$$\rho_n(ab) = a(y)\rho_n(b) + b(y)\rho_n(a).$$

This proves (18) and the properties of ρ_n in general.

Remark 2.2 Formula (18) can be viewed as a nonlocal substitute of the hamiltonian property of H_n .

2.3. Extension by $\mathbb{C}[B_-]$. Let $\mathbb{C}[B_-]$ be the ring of algebraic functions on B_- . Let $\tilde{\pi}_0 = \pi_0 \otimes \mathbb{C}[B_-]$. We extend the actions of \mathfrak{n}_+ and \mathfrak{a}_- on π_0 , to actions of \mathfrak{g} and \mathfrak{a}_- on $\tilde{\pi}_0$, in the following way.

Recall the identification as rings and \mathfrak{n}_+ -modules, of π_0 with $\mathbb{C}[N_+/A_+]$. Moreover, the \mathfrak{a}_- -action on π_0 is identified with its action by vector fields on $B_-\backslash G/A_+$ from the right, where N_+/A_+ is an open subset (recall that G is the group corresponding to \mathfrak{g}). The ring $\tilde{\pi}_0$ is then identified with the ring of functions on $B_- \times N_+/A_+$.

We define on it the actions of \mathfrak{g} and \mathfrak{a}_{-} as follows: the value of the vector field generated by $x \in \mathfrak{g}$ at $(b_{-}, n_{+}A_{+})$ is

(19)
$$(r(\operatorname{Ad}(b_{-}^{-1})(x))_{-}, \ell(\operatorname{Ad}(b_{-}^{-1})(x))_{+})$$

and the value of the vector field ∂_n (denoted by ∂ for n = 1) generated by p_{-n} , $n \in I$ is

(20)
$$(r(\operatorname{Ad}(n_+)(p_{-n})_-), \ell(\operatorname{Ad}(n_+)(p_{-n})_+));$$

here r(y) and $\ell(y)$ denote the right and left vector fields generated by a Lie algebra element y. The proof of these formulas is analogous to the proof of Lemma 1 in [10]. In particular for n = 1 we have according to Lemma 2 of [10]

In particular, for n = 1 we have according to Lemma 2 of [10]

(21)
$$\partial n_{+} = (n_{+}p_{-1}n_{+}^{-1})_{+}n_{+} = -(n_{+}p_{-1}n_{+}^{-1})_{-}n_{+} + (n_{+}p_{-1}n_{+}^{-1})n_{+} = -(p_{-1} + \sum_{i} u_{i}h_{i}^{\vee})n_{+} + n_{+}p_{-1}.$$

We also have:

(22)
$$\partial b_{-} = b_{-}(n_{+}p_{-1}n_{+}^{-1})_{-} = b_{-}(p_{-1} + \sum_{i} u_{i}h_{i}^{\vee}).$$

The last two formulas should be considered in an arbitrary representation of G of the form $V((\lambda))$, where V is finite-dimensional (see [10]).

Set now

(23)
$$P(b_{-} \otimes u_{i}) = -\partial_{y} \Big[\operatorname{Ad}(b_{-}(y))(h_{i})b_{-}(x)\partial_{x}^{-1}\delta_{xy} \Big].$$

By this formula we mean the following. In any representation of B_{-} of the form $V((\lambda))$, an element b_{-} of B_{-} can be viewed as a matrix $(b_{-,kl})$ whose entries $b_{-,kl}$ are Taylor series in λ^{-1} with coefficients in the ring $\mathbb{C}[B_{-}]$. Such functions in fact generate $\mathbb{C}[B_{-}]$. The left hand side of formula (23) is the matrix whose entries are $P(b_{-,kl} \otimes u_i)$. The right hand side of the formula is also a matrix of the same size, whose entries are elements of $\mathcal{R}_2(\mathbb{C}[B_{-}])$. Via Leibnitz rule, formula (23) defines $P(f \otimes u_i) \in \mathcal{R}_2(\mathbb{C}[B_{-}])$ for any $f \in \mathbb{C}[B_{-}]$. We interpret similarly formulas below for $P(b_{-} \otimes b_{-}), P(g \otimes g)$, etc.

Remark 2.3 In view of (22), we consider $b_{-}(x)$ as the ordered exponential

$$P \exp \int_{-\infty}^{x} (p_{-1} + \sum_{i=1}^{l} u_i(z)h_i^{\vee}) dz.$$

Note that in any representation $b_{-}(x)$ is represented by the matrix

(24)
$$\left(\operatorname{Id} - \sum_{i=0}^{l} \frac{(\alpha_i, \alpha_i)}{2} f_i S_i + \dots \right) \exp\left(\sum_{i=1}^{l} h_i^{\vee} \varphi_i(x) \right),$$

where $\varphi_i(x) = \int_{-\infty}^x u_i(y) dy$, and S_i is the *i*th "half screening"

$$S_i = \int_{-\infty}^x e^{-\varphi_i(y)} dy$$

The reason for this terminology is that if in the last formula we integrate over a closed contour, we obtain the classical limit of the screening operator of conformal field theory; the sum of the screening operators coincides with the hamiltonian of the affine Toda field theory. The other terms in the first factor of formula (24) can be expressed as consecutive Poisson brackets of the S_i 's.

Poisson bracket (23) can be informally obtained as follows. Since

$$P((p_{-1} + \sum_{i=1}^{l} u_i h_j^{\vee}) \otimes \varphi_j) = -h_j \delta_{xy},$$

we can write

$$\begin{aligned} \{b_{-}(x),\varphi_{j}(y)\} &= \int_{-\infty}^{x} dz P \exp \int_{-\infty}^{z} (p_{-1} + \sum_{i=1}^{l} u_{i}h_{i}^{\vee})(-h_{j}\delta_{zx}) \\ P \exp \int_{z}^{x} (p_{-1} + \sum_{i=1}^{l} u_{i}h_{i}^{\vee}) \\ &= -\left(P \exp \int_{-\infty}^{y} (p_{-1} + \sum_{i=1}^{l} u_{i}h_{i}^{\vee})\right)h_{j}\left(P \exp \int_{y}^{x} (p_{-1} + \sum_{i=1}^{l} u_{i}h_{i}^{\vee})\right)1_{y < x} \\ &= -b_{-}(y)h_{j}b_{-}(y)^{-1}b_{-}(x)1_{y < x},\end{aligned}$$

where $1_{y < x}$ is the function of x, y equal to 1 when y < x and to 0 else. For y fixed, this is a shifted Heaviside function in x; applying ∂_x to it gives $\delta(x - y)$, so that we identify $1_{y < x}$ with $\partial_x^{-1} \delta_{xy}$. Formula (23) is obtained by applying ∂_y to this identity. \Box

Lemma 2.1. Definition (23) is compatible with the ∂ -linearity condition (4) on P.

Proof. Both left and right hand sides of (23) satisfy the same identity

$$\partial_x(\text{lhs}) = (\text{lhs})(p_{-1} + \sum_j u_j(x)h_j^{\vee}) - b_-(x)h_i\partial_y\delta_{xy},$$

and

$$\partial_x(\mathbf{rhs}) = (\mathbf{rhs})(p_{-1} + \sum_j u_j(x)h_j^{\vee}) - b_-(x)h_i\partial_y\delta_{xy}.$$

Let us define for $b_{-} \in B_{-}$, and $x \in \mathfrak{g}$, $r(x)(b_{-}) = (\operatorname{Ad} b_{-}(x))_{-}b_{-}$ in any representation of G (recall that x_{-} stands for the projection of $x \in \mathfrak{g}$ on the second factor of the decomposition $\mathfrak{g} = \mathfrak{n}_{+} \oplus \mathfrak{b}_{-}$). This formula defines the right action of \mathfrak{g} on B_{-} viewed as an open subset of $N_{+} \setminus G$.

Let Δ_+ be the set of positive roots of $\tilde{\mathfrak{g}}$, and let e^{α} , e_{α} , $\alpha \in \Delta_+$, be dual bases of \mathfrak{n}_+ , \mathfrak{n}_- for the inner product \langle,\rangle .

Let

(25)
$$R^{+} = \frac{1}{2} \left(\sum_{\alpha} e^{\alpha} \otimes e_{\alpha} + \sum_{i} h_{i} \otimes h_{i}^{\vee} - \sum_{\alpha} e_{\alpha} \otimes e^{\alpha} \right),$$

(26)
$$R^{-} = \frac{1}{2} \left(\sum_{\alpha} e^{\alpha} \otimes e_{\alpha} - \sum_{i} h_{i} \otimes h_{i}^{\vee} - \sum_{\alpha} e_{\alpha} \otimes e^{\alpha} \right),$$

Let us define, in the tensor product of any pair of representations of G,

(27)

$$P(b_{-} \otimes b_{-}) = \{ ((r \otimes r)(R^{-})(b_{-}(x) \otimes b_{-}(x))) (1 \otimes b_{-}(x)^{-1}b_{-}(y)) \\ - ((r \otimes r)(R^{+})(b_{-}(y) \otimes b_{-}(y))) (b_{-}(y)^{-1}b_{-}(x) \otimes 1) \} \partial_{x}^{-1} \delta_{xy},$$

where pr_{-} is the projection $\mathfrak{g} \to \mathfrak{b}_{-}$ along \mathfrak{n}_{+} . This formula can be derived using the argument of Remark 2.3. A similar formula has been obtained by Faddeev and Takhtajan [12] (in the rational case) and used by Bazhanov, Lukyanov and Zamolodchikov [4] (see also [3]).

Formula (27) can also be written as

(28)
$$P(b_{-} \otimes b_{-}) = (pr_{-} \otimes pr_{-})[\mathrm{Ad}^{\otimes 2}(b_{-}(x))(R^{-}) - \mathrm{Ad}^{\otimes 2}(b_{-}(y))(R^{+})] \\ (b_{-}(x) \otimes b_{-}(y))\partial_{x}^{-1}\delta_{xy}.$$

It satisfies the antisymmetry condition (5), because

$$R^{-} = -R^{+(21)}$$

Lemma 2.2. Formula (27) is compatible with the ∂ -linearity condition (4) on P.

Proof. The l.h.s. of (27) satisfies

$$\partial_y(\text{lhs}) = (\text{lhs})(1 \otimes (p_{-1} + \sum_i u_i(y)h_i^{\vee})) \\ + \sum_i \{b_-(x)h_i\delta_{xy} - \text{Ad}(b_-(y))([p_{-1}, h_i])b_-(x)\} \otimes b_-(y)h_i^{\vee}\partial_x^{-1}\delta_{xy},$$

and the r.h.s. satisfies

$$\begin{aligned} \partial_y(\mathbf{rhs}) &= \left\{ [(r \otimes r)(R^-)(b_-(x) \otimes b_-(x))] [1 \otimes b_-(x)^{-1}b_-(y)(p_{-1} \\ &+ \sum_i h_i^{\vee} u_i(y))] \right. \\ &- \left[(r \otimes r)(R^+) [(b_-(y) \otimes b_-(y))(p_{-1} \otimes 1 + 1 \otimes p_{-1} - \sum_i u_i(y)(h_i^{\vee} \otimes 1 \\ &+ 1 \otimes h_i^{\vee})] \right] (b_-(y)^{-1}b_-(x) \otimes 1) \\ &+ [(r \otimes r)(R^+)(b_-(y) \otimes b_-(y))] [(p_{-1} + \sum_i u_i(y)h_i^{\vee})b_-(y)^{-1}b_-(x) \otimes 1] \right\} \\ &= \frac{\partial_x^{-1} \delta_{xy}}{\partial_x^{-1} \delta_{xy}} \\ &- [(r \otimes r)(R^- - R^+) \left(b_-(x) \otimes b_-(x) \right)] \delta_{xy} \end{aligned}$$

The δ_{xy} -terms coincide because $R^+ - R^- = \sum_i h_i \otimes h_i^{\vee}$. The $\partial_x^{-1} \delta_{xy}$ -terms containing $u_i(y)$ are coincide, since R^+ is invariant by the conjugation by \mathfrak{h} .

The identification of the remaining terms follows from the formula

(29)
$$[p_{-1} \otimes 1 + 1 \otimes p_{-1}, R^+] = \sum_i [p_{-1}, h_i] \otimes h_i^{\vee}.$$

Let $\rho : \mathfrak{g} \to \mathfrak{g}$ be the linear map defined by $\rho(x) = \langle R^+, 1 \otimes x \rangle$. Formula (29) means that $[\mathrm{ad}(p_{-1}), \rho]$ is the linear endomorphism of \mathfrak{g} , equal to 0 on \mathfrak{n}_+ and \mathfrak{n}_- and to $\mathrm{ad}(p_{-1})$ on \mathfrak{h} , which can be easily verified. \Box

Remark 2.4 It follows from the proof that the brackets (27) can also be expressed replacing R^+ and R^- by $R^+ + \kappa \sum_i \epsilon^i \otimes \epsilon_i$ and $R^- + \kappa \sum_i \epsilon^i \otimes \epsilon_i$, (ϵ^i) , (ϵ_i) dual bases of \mathfrak{g} . This follows also from the fact that $[\sum_i \epsilon^i \otimes \epsilon_i, b_- \otimes b_-] = 0$ for any $b_- \in B_-$. We prefer the form given here because then (5) is manifestly satisfied. \Box

Let us extend P to $\mathbb{C}[B_-]^{\otimes 2}$ by linearity, to $\mathbb{C}[B_-] \otimes \pi_0$ by formulas (23), (4) and Leibnitz rule, to $\pi_0 \otimes \mathbb{C}[B_-]$ by antisymmetry, and finally to $(\mathbb{C}[B_-] \otimes \pi_0)^{\otimes 2}$ by the Leibnitz rule.

Proposition 2.4. The operation P on $\mathbb{C}[B_{-}] \otimes \pi_0$ defined by formulas (11), (23), (27) and the rules above, satisfies the nonlocal VPA axioms.

Proof. By construction, P satisfies the ∂ -linearity axiom. The Leibnitz rule is satisfied for the brackets involving b_{-} 's since the right hand sides of (23) and (27) respect the tensor structure. For the other brackets, the Leibnitz rule is satisfied by construction. As was already mentioned, the antisymmetry of P follows from $R^{+} = R^{-(21)}$. Let us check now the Jacobi identity. For the $\mathbb{C}[B_{-}]$ part it follows from the following lemma.

Lemma 2.3. Let (R, ∂) be a differential ring. Let R_0 be a subring of R, such that R is spanned by elements of the form $\partial^i a_i$, where $a_i \in R_0$.

Let $\sum_i \varpi_i \otimes \varpi'_i \in S^2(\text{Der}(R_0))$, and let us assume that there exists a linear map $P: R \otimes R \to \mathcal{R}_2$, satisfying the Leibnitz rule and the ∂ -linearity, such that

$$P(a \otimes b) = \sum_{i} (\varpi_i a)(x)(\varpi'_i b)(y) \partial_x^{-1} \delta_{xy},$$

for $a, b \in R_0$. Then P satisfies the Jacobi identity.

Proof. It is enough to check it for elements of R_0 . We have

$$P_{xy,z}(P(a \otimes b) \otimes c) = \sum_{i,j} (\varpi_j \varpi_i a)(x)(\varpi'_i b)(y)(\varpi'_j c)(z) \varpi_x^{-1} \partial_x^{-1} \delta_{xyz} + (\varpi_i a)(x)(\varpi_j \varpi'_i b)(y)(\varpi'_j c)(z) \partial_x^{-1} \partial_y^{-1} \delta_{xyz};$$

the first term is cancelled by

$$\sum_{i,j} (\varpi_i c)(z) (\varpi_j \varpi'_i a)(x) (\varpi'_j b)(y) \partial_z^{-1} \partial_x^{-1} \delta_{xyz}$$

(which is obtained from the second by a cyclic permutation).

The Jacobi identity in the case of the proposition is then proved as follows: introduce the variables φ_i , $i = 1, \ldots, l$; let $\tilde{R} = \mathbb{C}[B_-] \otimes \mathbb{C}[\varphi_i^{(k)}]$, and identify R with a subalgebra of \tilde{R} by $\partial \varphi_i = u_i$. Extend the bracket P to \tilde{R} , by the rules

$$P(\varphi_i \otimes \varphi_j) = -(\alpha_i, \alpha_j)\partial_x^{-1}\delta_{xy}$$

and

$$P(b_{-} \otimes \varphi_i) = -\operatorname{Ad}(b_{-}(y))(h_i)b_{-}(x)\partial_x^{-1}\delta_{xy}$$

Let now $R_0 = \mathbb{C}[B_-] \otimes \mathbb{C}[\varphi_i]$; the restriction of P to R_0 is of the form given in the lemma, so P satisfies the Jacobi identity.

2.4. Extension by $\mathbb{C}[B_-]$ and half IM's. Consider now the algebra $\overline{\pi}_0 = \pi_0 \otimes \mathbb{C}[F_n] \otimes \mathbb{C}[B_-]$. We extend the actions of ∂ and of the flows ∂_n on it in the way compatible with the embeddings $\pi_0^+ \subset \overline{\pi}_0$ and $\widetilde{\pi}_0 \subset \overline{\pi}_0$. We also define a nonlocal VPA structure on $\overline{\pi}_0$, in such a way that the above embeddings are nonlocal VPA morphisms, and we define the brackets $P(F_n \otimes b_-)$ by

(30)
$$P(F_n \otimes b_-) = \partial_x^{-1} P(H_n \otimes b_-);$$

we then extend P to $\bar{\pi}_0^{\otimes 2}$ by antisymmetry and the Leibnitz rule (it is easy to check that (30) is compatible with the Leibnitz rule for the products of matrix elements of b_{-}).

Proposition 2.5. The operation P defined by formulas (11), (16), (17), (23), (27), (30), antisymmetry and the Leibnitz rule, defines a nonlocal VPA structure on $\bar{\pi}_0$.

Proof. The same as in Prop. 2.1.

The rest of this section is devoted to proving that the image of P actually lies in $\mathcal{R}_2^{-1}(\bar{\pi}_0)$ (the definition of this space is in Rem. 1.3). It is enough to prove it for $P(F_n \otimes b_-)$.

Lemma 2.4. For certain $A_{ni} \in \mathcal{R}_2^+(\pi_0)$, we have

(31)
$$P(F_n \otimes (p_{-1} + \sum_{i=1}^l u_i h_i^{\vee})) = \sum_{i=1}^l n(\partial_n u_i)(y) h_i^{\vee} \partial_x^{-1} \delta_{xy} + A_{ni} h_i^{\vee}.$$

Proof. We have $\{\int_{-\infty}^{\infty} H_n, u_i\} = n\partial_n(u_i)$, so that $P(H_n \otimes u_i) = n\partial_n u_i(y)\delta_{xy} + \partial_x A_{ni}$, with $A_{ni} \in \mathcal{R}^+_2(\pi_0)$. Now

$$P(F_n \otimes u_i) = \partial_x^{-1} P(H_n \otimes u_i) = n(\partial_n u_i(y)) \partial_x^{-1} \delta_{xy} + A_{ni},$$

and the statement follows.

Lemma 2.5. $\partial_n(\sum_{i=1}^l u_i(x)h_i^{\vee})$ is equal to the Cartan component of $-[p_{-1}, \operatorname{Ad}(n_+(x))(p_{-n})].$

Proof. We will give two proofs of this fact. Observe that $[p_{-1}, (\log n_+(x))_1] = -\sum_i u_i(x)h_i^{\vee}$; also

$$\partial_n((\log n_+)_1) = (\operatorname{Ad}(n_+)(p_{-n}))_1$$

for any $n_+ \in N_+$ (here log stands for the inverse of the exponential map $\exp: \mathfrak{n}_+ \to N_+$, which is an isomorphism, and the index 1 means the component of principal degree one of an element of $\tilde{\mathfrak{g}}$). Hence

$$\partial_n(\sum_i u_i(x)h_i^{\vee}) = -\partial_n([p_{-1}, (\log n_+(x))_1]) = -[p_{-1}, \partial_n(\log n_+(x))_1] = -[p_{-1}, (\operatorname{Ad}(n_+(x))(p_{-n}))_1].$$

Alternatively, this statement is a formulation of the Cartan part of the equation

$$[\partial + p_{-1} + \sum_{i} u_{i} h_{i}^{\vee}, \partial_{n} - (\mathrm{Ad}(n_{+})(p_{-n}))_{+}] = 0,$$

which follows from the zero-curvature equation (see e.g. formula (10) of [10]) and the fact that $\operatorname{Ad}(n_+)(p_{-n})$ commutes with $\partial + p_{-1} + \sum_i u_i h_i^{\vee}$ (see formula (13) of [10]).

Let us set $A_{ni} = \sum_{k \ge 0} A_{ni}^{(k)}(y) \partial_x^k \delta_{xy}$, with $A_{ni}^{(k)} \in \pi_0$.

Lemma 2.6. Let $\mathcal{Z} \in \mathcal{R}_2(\bar{\pi}_0)$ be such that

(32)
$$\partial_y \mathcal{Z} = \mathcal{Z}(p_{-1} + \sum_{i=1}^l u_i(y)h_i^{\vee}) + b_-(y) \Big(\sum_{i=1}^l n(\partial_n u_i)(y)h_i^{\vee}\partial_x^{-1}\delta_{xy} + A_{ni}h_i^{\vee}\Big).$$

Then

(33)

$$\begin{aligned} \mathcal{Z} &= n[b_{-}(y)(\mathrm{Ad}(n_{+}(y))(p_{-n}))_{-} - \mathrm{Ad}(b_{-}(x))(\mathrm{Ad}(n_{+}(x))(p_{-n}))_{-}b_{-}(y)]\partial_{x}^{-1}\delta_{xy} \\ &+ \mathrm{Ad}(b_{-}(x))\Big(\sum_{i=1}^{l}A_{ni}^{(0)}(x)h_{i}^{\vee}\Big)b_{-}(y)\partial_{y}^{-1}\delta_{xy} \\ &- \sum_{k\geq 1}\partial_{x}^{k}[\mathrm{Ad}(b_{-}(x))\Big(\sum_{i=1}^{l}A_{ni}^{(k+1)}(x)h_{i}^{\vee}\Big)b_{-}(y)\delta_{xy}]. \end{aligned}$$

Proof. Let

$$\mathcal{Z}_0 = n[b_-(y)(\mathrm{Ad}(n_+(y))(p_{-n})) - \mathrm{Ad}(b_-(x))(\mathrm{Ad}(n_+(x))(p_{-n})) - b_-(y)]\partial_x^{-1}\delta_{xy}.$$

Then

$$\begin{aligned} \partial_{y} \mathcal{Z}_{0} - \mathcal{Z}_{0}(p_{-1} + \sum_{i=1}^{l} u_{i}(y)h_{i}^{\vee}) &= n \Big\{ b_{-}(y)[p_{-1} + \sum_{i=1}^{l} u_{i}(y)h_{i}^{\vee}, (\operatorname{Ad}(n_{+}(y))(p_{-n}))_{-}] \\ &- b_{-}(y) \Big([p_{-1} + \sum_{i=1}^{l} u_{i}(y)h_{i}^{\vee}, \operatorname{Ad}(n_{+}(y))(p_{-n})] \Big)_{-} \Big\} \partial_{x}^{-1} \delta_{xy} \\ &= -nb_{-}(y)[p_{-1}, \Big(\operatorname{Ad}(n_{+}(y))(p_{-n}) \Big)_{1}] \partial_{x}^{-1} \delta_{xy} \\ &= nb_{-}(y) \partial_{n} (\sum_{i=1}^{l} u_{i}(y)h_{i}^{\vee}) \partial_{x}^{-1} \delta_{xy}, \end{aligned}$$

where the last equality follows from Lemma 2.5. We then have

$$\partial_y (\mathcal{Z} - \mathcal{Z}_0) - (\mathcal{Z} - \mathcal{Z}_0)(p_{-1} + \sum_{i=1}^l u_i(y))h_i^{\vee}) = b_-(y)\sum_{i=1}^l A_{ni}h_i^{\vee},$$

so that

$$\partial_y [(\mathcal{Z} - \mathcal{Z}_0)b_-(y)^{-1}] = \mathrm{Ad}(b_-(y))[\sum_{i=1}^l A_{ni}h_i^{\vee}],$$

and the result.

Proposition 2.6.

$$P(F_n \otimes b_-) = n[b_-(y)(\operatorname{Ad}(n_+(y))(p_{-n}))_- - \operatorname{Ad}(b_-(x))(\operatorname{Ad}(n_+(x))(p_{-n}))_- b_-(y)]\partial_x^{-1}\delta_{xy} + \operatorname{Ad}(b_-(x))\Big(\sum_{i=1}^l A_{ni}^{(0)}(x)h_i^{\vee}\Big)b_-(y)\partial_y^{-1}\delta_{xy} - \sum_{k\geq 0} \partial_x^k[\operatorname{Ad}(b_-(x))\Big(\sum_{i=1}^l A_{ni}^{(k+1)}(x)h_i^{\vee}\Big)b_-(y)\delta_{xy}]$$

and so belongs to $\mathcal{R}_2^{-1}(\tilde{\pi}_0)$.

Proof. Indeed, $P(F_n \otimes b_-)$ defined by formula (30) satisfies (32), by the Leibnitz rule and Lemma 2.4, and we apply to it Lemma 2.6.

Remark 2.5 It follows from Thm. 3.1 that the extension of ∂_n to $\bar{\pi}_0$, defined again as $r(p_{-n})$, is an infinitesimal automorphism of the nonlocal VPA structure of $\bar{\pi}_0$. \Box

3. Geometric interpretation of the Poisson structures.

3.1. Determination of the nonlocal terms. In 2.3, we have defined a nonlocal VPA $\bar{\pi}_0$. It is isomorphic as a differential algebra to $\mathbb{C}[B_- \times N_+]$ with the derivation ∂ defined by the right action of p_{-1} on $B_- \times N_+$. We will now describe P in these geometric terms.

Let us set $t = \sum_{\alpha} e^{\alpha} \otimes e_{\alpha} + e_{\alpha} \otimes e^{\alpha} + \sum_{i} h_{i} \otimes h_{i}^{\vee}$. Recall that p_{-n} is a basis of \mathfrak{a}_{-} dual to p_{n} with respect to the inner product \langle , \rangle .

Lemma 3.1.

(34)
$$P(b_{-} \otimes b_{-}) = -\ell^{\otimes 2}(t)(b_{-}(x) \otimes b_{-}(y))\partial_{x}^{-1}\delta_{xy}$$

Proof. We have

$$(pr_{-} \otimes pr_{-})(\mathrm{Ad}^{\otimes 2}b_{-}(x)R^{-} - \mathrm{Ad}^{\otimes 2}(b_{-}(y))(R^{+})) = (pr_{-} \otimes pr_{-})(\mathrm{Ad}^{\otimes 2}(b_{-}(x)))$$

$$\sum_{\alpha} e^{\alpha} \otimes e_{\alpha}) + (pr_{-} \otimes pr_{-})(\mathrm{Ad}^{\otimes 2}(b_{-}(y))\sum_{\alpha} e_{\alpha} \otimes e^{\alpha}) - \sum_{i} h_{i} \otimes h_{i}^{\vee}$$

$$= \sum_{\alpha} (\mathrm{Ad}(b_{-}(x))(e^{\alpha}))_{-} \otimes \mathrm{Ad}(b_{-}(x))(e_{\alpha}) + \sum_{\alpha} \mathrm{Ad}(b_{-}(y))(e_{\alpha}) \otimes (\mathrm{Ad}(b_{-}(y))(e^{\alpha}))_{-}$$

$$- \sum_{i} h_{i} \otimes h_{i}^{\vee}$$

$$= - \mathrm{Ad}(b_{-}(x))(\mathrm{Ad}(b_{-}(x)^{-1})(e^{\alpha}))_{-} \otimes e_{\alpha} - e_{\alpha} \otimes \mathrm{Ad}(b_{-}(y))(\mathrm{Ad}(b_{-}(y)^{-1})(e^{\alpha}))_{-}$$

$$- \sum_{i} h_{i} \otimes h_{i}^{\vee}.$$

The first equality is obtained using the following arguments: $R^- = \sum_{\alpha} e^{\alpha} \otimes e_{\alpha} - \frac{1}{2}t$, $R^+ = -\sum_{\alpha} e_{\alpha} \otimes e^{\alpha} + \frac{1}{2}t$; t is B_- -invariant, and $(pr_- \otimes pr_-)t = \sum_i h_i \otimes h_i^{\vee}$; the second equality is straightforward, and the the last one is because for $b_- \in B_-$, we have

(35)
$$\sum_{\alpha} (\operatorname{Ad}(b_{-})(e^{\alpha}))_{-} \otimes \operatorname{Ad}(b_{-})(e_{\alpha}) = -\sum_{\alpha} \operatorname{Ad}(b_{-})(\operatorname{Ad}(b_{-}^{-1})e^{\alpha})_{-} \otimes e_{\alpha}.$$

Formula (35) can be shown as follows: let $\xi \in \mathfrak{n}_+$, then

$$\langle \text{lhs of } (35), 1 \otimes \xi \rangle = (\text{Ad } b_{-}(\text{Ad } b_{-}^{-1}(\xi))_{+})_{-},$$

and

$$\langle \text{rhs of } (35), 1 \otimes \xi \rangle = - \operatorname{Ad} b_{-} (\operatorname{Ad} b_{-}^{-1}(\xi))_{-},$$

with $x_{+} = x - x_{-}$.

Formula (34) then follows from (28).

Lemma 3.2 ([10], **Prop. 6**). In any representation of G, $n_+(x)$ has the form

(36)
$$\bar{n}_{+}(x) \exp\left(\sum_{n \in I} \frac{1}{n} p_n F_n(x)\right),$$

where $\bar{n}_{+}(x)$ is a matrix of polynomials with entries in $\mathbb{C}[u_{i}^{(n)}]$.

Recall from [10], Lemma 1, that in any representation of N_+ we have the following formula for the right action of $x \in \mathfrak{g}$ on $N_+ \subset B_- \backslash G$:

$$r(x)n_{+} = (\mathrm{Ad}(n_{+})(x))_{+}n_{+}.$$

Lemma 3.3.

(37)
$$P(n_+ \otimes n_+) = r(a)[n_+(x) \otimes n_+(y)]\partial_x^{-1}\delta_{xy} + \text{ local terms},$$

where $a = \sum_{n \in I \cup (-I)} p_n \otimes p_{-n}$.

Proof. Let us write n_+ in the form (36) and compute the nonlocal part of $P(n_+ \otimes n_+)$. It comes from three different terms: $P(F_n \otimes \bar{n}_+), P(\bar{n}_+ \otimes F_n), P(F_n \otimes F_m)$. We have

 $P(F_n \otimes n_+) = n\partial_n n_+(y)\partial_x^{-1}\delta_{xy} + \rho_n(n_+),$

by (18). Set $F = \sum_{n \in I} \frac{1}{n} p_n F_n$. We then have $\bar{n}_+ = n_+ e^{-F}$. The Leibnitz rule gives

$$P(F_n \otimes \bar{n}_+) = n\partial_n n_+(y)e^{-F(y)}\partial_x^{-1}\delta_{xy} + \rho_n(n_+)(y)e^{-F(y)} + (1 \otimes \bar{n}_+(y))$$
$$[\sum_{m \in I} -\frac{1}{m}(1 \otimes p_m)P(F_n \otimes F_m)].$$

Using (17), we obtain

$$P(F_{n} \otimes \bar{n}_{+}) \in n\partial_{n}n_{+}(y)e^{-F(y)}\partial_{x}^{-1}\delta_{xy} + \rho_{n}(n_{+})e^{-F(y)} + (1 \otimes \bar{n}_{+}(y)) \cdot \left(\sum_{m \in I} -\frac{1}{m}(1 \otimes p_{m})(mH_{n,m}(x) + nH_{n,m}(y))\partial_{x}^{-1}\right)\delta_{xy} + \mathcal{R}_{2}^{+}(\pi_{0}),$$

so that

(38)

(39)

$$P(F_n \otimes \bar{n}_+) \in n[(\operatorname{Ad} \bar{n}_+(p_{-n}))_+ \bar{n}_+ - \bar{n}_+ \sum_{m \in I} \frac{1}{m} p_m H_{n,m}](y) \partial_x^{-1} \delta_{xy} + \rho_n(n_+) e^{-F(y)} - \sum_{m \in I} H_{n,m}(x) (\bar{n}_+(y)p_m) \partial_x^{-1} \delta_{xy} + \mathcal{R}_2^+(\pi_0).$$

Let us show that

$$\mathcal{W} = \rho_n(n_+)e^{-F(y)} - \sum_{m \in I} H_{n,m}(x)(\bar{n}_+(y)p_m)\partial_x^{-1}\delta_{xy}$$

belongs to $\mathcal{R}_2^+(\pi_0)$. Apply ∂_x to (38). The l.h.s. of the resulting identity belongs to $\mathcal{R}_2^+(\pi_0)$, as well as $\partial_x (n[(\operatorname{Ad} \bar{n}_+(p_{-n}))_+ \bar{n}_+ - \bar{n}_+ \sum_{m \in I} \frac{1}{m} p_m H_{n,m}](y) \partial_x^{-1}) \delta_{xy}$. It follows that

$$\partial_x \mathcal{W} \in \mathcal{R}_2 + (\pi_0).$$

On the other hand, in view of Prop. 2.3, we can write the nonlocal part of \mathcal{W} as $\sum_{m \in I} H_{n,m}(x) w_m(y) \partial_x^{-1} \delta_{xy}$. The result of the action of ∂_x on this nonlocal part should be local; since all $\partial H_{n,m}$, $m \in I$, are independent, it follows that all w_m 's vanish, so that

$$\mathcal{W} \in \mathcal{R}_2^+(\pi_0).$$

Formula (38) becomes

$$P(F_n \otimes \bar{n}_+) \in n\left((\operatorname{Ad} \bar{n}_+(p_{-n}))_+ \bar{n}_+ - \bar{n}_+ \sum_{m \in I} \frac{1}{m} p_m H_{n,m} \right) (y) \partial_x^{-1} \delta_{xy} + \mathcal{R}_2^+(\pi_0).$$

We derive from this a similar statement on $P(\bar{n}_+ \otimes F_n)$. Combining these with (17), and the fact that $P(\bar{n}_+ \otimes \bar{n}_+)$ is contained in $\mathcal{R}_2^+(\pi_0)$, we obtain (37).

Introduce the following notation. For any tensor $\gamma = \sum_i \gamma_i \otimes \gamma'_i$, and two operators a and b, we write $a(\gamma^{(1)}) \otimes b(\gamma^{(2)})$ for $\sum_i a(\gamma_i) \otimes b(\gamma'_i)$.

Lemma 3.4.

(40)
$$P(b_{-} \otimes n_{+}) = -b_{-}(x) [\operatorname{Ad}(b_{-}^{-1}(x))(t^{(1)})]_{-} \otimes [\operatorname{Ad}(b_{-}^{-1}(y))(t^{(2)})]_{+} n_{+}(y) \partial_{x}^{-1} \delta_{xy} + \sum_{n \in I} b_{-}(x) (\operatorname{Ad}(n_{+}(x))(p_{-n}))_{-} \otimes n_{+}(y) p_{n} \partial_{x}^{-1} \delta_{xy} + \text{ local terms;}$$

this equation should be understood in the tensor product of two representations of \mathfrak{g} .

The proof is given in Sect. 5.

3.2. Determination of VPA structures.

Theorem 3.1. The nonlocal VPA structure of $\bar{\pi}_0$ is expressed, via the identification of $\bar{\pi}_0$ with $\mathbb{C}[B_- \times N_+]$, by the formula

(41)
$$P(g \otimes g) = [-\ell^{\otimes 2}(t)(g(x) \otimes g(y)) + r^{\otimes 2}(a)(g(x) \otimes g(y))]\partial_x^{-1}\delta_{xy} + \sum_{n \ge 0} r^{\otimes 2} \Big((\operatorname{ad} p_{-1})^{-n-1} \otimes 1)(t-a) \Big) (g(x) \otimes g(y))\partial_x^n \delta_{xy}.$$

Proof. Let us denote by g(x) the pair $(b_-(x), n_+(x))$; it lies in the variety $B_- \times N_+$. This variety is endowed with left and right actions of \mathfrak{g} , that we denote ℓ and r. They are defined as follows. The mapping $B_- \times N_+ \to G$, associating to (b_-, n_+) the product b_-n_+ embeds $B_- \times N_+$ in G as a Schubert cell. The left and right actions of \mathfrak{g} on G can be restricted to $B_- \times N_+$; so that $\ell(x)$ is the vector field equal at (b_-, n_+) to the sum of $r((\operatorname{Ad} b_-(x))_-)$ on the first component of the product, and $\ell((\operatorname{Ad} b_-(x))_+)$ on the second one, according to

$$x \cdot b_{-}n_{+} = b_{-}[(\mathrm{Ad}(b_{-})(x))_{-} + (\mathrm{Ad}(b_{-})(x))_{+}]n_{+}.$$

Likewise, r(x) is the vector field equal at (b_-, n_+) to the sum of $r((\operatorname{Ad} n_+(x))_-)$ on the first component of the product, and $\ell((\operatorname{Ad} n_+(x))_+)$ on the second one, since

$$b_{-}n_{+} \cdot x = b_{-}[(\operatorname{Ad}(n_{+})(x))_{-} + (\operatorname{Ad}(n_{+})(x))_{+}]n_{+}.$$

Formulas (27) and (53) then imply

(42)

$$P(g \otimes g) = \left[-\ell^{\otimes 2}(t)(g(x) \otimes g(y)) + r^{\otimes 2}(a)(g(x) \otimes g(y))\right]\partial_x^{-1}\delta_{xy} + \text{ local terms.}$$

The action of ∂ on g coincides with $r(p_{-1})$, due to formulas (22) and (21), and we obtain

(43)
$$\partial g = r(p_{-1})g.$$

Therefore $P(g \otimes g)$ satisfies the differential equations

$$\partial_x P(g \otimes g) = (r(p_{-1}) \otimes 1) P(g \otimes g), \quad \partial_y P(g \otimes g) = (1 \otimes r(p_{-1})) P(g \otimes g).$$

We will use these equations to determine $P(g \otimes g)$ completely. Let us first determine the local part of the r.h.s. of (42). Denote it by \mathcal{Y} .

We have

(44)
$$\partial_x \mathcal{Y} - (r(p_{-1}) \otimes 1) \mathcal{Y} = r^{\otimes 2} (t-a) (g(x) \otimes g(x)) \delta_{xy},$$

(45)
$$\partial_y \mathcal{Y} - (1 \otimes r(p_{-1})) \mathcal{Y} = -r^{\otimes 2} (t-a) (g(x) \otimes g(x)) \delta_{xy}$$

(we have used the N_+ - and B_- -invariances of t to replace $\ell^{\otimes 2}(t)$ by $r^{\otimes 2}(t)$, and that $(g(x) \otimes g(y))\delta_{xy} = (g(x) \otimes g(x))\delta_{xy}$).

Recall that $\mathfrak{g} = \mathfrak{a} \oplus \operatorname{Im}(\operatorname{ad} p_{-1})$; this is an orthogonal decomposition in \mathfrak{g} ; ad p_{-1} is an automorphism of $\operatorname{Im}(\operatorname{ad} p_{-1})$, and t - a belongs to $(\operatorname{Im}(\operatorname{ad} p_{-1}))^{\otimes 2}$.

Then

(46)
$$\mathcal{Y}_0 = \sum_{n \ge 0} r^{\otimes 2} \Big((\operatorname{ad} p_{-1})^{-n-1} \otimes 1)(t-a) \Big) (g(x) \otimes g(y)) \partial_x^n \delta_{xy}$$

is a solution to (44); to see it, one should apply the Leibnitz rule, formula (43) and note the cancellation of all terms except the one in δ_{xy} . It is also a solution of (45) because $((\operatorname{ad} p_{-1})^n \otimes 1)(t-a) = (-1)^n (1 \otimes (\operatorname{ad} p_{-1})^n)(t-a)$. Indeed, this identity can be proved by writing $t - a = \sum_{\alpha} \hat{e}^{\alpha} \otimes \hat{e}_{\alpha}$, where \hat{e}^{α} , \hat{e}_{α} are dual bases of Im(ad $p_{-1})$, and using the anti-selfadjointness of ad p_{-1} .

Let P_0 be the operation defined by (42) and (46). It defines a nonlocal VPA structure on $(\mathbb{C}[B_- \times N_+], r(p_{-1}))$ by virtue of Prop. 1.1 (the elements of E_i in the second condition being $\ell^{\otimes 2}(t) - r^{\otimes 2}(t)$ for i = 0, and 0 for i > 0).

The theorem follows from the following lemma which is proved in Sect. 5.

Lemma 3.5. P_0 is equal to P.

Remark 3.1 The variety $B_- \times N_+$ can be considered as an open subset of G. One can show that formula (52) defines a nonlocal VPA structure on the whole group G.

It is easy to derive now a formula for the nonlocal VPA structure of $\mathbb{C}[N_+]$.

Corollary 3.1. The nonlocal VPA structure of π_0^+ is expressed, via the identification of π_0^+ with $\mathbb{C}[N_+]$, by the formula

(47)
$$P(n_{+} \otimes n_{+}) = r^{\otimes 2}(a)(n_{+}(x) \otimes n_{+}(y))\partial_{x}^{-1}\delta_{xy} + \sum_{n \geq 0} r^{\otimes 2} \Big((\operatorname{ad} p_{-1})^{-n-1} \otimes 1)(t-a) \Big) (n_{+}(x) \otimes n_{+}(y))\partial_{x}^{n}\delta_{xy}.$$

We now reformulate (47) so as to make it clear that the A_+ -invariant functions of N_+ only have local Poisson brackets expressed in terms of A_+ -invariant functions of N_+ . Let us denote

$$(t-a)_k = ((\operatorname{ad} p_{-1})^{-k-1} \otimes 1)(t-a).$$

Corollary 3.2.

(48)
$$P(n_{+} \otimes n_{+}) = r^{\otimes 2}(a)(n_{+}(x) \otimes n_{+}(y))\partial_{x}^{-1}\delta_{xy} + \sum_{k \ge 0} \left\{ \operatorname{Ad}(\bar{n}_{+}(x)) \left[(\partial_{x} - \sum_{n \in I} \frac{1}{n} H_{n}(x) \operatorname{ad} p_{n})^{k} ((t-a)_{k}^{(1)} \right] \right\}_{+} n_{+}(x) \\ \otimes \left(\operatorname{Ad}(\bar{n}_{+}(y))(t-a)_{k}^{(2)} \right)_{+} n_{+}(y)\delta_{xy}.$$

Proof. Let $F(x) = \sum_{n \in I} \frac{1}{n} p_n F_n(x)$, and $\alpha \in \tilde{\mathfrak{g}}$. Then

(49)
$$\operatorname{Ad}(e^{F(x)-F(y)})(\alpha \partial_x^n \delta_{xy}) = [\partial_x - \operatorname{ad}(F'(x))]^n(\alpha \delta_{xy}).$$

Indeed, this is obvious for n = 0. Assume that it is true for n, and apply ∂_x to the corresponding identity. We find

$$\operatorname{Ad}(e^{F(x)-F(y)})(\alpha \partial_x^{n+1} \delta_{xy}) + [F'(x), \operatorname{Ad}(e^{F(x)-F(y)})(\alpha \partial_x^n \delta_{xy})] = \partial_x \{ [\partial_x - \operatorname{ad}(F'(x))]^n (\alpha \delta_{xy}) \};$$

but the l.h.s. of this identity is expressed as

$$\operatorname{Ad}(e^{F(x)-F(y)})(\alpha\partial_x^{n+1}\delta_{xy}) + [F'(x), (\partial_x - \operatorname{ad}(F'(x)))^n(\alpha\delta_{xy})],$$

and we obtain (49) at step n + 1.

Now formula (48) follows directly from (47) and (49). In fact, the local part of formula (47) can be rewritten as

$$\sum_{k\geq 0} \left(\operatorname{Ad}(n_+(x)\otimes n_+(y))(t-a)_k \right)_+ n_+(x) \otimes n_+(y) \partial_x^k \delta_{xy}.$$

After substituting $n_+(x) = \bar{n}_+(x)e^{F(x)}$ in this formula we obtain

$$\sum_{k\geq 0} \left(\operatorname{Ad}(\bar{n}_+(x)\otimes \bar{n}_+(y)) \operatorname{Ad}(e^{F(x)}\otimes e^{F(y)})(t-a)_k \right)_+ n_+(x)\otimes n_+(y)\partial_x^k \delta_{xy},$$

which is equal to

$$\sum_{k\geq 0} \left(\operatorname{Ad}(\bar{n}_+(x)\otimes \bar{n}_+(y)) (\operatorname{Ad}(e^{F(x)-F(y)}\otimes 1)(t-a)_k \right)_+ n_+(x)\otimes n_+(y) \partial_x^k \delta_{xy} \right)$$

by \mathfrak{a}_+ -invariance of $(t-a)_k$. Applying formula (49) we obtain (48).

Now let $V((\lambda))$, $W((\mu))$ be $\tilde{\mathfrak{g}}$ -modules, and $v \in V$, $w \in W$ be such that $\mathfrak{a}_+v = 0$, $\mathfrak{a}_+w = 0$. The matrix coefficients of $n_+(x)v$, $n_+(y)w$ are composed of functions on

 N_+/A_+ , or, equivalently, of elements of π_0 . Moreover, all elements of $\mathbb{C}[N_+/A_+]$ can be obtained this way. Formula (48) implies that

(50)
$$P(n_{+}v \otimes n_{+}w) = \sum_{k \ge 0} \left\{ \operatorname{Ad}(\bar{n}_{+}(x)) \left[(\partial_{x} - \sum_{n \in I} \frac{1}{n} H_{n}(x) \operatorname{ad} p_{n})^{k} ((t-a)_{k}^{(1)}] \right]_{+} n_{+}(x) v \right.$$
$$\otimes \left(\operatorname{Ad}(\bar{n}_{+}(y))(t-a)_{k}^{(2)} \right)_{+} n_{+}(y) w \delta_{xy};$$

since $\bar{n}_+(x)$, $\bar{n}_+(y)$ are matrices whose entries are elements of π_0 , (50) shows at the same time that the Poisson brackets of the entries of $n_+(x)v$ and $n_+(y)w$ are local and expressed in terms of functions of N_+/A_+ .

In the theory of the mKdV equations, an important role is played by the embedding of N_+/A_+ in $\tilde{\mathfrak{g}}$ as an N_+ -coadjoint orbit of an element of \mathfrak{a} . This motivates us to compute the Poisson brackets of the matrices $\operatorname{Ad}(n_+)(p_n)$. The result is a direct consequence of Cor. 3.1.

Corollary 3.3. For $n, m \in \pm I$,

$$P(\operatorname{Ad}(n_{+})(p_{n}) \otimes \operatorname{Ad}(n_{+})(p_{m})) =$$

$$\sum_{k \geq 0} \left[\left\{ \operatorname{Ad}(\bar{n}_{+}(x)) \left[(\partial_{x} - \sum_{n \in I} \frac{1}{n} H_{n}(x) \operatorname{ad} p_{n})^{k} ((t-a)_{k}^{(1)}) \right] \right\}_{+}, \operatorname{Ad}(n_{+}(x))(p_{n}) \right]$$

$$\otimes \left[\left(\operatorname{Ad}(\bar{n}_{+}(y))(t-a)_{k}^{(2)} \right)_{+}, \operatorname{Ad}(n_{+}(y))(p_{m}) \right] \delta_{xy}.$$

Here again, the locality of these brackets and the fact they are expressed in terms of A_+ -invariant functions on N_+ is manifest.

Remark 3.2 Formulas (47) and (50) give a geometric interpretation of the nonlocal VPA structures of π_0^+ and π_0 respectively.

We now give a formula for $P(u_i \otimes n_+)$. Due to the presence of nonlocal quantities F_n in n_+ , this bracket contains nonlocal as well as local terms.

Proposition 3.1.

$$P(u_i \otimes n_+) = n_+(y)\partial_x \Big(-(n_+(x)^{-1}h_i n_+(x))_{\mathfrak{a}} \partial_y^{-1} \delta_{xy} \\ +\sum_{n\geq 0} (-\operatorname{ad} \ p_{-1})^{-n-1} ((n_+(x)^{-1}h_i n_+(x))_{\operatorname{Im}(\operatorname{ad} p_{-1})})\partial_x^n \delta_{xy} \Big),$$

where the indices \mathfrak{a} and $\operatorname{Im}(\operatorname{ad} p_{-1})$ stand for the projections on the components of $\mathfrak{g} = \mathfrak{a} \oplus \operatorname{Im}(\operatorname{ad} p_{-1})$.

Proof. Consider again the variable $\varphi_i = \partial^{-1} u_i$. It is easier to compute $\mathcal{G} = P(\varphi_i \otimes n_+)$ first. We have

$$\partial_y \mathcal{G} = -(p_{-1} + \sum_i u_i(y)h_i^{\vee})\mathcal{G} + \mathcal{G}p_{-1} - h_i n_+(x)\partial_{xy}.$$

Let $\mathcal{H} = n_+(y)^{-1}\mathcal{G}$, then

$$\partial_y \mathcal{H} = [\mathcal{H}, p_{-1}] - n_+(x)^{-1} h_i n_+(x) \partial_{xy}.$$

A solution to this equation is

$$\mathcal{H}_0 = -(n_+(x)^{-1}h_in_+(x))_{\mathfrak{a}}\partial_y^{-1}\delta_{xy} + \sum_{n\geq 0}(-\operatorname{ad} p_{-1})^{-n-1}((n_+(x)^{-1}h_in_+(x))_{\operatorname{Im}(\operatorname{ad} p_{-1})})\partial_x^n\delta_{xy}.$$

On the other hand, the nonlocal part of \mathcal{H} coincides with that of \mathcal{H}_0 because it is equal to the nonlocal part of $\sum_{n \in I} \frac{1}{n} p_n P(\varphi_i \otimes F_n)$; but $P(\varphi_i \otimes F_n) = -n \partial_n \varphi_i(x) + \text{local terms}$, since $P(\varphi_i \otimes H_n) \in -n \partial_n \varphi_i(x) + \partial_y(\mathcal{R}_2(\pi_0))$; so this nonlocal part is expressed as $-\sum_{i \in I} \partial_n \varphi_i(x) p_n \partial_y^{-1} \delta_{xy}$. To establish the coincidence of the nonlocal parts of \mathcal{H} and \mathcal{H}_0 , it remains to check the identity

(51)
$$\partial_n \varphi_i(x) = \langle n_+^{-1}(x) h_i n_+(x), p_{-n} \rangle$$

which amounts to

$$\partial_n u_i(x) = -\langle h_i, [p_{-1} + \sum_i u_i(x)h_i^{\vee}, n_+(x)p_{-n}n_+(x)^{-1}] \rangle;$$

since by Lemma 2.5, $\partial_n(\sum_i u_i(x)h_i^{\vee})$ coincides with the Cartan part of

$$-[p_{-1}, n_+(x)p_{-n}n_+(x)^{-1}],$$

this is verified. So (51) holds, and the nonlocal parts of \mathcal{H} and \mathcal{H}_0 coincide.

So $\mathcal{H}_1 = \mathcal{H} - \mathcal{H}_0$ has no nonlocal terms, and satisfies $\partial_y \mathcal{H}_1 = [\mathcal{H}_1, p_{-1}]$; the arguments used to establish (62) then show that $\mathcal{H}_1 = 0$.

Remark 3.3 As we noted in the introduction, it is possible to derive (37) using Prop. 3.1 and a differential equation in the first variable, satisfied by $P(n_+ \otimes n_+)$. But we find the present use of B_- more natural.

3.3. Gelfand-Dickey-Dorfman structure. According to Sect. 1.5, the VPA $\pi_0 = \mathbb{C}[N_+/A_+]$ is endowed with a Gelfand-Dickey-Dorfman structure. We give here a geometric interpretation of it.

Proposition 3.2. Let $V((\lambda))$, $W((\mu))$ be $\tilde{\mathfrak{g}}$ -modules, and $v \in V$, $w \in W$ be such that $\mathfrak{a}_+v = 0$, $\mathfrak{a}_+w = 0$. We have:

$$\mathcal{V}_{n+v\otimes 1}(1\otimes n+w) = \sum_{k\geq 0} (-\partial)^k [(\mathrm{Ad}(n_+)(t-a)_k^{(1)}) + n+v] \otimes (\mathrm{Ad}(n_+)(t-a)_k^{(2)}) + n+w$$

Proof. We apply the definition (9) of \mathcal{V}_f to formula (47).

3.4. Compatible nonlocal VPA structures. Let us show how the nonlocal VPA structure on $\mathbb{C}[B_- \times N_+]$, defined in Thm. 3.1, can be embedded into an infinite family of compatible nonlocal VPA structures (we call such a family compatible, if any linear combination of these structures is again a nonlocal VPA structure).

Let us identify \mathfrak{g} with the subalgebra of the loop algebra $\overline{\mathfrak{g}} \otimes \mathbb{C}((\lambda))$ of a finite dimensional semisimple Lie algebra $\overline{\mathfrak{g}}$, consisting of the elements $x(\lambda)$ satisfying

$$x(\zeta\lambda) = x(\lambda)^{\sigma}$$

 σ an automorphism of $\bar{\mathfrak{g}}$ and ζ a root of unity of the same order r. There is an action of $\mathbb{C}((\lambda^r))$ on \mathfrak{g} . Let us denote by the same letter elements of $\mathbb{C}((\lambda^r))$ and the corresponding operators on \mathfrak{g} .

For $n \in \mathbb{Z}$, set

$$t_n = (\lambda^{rn} \otimes 1)t, \quad a_n = (\lambda^{rn} \otimes 1)a$$

Note that ad p_{-1} commutes with the action of λ^{rn} , so that

$$t_n - a_n \in \operatorname{Im}(\operatorname{ad} p_{-1})^{\otimes 2}$$

Let us define an operation P_n on $\mathbb{C}[B_- \times N_+]$, by the formula

(52)
$$P_n(g \otimes g) = \left[-\ell^{\otimes 2}(t_n)(g(x) \otimes g(y)) + r^{\otimes 2}(a_n)(g(x) \otimes g(y))\right]\partial_x^{-1}\delta_{xy} + \sum_{n \ge 0} r^{\otimes 2} \left((\operatorname{ad} p_{-1})^{-n-1} \otimes 1)(t_n - a_n)\right)(g(x) \otimes g(y))\partial_x^n\delta_{xy}.$$

Proposition 3.3. The formulae (52) define compatible nonlocal VPA structures on $\mathbb{C}[B_- \times N_+]$ (endowed with the derivation $r(p_{-1})$).

Proof. A combination of the brackets (52) corresponds to the same formula, with t_n and a_n replaced by t_f and a_f respectively, with $t_f = (f \otimes 1)t$ and $a_f = (f \otimes 1)a$, f a certain element of $\mathbb{C}[\lambda^r, \lambda^{-r}]$. The resulting bracket satisfies the conditions of Prop. 1.1; the elements of E_i in the second condition are $\ell^{\otimes 2}(t_f) - r^{\otimes 2}(t_f)$ for i = 0, and 0 for i > 0.

Remark 3.4 The variety $B_- \times N_+$ can be considered as an open subset of G. It has a compatible family of nonlocal VPA structures defined by (52). In the same way as in the proof of Prop. 3.3, one can show that formula (52) defines a compatible family of nonlocal VPA structures on the whole group G.

Remark 3.5 The extension to G of the nonlocal VPA structure defined in Thm. 3.1 is clearly left G-invariant. It follows that left G-translations provide symmetries of the mKdV hierarchy, respecting the Poisson structure. Infinitesimal left translations by elements of \mathfrak{n}_+ correspond to the Toda flows; left translations by elements of \mathfrak{b}_- do not change the variables u_i . A class of translations that would be interesting to study further are left translations by elements of the affine Weyl group; they should mix local and nonlocal variables while respecting the Poisson structure. A. Orlov pointed out to us that they probably coincide with the Darboux transformations. \Box

4. The proof of Lemma 3.4 and Lemma 3.5

4.1. Proof of Lemma 3.4. Equation (40) is rewritten as

(53)
$$P(b_{-} \otimes n_{+}) = -\sum_{\beta} \bar{e}_{\beta} b_{-}(x) \otimes (\operatorname{Ad}(b_{-}(y)^{-1})(\bar{e}^{\beta}))_{+} n_{+}(y) \partial_{x}^{-1} \delta_{xy} \\ + \sum_{n \in I} b_{-}(x) (\operatorname{Ad}(n_{+}(x))(p_{-n}))_{-} \otimes n_{+}(y) p_{n} \partial_{x}^{-1} \delta_{xy} + \text{ local terms},$$

with $\sum \bar{e}^{\beta} \otimes \bar{e}_{\beta} = \sum e^{\alpha} \otimes e_{\alpha} + \sum_{i} h_{i} \otimes h_{i}^{\vee}$. The differential equations satisfied by the left and right hand sides of (53) are

$$\begin{aligned} \partial_y(\text{lhs of } (53)) &= (\text{lhs of } (53))(1 \otimes p_{-1}) - 1 \otimes (p_{-1} + \sum_i u_i(y)h_i^{\vee}) \\ (\text{lhs of } (53)) &- \sum_i \text{Ad}(b_-(y))([p_{-1}, h_i])b_-(x) \otimes h_i^{\vee} n_+(y)\partial_x^{-1}\delta_{xy} \\ &+ \text{ local terms,} \end{aligned}$$

using (21) and (23), and

$$\partial_{y}(\text{rhs of } (53)) = (\text{rhs of } (53))(1 \otimes p_{-1}) - 1 \otimes (p_{-1} + \sum_{i} u_{i}(y)h_{i}^{\vee}) \\ (\text{rhs of } (53)) - \sum_{\beta} \bar{e}_{\beta}b_{-}(x) \otimes \{ [p_{-1} + \sum_{i} h_{i}^{\vee}u_{i}(y), (\operatorname{Ad}(b_{-}^{-1}(y))(\bar{e}^{\beta}))_{+}] \\ -([p_{-1} + \sum_{i} h_{i}^{\vee}u_{i}(y), \operatorname{Ad}(b_{-}^{-1}(y))(\bar{e}^{\beta})])_{+} \} \\ n_{+}(y)\partial_{x}^{-1}\delta_{xy} + \text{ local terms},$$

using (21) and (22). Since

(54)
$$\sum_{i} \operatorname{Ad}(b_{-}(y))([p_{-1}, h_{i}]) \otimes h_{i}^{\vee} = \sum_{\beta} \bar{e}_{\beta} \otimes [p_{-1}, (\operatorname{Ad}(b_{-}(y)^{-1})(\bar{e}^{\beta}))_{1}],$$

these two equations coincide. In (54), we denote by x_1 the part of $x \in \mathfrak{g}$, of principal degree 1. (54) is proved by pairing its right and left hand sides with $1 \otimes h_i$, $i = 1, \ldots, l$. It follows that the difference of the two sides of (53) satisfies

(55)
$$\frac{\partial_y(\text{lhs of } (53) - \text{rhs of } (53))}{-(1 \otimes (p_{-1} + \sum_i u_i(y)h_i^{\vee}))(\text{lhs of } (53) - \text{rhs of } (53)) + \text{local terms.} }$$

On the other hand, we have

$$\partial_x(\text{lhs of } (53)) = (\text{lhs of } (53))((p_{-1} + \sum_i u_i(x)h_i^{\vee}) \otimes 1) \\ -\sum_{i,k} [b_-(x)h_i^{\vee} \partial_k u_i(x) \otimes n_+(y)p_k] \partial_x^{-1} \delta_{xy} + \text{ local terms,}$$

and

$$\partial_{x}(\text{rhs of } (53)) = (\text{rhs of } (53))((p_{-1} + \sum_{i} u_{i}(x)h_{i}^{\vee}) \otimes 1) \\ + \sum_{n \in I} b_{-}(x)\{[p_{-1} + \sum_{i} h_{i}^{\vee} u_{i}(x), (\operatorname{Ad}(n_{+}(x))(p_{-n}))_{-}] \\ -([p_{-1} + \sum_{i} h_{i}^{\vee} u_{i}(x), \operatorname{Ad}(n_{+}(x))(p_{-n})])_{-} \\ \otimes n_{+}(y)p_{n}\partial_{x}^{-1}\delta_{xy}\} + \text{ local terms.}$$

We have the equality

$$[p_{-1}, (\mathrm{Ad}(n_+(x))(p_{-n}))_1] = \sum_i h_i^{\vee} \partial_n u_i(x),$$

because of Lemma 2.5.

Therefore the right hand sides of the last two formulas coincide up to local terms, and

(56)
$$\partial_x (\text{lhs of } (53) - \text{rhs of } (53)) = (\text{lhs of } (53) - \text{rhs of } (53)) \\ (1 \otimes (p_{-1} + \sum_i u_i(x)h_i^{\vee})) + \text{local terms}.$$

Let $\mathcal{X} = (1 \otimes n_+(y)^{-1})$ (lhs of (53) – rhs of (53)) $(1 \otimes b_-(x)^{-1})$, then

 $\partial_x \mathcal{X} = \text{local terms}$

by (56), and

 $\partial_y \mathcal{X} = [\mathcal{X}, 1 \otimes p_{-1}] + \text{local terms}$

by (21) and (55). The first equation gives

$$\partial_x \mathcal{X} = \sum_{n \ge 0} \mathcal{X}_n(y) \partial_x^n \delta_{xy},$$

 \mathbf{SO}

 $\mathcal{X} = \mathcal{X}_0(y)\partial_x^{-1}\delta_{xy} + \text{local terms},$

and the second equation gives us

$$\partial_y \mathcal{X}_0(y) = [\mathcal{X}_0(y), 1 \otimes p_{-1}].$$

Let ξ be any element of the dual to \mathfrak{b}_{-} , and $\mathcal{X}_{\xi}(y) = (\xi \otimes 1)(\mathcal{X}_{0}(y))$. Then $\mathcal{X}_{\xi}(y)$ has values in \mathfrak{n}_{+} and satisfies

(57)
$$\partial_y \mathcal{X}_{\xi}(y) = [\mathcal{X}_{\xi}(y), p_{-1}];$$

we then follow the proof of [10], lemma 3, to conclude that $\mathcal{X}_{\xi}(y)$ is constant and lies in \mathfrak{a}_+ . Recall how this can be done: decompose $\mathcal{X}_{\xi}(y)$ in its homogeneous principal components $\sum_i \mathcal{X}_{\xi,i}(y)$, and each component along the decomposition $\operatorname{Im}(\operatorname{ad} p_{-1}) \oplus \mathfrak{a}$, as $\mathcal{X}_{\xi,i}^1(y) + \mathcal{X}_{\xi,i}^2(y)$; let *i* be the smallest index, such that $\mathcal{X}_{\xi,i}^1(y)$ is not zero; the equation implies that $\partial_y \mathcal{X}_{\xi}(y)$ has a nonzero component of degree i - 1 in Im ad p_{-1} , hence a contradiction. So $\mathcal{X}_{\xi}(y)$ lies in \mathfrak{a} ; (57) then implies that it is constant. We finally obtain:

lhs of (53) – rhs of (53) =
$$\sum_{n \in I} x_n b_-(x) \otimes n_+(y) p_n \partial_x^{-1} \delta_{xy} + \text{ local terms},$$

with $x_n \in \mathfrak{b}_-$. But there is only one possibility, $x_n = 0$, which is compatible with the following invariance property of P.

Recall that for $\xi \in \mathfrak{b}_{-}$, $\ell(\xi)$ is the derivation of the algebra $\overline{\pi}_{0}$ defined by the action of the left translation by ξ , on $B_{-} \times N_{+}$. Since $\ell(\xi)$ commutes with ∂ , it induces an endomorphism (also denoted by $\ell(\xi)$) of $\mathcal{R}_{2}(\overline{\pi}_{0})$, according to the rules used in 2.1 in the case of ∂_{n} . We then have:

Lemma 4.1. For $a, b \in \overline{\pi}_0, \xi \in \mathfrak{b}_-$,

(58)
$$P(\ell(\xi)a \otimes b) + P(a \otimes \ell(\xi)b) = \ell(\xi)P(a \otimes b).$$

Proof. For the brackets $P(b_{-} \otimes b_{-})$, this follows from (27) and the invariance of t. We also have

$$P(\ell(\xi)b_{-} \otimes u_{i}) = \partial_{y}([\xi, \operatorname{Ad}(b_{-}(y))(h_{i})]b_{-}(x)\partial_{x}^{-1}\delta_{xy} + \operatorname{Ad}(b_{-}(y))(h_{i})\xi b_{-}(x)$$
$$\partial_{x}^{-1}\delta_{xy}) = \ell(\xi)P(b_{-} \otimes u_{i})$$

so that

$$P(H_n \otimes \ell(\xi)b_-) = \ell(\xi)P(H_n \otimes b_-)$$

(in this equality, the second $\ell(\xi)$ is $\ell(\xi) \otimes 1$ acting on $\operatorname{End}(V)((\xi)) \otimes \mathcal{R}_2(\bar{\pi}_0)$) and $(\ell(\xi) \otimes 1)r_n = (1 \otimes \ell(\xi))r_n$ (equality in $\operatorname{End}(V)((\xi)) \otimes \mathcal{R}_2(\bar{\pi}_0)$; so that

$$P(F_n \otimes \ell(\xi)b_-) = \ell(\xi)P(F_n \otimes b_-).$$

Finally, the elements of π_0^+ are invariant under $\ell(\xi)$, so the identity is trivially satisfied for their Poisson brackets.

Now (53) follows.

4.2. Proof of Lemma 3.5. The operation P_0 is defined by the identities

$$P_0(b_- \otimes b_-) = -\ell^{\otimes 2}(t)(b_-(x) \otimes b_-(y))\partial_x^{-1}\delta_{xy}$$

(59)
$$P_{0}(b_{-} \otimes n_{+}) = [-b_{-}(b_{-}^{-1}t^{(1)}b_{-})_{-}(x) \otimes (b_{-}^{-1}t^{(2)}b_{-})_{+}(y)n_{+}(y) + b_{-}(n_{+}a^{(1)}n_{+}^{-1})_{-}(x) \otimes (n_{+}a^{(2)}n_{+}^{-1})_{+}n_{+}(y)]\partial_{x}^{-1}\delta_{xy} + \sum_{n\geq0}[b_{-}(n_{+}(t-a)_{n}^{(1)}n_{+}^{-1})_{-}(x) \otimes (n_{+}(t-a)_{n}^{(2)}n_{+}^{-1})_{+}n_{+}(y)]\partial^{n}\delta_{xy},$$

where we denote $((ad p_{-1})^{-n-1} \otimes 1)(t-a)$ by $(t-a)_n$, any element $\alpha \in \mathfrak{g} \otimes \mathfrak{g}$ is decomposed as $\sum \alpha^{(1)} \otimes \alpha^{(2)}$, and

(60)
$$P_{0}(n_{+} \otimes n_{+}) = r^{\otimes 2}(a)(n_{+}(x) \otimes n_{+}(y))\partial_{x}^{-1}\delta_{xy} + \sum_{n \geq 0} r^{\otimes 2} \Big((\operatorname{ad} p_{-1})^{-n-1} \otimes 1)(t-a) \Big) (n_{+}(x) \otimes n_{+}(y))\partial_{x}^{n}\delta_{xy}.$$

By construction, $P_0(g \otimes g)$ satisfies the identities

(61)

$$\partial_x P_0(g \otimes g) = (r(p_{-1}) \otimes 1) P_0(g \otimes g), \quad \partial_y P_0(g \otimes g) = (1 \otimes r(p_{-1})) P_0(g \otimes g).$$

Clearly, $P_0(b_- \otimes b_-)$ coincides with $P(b_- \otimes b_-)$. $\mathcal{B} = P(b_- \otimes n_+)$ satisfies the equation

$$\partial_{y}\mathcal{B} + (1 \otimes (p_{-1} + \sum_{i} u_{i}h_{i}^{\vee}))\mathcal{B} - \mathcal{B}(1 \otimes p_{-1}) = \sum_{i} (\partial_{y} \operatorname{Ad}(b_{-}(y))(h_{i})b_{-}(x)\partial_{x}^{-1} \otimes h_{i}^{\vee}(h_{+}(y)) \partial_{x}y)$$

by virtue of (23) and (21). Let us determine an equation satisfied by $\mathcal{B}_0 = P_0(b_- \otimes n_+)$. Let $\mathcal{E} = P_0(b_- \otimes b_-)$. \mathcal{E} satisfies the equation

$$\partial_y \mathcal{E} = \mathcal{E}(1 \otimes (p_{-1} + \sum_i u_i h_i^{\vee})) - \partial_y [\operatorname{Ad}(b_-(y))(h_i)b_-(x)\partial_x^{-1} \otimes b_-(y)]\delta_{xy}.$$

Note that due to (61), $P_0(b_- \otimes g)$ satisfies

 $\partial_y P_0(b_- \otimes g) = (1 \otimes r(p_{-1})) P_0(b_- \otimes g).$

This implies, writing \mathcal{B}_0 as $P_0(b_- \otimes b_-^{-1}g) [B_- \times N_+$ has a left B_- -action, defined as the product of the left action of B_- on itself and of the trivial one, that we use here], that \mathcal{B}_0 satisfies

$$\partial_{y}\mathcal{B}_{0} + (1 \otimes (p_{-1} + \sum_{i} u_{i}h_{i}^{\vee}))\mathcal{B}_{0} - \mathcal{B}_{0}(1 \otimes p_{-1}) = \sum_{i} (\partial_{y} \operatorname{Ad}(b_{-}(y))(h_{i})b_{-}(x)\partial_{x}^{-1} \otimes h_{i}^{\vee}(h_{+}(y)) \partial_{x}y).$$

Let us set $\mathcal{B}_1 = (1 \otimes n_+^{-1}(y))(\mathcal{B} - \mathcal{B}_0)$, we obtain

$$\partial_y \mathcal{B}_1 + [1 \otimes p_{-1}, \mathcal{B}_1] = 0$$

The nonlocal parts of \mathcal{B} and \mathcal{B}_0 coincide, so that \mathcal{B}_1 contains only local terms; write $\mathcal{B}_1 = \sum_{n\geq 0} \mathcal{B}_1^{(n)}(x) \partial_x^n \delta_{xy}$ (each $\mathcal{B}_1^{(n)}$ belongs to the tensor product of the tangent space $T_{b_-(x)}B_-$ to B_- at $b_-(x)$ with \mathfrak{n}_+), we then get

$$[1 \otimes p_{-1}, \mathcal{B}_1^{(0)}(x)] = 0, \quad \mathcal{B}_1^{(n)} = [1 \otimes p_{-1}, \mathcal{B}_1^{(n+1)}(x)]$$

for $n \geq 0$, so $\mathcal{B}_1^{(0)}(x)$ belongs to $T_{b_-(x)}B_- \otimes \mathfrak{a}$ by the first equation and to $T_{b_-(x)}B_- \otimes \mathfrak{a}$ Im(ad p_{-1}) by the second one (specialized to n = 0), so that it is zero; repeating this argument for $\mathcal{B}_1^{(1)}$, we find it to vanish as well, etc. So $\mathcal{B}_1 = 0$ and

(62)
$$P(b_- \otimes n_+) = P_0(b_- \otimes n_+).$$

Now, $\mathcal{B} = P(b_- \otimes n_+)$ satisfies the equation

(63)
$$\partial_x \mathcal{B} = \mathcal{B}((p_{-1} + \sum_i u_i(x)h_i^{\vee}) \otimes 1) + \sum_i b_-(x) \otimes (h_i^{\vee} P(u_i \otimes n_+)).$$

On the other hand, let $C = P(n_+ \otimes n_+)$ and $C_0 = P_0(n_+ \otimes n_+)$. C satisfies the equation

$$\partial_x \mathcal{C} = -((p_{-1} + \sum_i u_i(x)h_i^{\vee}) \otimes 1)\mathcal{C} + \mathcal{C}(p_{-1} \otimes 1) - \sum_i h_i^{\vee} n_+(x) \otimes P(u_i \otimes n_+).$$

Let us determine an equation satisfied by \mathcal{C}_0 . Due to (61), $P_0(g \otimes n_+)$ satisfies

$$\partial_x P_0(g \otimes n_+) = P_0(g \otimes n_+)(p_{-1} \otimes 1),$$

and writing $P_0(n_+ \otimes n_+)$ as $P_0(b_-^{-1}g \otimes n_+)$ (using the same left B_- -action as above) and using (63), we get

$$\partial_x \mathcal{C}_0 = -((p_{-1} + \sum_i u_i(x)h_i^{\vee}) \otimes 1)\mathcal{C}_0 + \mathcal{C}_0(p_{-1} \otimes 1) - \sum_i h_i^{\vee} n_+(x) \otimes P(u_i \otimes n_+).$$

 \mathcal{C} and \mathcal{C}_0 satisfy the same equation, so that $\mathcal{C}_1 = (n_+(x)^{-1} \otimes 1)(\mathcal{C} - \mathcal{C}_0)$ (which belongs to the tensor product of \mathfrak{n}_+ with the tangent space to N_+ at $n_+(y)$) satisfies

$$\partial_x \mathcal{C}_1 = [\mathcal{C}_1, p_{-1} \otimes 1].$$

Since the nonlocal parts of P_0 and P coincide, C_1 contains no nonlocal terms. We can use the same arguments as in the case of \mathcal{B}_1 , to conclude that $C_1 = 0$ and

(64)
$$P(n_+ \otimes n_+) = P_0(n_+ \otimes n_+).$$

Lemma 3.5 now follows from (62), (64).

References

- M. Adler, P. van Moerbeke, Compatible Poisson structures and the Virasoro algebra, Commun. Pure and Appl. Math. 47 (1994), 5-37.
- 2. A.V. Antonov, A.A. Belov, B.L. Feigin, Geometric description of the local integrals of motion of the Maxwell-Bloch equation, Mod. Phys. Lett. A 10 (1995), 1209-24.
- O. Babelon, L. Bonora, F. Toppan, Exchange algebra and the Drinfeld-Sokolov theorem, Commun. Math. Phys. 140 (1991), 93-117.
- V.V. Bazhanov, S.L. Lukyanov, A.B. Zamolodchikov, Integrable structure of conformal field theory, quantum KdV theory and thermodynamic Bethe Ansatz, Comm. Math. Phys. 177 (1996) 381-98; hep-th/9412229.
- 5. A.A. Beilinson, V.G. Drinfeld, Chiral algebras, preprint.
- R. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, Proc. Natl. Acad. Sci. USA 83 (1986), 3068-71.
- V.G. Drinfeld, V.V. Sokolov, Lie algebras and equations of Korteweg-de Vries type, Sov. Math. Dokl. 23 (1981) 457-62; J. Sov. Math. 30 (1985) 1975-2035.
- B. Enriquez, Nilpotent action on KdV variables and 2-dimensional Drinfeld-Sokolov reduction, Theor. Math. Phys. 98:3 (1994), 256-8, 375-8 (Russian edition).
- B. Enriquez, B. Feigin, Integrals of motion of the classical lattice sine-Gordon system, Theor. Math. Phys. 103:3 (1995), 738-56, 507-28 (Russian edition); hep-th/9409075.
- 10. B. Enriquez, E. Frenkel, Equivalence of two approaches to the mKdV hierarchies, preprint q-alg/9606004.
- B. Enriquez, A. Orlov, V. Rubtsov, Higher Hamiltonian structures on the KdV phase space (the sl₂ case), JETP Lett., 58:8 (1993), 658-64, 677-83 (Russian edition); hep-th/9309038.
- L.D. Faddeev, L.A. Takhtajan, Hamiltonian Methods in Soliton Equations, Springer-Verlag, 1992.
- B. Feigin, E. Frenkel, Integrals of motions and quantum groups, Lect. Notes in Math. 1620 (1995), 349-418, Springer-Verlag; hep-th/9310022.
- B. Feigin, E. Frenkel, Kac-Moody groups and integrability of soliton equations, Invent. Math. 120 (1995), 379-408; hep-th/9311171.
- 15. B. Feigin, E. Frenkel, Non-linear Schrödinger equations and Wakimoto modules, to appear.
- 16. M.A.I. Flohr, On fusion rules in logarithmic conformal field theories, Preprint hep-th/9605151.
- I. Frenkel, J. Lepowsky, A. Meurman, Vertex Operator Algebras and the Monster, Academic Press, 1988.
- I.M. Gelfand, I.Ya. Dorfman, Hamiltonian operators and algebraic structures related to them, Funct. Anal. Appl. 13:4 (1979), 13-30.
- 19. V. Gurarie, Logarithmic operators in conformal field theory, Nucl. Phys. B 410 (1993), 535-49.
- 20. V.G. Kac, Infinite-dimensional Lie Algebras, 3rd edition, Cambridge Univ. Press, 1990.
- D.R. Lebedev, A.O. Radul, The generalized internal long waves equations: construction, hamiltonian structure and conservation laws, Comm. Math. Phys. 91 (1983), 543-55.
- A.O. Radul, A description of Poisson brackets on the space of nonlocal functionals, Funct. Anal. Appl. 19:2 (1985), 153-6, 85-7 (Russian edition).

32 BENJAMIN ENRIQUEZ AND EDWARD FRENKEL

23. V.E. Zakharov, A.B. Shabat, Integration of nonlinear equations of mathematical physics by the method of inverse scattering II, Funct. Anal. Appl. 13:3 (1979), 166-74, 13-22 (Russian edition).

Centre de Mathématiques, URA 169 du CNRS, Ecole Polytechnique, 91 128 Palaiseau, France

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA 02138, USA