### NONLINEAR DEFORMED su(2) ALGEBRAS INVOLVING TWO DEFORMING FUNCTIONS \*

D. BONATSOS, P. KOLOKOTRONIS

Institute of Nuclear Physics, NCSR Demokritos, GR-15310 Aghia Paraskevi, Attiki, Greece

C. DASKALOYANNIS

Department of Physics, Aristotle University of Thessaloniki, GR-54006 Thessaloniki, Greece

#### A. LUDU

Department of Theoretical Physics, Faculty of Physics, University of Bucharest, Bucharest-Magurele, P.O. Box MG-5211, Romania

#### C. QUESNE<sup>†</sup>

Physique Nucléaire Théorique et Physique Mathématique, Université Libre de Bruxelles, Campus de la Plaine CP229, Boulevard du Triomphe, B-1050 Brussels, Belqium

#### Abstract

The most common nonlinear deformations of the su(2) Lie algebra, introduced by Polychronakos and Roček, involve a single arbitrary function of  $J_0$  and include the quantum algebra  $\operatorname{su}_q(2)$  as a special case. In the present contribution, less common nonlinear deformations of  $\operatorname{su}(2)$ , introduced by Delbecq and Quesne and involving two deforming functions of  $J_0$ , are reviewed. Such algebras include Witten's quadratic deformation of  $\operatorname{su}(2)$  as a special case. Contrary to the former deformations, for which the spectrum of  $J_0$  is linear as for  $\operatorname{su}(2)$ , the latter give rise to exponential spectra, a property that has aroused much interest in connection with some physical problems. Another interesting algebra of this type, denoted by  $\mathcal{A}_q^+(1)$ , has two series of (N + 1)-dimensional unitary irreducible representations, where  $N = 0, 1, 2, \ldots$  To allow the coupling of any two such representations, a generalization of the standard Hopf axioms is proposed. The resulting algebraic structure, referred to as a two-colour quasitriangular Hopf algebra, is described.

<sup>\*</sup>Presented at the 5th International Colloquium "Quantum Groups and Integrable Systems", Prague, 20–22 June 1996

<sup>&</sup>lt;sup>†</sup>Directeur de recherches FNRS

### 1 Introduction

Quantized universal enveloping algebras, also called q-algebras, refer to some specific deformations of (the universal enveloping algebra of) Lie algebras, to which they reduce when the deforming parameter q (or set of deforming parameters) goes to one [1]. The simplest example of q-algebra,  $su_q(2) \equiv U_q(su(2))$ , was first introduced by Sklyanin, and by Kulish and Reshetikhin [2]. It has found a lot of applications in various branches of physics since its realization in terms of q-bosonic operators was proposed by Biedenharn and Macfarlane [3].

The  $su_q(2)$  algebra is a special case of more general deformations of su(2), independently introduced by Polychronakos and Roček [4]. They involve one arbitrary function  $f(J_0)$  in the commutator of  $J_+$  with  $J_-$ , and their representation theory is characterized by a rich variety of phenomena, whose interest in particle physics has been stressed.

More recently, deformations of su(2) involving two deforming functions  $F(J_0)$ and  $G(J_0)$  in the commutator of  $J_+$  with  $J_-$  and in that of  $J_0$  with  $J_+$  or  $J_-$ , respectively, have been proposed by Delbecq and Quesne [5, 6, 7]. It is the purpose of the present contribution to review the construction and representation theory of such algebras, and to show how the problem of endowing some of them with a Hopf algebraic structure can be addressed [8].

## 2 Nonlinear deformed su(2) algebras

Polychronakos-Roček algebras (PRA's) are associative algebras over  $\mathbb{C}$ , generated by three operators  $j_0 = (j_0)^{\dagger}$ ,  $j_+$ , and  $j_- = (j_+)^{\dagger}$ , satisfying the commutation relations [4]

$$[j_0, j_+] = j_+, \qquad [j_0, j_-] = -j_-, \qquad [j_+, j_-] = f(j_0), \tag{1}$$

where f(z) is a real, parameter-dependent function of z, holomorphic in the neighbourhood of zero, and going to 2z for some values of the parameters. These algebras have a Casimir operator given by  $c = j_-j_+ + h(j_0) = j_+j_- + h(j_0) - f(j_0)$ , in terms of another real function h(z), related to f(z) through the equation h(z) - h(z-1) = f(z). An explicit expression for h(z) has been given by Delbecq and Quesne [5] in terms of Bernoulli polynomials and Bernoulli numbers.

For all PRA's, the spectrum of  $j_0$  is linear as in the special case of the q-algebra  $\operatorname{su}_q(2)$ . The latter corresponds to  $f(j_0) = [2j_0]_q$ , and  $h(j_0) = [j_0]_q[j_0 + 1]_q$ , with  $[x]_q \equiv (q^x - q^{-x})/(q - q^{-1})$ , and  $q \in \mathbb{R}^+$  (the case where q is a phase will not be considered here, as throughout the present work we shall restrict the parameters to real values) [3].

Delbecq-Quesne algebras (DQA's) differ from PRA's by the replacement of (1) by [5]

$$[J_0, J_+] = G(J_0)J_+, \qquad [J_0, J_-] = -J_-G(J_0), \qquad [J_+, J_-] = F(J_0), \qquad (2)$$

where  $J_0 = (J_0)^{\dagger}$ ,  $J_- = (J_+)^{\dagger}$ , and the commutators involve two real, parameterdependent functions of z, F(z) and G(z), holomorphic in the neighbourhood of zero, and going to 2z and 1 for some values of the parameters, respectively. These functions are further restricted by the assumption that the algebras have a Casimir operator given by  $C = J_-J_+ + H(J_0) = J_+J_- + H(J_0) - F(J_0)$ , in terms of some real function H(z), holomorphic in the neighbourhood of zero. The latter restriction implies that F(z), G(z), and H(z) satisfy the consistency condition H(z) - H(z - G(z)) = F(z).

Since for  $G(J_0) = 1$ , DQA's reduce to PRA's, the first significant case corresponds to  $G(J_0) = 1 + (1 - q)J_0$ , where  $q \in \mathbb{R}^+$ . In such a case, it has been shown that there exist  $(\lambda - 1)$ -parameter algebras  $\mathcal{A}^+_{\alpha_2\alpha_3...\alpha_{\lambda-1}q}(\lambda, 1)$ , for which the functions  $F(J_0)$  and  $H(J_0)$  are polynomials of degree  $\lambda$  in  $J_0$  [5]. In particular, for  $\lambda = 2$  and 3, one finds the algebras  $\mathcal{A}^+_q(2,1)$  [6] and  $\mathcal{A}^+_{p,q}(3,1)$  [7], for which  $F(J_0) = 2J_0(1 + (1 - q)J_0)$ ,  $H(J_0) = 2(1 + q)^{-1}J_0(J_0 + 1)$ , and  $F(J_0) = 2J_0(1 + (1 - q)J_0)$ ,  $H(J_0) = 2((1 + q)(1 + q + q^2))^{-1}J_0(J_0 + 1)(1 + (p+q)q - (1 - p)(1 + q)J_0)$ , respectively. The former is equivalent to Witten's first deformation of su(2) [9].

The representation theory of the DQA's can be dealt with as that of  $\operatorname{su}_q(2)$ , or more generally of the PRA's. Considering the case where  $G(J_0) = 1 + (1 - q)J_0$ , and denoting by  $|cm\rangle$  a simultaneous eigenvector of the commuting Hermitian operators C and  $J_0$ , associated with the eigenvalues c and m respectively, it can be proved [5] that  $J^n_+|cm\rangle$  (resp.  $J^n_-|cm\rangle$ ),  $n \in \mathbb{N}^+$ , is either the null vector or a simultaneous eigenvector of C and  $J_0$ , corresponding to the eigenvalues c and  $mq^{-n} - (1 - q^{-n})/(1 - q)$  (resp.  $mq^n - (1 - q^n)/(1 - q)$ ). Hence, the spectrum of  $J_0$  is exponential, instead of linear as for the PRA's.

Moreover, if the starting m value belongs to the interval  $((q-1)^{-1}, +\infty)$ (resp.  $(-\infty, (q-1)^{-1})$ ), then all the  $J_0$  eigenvalues obtained by successive applications of  $J_+$  or  $J_-$  upon  $|cm\rangle \neq 0$  belong to the same interval and  $J_+$  (resp.  $J_-$ ) is a raising generator, whereas if  $m = (q-1)^{-1}$ , then neither  $J_+$  nor  $J_-$  change the  $J_0$  eigenvalue. The unirreps therefore separate into two classes according to whether the eigenvalues of  $J_0$  are contained in the interval  $(-\infty, (q-1)^{-1})$ , or in the interval  $((q-1)^{-1}, +\infty)$ .

In general, they may fall into one out of four categories: (i) infinite-dimensional unirreps with a lower bound -j, (ii) infinite-dimensional unirreps with an upper bound J, (iii) infinite-dimensional unirreps with neither lower nor upper bounds, and (iv) finite-dimensional unirreps with both lower and upper bounds, -j and J (where in general  $j \neq J$ ). In addition, there is a trivial one-dimensional unirrep corresponding to  $m = (q-1)^{-1}$ .

The exponential character of the  $J_0$  spectrum in DQA representations may be of interest in various physical problems, wherein such spectra have been encountered, such as alternative Hamiltonian quantizations, exactly solvable potentials, q-deformed supersymmetric quantum mechanics, and q-deformed interacting boson models [10].

# **3** The algebra $\mathcal{A}_a^+(1)$

Another example of DQA, for which the function  $G(J_0)$  is linear, has been recently constructed [8]. Contrary to those considered in the previous section, this algebra, denoted by  $\mathcal{A}_q^+(1)$ , is defined in terms of functions  $F(J_0)$  and  $H(J_0)$  that are not polynomials, but infinite series in  $J_0$ ,

$$F(J_0) = -\frac{(G(J_0))^2 - (G(J_0))^{-2}}{q - q^{-1}},$$
  

$$H(J_0) = \frac{q^{-1}(G(J_0))^2 + q(G(J_0))^{-2} - q - q^{-1}}{(q - q^{-1})^2},$$
(3)

with  $G(J_0) = 1 + (1 - q)J_0$ . Since the transformation  $q \to q^{-1}$ ,  $J_0 \to -qJ_0$ ,  $J_{\pm} \to J_{\mp}$  is an automorphism of  $\mathcal{A}_q^+(1)$ , the parameter values may be restricted to the range 0 < q < 1.

 $\mathcal{A}_q^+(1)$  can be obtained from  $\mathrm{su}_q(2)$ , a special case of PRA, by using a two-valued map  $P_\delta : \mathrm{su}_q(2) \to \mathcal{A}_q^+(1), \, \delta = \pm 1$ , defined by

$$J_0 = p_{\delta}(j_0), \qquad J_+ = j_+, \qquad J_- = j_-, \qquad \text{where} \qquad p_{\delta}(z) \equiv \frac{1 - \delta q^{-z}}{q - 1}.$$
 (4)

Such a generator map is well defined: it can indeed be easily checked that if  $j_0$ ,  $j_+$ ,  $j_-$  satisfy the  $su_q(2)$  commutation relations, then  $J_0$ ,  $J_+$ ,  $J_-$ , given in (4), fulfil those of  $\mathcal{A}_q^+(1)$ . The functions  $p_{\delta}(z)$ ,  $\delta = \pm 1$ , defined in (4), are entire and invertible functions, with  $g(z) = p_{\delta}^{-1}(z) = \ln(G^2(z)) / \ln(q^{-2})$ . If  $z \in \mathbb{R}$ , the range of  $p_{\delta}$  (and consequently the domain of  $p_{\delta}^{-1}$ ) is the interval  $(-\infty, (q-1)^{-1})$  or  $((q-1)^{-1}, +\infty)$  according to whether  $\delta = -1$  or  $\delta = +1$ . The function g(z) is well-behaved everywhere on  $\mathbb{R}$ , except in the neighbourhood of the point  $z = (q-1)^{-1}$ .

It should be stressed that the use of  $P_{\delta}$ ,  $\delta = \pm 1$ , implies an extension of the well-known deforming functional technique [11] for two reasons: first because here a map between two deformed algebras,  $\operatorname{su}_q(2)$  and  $\mathcal{A}_q^+(1)$ , is considered instead of a map between a Lie algebra and a deformed one, as in the original method; and second because use is made of a two-valued functional, whose inverse is singular, instead of a single-valued one.

It can be easily shown [8] that  $\mathcal{A}_{q}^{+}(1)$  has no infinite-dimensional unirrep, but has, for any  $N = 0, 1, 2, \ldots$ , two (N + 1)-dimensional unirreps, which may be distinguished by  $\delta = \pm 1$ . The corresponding spectrum of  $J_0$  is given by  $m^{\delta} = (1 - \delta q^{-(N-2n)/2})/(q-1), n = 0, 1, \ldots, N$ , with maximum and minimum eigenvalues  $J^{\delta} = (1 - \delta q^{-\delta N/2})/(q-1)$ , and  $-j^{\delta} = (1 - \delta q^{\delta N/2})/(q-1)$ respectively. The unirrep specified by  $J^+$  (resp.  $J^-$ ) is entirely contained in the interval  $((q-1)^{-1}, +\infty)$  (resp.  $(-\infty, (q-1)^{-1})$ ). For both unirreps, the eigenvalue of the Casimir operator is given by  $\langle C \rangle = H(\gamma^{\delta})$ , where  $\gamma^{\delta} = (1 - \delta q^{-N/2})/(q-1)$ .

In the carrier space  $V^{J^{\delta}}$  of the unirrep characterized by  $J^{\delta}$ , whose basis vectors are specified by the values of  $J^{\delta}$  and  $m^{\delta}$ , the  $\mathcal{A}_{q}^{+}(1)$  generators are represented by

some linear operators  $\Phi^{J^{\delta}}(A), A \in \mathcal{A}_q^+(1)$ , defined by

$$\Phi^{J^{\delta}}(J_{0}) | J^{\delta}, m^{\delta} \rangle = m^{\delta} | J^{\delta}, m^{\delta} \rangle = \left(\frac{N}{2} - n\right) | J^{\delta}, m^{\delta} \rangle,$$

$$\Phi^{J^{\delta}}(J_{-}) | J^{\delta}, m^{\delta} \rangle = \sqrt{H(\gamma^{\delta}) - H(qm^{\delta} - 1)} | J^{\delta}, qm^{\delta} - 1 \rangle$$

$$= \sqrt{[n+1]_{q}[N-n]_{q}} | J^{\delta}, qm^{\delta} - 1 \rangle,$$

$$\Phi^{J^{\delta}}(J_{+}) | J^{\delta}, m^{\delta} \rangle = \sqrt{H(\gamma^{\delta}) - H(m^{\delta})} | J^{\delta}, q^{-1}(m^{\delta} + 1) \rangle$$

$$= \sqrt{[n]_{q}[N-n+1]_{q}} | J^{\delta}, q^{-1}(m^{\delta} + 1) \rangle.$$
(5)

The generator mapping  $P_{\delta}$  can be used to transfer the quasitriangular Hopf structure of  $\operatorname{su}_q(2)$  to  $\mathcal{A}_q^+(1)$  [8]. One gets in this way a double quasitriangular Hopf structure, with comultiplication, counit, antipode maps, and universal  $\mathcal{R}$ matrix given by

$$\begin{aligned} \Delta_{\delta} \left( J_{0} \right) &= (q-1)^{-1} \left( 1 \otimes 1 - \delta G(J_{0}) \otimes G(J_{0}) \right), \\ \Delta_{\delta} \left( J_{\pm} \right) &= \delta \left( J_{\pm} \otimes (G(J_{0}))^{-1} + G(J_{0}) \otimes J_{\pm} \right), \\ \epsilon_{\delta} (J_{0}) &= (1-\delta)(q-1)^{-1}, \quad \epsilon_{\delta} (J_{\pm}) = 0, \\ S_{\delta} (J_{0}) &= -J_{0} (G(J_{0}))^{-1}, \quad S_{\delta} (J_{\pm}) = -q J_{\pm}, \quad S_{\delta} (J_{\pm}) = -q^{-1} J_{\pm}, \\ \mathcal{R}^{\delta} &= q^{2 \log_{q} (\delta G(J_{0})) \otimes \log_{q} (\delta G(J_{0}))} \sum_{n=0}^{\infty} \frac{(1-q^{-2})^{n}}{[n]_{q}!} q^{n(n-1)/2} \\ &\times \left( (G(J_{0}))^{-1} J_{\pm} \otimes G(J_{0}) J_{\pm} \right)^{n}, \end{aligned}$$
(6)

respectively. Both  $(\Delta_+, \epsilon_+, S_+, \mathcal{R}^+)$ , and  $(\Delta_-, \epsilon_-, S_-, \mathcal{R}^-)$  satisfy the Hopf and quasitriangularity axioms, but the former are only valid for the representations of  $\mathcal{A}_q^+(1)$  with eigenvalues of  $J_0$  in the interval  $((q-1)^{-1}, +\infty)$ , whereas the latter act in  $(-\infty, (q-1)^{-1})$ .

# 4 Two-colour quasitriangular Hopf structure of $\mathcal{A}_q^+(1)$

The double Hopf structure considered in the previous section allows one to couple any two  $\mathcal{A}_q^+(1)$  unirreps characterized by  $J_1^+$  and  $J_2^+$  (resp.  $J_1^-$  and  $J_2^-$ ), and with respective carrier spaces  $V^{J_1^+}$  and  $V^{J_2^+}$  (resp.  $V^{J_1^-}$  and  $V^{J_2^-}$ ), to obtain a reducible representation of the same in  $V^{J_1^+} \otimes V^{J_2^+}$  (resp.  $V^{J_1^-} \otimes V^{J_2^-}$ ). No coupling of two unirreps of the types  $J_1^+$  and  $J_2^-$ , or  $J_1^-$  and  $J_2^+$ , is however possible.

To allow such types of couplings, it is necessary to extend the double Hopf structure of  $\mathcal{A}_q^+(1)$  [8]. This can be accomplished by considering the 'transmutation' operators  $T^{J^{\delta}}: V^{J^{\delta}} \to V^{J^{-\delta}}$ , which change the basis states of an (N+1)-dimensional unirrep, characterized by  $J^{\delta}$ , into those of its partner with the same dimension, specified by  $J^{-\delta}$ , i.e.,  $T^{J^{\delta}} | J^{\delta}, m^{\delta} \rangle = | J^{-\delta}, m^{-\delta} \rangle$ . By applying  $T^{J^{\delta}}$  on both sides of (5), we obtain that for any  $\mathcal{A}_{q}^{+}(1)$  generator A,  $T^{J^{\delta}} \Phi^{J^{\delta}}(A) T^{J^{-\delta}} = \Phi^{J^{-\delta}}(\sigma(A))$ , where  $\sigma : \mathcal{A}_{q}^{+}(1) \to \mathcal{A}_{q}^{+}(1)$ , defined by  $\sigma(J_{0}) = 2(q-1)^{-1} - J_{0}$ , and  $\sigma(J_{\pm}) = J_{\pm}$ , is an involutive automorphism of the algebra  $\mathcal{A}_{q}^{+}(1)$ . This clearly shows that at the algebra level, the operator  $\sigma$  is responsible for the transmutation. Defining now  $\sigma_{\delta} : \mathcal{A}_{q}^{+}(1) \to \mathcal{A}_{q}^{+}(1)$ ,  $\delta = \pm 1$ , as  $\sigma_{+} = \mathrm{id}$ , and  $\sigma_{-} = \sigma$ , we note that the basic mapping  $P_{\delta}$ , defined in (4), satisfies the relation  $\sigma_{\zeta \eta} \circ P_{\eta} = P_{\zeta}$ , where  $\zeta$ ,  $\eta = \pm 1$ .

The comultiplication and antipode maps, as well as the double  $\mathcal{R}$ -matrix of equation (6) can be extended by setting

$$\Delta_{\delta}^{\zeta,\eta}(A) = (\sigma_{\zeta\delta} \otimes \sigma_{\eta\delta}) \circ \Delta_{\delta}(A), \qquad S_{\delta}^{\zeta}(A) = \sigma_{\zeta\delta} \circ S_{\delta}(A),$$
  
$$\mathcal{R}^{\zeta,\eta} = (\sigma_{\zeta\delta} \otimes \sigma_{\eta\delta}) \left(\mathcal{R}^{\delta}\right), \qquad (7)$$

where  $\zeta$ ,  $\eta$ ,  $\delta = \pm 1$ , while the counit map  $\epsilon_{\delta}$ , defined in the same equation, is left unchanged. The results can be written as

$$\begin{split} \Delta_{\delta}^{\zeta,\eta} (J_0) &= (q-1)^{-1} \left( 1 \otimes 1 - \delta \zeta \eta G(J_0) \otimes G(J_0) \right), \\ \Delta_{\delta}^{\zeta,\eta} (J_{\pm}) &= \eta J_{\pm} \otimes (G(J_0))^{-1} + \zeta G(J_0) \otimes J_{\pm}, \\ S_{\delta}^{\zeta} (J_0) &= (q-1)^{-1} \left( 1 - \zeta \delta \left( G(J_0) \right)^{-1} \right), \\ S_{\delta}^{\zeta} (J_{\pm}) &= -q^{\pm 1} J_{\pm}, \\ \mathcal{R}^{\zeta,\eta} &= q^{2 \log_q(\zeta G(J_0)) \otimes \log_q(\eta G(J_0))} \\ &\times \sum_{n=0}^{\infty} \frac{(1 - q^{-2})^n}{[n]_q!} q^{n(n-1)/2} \left( (\zeta G(J_0))^{-1} J_{\pm} \otimes \eta G(J_0) J_{\pm} \right)^n. \end{split}$$
(8)

It can be easily shown [8] that the generalized comultiplication, counit, and antipode maps,  $\Delta_{\delta}^{\zeta,\eta}$ ,  $\epsilon_{\delta}$ ,  $S_{\delta}^{\zeta}$ , defined in (6) and (8), transform under  $\sigma_{\delta}$  as

$$(\sigma_{\mu\zeta} \otimes \sigma_{\nu\eta}) \circ \Delta_{\delta}^{\zeta,\eta} = \Delta_{\rho}^{\mu,\nu} \circ \sigma_{\rho\delta}, \qquad \epsilon_{\delta} \circ \sigma_{\delta\zeta} = \epsilon_{\zeta}, \qquad \sigma_{\zeta\eta} \circ S_{\delta}^{\eta} = S_{\mu}^{\zeta} \circ \sigma_{\mu\delta}, \quad (9)$$

and satisfy the following generalized coassociativity, counit, and antipode axioms,

$$(\Delta_{\mu}^{\zeta,\eta} \otimes \mathrm{id}) \circ \Delta_{\delta}^{\mu,\nu}(A) = (\mathrm{id} \otimes \Delta_{\rho}^{\eta,\nu}) \circ \Delta_{\delta}^{\zeta,\rho}(A), (\epsilon_{\zeta} \otimes \sigma_{\eta\delta}) \circ \Delta_{\delta}^{\zeta,\eta}(A) = (\sigma_{\zeta\delta} \otimes \epsilon_{\eta}) \circ \Delta_{\delta}^{\zeta,\eta}(A) = A, m \circ \left(S_{\zeta}^{\mu} \otimes \sigma_{\mu\eta}\right) \circ \Delta_{\delta}^{\zeta,\eta}(A) = m \circ (\sigma_{\mu\zeta} \otimes S_{\eta}^{\mu}) \circ \Delta_{\delta}^{\zeta,\eta}(A) = \iota \circ \epsilon_{\delta}(A),$$
(10)

where A denotes any element of  $\mathcal{A}_q^+(1)$ , m and  $\iota$  are the multiplication and unit maps of  $\mathcal{A}_q^+(1)$ ,  $\delta$ ,  $\zeta$ ,  $\eta$ ,  $\mu$ ,  $\nu$ ,  $\rho$  take any values in the set  $\{-1, +1\}$ , and no summation over repeated indices is implied. Moreover,  $\Delta_{\delta}^{\zeta,\eta}$  and  $\epsilon_{\delta}$  are algebra homomorphisms, while  $S_{\delta}^{\zeta}$  is both an algebra and a coalgebra antihomomorphism.

By using the generalized coproduct  $\Delta_{\delta}^{\zeta,\eta}$ , it is now possible to couple any  $(N_1 + 1)$ - and  $(N_2 + 1)$ -dimensional unirreps of  $\mathcal{A}_q^+(1)$ , specified by  $J_1^{\zeta}$  and  $J_2^{\eta}$  respectively, to provide two reducible representations in  $V^{J_1^{\zeta}} \otimes V^{J_2^{\eta}}$ , which are characterized by  $\delta = +1$  and  $\delta = -1$ , respectively. They can be decomposed into a direct sum of (N + 1)-dimensional unirreps, specified by  $J^{\delta}$ , by using some Wigner coefficients  $\langle J_1^{\zeta} m_1^{\zeta}, J_2^{\eta} m_2^{\eta} | J^{\delta} m^{\delta} \rangle_{DQ}$ , given in terms of  $\mathrm{su}_q(2)$  Wigner coefficients by the relation

$$\left\langle J_{1}^{\zeta} m_{1}^{\zeta}, J_{2}^{\eta} m_{2}^{\eta} \middle| J^{\delta} m^{\delta} \right\rangle_{DQ} = \left\langle \frac{N_{1}}{2} \frac{N_{1}}{2} - n_{1}, \frac{N_{2}}{2} \frac{N_{2}}{2} - n_{2} \middle| \frac{N}{2} \frac{N}{2} - n \right\rangle_{q}.$$
(11)

The carrier space of the unirrep  $J^{\delta}$  in  $V^{J_1^{\zeta}} \otimes V^{J_2^{\eta}}$  is therefore spanned by the states

$$\left|J_{1}^{\zeta}J_{2}^{\eta}J^{\delta}m^{\delta}\right\rangle = \sum_{m_{1}^{\zeta},m_{2}^{\eta}} \left\langle J_{1}^{\zeta}m_{1}^{\zeta}, J_{2}^{\eta}m_{2}^{\eta}\right| J^{\delta}m^{\delta} \right\rangle_{DQ} \left|J_{1}^{\zeta},m_{1}^{\zeta}\right\rangle \otimes \left|J_{2}^{\eta},m_{2}^{\eta}\right\rangle.$$
(12)

Turning now to the generalized universal  $\mathcal{R}$ -matrix defined in (7) or (8), it can be easily shown [8] that its four pieces  $\mathcal{R}^{\zeta,\eta}$ ,  $\zeta$ ,  $\eta = \pm 1$ , are invertible and satisfy the properties

$$(\sigma_{\mu\zeta} \otimes \sigma_{\nu\eta}) \left( \mathcal{R}^{\zeta,\eta} \right) = \mathcal{R}^{\mu,\nu}, \quad \tau \circ \Delta^{\eta,\zeta}_{\delta}(A) = \mathcal{R}^{\zeta,\eta} \Delta^{\zeta,\eta}_{\delta}(A) \left( \mathcal{R}^{\zeta,\eta} \right)^{-1}, \left( \Delta^{\lambda,\mu}_{\zeta} \otimes \sigma_{\nu\eta} \right) \left( \mathcal{R}^{\zeta,\eta} \right) = \mathcal{R}^{\lambda,\nu}_{13} \mathcal{R}^{\mu,\nu}_{23}, \quad \left( \sigma_{\lambda\zeta} \otimes \Delta^{\mu,\nu}_{\eta} \right) \left( \mathcal{R}^{\zeta,\eta} \right) = \mathcal{R}^{\lambda,\nu}_{13} \mathcal{R}^{\lambda,\mu}_{12}, (13)$$

for any  $A \in \mathcal{A}_q^+(1)$ . From these results, or more simply from the corresponding properties fulfilled by  $\mathcal{R}^{\delta}$ , one obtains the relations

$$\mathcal{R}_{12}^{\zeta,\eta} \mathcal{R}_{13}^{\zeta,\mu} \mathcal{R}_{23}^{\eta,\mu} = \mathcal{R}_{23}^{\eta,\mu} \mathcal{R}_{13}^{\zeta,\eta} \mathcal{R}_{12}^{\zeta,\eta}, 
(\epsilon_{\zeta} \otimes \mathrm{id}) \left( \mathcal{R}^{\zeta,\eta} \right) = (\mathrm{id} \otimes \epsilon_{\eta}) \left( \mathcal{R}^{\zeta,\eta} \right) = 1, 
(S_{\zeta}^{\lambda} \otimes \sigma_{\mu\eta}) \left( \mathcal{R}^{\zeta,\eta} \right) = \left( \sigma_{\lambda\zeta} \otimes (S_{\eta}^{\mu})^{-1} \right) \left( \mathcal{R}^{\zeta,\eta} \right) = \left( \mathcal{R}^{\lambda,\mu} \right)^{-1}.$$
(14)

The first relation in (14) shows that the generalized universal  $\mathcal{R}$ -matrix is a solution of the coloured YBE [12], where the 'colour' parameters  $\zeta$ ,  $\eta$ ,  $\mu$  take discrete values in the set  $\{-1, +1\}$ . We may therefore call  $\left(\mathcal{A}_q^+(1), +, m, \iota, \Delta_{\delta}^{\zeta,\eta}, \epsilon_{\delta}, S_{\delta}^{\zeta}, \mathcal{R}^{\zeta,\eta}; \mathbb{C}\right)$  a two-colour quasitriangular Hopf algebra over  $\mathbb{C}$  [8]. As will be shown elsewhere [13], this type of algebraic structure admits generalizations, which will be referred to as coloured quasitriangular Hopf algebras.

### References

- Drinfeld V. G.: in Proc. Int. Congress of Mathematicians (Berkeley, CA, 1986) (Ed. A. M. Gleason), AMS, Providence, RI, 1987, p. 798. Jimbo M.: Lett. Math. Phys. 10 (1985) 63.
- [2] Sklyanin E. K.: Funct. Anal. Appl. 16 (1982) 262.
   Kulish P. P. and Reshetikhin N. Y.: J. Sov. Math. 23 (1983) 2435.
- [3] Biedenharn L. C.: J. Phys. A 22 (1989) L873.
   Macfarlane A. J.: J. Phys. A 22 (1989) 4581.
- [4] Polychronakos A. P.: Mod. Phys. Lett. A 5 (1990) 2325.
   Roček M.: Phys. Lett. B 255 (1991) 554.
- [5] Delbecq C. and Quesne C.: J. Phys. A 26 (1993) L127.
- [6] Delbecq C. and Quesne C.: Phys. Lett. B 300 (1993) 227.
- [7] Delbecq C. and Quesne C.: Mod. Phys. Lett. A 8 (1993) 961.
- [8] Bonatsos D., Daskaloyannis C., Kolokotronis P., Ludu A., and Quesne C.: A nonlinear deformed su(2) algebra with a two-colour quasitriangular Hopf structure, Preprint, Université Libre de Bruxelles, 1996.
- [9] Witten E.: Nucl. Phys. B 330 (1990) 285.
- [10] Fairlie D. B. and Nuyts J., J. Phys. A 24 (1991) L1001.
  Spiridonov V., Phys. Rev. Lett. 69 (1992) 398; Mod. Phys. Lett. A 7 (1992) 1241.
  Gupta R. K. and Ludu A., Phys. Rev. C 48 (1993) 593.
- [11] Curtright T. L. and Zachos C. K., Phys. Lett. B 243 (1990) 237.
   Curtright T. L., Ghandour G. I., and Zachos C. K., J. Math. Phys. 32 (1991) 676.
- [12] Murakami J.: in Proc. Int. Conf. of Euler Mathematical School: Quantum Groups (Leningrad, 1990), Lecture Notes in Physics, Springer, Berlin, 1991, p. 350.
  Akutsu Y. and Deguchi T.: Phys. Rev. Lett. 67 (1991) 777.
  Ge M.-L., Liu X.-F., and Sun C.-P.: Phys. Lett. A 155 (1991) 137.
  Burdík Č. and Hellinger P.: J. Phys. A 25 (1992) L1023.
  Kundu A. and Basu-Mallick B.: J. Phys. A 25 (1992) 6307.
- [13] Quesne C.: Coloured Hopf algebras, in preparation.