3-State Potts model and automorphism of vertex operator algebra of order 3

Masahiko Miyamoto

Institute of Mathematics University of Tsukuba Tsukuba 305, Japan

Abstract

In [M2], the author has defined an automorphism τ of a vertex operator algebra (VOA) of order 2 using a sub VOA isomorphic to Ising model $L(\frac{1}{2}, 0)$. We here define an automorphism of VOA of order 3 by using a sub VOA isomorphic to a direct sum of 3-state Potts models $L(\frac{4}{5}, 0)$ and an its module $L(\frac{4}{5}, 3)$. If V is the moonshine VOA V^{\natural} , the defined automorphism is a 3A element of the monster simple group.

1 Introduction

In the research of the Griess algebra V_2^{\natural} , Conway [C] found several idempotents called axes of the Griess algebra corresponding to elements of the monster simple group $\mathbf{M} = Aut(V^{\natural})$. It was discovered that the Griess algebra is the second primary part $(V^{\natural})_2$ of the moonshine VOA V^{\natural} constructed in [FLM].

It was proved in [M2] that idempotents in the Griess algebra are halves of conformal vectors (or Virasoro elements of sub VOAs). In particular, every idempotent in the Conway's list is a half of the Virasoro element of a sub VOA isomorphic to one of the minimal discrete series of Virasoro VOA L(n, 0) with central charge n for some 0 < n < 1. For example, a 2A-involution of the monster simple group is a half of the Virasoro element of a sub VOA isomorphic to the Ising model $L(\frac{1}{2}, 0)$ with central charge $\frac{1}{2}$. We note that a sub VOA isomorphic to the Ising model $L(\frac{1}{2}, 0)$ with central charge $\frac{1}{2}$. We note that a sub VOA W in this paper does not usually have the same Virasoro element of V. Conversely, the author showed that if a VOA V contains a sub VOA W isomorphic to the Ising model $L(\frac{1}{2}, 0)$, then it defines an automorphism τ_W of V of order at most 2, which is a 2A-element of the monster simple group if V is the monshine VOA V^{\natural} .

is very simple and done as follows:

Let e be a Virasoro element of W. As we will prove the general case in Theorem 5.1, V is a direct sum of (possibly infinite number of) irreducible W-modules since $W \cong L(\frac{1}{2}, 0)$ is rational. It is well known that $L(\frac{1}{2}, 0)$ has the exactly three irreducible modules $L(\frac{1}{2}, 0)$, $L(\frac{1}{2}, \frac{1}{2})$ and $L(\frac{1}{2}, \frac{1}{16})$. We can define an automorphism τ_e by

$$\tau_e: \begin{cases} 1 & on \quad W\text{-submodules isomorphic to } L(\frac{1}{2}, 0) \text{ or } L(\frac{1}{2}, \frac{1}{2}) \\ -1 & on \quad W\text{-submodules isomorphic to } L(\frac{1}{2}, \frac{1}{16}) \end{cases}$$

Throughout this paper, we will use the similar notation in order to define an endomorphism of V by a sub VOA W and we will omit "W-submodules isomorphic to" from the definition of automorphisms in order to simplify the notation. In the Conway's list, an idempotent for a 3A element is a half of the Virasoro element of a sub VOA isomorphic to $L(\frac{4}{5}, 0)$ with central charge $\frac{4}{5}$. So it is natural for us to expect an automorphism g (of order 3) defined by a sub VOA isomorphic to $L(\frac{4}{5}, 0)$, where $L(\frac{4}{5}, 0)$ is the third of the discrete series of minimal Virasoro vertex operator algebras called 3-state Potts model. It is a rational VOA and has the exactly ten irreducible modules :

$$L(\frac{4}{5},0), L(\frac{4}{5},\frac{1}{8}), L(\frac{4}{5},\frac{2}{3}), L(\frac{4}{5},\frac{13}{8}), L(\frac{4}{5},3), L(\frac{4}{5},\frac{2}{5}), L(\frac{4}{5},\frac{1}{40}), L(\frac{4}{5},\frac{1}{15}), L(\frac{4}{5},\frac{21}{40}), L(\frac{4}{5},\frac{7}{5}).$$

As we showed in [M2], if a VOA V contains a sub VOA $W \cong L(\frac{4}{5}, 0)$, then we can define an automorphism σ_W of at most 2 given by

$$\sigma_W : \begin{cases} 1 & on \quad L(\frac{4}{5}, 0), L(\frac{4}{5}, 3), L(\frac{4}{5}, \frac{13}{8}), L(\frac{4}{5}, \frac{2}{5}), L(\frac{4}{5}, \frac{1}{15}), L(\frac{4}{5}, \frac{7}{5}) \\ -1 & on \quad L(\frac{4}{5}, \frac{1}{8}), L(\frac{4}{5}, \frac{13}{8}), L(\frac{4}{5}, \frac{1}{40}), L(\frac{4}{5}, \frac{21}{40}) \end{cases}$$

We note that we can't observe this automorphism in the moonshine VOA. Namely, V^{\ddagger} does not contain any submodules of the second lines. Under such a situation, we want to define a 3A automorphism τ_W of V by a sub VOA W isomorphic to $L(\frac{4}{5}, 0)$. It is clear that it is not enough to think of only $L(\frac{4}{5}, 0)$ because there is no difference between the eigenspaces of τ with eigenvalues $e^{2\pi i/3}$ and $e^{4\pi i/3}$. Namely, let V^1 and V^2 be eigenspaces of τ with eigenvalue $e^{2\pi i/3}$ and $e^{4\pi i/3}$, then they are isomorphic as $L(\frac{4}{5}, 0)$ -modules. However, Dong and Mason [DM] showed a wonderful result that V^1 and V^2 are not isomorphic as $V^{<\tau_W>}$ -modules, where $V^{<\tau_W>}$ is the space of τ_W -invariants. So we have to think of a bigger sub VOA. What is the difference between $V^{<\tau_W>}$ and $L(\frac{4}{5}, 0)$? Recently, Kitazume, Yamada and the author have constructed a new class of VOAs by using codes over \mathbb{Z}_3 in [KMY]. The interesting point is that they used a VOA isomorphic to $L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$ as a sub VOA corresponding to $0 \in \mathbb{Z}_3$.

This is our key point and the main result in this paper is to show that if V contains a sub VOA W isomorphic to $L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$, we can define a triality automorphism τ_W of V (of order 3 or possibly 1).

In order to define the automorphism, we need quote several results from [KMY]. They classified the irreducible modules of $W(0) = L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$. Namely,

Theorem 1.1 ([KMY]) $L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$ is a rational VOA and it has the exactly six irreducible modules:

$$W(0), W(\frac{2}{5}), W(\frac{2}{3}, +), W(\frac{1}{15}, +), W(\frac{2}{3}, -), W(\frac{1}{15}, -).$$

Here h in W(h) or $W(h, \pm)$ denotes the lowest degree and W(k, -) is a contragredient (dual) module of W(k, +) for $k = \frac{2}{3}, \frac{1}{15}$. In particular,

$$\begin{split} W(0) &\cong L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3), \\ W(\frac{2}{5}) &\cong L(\frac{4}{5}, \frac{2}{5}) \oplus L(\frac{4}{5}, \frac{7}{5}), \\ W(\frac{2}{3}, +) &\cong L(\frac{4}{5}, \frac{2}{3}), \\ W(\frac{2}{3}, -) &\cong L(\frac{4}{5}, \frac{2}{3}), \\ W(\frac{1}{15}, +) &\cong L(\frac{4}{5}, \frac{1}{15}), \\ W(\frac{1}{15}, -) &\cong L(\frac{4}{5}, \frac{1}{15}), \end{split}$$

as $L(\frac{4}{5}, 0)$ -modules.

Using the notation in the above theorem, we have:

Theorem A If a VOA V contains a sub VOA W isomorphic to $L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$, then an endomorphism τ_W of V defined by

$$\tau_W : \begin{cases} 1 & on \quad W(0) \text{ and } W(\frac{2}{5}) \\ e^{2\pi i/3} & on \quad W(\frac{2}{3}, +) \text{ and } W(\frac{1}{15}, +) \\ e^{4\pi i/3} & on \quad W(\frac{2}{3}, -) \text{ and } W(\frac{1}{15}, -) \end{cases}$$

is an automorphism of V.

In order to tell the difference between W(h, +) and W(h, -) for $h = \frac{2}{3}, \frac{1}{15}$, we have to explain the actions of the lowest degree vector of $L(\frac{4}{5}, 3)$ since both $W(h, \pm)$ are isomorphic to $L(\frac{4}{5}, h)$ as $L(\frac{4}{5}, 0)$ -modules. However, we will take an easier way to avoid such a complicated job. We only note that if we fix $W(\frac{2}{3}, +)$, then $W(\frac{1}{15}, \pm)$ and $W(\frac{2}{3}, -)$ are uniquely determined by the fusion rule

$$W(\frac{2}{3},\pm) \times W(\frac{2}{5}) = W(\frac{1}{15},\pm).$$

We don't distinguish between $W(\frac{2}{3}, +)$ and $W(\frac{2}{3}, -)$, but if we switch them then we shall define τ_W^{-1} .

Let T be a sub VOA of W isomorphic to $L(\frac{4}{5}, 0)$. If $\tau = 1$, then all T-submodules of V are isomorphic to $L(\frac{4}{5}, 0)$, $L(\frac{4}{5}, 3)$, $L(\frac{4}{5}, \frac{2}{5})$ or $L(\frac{4}{5}, \frac{7}{5})$. In this case, we can define another automorphism μ_T of V as follows:

Theorem B Assume that V contains a sub VOA T isomorphic to $L(\frac{4}{5}, 0)$ and all T-submodules of V are isomorphic to $L(\frac{4}{5}, 0)$, $L(\frac{4}{5}, 3)$, $L(\frac{4}{5}, \frac{2}{5})$ or $L(\frac{4}{5}, \frac{7}{5})$. Then the endomorphism μ_T defined by

$$\mu_T : \begin{cases} 1 & on \quad L(\frac{4}{5}, 0) \text{ and } L(\frac{4}{5}, \frac{7}{5}) \\ -1 & on \quad L(\frac{4}{5}, 3) \text{ and } L(\frac{4}{5}, \frac{2}{5}) \end{cases}$$

is an automorphism of V.

The proofs of these theorems are based on Theorem 2.1 (Proposition 4.4 in [M1]). Namely, it is sufficient to show that the fusion rules among the irreducible $L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$ modules commutes with τ_W . For an example, we know the fusion rules among irreducible $L(\frac{4}{5}, 0)$ -modules (Table A), which proves Theorem B. Therefore, the main thing we will do in this paper is to determine the fusion rules among the irreducible $L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$ modules.

In this paper, we often view V as a W-module (or an infinite direct sum of W-modules) if V contains a sub VOA W. This is not obvious since one of the axioms of VOA-modules expects the grade keeping operator e_1 of Virasoro element e of W to act on V diagonally. We will prove in §4 that this is generally true for a rational sub VOA W.

2 Preliminary results and a generalized VOA constructed from a lattice

Throughout this paper, W(0) denotes a VOA isomorphic to $L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$. Since we will treat only a rational VOA V isomorphic to $L(\frac{4}{5}, 0)$ or W(0), the tensor products of two V-modules M^1 and M^2 are always well-defined and it is isomorphic to $\oplus_U N_{M^1,M^2}^U U$, where U runs over the all irreducible V-modules. Therefore, it is equal to the fusion rule in our case and so we will use the same notation $M^1 \times M^2$ to denote the tensor product.

Since $L(\frac{4}{5}, 0) \subseteq W(0)$, all W(0)-modules are $L(\frac{4}{5}, 0)$ -modules. Using this fact, we will give an upperbound of the fusion rules of W(0)-modules. Using exactly the same proof, we can modify Proposition 11.9 in [DL] into the following theorem.

Theorem 2.1 ([DL]) Let W^1, W^2, W^3 be V-modules and assume that W^1, W^2 have no proper submodules containing v^1 and v^2 , respectively. Let $I \in I\begin{pmatrix} W^3 \\ W^1 & W^2 \end{pmatrix}$. If $I(v^1, z)v^2 = 0$, then $I(\cdot, z) = 0$.

In the case where $W^1 = W(h, +) \oplus W(h, -)$ for $h = \frac{1}{15}, \frac{2}{3}, W^1$ has no proper submodule containing $U^1 = \{(v, \phi(v)) \in W(h, +) \oplus W(h, -)\}$, where $\phi : W(h, +) \to W(h, -)$ is a $L(\frac{4}{5}, 0)$ -isomorphism. Therefore, we have the following theorem:

Lemma 2.1 The maps

$$\begin{split} \phi^{1} : & I_{W(0)} \begin{pmatrix} W^{3} \\ W(i) & W(j) \end{pmatrix} \to I_{L(\frac{4}{5},0)} \begin{pmatrix} W^{3} \\ L(\frac{4}{5},i) & L(\frac{4}{5},j) \end{pmatrix} \end{pmatrix}, \\ \phi^{2} : & I_{W(0)} \begin{pmatrix} W^{3} \\ W(h,+) \oplus W(h,-) & W(k,+) \oplus W(k,-) \end{pmatrix} \to I_{L(\frac{4}{5},0)} \begin{pmatrix} W^{3} \\ L(\frac{4}{5},h) & L(\frac{4}{5},k) \end{pmatrix} \\ and \\ and \\ \end{split}$$

$$\phi^3 : I_{W(0)} \begin{pmatrix} W^3 \\ W(h, +) \oplus W(h, -) & W(i) \end{pmatrix} \to I_{L(\frac{4}{5}, 0)} \begin{pmatrix} W^3 \\ L(\frac{4}{5}, h) & L(\frac{4}{5}, j) \end{pmatrix}$$

induced by the restrictions are all injective for $i, j = 0, \frac{2}{5}$ and $h, k = \frac{2}{3}, \frac{1}{15}$.

Throughout this paper, $N_{W^1,W^2}^{W^3}$ denotes dim $I\begin{pmatrix} W^3\\ W^1 & W^2 \end{pmatrix}$. It is known that

$$N^{W^3}_{W^1,W^2} = N^{W^3}_{W^2,W^1} = N^{(W^1)'}_{W^2,(W^3)'}$$

where W' denotes the contragredient (dual) module of W, (see Proposition 5.5.2 in [FHL]). We note that $N_{W(0),W^1}^{W^1} = 1$ and $N_{W(0),W^1}^{W^2} = 0$ for $W^1 \not\cong W^2$. Let $k' = 3, \frac{7}{5}$ for $k = 0, \frac{2}{5}$, respectively. Namely, $W(k) \cong L(\frac{4}{5}, k) \oplus L(\frac{4}{5}, k')$ as $L(\frac{4}{5}, 0)$ -modules. By the above lemma and the fusion rules of the irreducible $L(\frac{4}{5}, 0)$ -modules (see Table A), we have the following lemma.

Lemma 2.2

$$\begin{split} N_{W(i),W(j)}^{W(k)} &\leq N_{L(\frac{4}{5},k)\oplus L(\frac{4}{5},k')}^{L(\frac{4}{5},k)\oplus L(\frac{4}{5},k')} \leq 1\\ N_{W(i),W(j)}^{W(k,\pm)} &\leq N_{L(\frac{4}{5},i),L(\frac{4}{5},j)}^{L(\frac{4}{5},k)} = 0\\ N_{W(i,+)\oplus W(i,-),W(j,+)\oplus W(j,-)}^{W(k)} &\leq N_{L(\frac{4}{5},i),L(\frac{4}{5},j)}^{L(\frac{4}{5},k)\oplus L(\frac{4}{5},k')} \leq 2\\ N_{W(i,+)\oplus W(i,-),W(j,+)\oplus W(j,-)}^{W(k,\pm)} &\leq N_{L(\frac{4}{5},i),L(\frac{4}{5},j)}^{L(\frac{4}{5},k)} \leq 1 \end{split}$$

Let's explain how we determine the fusion rules of W(0)-modules. Since all VOAs in this paper are rational, we identify the fusion product and the tensor product. Namely, we will see $W^1 \times W^2$ as a W(0)-module for W(0)-modules W^1 and W^2 . Let T be a sub VOA of W(0) isomorphic to $L(\frac{4}{5}, 0)$. For a W(0)-module W, it is also a T-module and we denote them by $(W)_T$. As W(0)-modules, we have a fusion product $W^1 \times W^2$. Also viewing W^1 and W^2 as $L(\frac{4}{5}, 0)$ -modules, we have a fusion product $(W^1)_T \times (W^2)_T$. The above two lemmas tell that there is an injective W(0)-homomorphism π of $(W^1 \times W^2)$ into $(W^1)_T \times (W^2)_T$. We note that all fusion rules $N_{L(\frac{4}{5},i),L(\frac{4}{5},h)}^{L(\frac{4}{5},i)}$ are less than or equal to 1. We will next show $N_{W^2,W^3}^{W^1} \neq 0$ for desired W(0)-modules W^1, W^2, W^3 so that the above injection π of $(W^1 \times W^2)$ into $(W^1)_T \times (W^2)_T$ is an isomorphism. Therefore, we can determine the fusion rules.

In order to show $N_{W^2,W^3}^{W^1} \neq 0$ for some W^1, W^2, W^3 , we will use a (generalized) VOA V_L constructed from a lattice $L = \frac{\sqrt{2}}{3}A_2$. First, we quote a construction of V_S for $S \subseteq \mathbb{R}L$ for [FLM]. We note that we don't need a group extension here since all the values of inner products of elements of L are in $2\mathbb{Z}[1/3]$.

Let L be a lattice. Viewing $H = \mathbb{C}L$ as a commutative Lie algebra with a bilinear form \langle , \rangle , we define the affine Lie algebra

$$\hat{H} = H[t, t^{-1}] + \mathbb{C}k$$

associated with H and the symmetric tensor algebra $M(1) = S(\hat{H}^-)$ of \hat{H}^- , where $\hat{H}^- = H[t^{-1}]t$. As in [FLM], we shall define the Fock space $V_L = \bigoplus_{a \in L} M(1) \mathbf{e}^{\mathbf{a}}$ with the vacuum $\mathbf{1} = \mathbf{e}^{\mathbf{0}}$ and the vertex operators Y(*, z) as follows:

The vertex operator of \mathbf{e}^a is given by

$$Y(\mathbf{e}^{\mathbf{a}}, \mathbf{z}) = \exp\left(\sum_{\mathbf{n}\in\mathbb{Z}_{+}}\frac{\mathbf{a}(-\mathbf{n})}{\mathbf{n}}\mathbf{z}^{\mathbf{n}}\right)\exp\left(\sum_{\mathbf{n}\in\mathbb{Z}_{+}}\frac{\mathbf{a}(\mathbf{n})}{-\mathbf{n}}\mathbf{z}^{-\mathbf{n}}\right)\mathbf{e}^{\mathbf{a}}\mathbf{z}^{\mathbf{a}}.$$

and that of $a(-1)\mathbf{e}^{\mathbf{0}}$ is

$$Y(a(-1)\mathbf{e}^0, \mathbf{z}) = \mathbf{a}(\mathbf{z}) = \sum \mathbf{a}(\mathbf{n})\mathbf{z}^{-\mathbf{n}-1}.$$

The vertex operators of other elements are defined by the n-th normal product:

$$Y(a(n)v, z) = a(z)_n Y(v, z) = Res_x \{ (x - z)^n a(x) Y(v, z) - (-z + x)^n Y(v, z) a(x) \}.$$

Here the operator of $a \otimes t^n$ on $M(1)\mathbf{e}^{\mathbf{b}}$ are denoted by a(n) and

$$a(n)\mathbf{e}^{\mathbf{b}} = \mathbf{0} \text{ for } \mathbf{n} > \mathbf{0},$$

$$a(0)\mathbf{e}^{\mathbf{b}} = < \mathbf{a}, \mathbf{b} > \mathbf{e}^{\mathbf{b}},$$

$$e^{a}\mathbf{e}^{\mathbf{b}} = \mathbf{e}^{\mathbf{a}+\mathbf{b}},$$

$$z^{a}\mathbf{e}^{\mathbf{b}} = \mathbf{e}^{\mathbf{b}}\mathbf{z}^{<\mathbf{a},\mathbf{b}>},$$

$$(x+y)^{n} = \sum_{i=0}^{\infty} {n \choose i} x^{n-i} y^{i} \text{ and }$$

$${n \choose i} = \frac{n(n-1)\cdots(n-i+1)}{i!}.$$

We note that the above definition of vertex operator is very general, that is, it is well defined for any $v \in V_{\mathbb{R} \otimes L}$ and so we may think

$$Y(v,z) \in End(V_{\mathbb{R}\otimes L})((z,z^{-1}))$$

for $v \in V_{\mathbb{R}\otimes_{\mathbb{Z}}L}$, where $P((z, z^{-1}))$ denotes $\{\sum_{n \in \mathbb{C}} a_n z^n : a_n \in P\}$ for any set P. In particular,

$$Y(v,z)u \in V_{\mathbb{R}\otimes L}[[z,z^{-1}]]z^{\langle a,b\rangle}$$

for $v \in M(1)e^a$ and $u \in M(1)e^b$ and $a, b \in \mathbb{R}L$. Set $\mathbf{1} = \mathbf{e}^{\mathbf{0}}$. It is worth to note that if we set $Y(v, z) = \sum_{n \in \mathbb{R}} v_n z^{-n-1}$, then $v_{-1} \mathbf{e}^{\mathbf{0}} = \mathbf{v}$ for any $v \in \mathbb{R} \otimes L$. Also, for the Virasoro element, we set

$$w = \sum v^i (-1)^1 e^0$$

where $\{v^1, ..., v^k\}$ is an orthonormal basis of $\mathbb{R}L$.

For any subset S of $\mathbb{R}L$, we can define

$$V_S = \bigoplus_{a \in S} M(1) \mathbf{e}^{\mathbf{a}}.$$

The followings are obtained in Chapter 4 of [FLM].

Lemma 2.3 ([FLM])

$$\begin{split} & [\sum a(n)x^{-n-1}, Y(\mathbf{e}^{\mathbf{b}}, \mathbf{z})] \sim \mathbf{0} & \text{for any } a, b \in L, \\ & Y(\mathbf{e}^{\mathbf{a}}, \mathbf{x})\mathbf{Y}(\mathbf{e}^{\mathbf{b}}, \mathbf{z}) \sim \mathbf{Y}(\mathbf{e}^{\mathbf{b}}, \mathbf{z})\mathbf{Y}(\mathbf{e}^{\mathbf{a}}, \mathbf{x}) & \text{for } < a, b \geq \equiv 2 \pmod{2}, \\ & Y(\mathbf{e}^{\mathbf{a}}, \mathbf{x})\mathbf{Y}(\mathbf{e}^{\mathbf{b}}, \mathbf{z}) \sim -\mathbf{Y}(\mathbf{e}^{\mathbf{b}}, \mathbf{z})\mathbf{Y}(\mathbf{e}^{\mathbf{a}}, \mathbf{z}) & \text{for } < a, b \geq \equiv 1 \pmod{2}. \end{split}$$

where $a(x,z) \sim b(x,z)$ means $(z-x)^m(a(x,z)-b(x,z)) = 0$ for some $m \in \mathbb{Z}$. Especially, if $\langle a,b \rangle \in 2\mathbb{Z}$, then $[Y(v,z), Y(u,x)] \sim 0$ for $v \in M(1)e^a, u \in M(1)e^b$.

In [KMY], they studied the structure of the VOA $M^0 = V_L$ and its modules for the lattice $L = \sqrt{2}A_2$. Namely, let $\langle x, y \rangle = -2$, $\langle x, x \rangle = \langle y, y \rangle = 4$ and set $L = \mathbb{Z}x + \mathbb{Z}y$ be a lattice (of type $\sqrt{2}A_2$). It is easy to see that $M^1 = V_{\frac{x+2y}{3}+L}$ and $M^2 = V_{\frac{2x+y}{3}+L}$ are V_L -modules. Set

$$M = M^0 \oplus M^1 \oplus M^2.$$

We note that M is closed under the operators u_n of $u \in M$. It is proved by [DLMN] that the Virasoro element w of V_L is an orthogonal sum of three conformal vectors w^1 , w^2 , and w^3 with central charges $\frac{1}{2}$, $\frac{7}{10}$, and $\frac{4}{5}$, respectively. Namely, V_L contains a sub VOA T isomorphic to $L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, 0) \otimes L(\frac{4}{5}, 0)$. Viewing V as a T-module, it is a direct sum of irreducible modules of T and each irreducible T-module is isomorphic to $L(\frac{1}{2}, h_1) \otimes L(\frac{7}{10}, h_2) \otimes L(\frac{4}{5}, h_3)$ for some h_1, h_2, h_3 .

In the following argument, we recall the study in [KMY].

It is clear that $(V_L)_1 = \mathbb{C}x(-1)e^0 + \mathbb{C}y(-1)e^0$. The sum of all subspaces of $M^0 = V_L$ isomorphic to $L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, k^1) \otimes L(\frac{4}{5}, k^2)$ for any k^1, k^2 is isomorphic to a direct sum of

$$T^{1} = L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, 0) \otimes \left(L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)\right)$$

and

$$T^{2} = L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, \frac{3}{5}) \otimes \left(L(\frac{4}{5}, \frac{2}{5}) \oplus L(\frac{4}{5}, \frac{7}{5})\right).$$

Also, the sum of subspaces of M^1 isomorphic to $L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, k^1) \otimes L(\frac{4}{5}, k^2)$ is a direct sum of

$$T^{3+} = L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, 0) \otimes L(\frac{4}{5}, \frac{2}{3})$$
$$T^{4+} = L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, \frac{3}{5}) \otimes L(\frac{4}{5}, \frac{1}{15})$$

Similarly, the sum of subspaces of M^2 isomorphic to $L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, k^1) \otimes L(\frac{4}{5}, k^2)$ is a direct sum of

$$T^{3-} = L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, 0) \otimes L(\frac{4}{5}, \frac{2}{3})$$
$$T^{4-} = L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, \frac{3}{5}) \otimes L(\frac{4}{5}, \frac{1}{15}).$$

It is easy to see that $T^{n\pm}$ are contragredient (dual) modules of $T^{n\mp}$ by the natural inner product of $V_{\mathbb{Z}(\frac{x+2y}{2})+L}$.

Since $\langle L, \frac{x+2y}{3} + L \rangle \subseteq 2\mathbb{Z}$, for $v \in M^g$ with $g \in \mathbb{Z}_3$, the vertex operator Y(v, z) satisfies L(-1)-derivative property and the local commutativity with all vertex operators Y(u, z) of $u \in V_L$. By applying it to M^h for $h \in \mathbb{Z}_3$, we have an intertwining operator $Y(*, z) \in I\begin{pmatrix} M^{h+g} \\ M^g & M^h \end{pmatrix}$.

When we view these intertwining operators as intertwining operators among $L(\frac{1}{2},0) \otimes L(\frac{7}{10},0) \otimes \{L(\frac{4}{5},0) \oplus L(\frac{4}{5},3)\}$ -submodules and then as intertwining operators among $L(\frac{4}{5},0) \oplus L(\frac{4}{5},3)$ -modules, the following theorem is very useful.

Theorem 2.2 ([M1]) Let W be a sub VOA of V which may have a different Virasoro element $e \in V_2$. Assume that W is rational. Let M^1 and M^2 be irreducible W-submodules of V. Set

$$M(M^1, M^2) = \sum_{v \in M^1, u \in M^2, m \in \mathbb{Z}} \mathbb{C}v(m)u.$$

Then $M(M^1, M^2)$ is a W-module and we have $I_W\binom{M^3}{M^1 M^2} \neq 0$ for any irreducible W-submodule M^3 of $M(M^1, M^2)$. In particular, let $\{W^i : i \in I\}$ be the set of distinct

irreducible W-modules and $V = \bigoplus V^i$ be the decomposition into the direct sum of homogeneous W-modules V^i , where V^i is the sum of all irreducible W-submodules of V isomorphic to W^i . For $u \in V$, let $u = \sum u^i$, where $u^i \in V^i$. If there are $v \in V^i$, $u \in V^j$, $n \in \mathbb{Z}$ such that $(v_n u)^k \neq 0$, then $I\begin{pmatrix}V^k\\V^k\\V^i&V^j\end{pmatrix} \neq 0$.

We should note that by Theorem 2.2 and the fusion rules $L(\frac{1}{2}, 0) \times L(\frac{1}{2}, 0) = L(\frac{1}{2}, 0)$, $T^1 \oplus T^2 \oplus T^{3+} \oplus T^{3-} \oplus T^{4+} \oplus T^{4-}$ is closed by the products.

Since W(h, -) and W(i) are contragredient (dual) modules of W(h, +) and W(i), respectively, we have:

$$N_{W(h,\pm),W(h,\pm)}^{W(0)} \neq 0$$
 and $N_{W(i),W(i)}^{W(0)} \neq 0.$ (2.1)

for $h = 0, \frac{2}{5}$ and $i = \frac{2}{3}, \frac{1}{15}$.

It is easy to check that

$$(T^{2})_{1} = \mathbb{C}(x+2y)(-1)e^{0}$$

$$(T^{1})_{2} = \mathbb{C}(3x(-1)^{2}e^{0} + (x+2y)(-1)^{2}e^{0})$$

$$(T^{3+})_{2/3} = \mathbb{C}(e^{(x+2y)/3} + e^{(x-y)/3} + e^{(-2x-y)/3})$$

$$(T^{4+})_{2/3} = \mathbb{C}(2e^{(x+2y)/3} - e^{(x-y)/3} - e^{(-2x-y)/3})$$

Since

$$((x+2y)(-1)e^{0})_{-1}(x+2y)(-1)e^{0} = (x+2y)(-1)^{2}e^{0} \notin (T^{1})_{2}$$

we have

$$N_{W(\frac{2}{5}),W(\frac{2}{5}),W(\frac{2}{5})}^{W(\frac{2}{5})} \neq 0.$$
(2.2)

Also, since

$$(x+2y)(-1)\{\lambda e^{(x+2y)/3} + \mu(e^{(x-y)/3} + e^{(-2x-y)/3})\} = 4\lambda e^{(x+2y)/3} - 2\mu(e^{(x+2y)/3} + e^{(-2x-y)/3}),$$

we have :

$$N_{W(\frac{2}{5}),W(\frac{2}{3},+)}^{W(\frac{1}{15},+)} = N_{W(\frac{2}{3},+),W(\frac{1}{15},-)}^{W(\frac{2}{3},-)} = N_{W(\frac{2}{3},-)}^{W(\frac{2}{3},-)} \neq 0$$
(2.3)

and

$$N_{W(\frac{2}{5}),W(\frac{1}{15},+)}^{W(\frac{1}{15},+)} = N_{W(\frac{1}{15},+),W(\frac{1}{15},-)}^{W(\frac{2}{5})} \neq 0.$$
(2.4)

Similarly,

$$N_{W(\frac{2}{5}),W(\frac{2}{3},-)}^{W(\frac{1}{15},-)} = N_{W(\frac{2}{3},-),W(\frac{1}{15},+)}^{W(\frac{2}{5})} = N_{W(\frac{2}{3},+)}^{W(\frac{2}{3},+)} \neq 0$$
(2.5)

and

$$N_{W(\frac{2}{5}),W(\frac{1}{15},-)}^{W(\frac{1}{15},-)} = N_{W(\frac{1}{15},-),W(\frac{1}{15},+)}^{W(\frac{2}{5})} \neq 0.$$
(2.6)

It follows from the direct calculations that $e^{(x+2y)/3} + e^{(x-y)/3} + e^{(-2x-y)/3}$ and $2e^{(x+2y)/3} - (e^{(x-y)/3} + e^{(-2x-y)/3})$ are lowest degree vectors of $L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, 0) \otimes L(\frac{4}{5}, \frac{2}{3}) \subseteq M^1$ and $L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, \frac{3}{5}) \otimes L(\frac{4}{5}, \frac{1}{15}) \subseteq M^1$, respectively. Similarly, $e^{(-x-2y)/3} + e^{(-x+y)/3} + e^{(2x+y)/3}$ and $2e^{(-x-2y)/3} - (e^{(-x+y)/3} + e^{(2x+y)/3})$ are lowest degree vectors of $L(\frac{1}{2}, 0) \otimes L(\frac{7}{10}, 0) \otimes L(\frac{7}{10}, 0) \otimes L(\frac{7}{10}, \frac{3}{5}) \otimes L(\frac{4}{5}, \frac{1}{15}) \subseteq M^2$, respectively. Also, for $u = \alpha e^{(x+2y)/3} + \beta (e^{(x-y)/3} + e^{(-2x-y)/3})$ and $v = \lambda e^{(x+2y)/3} + \mu (e^{(x-y)/3} + e^{(-2x-y)/3})$, we have $u_{-1/3}v = 2\beta \mu e^{(x+2y)/3} + \beta\lambda + \alpha \mu (e^{(x-y)/3} + e^{(-2x-y)/3})$, where $u_{-1/3}$ is the grade keeping operator of u. Hence, we have

$$N_{W(\frac{2}{3},\pm),W(\frac{2}{3},\pm)}^{W(\frac{2}{3},\pm)} \neq 0,$$
(2.7)

$$N_{W(\frac{1}{15},\pm),W(\frac{1}{15},\pm)}^{W(\frac{2}{3},\mp)} = N_{W(\frac{1}{15},\pm),W(\frac{2}{3},\pm)}^{W(\frac{1}{15},\mp)} \neq 0,$$
(2.8)

$$N_{W(\frac{1}{15},\pm),W(\frac{1}{15},\pm)}^{W(\frac{1}{15},\mp)} \neq 0.$$
(2.9)

3 Fusion rule

We first list the fusion rules among $L(\frac{4}{5}, 0)$ -modules $L(\frac{4}{5}, 0)$, $L(\frac{4}{5}, 3)$, $L(\frac{4}{5}, 2/5)$, $L(\frac{4}{5}, 7/5)$, $L(\frac{4}{5}, \frac{2}{3})$ and $L(\frac{4}{5}, \frac{1}{15})$, which are the only irreducible modules we need in this section. For the fusion rules for the remaining cases, see Appendix.

0	$\frac{2}{5}$	$\frac{7}{5}$	$\frac{1}{15}$	3	$\frac{2}{3}$
$\frac{2}{5}$	$0:\frac{7}{5}$	$\frac{2}{5}:3$	$\frac{1}{15}:\frac{2}{3}$	$\frac{7}{5}$	$\frac{1}{15}$
$\frac{\frac{2}{5}}{\frac{7}{5}}$	$\frac{2}{5}:3$	$0:\frac{7}{5}$	$\frac{2}{3}:\frac{1}{15}$	$\frac{2}{5}$	$\frac{1}{15}$
$\frac{1}{15}$	$\frac{1}{15}:\frac{2}{3}$	$\frac{2}{3}:\frac{1}{15}$	$0:\frac{7}{5}:\frac{2}{3}:\frac{1}{15}:3:\frac{2}{5}$	$\frac{1}{15}$	$\frac{2}{5}:\frac{1}{15}:\frac{7}{5}$
3	$\frac{7}{5}$	$\frac{2}{5}$	$\frac{1}{15}$	0	$\frac{2}{3}$
$\frac{2}{3}$	$\frac{1}{15}$	$\frac{1}{15}$	$\frac{2}{5}:\frac{1}{15}:\frac{7}{5}$	$\frac{2}{3}$	$0:\frac{2}{3}:3$

Table A

In the table, the number h denotes $L(\frac{4}{5}, h)$ and $h: \dots: k$ denotes $L(\frac{4}{5}, h) + \dots + L(\frac{4}{5}, k)$.

By (2.5) and (2.6), $N_{W(h,\pm),W(h,\pm)}^{W(h,\mp)} \neq 0$. Hence, by the fusion rules $L(\frac{4}{5},\frac{2}{3}) \times L(\frac{4}{5},\frac{2}{3}) = L(\frac{4}{5},0) + L(\frac{4}{5},3) + L(\frac{4}{5},\frac{2}{3})$ of $L(\frac{4}{5},0)$ -modules and (2.1) and (2.7), we have

$$W(\frac{2}{3}, \pm) \times W(\frac{2}{3}, \pm) = W(\frac{2}{3}, \mp)$$

$$W(\frac{2}{3}, \pm) \times W(\frac{2}{3}, \mp) = W(0).$$
(3.1)

Similarly, by the fusion rules $L(\frac{4}{5}, \frac{1}{15}) \times L(\frac{4}{5}, \frac{1}{15}) = L(\frac{4}{5}, 0) + L(\frac{4}{5}, 3) + L(\frac{4}{5}, \frac{2}{3}) + L(\frac{4}{5}, \frac{2}{5}) + L(\frac{4}{5}, \frac{7}{5})$ of $L(\frac{4}{5}, 0)$ -modules and (2.1),(2.4),(2.8) and (2.9), we have

$$W(\frac{1}{15}, \pm) \times W(\frac{1}{15}, \pm) = W(\frac{1}{15}, \mp) + W(\frac{2}{3}, \mp)$$

$$W(\frac{1}{15}, \pm) \times W(\frac{1}{15}, \mp) = W(0) + W(\frac{2}{5})$$
(3.2)

By the fusion rules $L(\frac{4}{5},\frac{2}{5}) \times L(\frac{4}{5},\frac{2}{3}) = L(\frac{4}{5},\frac{1}{15})$ and (2.3) and (2.5), we have

$$W(\frac{2}{5}) \times W(\frac{2}{3}, \pm) = W(\frac{1}{15}, \pm).$$
 (3.3)

By the fusion rules $L(\frac{4}{5}, \frac{2}{5}) \times L(\frac{4}{5}, \frac{1}{15}) = L(\frac{4}{5}, \frac{1}{15}) + L(\frac{4}{5}, \frac{2}{3})$ and (2.3) ~ (2.6), we have

$$W(\frac{2}{5}) \times W(\frac{1}{15}, \pm) = W(\frac{1}{15}, \pm) + W(\frac{2}{3}, \pm).$$
 (3.4)

By the fusion rules $L(\frac{4}{5},\frac{2}{5}) \times L(\frac{4}{5},\frac{2}{5}) = L(\frac{4}{5},0) + L(\frac{4}{5},\frac{7}{5})$ and (2.1) and (2.2), we have

$$W(\frac{2}{5}) \times W(\frac{2}{5}) = W(0) + W(\frac{2}{5}).$$
 (3.5)

By the fusion rules $L(\frac{4}{5}, \frac{2}{3}) \times L(\frac{4}{5}, \frac{1}{15}) = L(\frac{4}{5}, \frac{2}{5}) + L(\frac{4}{5}, \frac{7}{5}) + L(\frac{4}{5}, \frac{1}{15})$ and (2.3),(2.5) and (2.8), we have

$$W(\frac{2}{3},\pm) \times W(\frac{1}{15},\pm) = W(\frac{1}{15},\mp)$$

$$W(\frac{2}{3},\pm) \times W(\frac{1}{15},\mp) = W(\frac{2}{5}).$$
(3.6)

We put the above fusion rules in the following table.

Table B

W(0)	$W(\frac{2}{5})$	$W(\frac{2}{3}, +)$	$W(\frac{1}{15}, +)$	$W(\frac{2}{3}, -)$	$W(\frac{1}{15})$
$W(\frac{2}{5})$	$W(0): W(\frac{2}{5})$	$W(\frac{1}{15}, +)$	$W(\frac{1}{15},+):W(\frac{2}{3},+)$	$W(\frac{1}{15}, -)$	$W(\frac{1}{15},-):W(\frac{2}{3},-)$
$W(\frac{2}{3}, +)$	$W(\frac{1}{15}, +)$	$W(\frac{2}{3}, -)$	$W(\frac{1}{15}, -)$	W(0)	$W(\frac{2}{5})$
$W(\frac{1}{15}, +)$	$W(\frac{1}{15},+):W(\frac{2}{3},+)$	$W(\frac{1}{15}, -)$	$W(\frac{1}{15},-):W(\frac{2}{3},-)$	$W(\frac{2}{5})$	$W(0): W(\frac{2}{5})$
$W(\frac{2}{3}, -)$	$W(\frac{1}{15}, -)$	W(0)	$W(\frac{2}{5})$	$W(\frac{2}{3}, +)$	$W(\frac{1}{15}, +)$
$W(\frac{1}{15}, -)$	$W(\frac{1}{15},-):W(\frac{2}{3},-)$	$W(\frac{2}{5})$	$W(0): W(\frac{2}{5})$	$W(\frac{1}{15}, +)$	$W(\frac{1}{15},+):W(\frac{2}{3},+)$

4 Automorphisms

As we showed in [M2], if a VOA contains $L(\frac{4}{5}, 0)$, then we have an automorphism σ of at most 2 given by

$$\sigma: \begin{cases} 1 & on \quad L(\frac{4}{5},0), L(\frac{4}{5},3), L(\frac{4}{5},\frac{2}{3}), L(\frac{4}{5},\frac{2}{5}), L(\frac{4}{5},\frac{1}{15}), L(\frac{4}{5},\frac{7}{5}) \\ -1 & on \quad L(\frac{4}{5},\frac{1}{8}), L(\frac{4}{5},\frac{13}{8}), L(\frac{4}{5},\frac{1}{40}), L(\frac{4}{5},\frac{21}{40}) \end{cases}$$

So we next think about the case $\sigma = 1$ or the space V^{σ} of σ -invariants. In this case, there are no $L(\frac{4}{5}, 0)$ -submodules isomorphic to $L(\frac{4}{5}, \frac{1}{8}), L(\frac{4}{5}, \frac{13}{8}), L(\frac{4}{5}, \frac{1}{40})$ or $L(\frac{4}{5}, \frac{21}{40})$. We next assume that V contains $L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$. We should note that if V contains $L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$, then there are no $L(\frac{4}{5}, 0)$ -submodules in V isomorphic to $L(\frac{4}{5}, \frac{1}{8}), L(\frac{4}{5}, \frac{13}{8}), L(\frac{4}{5}, \frac{13}{8}), L(\frac{4}{5}, \frac{13}{8})$

or $L(\frac{4}{5}, \frac{21}{40})$.

Theorem A If a VOA V contains $L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$, then an endomorphism τ of V defined by

$$\tau : \begin{cases} 1 & on \quad W(0) \text{ and } W(\frac{2}{5}) \\ e^{2\pi i/3} & on \quad W(\frac{2}{3}, +) \text{ and } W(\frac{1}{15}, +) \\ e^{4\pi i/3} & on \quad W(\frac{2}{3}, -) \text{ and } W(\frac{1}{15}, -) \end{cases}$$

is an automorphism of V.

[Proof] Replacing W(i) and W(h, +) and W(k, -) in the table (B) by 1 and $e^{2\pi i/3}$ and $e^{4\pi i/3}$, we have

1	1	$e^{2\pi i/3}$	$e^{2\pi i/3}$	$e^{4\pi i/3}$	$e^{4\pi i/3}$
1	1:1	$e^{2\pi i/3}$	$e^{2\pi i/3}:e^{2\pi i/3}$	$e^{4\pi i/3}$	$e^{4\pi i/3}:e^{4\pi i/3}$
$e^{2\pi i/3}$	$e^{2\pi i/3}$	$e^{4\pi i/3}$	$e^{4\pi i/3}$	1	1
$e^{2\pi i/3}$	$e^{2\pi i/3}:e^{2\pi i/3}$	$e^{4\pi i/3}$	$e^{4\pi i/3}$: $e^{4\pi i/3}$	1	1:1
$e^{4\pi i/3}$	$e^{4\pi i/3}$	1	1	$e^{2\pi i/3}$	$e^{2\pi i/3}$
$e^{4\pi i/3}$	$e^{4\pi i/3}:e^{4\pi i/3}$	1	1:1	$e^{2\pi i/3}$	$e^{2\pi i/3}:e^{2\pi i/3}$

which is compatible with the products. Hence, by Theorem 2.2, τ is an automorphism of V.

Q.E.D.

If $\tau_W = 1$, then all *T*-submodules of *V* are isomorphic to $L(\frac{4}{5}, 0)$, $L(\frac{4}{5}, 3)$, $L(\frac{4}{5}, \frac{2}{5})$ or $L(\frac{4}{5}, \frac{7}{5})$ for $T \subseteq W$ and $T \cong L(\frac{4}{5}, 0)$. In this case, we can define another automorphism μ_T of *V* as follows:

Theorem B Assume that V contains a sub VOA T isomorphic to $L(\frac{4}{5}, 0)$ and all T-submodules of V are isomorphic to $L(\frac{4}{5}, 0)$, $L(\frac{4}{5}, 3)$, $L(\frac{4}{5}, \frac{2}{5})$ or $L(\frac{4}{5}, \frac{7}{5})$. Then the endomorphism μ_T defined by

$$\mu_T : \begin{cases} 1 & on \quad L(\frac{4}{5}, 0) \text{ and } L(\frac{4}{5}, \frac{7}{5}) \\ -1 & on \quad L(\frac{4}{5}, 3) \text{ and } L(\frac{4}{5}, \frac{2}{5}) \end{cases}$$

is an automorphism of V.

5 V as a sub VOA-module

The notion of sub VOAs of V in this paper is not the same as in [FZ], where they expected sub VOA W to have the same Virasoro element with V. Our definition of sub VOAs is: $(W = \oplus W_n, Y^W, w_W, \mathbf{1}_W)$ is a sub VOA of V if

(1) $(W, Y^W, w_W, \mathbf{1}_W)$ is a VOA,

- (2) $W \subseteq V$ and $W_n = W \cap V_n$,
- (3) $\mathbf{1}_{\mathbf{W}} = \mathbf{1}_{\mathbf{V}}$ and
- (4) $Y^{W}(v, z) = Y(v, z)|_{W}$ for $v \in W$.

There are several definitions for VOA-modules, but we will include an infinite direct sum of irreducible modules as a VOA-module M. Namely, we don't assume dim $M_n < \infty$.

Let W be a sub VOA of V and e a Virasoro element of W. Different from the ordinary algebras, it is not obvious that V is a W-module. The problem is whether e_1 acts on V diagonally or not.

The purpose of this section is to show that V is a W-module for W in our cases.

Let V be a VOA and W a sub VOA. Let w and e be Virasoro elements of V and W, respectively.

In particular, $e \in V_2$ and so e_1 keeps the grade of V. By the assumption, $f_1 = w_1 - e_1$ acts on W as 0. Furthermore, for $v \in W Y(e_0v, z)|_W = \frac{d}{dz}Y(v, z)|_W = Y(w_0v, z)|_W$ and so $Y(w_0v - e_0v, z)|_W = 0$. In particular, $w_0v - e_0v = (w_0v - e_0v)_{-1}\mathbf{1} = 0$ and so, $f_0 = w_0 - e_0$ acts on W as 0. Hence we have:

Lemma 5.1 Both f_1 and f_0 commutes v_m on V for $v \in W$ and $m \in \mathbb{Z}$.

[**Proof**] It follows from
$$[f_0, v_m] = (f_0 v)_m = 0$$
 and $[f_1, v_m] = (f_0 v)_{m+1} + (f_1 v)_m = 0$.
Q.E.D.

The main purpose in this section is to prove the following theorem:

Theorem 5.1 If W is rational, then V is a W-module.

[Proof] Define a module vertex operator $Y^V(v, z)$ of $v \in W$ by the vertex operator of $v \in V$. Clearly, they satisfy the local commutativity and the e_0 -derivative property:

$$Y^{V}(e_{0}v, z) = Y^{V}((f_{0} + e_{0})v, z) = Y(w_{0}v, z) = \frac{d}{dz}Y(v, z)$$

for $v \in W$. Hence, what we have to do is to prove that V is a direct sum of eigenspaces of e_1 . Suppose false. Since f_1 commutes with all v_n for $v \in W$, the eigenspace V_{λ} and the generalized eigenspace $T_{\lambda} = \{v \in V | \exists n \in \mathbb{Z} \ (f_1 - \lambda)^n v = 0\}$ of f_1 with eigenvalue λ is invariant under the actions v_n of $v \in W$. We first prove that the eigenspace V_{λ} is a direct sum of irreducible W-modules. Since f_1 keeps the grade, it acts on V_n and so we have $V_{\lambda} = \bigoplus (V_{\lambda})_n$, where $(V_{\lambda})_n = V_n \cap V_{\lambda}$. Since $e_1 = (w_1 - f_1)$ acts on $(V_{\lambda})_n$ as $n - \lambda$, V_{λ} is a W-module. Since W is rational by the assumption, V_{λ} is a direct sum of irreducible W-modules.

Since f_1 acts on each finite dimensional homogenous spaces V_n , V is a direct sum of generalized eigenspaces of f_1 . Hence, there are λ and n such that $T_{\lambda} \cap V_n \neq V_{\lambda} \cap V_n$. Take n as a minimal one.

As we explained as above, v_n acts on T_{λ}/V_{λ} for $v \in W$ and the eigenspace X_{λ} of f_1 in T_{λ}/V_{λ} is a W-module. By the choice of n, $(X_{\lambda})_n \neq 0$. Let \overline{X} be an irreducible Wsubmodule of X_{λ} whose lowest degree is n. We should note that since V_{λ} is an eigenspace of f_1 and $e_1 = w_1 - f_1$, the lowest eigenspace of w_1 in V_{λ} is the lowest eigenspace of e_1 .

Hence there is an irreducible W-submodule \overline{B} of T_{λ}/V_{λ} whose lowest degree space \overline{B}_0 is in $(V_n + V_{\lambda})/V_{\lambda}$ since W is rational. Let B be its inverse image. Clearly, B contains V_{λ} and f_1 does not act on B diagonally. Let S is a submodule of V_{λ} generated by Wsubmodules which are not isomorphic to \overline{B} . Since $(f_1 - \lambda)B \neq 0$ and all submodule of $(f_1 - \lambda)B$ is isomorphic to \overline{B} , all composition factors of B/S is isomorphic to \overline{B} as W-modules and we have $S \neq V_{\lambda}$. In particular, $(B/S) \cap (V_m + S/S) = 0$ for all m < n. We next show that Zhu-algebra A(W) acts on the top module $(B_n + S)/S$ of B/S.

In order to prove the above assertion, we will use an idea for Zhu-algebra in [Z]. We will treat a general case for a while. Let A(V) = V/O(V) be the Zhu-algebra of V. For $v \in V$, o(v) denotes the grade keeping operator of v. For a homogeneous element $v \in V_m$, if $Y^M(v,z) = \sum v_i z^{-i-1}$ is a module vertex operator, then $o(v) = v_{m-1}$. It actually depends on the module M, but we write $o(v) = v_{m-1}$ formally.

Let R be the ring generated by all o(v) of $v \in V$. Let I be a subspace of R generated by the elements $\sigma = v_{i_1}^1 \cdots v_{i_r}^r \in R$ satisfying that $v_{i_s}^s \cdots v_{i_r}^r$ decreases the grade for some $1 \leq s \leq r$. We permit an infinite sum of such elements if it is well-defined in R. Clearly, I is a two-sided ideal of R. It is known that A(V) = R/I, see [Z].

Let's go back to the proof. Since W is rational, A(W) is a semi-simple. Let $\phi = v_{i_1}^1 \cdots v_{i_t}^t \in R$ and assume that $v_{i_s}^s \cdots v_{i_t}^t$ decreases the grade on W. Since the grade on W is the same as that on V, ϕ acts on $(B_n + S)/S$ trivially. Hence A(W) acts on $(B_n + S)/S$. Since A(W) is semi-simple, $(B_n + S)/S$ is a direct sum of irreducible A(W)-modules. By the definition of B and S, $(B_n + S)/S$ is a homogeneous A(W)-module. Since e_1 is in the center of A(W), e_1 acts on $(B_n + S)/S$ as a scalar times and so does f_1 , which contradicts to the facts that $(f_1 - \lambda)(B_n + S)/S \neq 0$.

Hence, V is a direct sum of eigenspaces of f_1 and so e_1 acts on V diagonally. This completes the proof of Theorem 5.1.

Q.E.D.

6 Appendix

6.1 Fusion rules of irreducible $L(\frac{4}{5}, 0)$ -modules

For $L(\frac{4}{5}, 0)$, the following fusion rules are known, see [W]. In the following table, the numbers h denote $L(\frac{4}{5}, h)$ and $h_1 : \ldots : h_t$ denotes $L(\frac{4}{5}, h_1) + \ldots + h(\frac{4}{5}, h_t)$.

0	$\frac{2}{5}$	$\frac{1}{40}$	$\frac{7}{5}$	$\frac{21}{40}$	$\frac{1}{15}$	3	$\frac{13}{8}$	$\frac{2}{3}$	$\frac{1}{8}$
$\frac{2}{5}$	$0:\frac{7}{5}$	$\frac{1}{8}:\frac{21}{40}$	$\frac{2}{5}:3$	$\frac{1}{40}:\frac{13}{8}$	$\frac{1}{15}:\frac{2}{3}$	$\frac{7}{5}$	$\frac{21}{40}$	$\frac{1}{15}$	$\frac{1}{40}$
$\frac{1}{40}$	$\frac{1}{8}:\frac{21}{40}$	$0:\frac{7}{5}:\frac{2}{3}:\frac{1}{15}$	$\frac{1}{40}:\frac{13}{8}$	$\frac{2}{5}:3:\frac{1}{15}:\frac{2}{3}$	$\frac{1}{40}:\frac{13}{8}:\frac{21}{40}:\frac{1}{8}$	$\frac{21}{40}$	$\frac{7}{5}:\frac{1}{15}$	$\frac{21}{40}$: $\frac{1}{40}$	$\frac{1}{15}:\frac{2}{5}$
$\frac{7}{5}$	$\frac{2}{5}:3$	$\frac{1}{40}:\frac{13}{8}$	$0:\frac{7}{5}$	$\frac{1}{8}:\frac{21}{40}$	$\frac{2}{3}:\frac{1}{15}$	$\frac{2}{5}$	$\frac{1}{40}$	$\frac{1}{15}$	$\frac{21}{40}$
$\frac{21}{40}$	$\frac{1}{40}:\frac{13}{8}$	$\frac{2}{5}:3:\frac{1}{15}:\frac{2}{3}$	$\frac{1}{8}:\frac{21}{40}$	$0:\frac{7}{5}:\frac{2}{3}:\frac{1}{15}$	$\frac{1}{8}:\frac{21}{40}:\frac{13}{8}:\frac{1}{40}$	$\frac{1}{40}$	$\frac{2}{5}:\frac{1}{15}$	$\frac{1}{40}$: $\frac{21}{40}$	$\frac{1}{15}:\frac{7}{5}$
$\frac{1}{15}$	$\frac{1}{15}:\frac{2}{3}$	$\frac{1}{40}$: $\frac{13}{8}$: $\frac{21}{40}$: $\frac{1}{8}$	$\frac{2}{3}:\frac{1}{15}$	$\frac{1}{8}:\frac{21}{40}:\frac{13}{8}:\frac{1}{40}$	$0:\frac{7}{5}:\frac{2}{3}:\frac{1}{15}:3:\frac{2}{5}$	$\frac{1}{15}$	$\frac{1}{40}:\frac{21}{40}$	$\frac{2}{5}:\frac{1}{15}:\frac{7}{5}$	$\frac{1}{40}$: $\frac{21}{40}$
3	$\frac{7}{5}$	$\frac{21}{40}$	$\frac{2}{5}$	$\frac{1}{40}$	$\frac{1}{15}$	0	$\frac{1}{8}$	$\frac{2}{3}$	$\frac{13}{8}$
$\frac{13}{8}$	$\frac{21}{40}$	$\frac{7}{5}:\frac{1}{15}$	$\frac{1}{40}$	$\frac{2}{5}:\frac{1}{15}$	$\frac{1}{40}:\frac{21}{40}$	$\frac{1}{8}$	$0:\frac{2}{3}$	$\frac{1}{8}:\frac{13}{8}$	$\frac{2}{3}:3$
$\frac{2}{3}$	$\frac{1}{15}$	$\frac{21}{40}$: $\frac{1}{40}$	$\frac{1}{15}$	$\frac{1}{40}$: $\frac{21}{40}$	$\frac{2}{5}:\frac{1}{15}:\frac{7}{5}$	$\frac{2}{3}$	$\frac{1}{8}:\frac{13}{8}$	$0:\frac{2}{3}:3$	$\frac{1}{8}:\frac{13}{8}$
$\frac{1}{8}$	$\frac{1}{40}$	$\frac{1}{15}:\frac{2}{5}$	$\frac{21}{40}$	$\frac{1}{15}:\frac{7}{5}$	$\frac{1}{40}:\frac{21}{40}$	$\frac{13}{8}$	$\frac{2}{3}:3$	$\frac{1}{8}:\frac{13}{8}$	$0:\frac{2}{3}$

Table (

References

- [B] R.E. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, Proc. Natl. Acad. Sci. USA 83 (1986), 3068–3071.
- [C] J.H. Conway, A simple construction of the Fischer-Griess monster group, Invent. Math. 79 (1985), no. 3, 513–540.
- [CS] J.H. Conway and N.J.A. Sloane: "Sphere Packings, Lattices and Groups", Springer-Verlag, 1988.
- [DL] C. Dong and J. Lepowsky, Generalized vertex algebras and relative vertex operators, Progress in Mathematics, 112. Birkhäuser Boston, Inc., Boston, 1993.

- [DLMN] C. Dong, H. Li, G. Mason and S. Norton, Associative subalgebras of the Griess algebra and related topics, preprint.
- [DM] C. Dong and G. Mason, On quantum Galois theory, *Duke Math. J.* 86 (1997), no.2, 305–321.
- [DMZ] C. Dong, G. Mason and Y. Zhu, Discrete series of the Virasoro algebra and the moonshine module, Proc. Symp. Pure. Math., American Math. Soc. 56 II (1994), 295-316.
- [FHL] I. Frenkel, Y.-Z. Huang and J. Lepowsky, "On axiomatic approaches to vertex operator algebras and modules", Memoirs Amer. Math. Soc. 104, 1993.
- [FLM] I.B. Frenkel, J. Lepowsky, and A. Meurman, Vertex Operator Algebras and the Monster, Pure and Applied Math., Vol. 134, Academic Press, 1988.
- [FZ] I. Frenkel and Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, *Duke Math. J.* 66 (1992), 123-168.
- [KMY] M. Kitazume, M. Miyamoto and H. Yamada, Ternary codes and VOA, preprint
- [Li1] Hai-sheng Li, An approach to tensor product theory for representations of a vertex operator algebra, to appear.
- [M1] M. Miyamoto, Griess algebras and conformal vectors in vertex operator algebras, J. Algebra **179**, (1996) 523-548
- [M2] M. Miyamoto, Binary codes and vertex operator (super)algebras, J. Algebra 181, (1996) 207-222
- [M3] M. Miyamoto, Representation theory of code VOA, J. Algebra, to appear.
- [W] W. Wang, Rationality of Virasoro vertex operator algebra, Duke Math. J. IMRN, Vol. 71, No. 1 (1993), 197-211.
- [Z] Y.-C. Zhu, Vertex operator algebra, elliptic functions and modular forms, Ph.D. thesis, Yale University, 1990.