

Discretizations of the Schrödinger equation with quantum algebra symmetry

Angel Ballesteros, Francisco J. Herranz¹ and Preeti Parashar

*Departamento de Física, Universidad de Burgos
Pza. Misael Bañuelos, E-09001-Burgos, Spain*

Abstract

Two quantum Hopf structures for the Schrödinger algebra as well as their corresponding differential-difference realizations are presented. For each case a (space or time) discretization of the Schrödinger equation is deduced and the quantum Schrödinger generators are shown to be symmetry operators.

¹Communication presented in the 5th Wigner Symposium, Vienna, 1997

1 A space discretization of the Schrödinger equation

The (1+1)-dimensional Schrödinger algebra [1, 2] \mathcal{S} is generated by the time translation H , space translation P , Galilean boost K , dilation D , conformal transformation C and mass operator M . The non-zero Lie brackets of \mathcal{S} are:

$$\begin{aligned} [D, P] &= -P & [D, K] &= K & [K, P] &= M & [K, H] &= P \\ [D, H] &= -2H & [D, C] &= 2C & [H, C] &= D & [P, C] &= -K. \end{aligned} \quad (1.1)$$

The usual differential realization of the Schrödinger generators in terms of the time and space coordinates (t, x) and two constants a, m , is given by:

$$\begin{aligned} H &= \partial_t & P &= \partial_x & M &= m & K &= -t\partial_x - mx \\ D &= 2t\partial_t + x\partial_x - a & C &= t^2\partial_t + tx\partial_x - at + \frac{m}{2}x^2. \end{aligned} \quad (1.2)$$

The heat/Schrödinger equation (SE) can be obtained from this representation as the Casimir $E = P^2 - 2MH$ of the Galilei subalgebra $\{K, H, P, M\}$ of \mathcal{S} :

$$(\partial_x^2 - 2m\partial_t)\phi(x, t) = 0. \quad (1.3)$$

We say that an operator S is a symmetry of E if S transforms solutions into solutions: $(ES)\phi(x, t) = (\Lambda S)\phi(x, t)$, where Λ is another operator. The Galilei generators are symmetries S of the SE with $\Lambda = S$. The dilation is also a symmetry, since (1.1) implies that $[E, D] = 2E$. Finally, we compute $[E, C] = -(KP + PK + 2MD)$ and by introducing the representation (1.2) with $a = -1/2$ we obtain that $[E, C] = 2tE$.

Let $U_z^{(s)}(\mathcal{S})$ be the quantum Schrödinger algebra [3] whose (nonstandard) classical r -matrix is $r = zP \wedge (D + \frac{1}{2}M)$; its coproduct and commutators are

$$\begin{aligned} \Delta(P) &= 1 \otimes P + P \otimes 1 & \Delta(M) &= 1 \otimes M + M \otimes 1 \\ \Delta(H) &= 1 \otimes H + H \otimes e^{-2zP} \\ \Delta(K) &= 1 \otimes K + K \otimes e^{zP} - zD' \otimes e^{zP}M \\ \Delta(D) &= 1 \otimes D + D \otimes e^{zP} + \frac{1}{2}M \otimes (e^{zP} - 1) \\ \Delta(C) &= 1 \otimes C + C \otimes e^{2zP} + \frac{z}{2}K \otimes e^{zP}D' - \frac{z}{2}D' \otimes e^{zP}(K + zD'M), \end{aligned} \quad (1.4)$$

$$\begin{aligned} [D, P] &= \frac{1}{z}(1 - e^{zP}) & [D, K] &= K & [K, P] &= Me^{zP} & [M, \cdot] &= 0 \\ [D, H] &= -2H & [D, C] &= 2C - \frac{z}{2}KD' & [H, P] &= 0 \\ [H, C] &= \frac{1}{2}(1 + e^{-zP})D' - \frac{1}{2}M + zKH & [K, C] &= -\frac{z}{2}K^2 \\ [P, C] &= -\frac{1}{2}(1 + e^{zP})K - \frac{z}{2}e^{zP}MD' & [K, H] &= \frac{1}{z}(1 - e^{-zP}), \end{aligned} \quad (1.5)$$

where $D' = D + \frac{1}{2}M$. We deduce the quantum SE with $U_z^{(s)}(\mathcal{S})$ as its quantum symmetry algebra. A differential-difference realization of (1.5) reads

$$\begin{aligned} P &= \partial_x & H &= \partial_t & M &= m \\ K &= -t\frac{1}{z}(1 - e^{-z\partial_x}) - mxe^{z\partial_x} & D &= 2t\partial_t + x\frac{1}{z}(e^{z\partial_x} - 1) - a \\ C &= t^2\partial_t e^{-z\partial_x} + tx\left(\frac{1}{z}\sinh z\partial_x - zm\partial_t e^{z\partial_x}\right) + \frac{1}{2}mx^2 e^{z\partial_x} \\ &\quad - \frac{1}{2}t\left\{a(1 + e^{-z\partial_x}) + (1 + \frac{1}{2}m)(1 - e^{-z\partial_x})\right\} \\ &\quad + \frac{1}{2}zm(1 + a - \frac{1}{2}m)xe^{z\partial_x}. \end{aligned} \quad (1.6)$$

At the level of commutation rules (1.5), the Galilei structure remains as a subalgebra and its deformed Casimir is $E_z = \frac{1}{z^2}(1 - e^{-zP})^2 - 2MH$. We define a space discretization of the SE as the action of E_z on $\phi(x, t)$ through (1.6), that is,

$$\left(\frac{1}{z^2}(1 - e^{-z\partial_x})^2 - 2m\partial_t\right)\phi(x, t) = 0. \quad (1.7)$$

By construction, the Galilei generators $\{K, H, P, M\}$ are symmetries of E_z . For the dilation we obtain from the commutation rules (1.5) that $[E_z, D] = 2E_z$, so it is also a symmetry. For the conformal transformation we have

$$[E_z, C] = -K(1 - e^{-2zP})/z + M - 2MD - 2zMKH. \quad (1.8)$$

By introducing the quantum realization (1.6) we get

$$[E_z, C] = \left\{t(e^{-z\partial_x} + 1) - zmx e^{z\partial_x}\right\} E_z + m(1 + 2a). \quad (1.9)$$

Therefore we have to set $a = -1/2$ in order to obtain C as a symmetry.

2 A time discretization of the Schrödinger equation

We consider now the quantum Schrödinger algebra [4] $U_z^{(t)}(\mathcal{S})$, associated with the (nonstandard) classical r -matrix $r = zH \wedge (D + \frac{1}{2}M) \equiv zH \wedge D'$:

$$\begin{aligned} \Delta(H) &= 1 \otimes H + H \otimes 1 & \Delta(M) &= 1 \otimes M + M \otimes 1 \\ \Delta(P) &= 1 \otimes P + P \otimes e^{-zH} \\ \Delta(K) &= 1 \otimes K + K \otimes e^{zH} - zD' \otimes e^{2zH}P \\ \Delta(D) &= 1 \otimes D + D \otimes e^{2zH} + \frac{1}{2}M \otimes (e^{2zH} - 1) \\ \Delta(C) &= 1 \otimes C + C \otimes e^{2zH} - \frac{z}{2}D' \otimes e^{2zH}M, \end{aligned} \quad (2.10)$$

$$\begin{aligned} [D, P] &= -P & [D, K] &= K & [K, P] &= M & [M, \cdot] &= 0 \\ [D, H] &= \frac{1}{z}(1 - e^{2zH}) & [D, C] &= 2C + z(D')^2 & [H, P] &= 0 \\ [H, C] &= D + \frac{1}{2}M(1 - e^{2zH}) & [K, C] &= \frac{z}{2}(DK + KD + KM) \\ [P, C] &= -K - \frac{z}{2}(DP + PD + PM) & [K, H] &= e^{2zH}P. \end{aligned} \quad (2.11)$$

A realization of $U_z^{(t)}(\mathcal{S})$ reads ($b = m/2 - 2$):

$$\begin{aligned} H &= \partial_t & P &= \partial_x & M &= m \\ K &= -(t + 2z)e^{2z\partial_t}\partial_x - mx & D &= (t + 2z)\frac{1}{z}(e^{2z\partial_t} - 1) + x\partial_x - a \\ C &= (t^2 - 2zbt)\frac{1}{2z}(e^{2z\partial_t} - 1) + tx\partial_x - at + \frac{m}{2}x^2 - 2z(b + 1)e^{2z\partial_t} \\ &\quad - \frac{z}{2}x^2\partial_x^2 - z(b - a + \frac{1}{2})x\partial_x - \frac{z}{2}(b - a)^2. \end{aligned} \quad (2.12)$$

The deformed Casimir for the Galilei sector is $E_z = P^2 - M\frac{1}{z}(1 - e^{-2zH})$; its action on $\phi(x, t)$ through (2.12) leads to a time discretization of the SE:

$$\left(\partial_x^2 - m\frac{1}{z}(1 - e^{-2z\partial_t})\right)\phi(x, t) = 0. \quad (2.13)$$

As in Sec. 1 the generators $\{K, H, P, M, D\}$ are easily shown to be symmetries of this SE. For the conformal transformation, we find that

$$[E_z, C] = -(KP + PK + 2MD e^{-2zH}) + M(M + 2)(1 - e^{-2zH}) - z(DP^2 + 2PD P + P^2 D + 2P^2 M)/2. \quad (2.14)$$

Now adopting the quantum realization (2.12):

$$[E_z, C] = 2(t + 2z - zx\partial_x)E_z - z\{m + 2(1 - a)\}\partial_x^2 + m(1 + 2ae^{-2z\partial_t}) + m(m + 2)(1 - e^{-2z\partial_t}), \quad (2.15)$$

and setting again $a = -\frac{1}{2}$ gives us $[E_z, C] = 2\left\{t + \frac{z}{2}(1 - m - 2x\partial_x)\right\}E_z$.

Finally, we recall that there exist other quantum SE [5, 6, 7] although without a known associated Hopf structure.

Acknowledgments

A.B. and F.J.H. have been partially supported by DGICYT (Project PB94-1115) from the Ministerio de Educación y Cultura de España and by Junta de Castilla y León (Projects CO1/396 and CO2/297). P.P. acknowledges A.E.C.I. for the award of a fellowship.

References

- [1] C.R. Hagen, *Phys. Rev. D* **5**, 377 (1972).
- [2] U. Niederer, *Helv. Phys. Acta* **45**, 802 (1972).
- [3] A. Ballesteros *et al.*, A Jordanian quantum two-photon/Schrödinger algebra, *J. Phys. A* to appear.
- [4] A. Ballesteros *et al.*, q-alg/9709009.
- [5] R. Floreanini and L. Vinet, *Lett. Math. Phys.* **32**, 37 (1994).
- [6] R. Floreanini and L. Vinet, *J. Math. Phys.* **36**, 3134 (1995).
- [7] V.K. Dobrev *et al.*, *J. Phys. A* **29**, 5909 (1996).