

Species Numbers versus Area in Chowdhury Ecosystems

Dietrich Stauffer* and Andrzej Pękalski**

Department of Genetics, Institute of Microbiology, University of Wrocław,
ul. Przybyszewskiego 63/77, PL-54148 Wrocław, Poland; visiting from Institute for Theoretical Physics, Cologne University, D-50923 Köln, Euroland.
e-mail: stauffer@thp.uni-koeln.de

**Institute of Theoretical Physics, University of Wrocław, pl. M.Borna
9, PL-51-204 Wrocław, Poland. e-mail: apekal@ift.uni.wroc.pl

Abstract: For a fixed number of food levels on a lattice, the Chowdhury model for ecosystem is simulated and found to give a number of different species which approaches exponentially a plateau if the considered area is increased.

The Chowdhury model [1] is one of several individual-based models for ecosystems with many (prey-predator) species [2, 3]. We use it here with the parameters of [4], that means a low speciation rate 0.001, and low diffusivity 0.001, and a fixed number (usually six) of trophic layers. We refer to [1, 4, 5] for more details on this model. Here we check if the number of different species which are found in a square of size $R \times R$ increases with a small power of the area R^2 , as summarized in [6].

So after some equilibration time (one sixth of the total number of iterations) we sum at each iteration over all different species found in a corner of size $R \times R$ of the $L \times L$ lattice, with $1 \leq R \leq L$ and $10 \leq L \leq 100$. To see better the trends with increasing R we look at the R -derivative of this number of species, that means we determine how many new species were found in a square of side length R which were not already found in its inner square of side length $R - 1$. A species in this asexual model is defined to consist of all animals occupying the corresponding ecological niches on the considered lattice sites.

Figure 1 shows that this number of new species decays roughly exponentially with increasing R (though for small R also a power law fits). Thus the total number of species approaches exponentially a plateau for large R , in contrast to the desired power law [6]. This $\exp(-R/\xi)$ decay with R is found to be independent of the simulation time t , Fig.2a, while the decay length ξ increases when the number of food levels increases from 3 to 7, Fig.2b.




Figure 1: Number of new species in R -squares which are not yet found in the inner squares of size $R - 1$, summed over 20,000 time steps (iterations) for lattices of linear dimension $L = 10$ (+), 20 (x), 50 (stars), and 100 (squares) from bottom to top (one sample each, except sum over ten samples for $L = 10$). The counted numbers are divided by the number $2R - 1$ of involved lattice sites and rounded down to integer values. Each lattice carried six food levels with 1 to 32 ecological niches.

An inhomogeneous ecosystem with different but fixed numbers of food levels thus might give a power law as a superposition of many exponential laws.

We thank PapaSmorf Cebrat for hospitality during our collaboration, and the Wrocław-Köln university partnership for partial support.

References

- [1] D. Chowdhury and D. Stauffer, J. Biosci. (India) 30, 377 (2005); D. Chowdhury, D. Stauffer and A.Kunwar, Phys. Rev. Lett. 90, 068101 (2003).

- [2] M.A. Droz and A. Pękalski, *Physica A* 336, 86 (2005).
- [3] V.Schwämmle and E. Brigatti, preprint q-bio/0509032.
- [4] K. Rohde and D. Stauffer, preprint for *Adv. Complex Syst.*
- [5] D. Stauffer, S. Moss de Oliveira, P.M.C. de Oliveira, J.S. Sa Martins, *Biology, Sociology, Geology by Computational Physicists*, Elsevier, Amsterdam 2006 in press.
- [6] M.A.F. Gomes, G.L. Vasconcelos, I.S. Tsang, I.R. Tsang, *Physica A* 271, 489 (1999).




Figure 2: As Fig.1 but in part a for $t = 2,400$ to $2,400,000$ iterations at $L = 50$, and in part b for three to seven food levels (from bottom to top) at $L = 50, t = 24,000$.