

# Conditions for multiplicativity of maximal $\ell_p$ -norms of channels for fixed integer $p$

Vittorio Giovannetti<sup>1</sup>, Seth Lloyd<sup>2</sup>, and Mary Beth Ruskai<sup>3</sup>

<sup>1</sup>*NEST-INFM & Scuola Normale Superiore, Piazza dei Cavalieri 7, I-56126, Pisa, Italy*

<sup>2</sup>*Massachusetts Institute of Technology – Department of Mechanical Engineering and Research Laboratory of Electronics  
77 Massachusetts Ave., Cambridge, MA 02139, USA*

<sup>3</sup>*Department of Mathematics, Tufts University Medford, Massachusetts 02155 USA*

(Dated: December 2, 2024)

We introduce a condition for memoryless quantum channels which, when satisfied guarantees the multiplicativity of the maximal  $\ell_p$ -norm with  $p$  a fixed integer. By applying the condition to qubit channel, it can be seen that it is not a necessary condition, although some known results for qubits can be recovered. For  $p = 2, 3, 4$  the condition is applied to the Werner-Holevo channel, which is known to violate multiplicativity for large  $p$ , and shown to hold when  $d \geq 2^{p-1}$ . Finally, a new class of channels is considered which generalize the depolarizing channel to maps which are combinations of the identity channel and a noisy one whose image is an arbitrary density matrix. It is shown that these channels are multiplicative for  $p = 2$ .

PACS numbers: 03.67.Hk, 03.67.-a, 03.65.Db, 42.50.-p

## I. INTRODUCTION

A noisy quantum channel can be described by [9, 26, 29] means of a completely-positive, trace-preserving (CPT) map  $\mathcal{E}$  which transforms the density matrices  $\gamma$  on the Hilbert space  $\mathcal{H}$  into the output states  $\mathcal{E}(\gamma)$ . Such maps can always be represented [8, 26, 29] in the form

$$\mathcal{E}(\gamma) = \sum_k A_k \gamma A_k^\dagger, \quad \sum_k A_k^\dagger A_k = \mathbb{1}, \quad (1)$$

with  $\{A_k\}$  called a set of Kraus operators associated with  $\mathcal{E}$ . When the channel is memoryless [6],  $m$  successive uses of are described by the map  $\mathcal{E}^{\otimes m}$ . It is natural to ask if entangled inputs can decrease the effects of noise for memoryless channels [5] in some way.

One measure of the effect of noise is the maximal  $\ell_p$ -norm of a channel, which is defined as

$$\nu_p(\mathcal{E}) \equiv \sup_{\gamma \in \mathcal{D}(\mathcal{H})} \|\mathcal{E}(\gamma)\|_p \quad p \geq 1, \quad (2)$$

where  $\|A\|_p \equiv (\text{Tr}|A|^p)^{1/p}$  is the  $p$ -norm of the operator  $A$  and where the supremum is taken over all  $\mathcal{D}(\mathcal{H})$ , the set of density matrices. The quantity  $\text{Tr}[\mathcal{E}(\gamma)^p]$  is a measure of the closeness of the output to a pure state, and  $\nu_p(\mathcal{E}) = 1$  if and only if some output state  $\mathcal{E}(\gamma)$  is pure. Because the Rényi entropy [30] can be written as  $S_p(\rho) = -\frac{1}{p-1} \log \|\rho\|_p^p$  one could define a maximal output Rényi entropy [2, 13, 14] satisfying  $(p-1)S_{p,\max}(\mathcal{E}) = -p \log \nu_p(\mathcal{E})$ .

Amosov, Holevo and Werner (AHW) conjectured [4] that  $\nu_p(\mathcal{E})$  is multiplicative for tensor product channels

$$\nu_p(\mathcal{E}^{\otimes m}) \equiv \sup_{\Gamma \in \mathcal{D}(\mathcal{H}^{\otimes m})} \|\mathcal{E}^{\otimes m}(\Gamma)\|_p = [\nu_p(\mathcal{E})]^m, \quad (3)$$

where  $\mathcal{E}^{\otimes m}$  is the CPT map which describes  $m$  successive memoryless uses of the channel  $\mathcal{E}$ , and where the maximization in the second term of Eq. (3) is now performed over the density matrices  $\Gamma \in \mathcal{D}(\mathcal{H}^{\otimes m})$ . The AHW conjecture requires that a product state  $\Gamma$  saturate the supremum of  $\nu_p(\mathcal{E}^{\otimes m})$  for the memoryless channel  $\mathcal{E}^{\otimes m}$  so that entangled input states  $\Gamma$  do not increase the output norm. One rational for the multiplicativity hypothesis [4] is the physical intuition that quantum coherence among successive channel uses should be degraded by the action of a memoryless channel. Since the  $\ell_p$ -norm “measures” the purity of the states emerging from the channel, one might expect separable inputs to perform better than entangled inputs. The multiplicativity of  $\nu_p(\mathcal{E}^{\otimes m})$  is equivalent to additivity for the minimum Rényi entropy with the same  $p$  [2, 13, 14]. Moreover, if (3) holds for  $p$  arbitrarily close to 1, then it implies [4] the additivity of the minimum output von Neumann entropy [24], another measure of output purity. This has been shown [33] to be related to a conjectured additivity property of the Holevo information [15], and to conjectures about additivity and superadditivity of the entanglement of formation [1, 27].

Subsequently, Werner and Holevo [34] showed that the general multiplicativity conjecture is false by producing a channel that violates (3) for  $p > 4.79$ . Nevertheless, one might still expect multiplicativity to hold for some range of  $p$ , most notably  $1 \leq p \leq 2$  and this would suffice for many applications in quantum information theory. However,

even the case  $p = 2$  is still not resolved. It is hence important to understand under which circumstances and for which values of  $p$  a given channel satisfies Eq. (3). Many authors have tackled this problem by discussing special situations for which the conjecture can be proved [2, 3, 13, 18, 19, 20, 21, 22, 23, 25, 28, 32]. In the case of a fixed integer  $p$ , we provide an upper bound for  $\nu_p(\mathcal{E}^{\otimes m})$ , and derive a pair of sufficient conditions, either of which ensures that  $\mathcal{E}$  satisfies the multiplicativity conjecture (3).

The material is organized as follows. In Section II we introduce some notation and present a linearization technique that allows one to compute the  $\ell_p$ -norm of integer order as the expectation value of an operator defined on an extended Hilbert space. In Section III we derive our upper bound and show how it leads to a sufficient condition for the multiplicativity of the  $\ell_p$ -norm. Then we apply our condition to several classes of channels. By considering qubit channels in the case  $p = 2$ , we show in Section IV A that our sufficient condition is not necessary. We also obtain new proofs of multiplicativity when the two shortest axes of the image ellipsoid (whether or not shifted) are equal. In Section IV B we prove multiplicativity when  $p = 2$  for a shifted depolarizing channel and further generalizations which do not seem to have been considered in the literature. Finally, in Section IV C we consider the Werner-Holevo channel [34] for  $p = 2, 3, 4$ , and obtain new results about multiplicativity when  $p = 3, 4$ . We also conjecture that the channel is multiplicative for *any*  $p$  when it acts on a space of dimension  $d \geq 2^{p-1}$ .

We include several appendices. The first reviews useful facts about operators, including Hilbert-Schmidt duality, shift and permutation operators, and double stochastic matrices. Appendix A also contains information about the notation, and the proof of an important identity. Appendix B discusses properties and alternative forms of the linearizing operators we use. Appendix C provides details needed for our analysis of the Werner-Holevo channel.

## II. LINEARIZATION OF $p$ -NORM FUNCTIONS

### A. Basic linearization strategy

In this section we present a method, introduced in [13], that allows one to compute the  $\ell_p$ -norm from the expectation value of an operator defined in an extended Hilbert space. For any  $p$ -integer, it is possible to find a linear operator  $X(\mathcal{E}, p)$  defined in the extended Hilbert space  $\mathcal{H}^{\otimes p}$  such that, for any density matrix  $\gamma \in \mathcal{H}$ , we have

$$\text{Tr}[\mathcal{E}(\gamma)]^p = \text{Tr}[\underbrace{(\gamma \otimes \gamma \otimes \cdots \otimes \gamma)}_{p\text{-times}} X(\mathcal{E}, p)] \quad (4)$$

where the trace in the left-hand side is computed with respect to an orthonormal basis of  $\mathcal{H}$ , while the trace in the right-hand side is computed with respect to an orthonormal basis of  $\mathcal{H}^{\otimes p}$ . In other words, we can represent the  $p$ -purity function  $\text{Tr}[\mathcal{E}(\gamma)^p]$  as the expectation value of  $X(\mathcal{E}, p)$  on  $p$  copies of  $\gamma$ . The operator  $X(\mathcal{E}, p)$  is not uniquely defined; in fact, it can be realized by the action of tensor products of the dual map of  $\mathcal{E}$  on any permutation operator acting on  $\mathcal{H}^{\otimes p}$  whose shortest cycle is length  $p$ .

To make this explicit, we need some notation, which is explained in more detail in Appendix A, particularly sections A 1 and A 2. We will use a hat to denote the dual, or adjoint, map  $\hat{\mathcal{E}}$  with respect to the Hilbert-Schmidt inner product. Let  $L_p$  and  $R_p$  denote the left and right cyclic shifts which can be defined by their action on an orthonormal product basis as

$$L_p |\xi_1 \xi_2 \cdots \xi_{p-1} \xi_p\rangle = |\xi_2 \cdots \xi_p \xi_1\rangle \quad (5a)$$

$$R_p |\xi_1 \xi_2 \cdots \xi_{p-1} \xi_p\rangle = |\xi_p \xi_1 \cdots \xi_{p-1}\rangle. \quad (5b)$$

where  $|\xi_1 \xi_2 \cdots \xi_{p-1} \xi_p\rangle = \otimes_{j=1}^p |\xi_j\rangle$  and  $\{|\xi_k\rangle\}$  is an orthonormal basis for  $\mathcal{H}$ . Then the operator

$$\Omega(\mathcal{E}, p) \equiv \hat{\mathcal{E}}^{\otimes p}(L_p) \quad (6)$$

satisfies (4). This follows from

$$\begin{aligned} \text{Tr} \gamma^{\otimes p} \Omega(\mathcal{E}, p) &= \text{Tr}(\gamma \otimes \gamma \otimes \cdots \otimes \gamma) \hat{\mathcal{E}}^{\otimes p}(L_p) \\ &= \text{Tr} [\mathcal{E}(\gamma) \otimes \mathcal{E}(\gamma) \otimes \cdots \otimes \mathcal{E}(\gamma)] L_p \\ &= \text{Tr}[\mathcal{E}(\gamma)]^p \end{aligned} \quad (7)$$

where the last step used (A12). It follows from (A14) that  $L_p$  could be replaced by another permutation; however, it is important to make a definite choice for later use.

In previous work [13, 14], a different realization of  $X(\mathcal{E}, p)$  was used which is valid only for pure states. Let

$$\Theta(\mathcal{E}, p) = \Omega(\mathcal{E}, p) R_p = \widehat{\mathcal{E}}^{\otimes p}(L_p) R_p \quad (8)$$

$$= \sum_{k_1, \dots, k_p} A_{k_1}^\dagger A_{k_2} \otimes A_{k_2}^\dagger A_{k_3} \otimes \dots \otimes A_{k_p}^\dagger A_{k_1} \quad (9)$$

where  $\{A_k\}$  form a set of Kraus operators for  $\mathcal{E}$  as in (1). The operator  $\Theta(\mathcal{E}, p)$  satisfies (4) when  $\gamma = |\psi\rangle\langle\psi|$  is a pure state. This relation is proved in Appendix B 1, and implicitly shows that it does not depend on the chosen Kraus representation (1) of  $\mathcal{E}$ . For  $p = 2$ , (9) and (8) were obtained earlier in Ref. [35].

In general, the operator  $\Omega(\mathcal{E}, p)$  will not be Hermitian. We have already observed that  $X(\mathcal{E}, p)$  is not unique and that whenever  $P_p$  is a permutation operator whose shortest cycle is length  $p$ , the operator  $\widehat{\mathcal{E}}^{\otimes p}(P_p)$  provide another realization. Since  $L_p = R_p^\dagger$ ,

$$[\widehat{\mathcal{E}}^{\otimes p}(L_p)]^\dagger = \widehat{\mathcal{E}}^{\otimes p}(L_p^\dagger) = \widehat{\mathcal{E}}^{\otimes p}(R_p). \quad (10)$$

This implies that  $\Omega(\mathcal{E}, 2)$  is Hermitian for  $p = 2$ , and that the operator

$$\frac{1}{2}[\Omega(\mathcal{E}, p) + [\Omega(\mathcal{E}, p)]^\dagger] = \frac{1}{2}[\widehat{\mathcal{E}}^{\otimes p}(L_p) + \widehat{\mathcal{E}}^{\otimes p}(R_p)] \quad (11)$$

gives a Hermitian realization of  $X(\mathcal{E}, p)$  for any  $p$ . However, we do not expect (11) to have the important multiplicity property (14) for repeated uses of the channel. Further discussion of other realizations  $X(\mathcal{E}, p)$  is given Appendices B 2 and B 3.

Linear operators satisfying (4) provide a useful tool for studying the  $p$ -purity functions, which are intrinsically non-linear objects; it reduces some associated problems to the analysis of the linear operator  $X(\mathcal{E}, p)$  acting on the extended Hilbert space  $\mathcal{H}^{\otimes p}$  obtained by adding  $p-1$  “fictitious” copies of the input Hilbert space  $\mathcal{H}$ . In Refs. [13, 14], this approach was used to obtain some additivity properties of Gaussian Bosonic channels. For  $p = 2$ , Eq. (4) was used in Ref. [7] to study the fidelity obtainable in continuous-variable teleportation with finite two-mode squeezing, and in Ref. [35] to analyze the purity of generic quantum channels.

## B. Tensor product maps

The results derived in the preceding section can also be applied when the basic CPT map is itself a tensor product. Then Eq. (4) becomes

$$\text{Tr}[\mathcal{E}^{\otimes m}(\Gamma)]^p = \text{Tr}[\underbrace{(\Gamma \otimes \Gamma \otimes \dots \otimes \Gamma)}_{p\text{-times}} X(\mathcal{E}^{\otimes m}, p)], \quad (12)$$

where  $\Gamma$  is a generic density matrix in the input Hilbert space  $\mathcal{H}^{\otimes m}$  and  $X(\mathcal{E}^{\otimes m}, p)$  is a linear operator on  $(\mathcal{H}^{\otimes m})^{\otimes p} = \mathcal{H}^{\otimes mp}$ . Following the strategy of Section II A, we now choose  $X(\mathcal{E}^{\otimes m}, p)$  to be the operator,

$$\Omega(\mathcal{E}^{\otimes m}, p) \equiv (\widehat{\mathcal{E}^{\otimes m}})^{\otimes p}(\mathbb{L}_p) = (\widehat{\mathcal{E}^{\otimes m}})^{\otimes p}(L_p^{\otimes m}) \quad (13)$$

The operator  $\mathbb{L}_p$  is described in more detail in Appendix A 3 where it is proved that  $\mathbb{L}_p = L_p^{\otimes m} = (L_{mp})^m$ . Using  $(\widehat{\mathcal{E}^{\otimes m}})^{\otimes p} = (\widehat{\mathcal{E}})^{\otimes mp}$ , we find

$$\Omega(\mathcal{E}^{\otimes m}, p) = \widehat{\mathcal{E}}^{\otimes mp}(L_p^{\otimes m}) = [\widehat{\mathcal{E}}^{\otimes p}(L_p)]^{\otimes m} = [\Omega(\mathcal{E}, p)]^{\otimes m}. \quad (14)$$

Equation (14) is a key result whose simplicity hides a great deal of subtlety. The essential point is that the linear operator  $X(\mathcal{E}^{\otimes m}, p)$  which satisfies (1) for the tensor product channel  $\mathcal{E}^{\otimes m}$  can be realized by the action of the dual of  $(\mathcal{E}^{\otimes m})$  on the permutation  $L_p^{\otimes m}$ .

### III. CONDITIONS FOR MULTIPLICATIVITY

#### A. Upper bound

We now use the singular value decomposition [16, 17] to observe that one can write

$$\Omega(\mathcal{E}, p) = \sum_j \xi_j |\mu_j\rangle\langle\nu_j| \quad (15)$$

where  $\{|\mu_j\rangle\}$  and  $\{|\nu_j\rangle\}$  denote orthonormal bases for  $\mathcal{H}^{\otimes p}$  and  $\xi_j > 0$  are the singular values of  $\Omega(\mathcal{E}, p)$ , i.e., the non-zero eigenvalues of  $|\Omega(\mathcal{E}, p)| \equiv \sqrt{[\Omega(\mathcal{E}, p)]^\dagger \Omega(\mathcal{E}, p)}$ . Before applying this, it will be convenient to introduce the convention of using bold uppercase Greek letters to denote tensor product vectors as in  $|\Psi\rangle \equiv |\psi\rangle^{\otimes p} = |\psi\rangle \otimes |\psi\rangle \otimes \cdots \otimes |\psi\rangle \in \mathcal{H}^{\otimes p}$ . Then

$$\begin{aligned} \text{Tr}[\mathcal{E}(|\psi\rangle\langle\psi|)]^p &= \langle\Psi|\Omega(\mathcal{E}, p)|\Psi\rangle \\ &= \sum_j \xi_j \langle\Psi|\mu_j\rangle\langle\nu_j|\Psi\rangle \\ &\leq \xi_{\max} \sum_j |\langle\Psi|\mu_j\rangle\langle\nu_j|\Psi\rangle| \\ &\leq \xi_{\max} \|\Psi\|^2 = \|\Omega(\mathcal{E}, p)\|_\infty \end{aligned} \quad (16)$$

where  $\xi_{\max} = \sup_j \xi_j = \|\Omega(\mathcal{E}, p)\|_\infty$  is the largest singular value of  $\Omega(\mathcal{E}, p)$ . Applying this analysis to multiple uses of the channel, one can similarly conclude that

$$\text{Tr}[\mathcal{E}^{\otimes m}(|\Psi\rangle\langle\Psi|)]^p \leq \|\Omega(\mathcal{E}^{\otimes m}, p)\|_\infty, \quad (17)$$

where  $|\Psi\rangle$  is now an arbitrary vector in  $\mathcal{H}^{\otimes m}$ . However, it follows from (14) that the singular values of  $\Omega(\mathcal{E}^{\otimes m}, p)$  are products of those of  $\Omega(\mathcal{E}, p)$  so that

$$\|\Omega(\mathcal{E}^{\otimes m}, p)\|_\infty = (\|\Omega(\mathcal{E}, p)\|_\infty)^m = (\xi_{\max})^m. \quad (18)$$

Combining (17) and (18), one finds

$$\text{Tr}[\mathcal{E}^{\otimes m}(|\Psi\rangle\langle\Psi|)]^p \leq (\|\Omega(\mathcal{E}, p)\|_\infty)^m = (\xi_{\max})^m. \quad (19)$$

Since, the supremum in (2) is attained using a pure state input and (19) holds for all pure inputs  $|\Psi\rangle$ , we conclude that the upper bound

$$\nu_p(\mathcal{E}^{\otimes m}) \leq (\xi_{\max})^{m/p}, \quad (20)$$

holds for all pairs of integers  $m$  and  $p$ .

#### B. Multiplicativity condition

The bound (20) leads to a sufficient condition for multiplicativity. We state this formally, and give a relate condition as a corollary.

**Theorem 1** *The channel  $\mathcal{E}$  has the multiplicativity property (3) if the largest singular value of  $\Omega(\mathcal{E}, p)$  satisfies*

$$\|\Omega(\mathcal{E}, p)\|_\infty = [\nu_p(\mathcal{E})]^p \quad (21)$$

**Corollary 2** *The channel  $\mathcal{E}$  has the multiplicativity property (3) if the largest singular value of  $\Omega(\mathcal{E}, p)$  is also an eigenvalue of  $\Omega(\mathcal{E}, p)$  with a product eigenvector of the form  $|\phi\rangle^{\otimes p}$*

To prove Theorem 1, observe that in the notation of the preceding section (21) can be written as  $\xi_{\max} = [\nu_p(\mathcal{E})]^p$ . Then (20) implies

$$\nu_p(\mathcal{E}^{\otimes m}) \leq (\xi_{\max})^{m/p} = [\nu_p(\mathcal{E})]^m. \quad (22)$$

On the other hand, one always has

$$\nu_p(\mathcal{E}^{\otimes m}) \geq \|\mathcal{E}^{\otimes m}(\gamma_{\max}^{\otimes m})\|_p = \|\mathcal{E}(\gamma_{\max})\|_p^m = [\nu_p(\mathcal{E})]^m$$

where  $\gamma_{\max}$  denotes the state which achieves the supremum for  $\nu_p(\mathcal{E})$ . Combining these inequalities gives  $\nu_p(\mathcal{E}^{\otimes m}) = [\nu_p(\mathcal{E})]^m$ . **QED**

To prove the corollary, observe that hypothesis holds if and only if there is a state  $|\phi\rangle$  in  $\mathcal{H}$  such that

$$\xi_{\max} = \langle \Phi | \Omega(\mathcal{E}, p) | \Phi \rangle = \text{Tr}[\mathcal{E}(|\phi\rangle\langle\phi|)]^p \quad (23)$$

where the second equality used (7) and our convention that  $|\Phi\rangle = |\phi\rangle^{\otimes p}$ . But it is always true that

$$\text{Tr}[\mathcal{E}(|\phi\rangle\langle\phi|)]^p \leq \sup_{\gamma} \text{Tr}[\mathcal{E}(\gamma)]^p \equiv [\nu_p(\mathcal{E})]^p \quad (24)$$

so that  $\xi_{\max} \leq [\nu_p(\mathcal{E})]^p$ . Combining this with (20) when  $m = 1$ , implies that  $\xi_{\max} = [\nu_p(\mathcal{E})]^p$  so that the hypothesis of Theorem 1 holds. **QED**

In Section IV A we will see that the condition in Theorem 1 is not necessary. There are unital qubit CPT maps, which are known to be multiplicative, but do not satisfy (21). Verifying the hypothesis of Corollary 2 requires that one find an eigenvector as well as the largest singular value of an operator, but does not require knowledge of  $\nu_p(\mathcal{E})$ ; condition (21) does require the latter, but does not require computation of any eigenvectors. In general, (21) seems easier to check. However, in the examples we analyzed, both conditions hold and the process of verifying one easily yields the other. It would be interesting to know if (21) implies that the singular value of  $\Omega(\mathcal{E}, p)$  is also an eigenvector with a product eigenvalue as in Corollary 2.

## IV. APPLICATIONS

### A. Qubit channels

#### 1. Notation

We illustrate our condition by looking at some examples of qubit channels, for which will use notation similar to that introduced in [25, 31]. Any  $2 \times 2$  matrix can be represented in the basis consisting the  $2 \times 2$  identity matrix  $\mathbb{1}$  and the three Pauli matrices which we often write as a formal vector  $\vec{\sigma} \equiv (\sigma_1, \sigma_2, \sigma_3)$ . In this basis a density matrix can be written as  $\gamma = \frac{1}{2}[\mathbb{1} + \vec{w} \cdot \vec{\sigma}]$  with  $\vec{w}$  in  $\mathbf{R}^3$  and  $|\vec{w}| \leq 1$ . The density matrix is pure if and only if  $|\vec{w}| = 1$ . Any linear map  $\Phi$  on a qubit, can be described by two real vectors  $\vec{s}, \vec{t} \in \mathbf{R}^3$  and by a  $3 \times 3$  real matrix  $T$ , through the expression

$$\Phi(z_0 \mathbb{1} + \vec{z} \cdot \vec{\sigma}) = (z_0 + \vec{s} \cdot \vec{z}) \mathbb{1} + (z_0 \vec{t} + T \cdot \vec{z}) \cdot \vec{\sigma}, \quad (25)$$

which holds for all  $z_0 \in \mathbb{C}$  and  $\vec{z} \in \mathbb{C}^3$ . This corresponds to representing  $\Phi$  in the basis  $\{\mathbb{1}, \vec{\sigma}\}$  by the  $4 \times 4$  matrix  $\begin{pmatrix} 1 & \vec{s}^t \\ \vec{t} & T \end{pmatrix}$  which we have written in block form (with the convention that  $\vec{t}$  corresponds to a column vector and  $\vec{s}^t$  a row vector, using the superscript  $t$  to denote transpose). It was shown in [24] that it suffices to consider  $T$  diagonal with real elements  $\{\lambda_1, \lambda_2, \lambda_3\}$ . [In essence, a variant of the SVD (which leads to negative as well as positive  $\lambda_k$ ) can be applied to  $T$  corresponding to rotations on the input and output bases respectively.]

In this notation,  $\Phi$  is trace preserving (TP) if and only if  $\vec{s} = 0$  and it is unital if and only if  $\vec{t} = 0$ . Additional conditions under which the map is positivity preserving or completely positive (CP) are more complex. A complete set of conditions for the map to be CPT was obtained in [31]. When  $t_1 = t_2 = 0$ , these CPT conditions reduce to  $(\lambda_1 \pm \lambda_2)^2 \leq (1 \pm \lambda_3)^2 - t_3^2$ , as shown in Refs. [12, 31]. Since the dual map of  $\Phi$  is represented by the adjoint matrix, it satisfies,

$$\widehat{\Phi}(z_0 \mathbb{1} + \vec{z} \cdot \vec{\sigma}) = (z_0 + \vec{t} \cdot \vec{z}) \mathbb{1} + (z_0 \vec{s} + T^t \cdot \vec{z}) \cdot \vec{\sigma}. \quad (26)$$

Since  $\mathcal{H}$  is now 2-dimensional, the left shift  $L_2$  is simply the SWAP operator  $S$  which satisfies

$$S = \frac{1}{2}[\mathbb{1} \otimes \mathbb{1} + \sigma_1 \otimes \sigma_1 + \sigma_2 \otimes \sigma_2 + \sigma_3 \otimes \sigma_3] \quad (27)$$

It is then straightforward to use (6) to show that

$$\Omega(\Phi, 2) = \hat{\Phi}^{\otimes 2}(S) = \frac{1}{2} \left[ (1 + |\vec{t}|^2) \mathbb{1} \otimes \mathbb{1} + \sum_{j=1}^3 \lambda_j^2 \sigma_j \otimes \sigma_j + \sum_{j=1}^3 \lambda_j t_j (\mathbb{1} \otimes \sigma_j + \sigma_j \otimes \mathbb{1}) \right]. \quad (28)$$

## 2. Unital maps

For qubit channels the conjecture (2) has been extensively studied in [18, 19, 22, 24]. Multiplicativity has been proven for all  $p$  for unital qubit channels [18] and for  $p = 2$  for all qubit channels (Theorem 2 of [18]). Here we will use the case  $p = 2$  to illustrate the multiplicativity-criterion presented in Section III B.

It will be useful to choose the subscript “max” in  $\{1, 2, 3\}$  so that  $|\lambda_{\max}| = \max_k |\lambda_k|$ . For unital qubits maps, the maximum  $\ell_2$ -norm of  $\Phi$  can be achieved with an input state of the form  $\frac{1}{2}[\mathbb{1} \pm \sigma_{\max}]$  for which the output  $\frac{1}{2}[\mathbb{1} \pm \lambda_{\max} \sigma_{\max}]$  has eigenvalues  $\frac{1}{2}[1 \pm \lambda_{\max}]$  and

$$\nu_2(\Phi) = \frac{1}{\sqrt{2}} \sqrt{1 + \lambda_{\max}^2} \quad (29)$$

When  $\Phi$  is unital,  $\vec{t} = 0$  and the third term in the expression (28) vanishes. It then follows that in the product basis  $\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$ , the operator  $\Omega(\Phi, 2)$  is represented by the matrix

$$\frac{1}{2} \begin{pmatrix} 1 + \lambda_3^2 & 0 & 0 & \lambda_1^2 - \lambda_2^2 \\ 0 & 1 - \lambda_3^2 & \lambda_1^2 + \lambda_2^2 & 0 \\ 0 & \lambda_1^2 + \lambda_2^2 & 1 - \lambda_3^2 & 0 \\ \lambda_1^2 - \lambda_2^2 & 0 & 0 & 1 + \lambda_3^2 \end{pmatrix}. \quad (30)$$

This is easily seen to have two non-zero  $2 \times 2$  blocks. The “inner” block has eigenvalues  $\frac{1}{2}[1 - \lambda_3^2 \pm (\lambda_1^2 - \lambda_2^2)]$  with eigenvectors  $\frac{1}{\sqrt{2}}(0, 1, \pm 1, 0)^t$  corresponding to the Bell states  $\frac{1}{\sqrt{2}}(|01\rangle \pm |10\rangle)$ . The “outer” block has eigenvalues  $\frac{1}{2}[1 + \lambda_3^2 \pm (\lambda_1^2 + \lambda_2^2)]$  with eigenvectors  $\frac{1}{\sqrt{2}}(1, 0, 0, \pm 1)^t$  corresponding to the Bell states  $\frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle)$ . Since  $\Omega(\Phi, 2)$  is Hermitian, its singular values are simply the absolute values of the eigenvalues above.

When the  $|\lambda_k|$  are distinct for  $k = 1, 2, 3$ , the singular values of  $\Omega(\Phi, 2)$  are all distinct and correspond to maximally entangled, rather than product, states. Moreover, one of the singular values is always strictly greater than  $\nu_2(\Phi)$ . For example, when  $|\lambda_{\max}| = |\lambda_3|$ , one of the “inner” eigenvalues equals  $\nu_2(\Phi)^2 + \frac{1}{2}(\lambda_1^2 - \lambda_2^2)$  which is strictly greater than (29) unless  $|\lambda_1| = |\lambda_2|$ . Therefore, although  $\Phi$  is multiplicative, it does not satisfy (21). This establishes that (21) is not a necessary condition for multiplicativity.

Now consider the case  $\lambda_3 > \lambda_1 = \lambda_2 \geq 0$ ; such channels are sometimes called “two-Pauli” channels [5]. The image of the Bloch sphere is an ellipsoid shaped like an American football. For these channels, the “outer” block in (30) is diagonal, its (degenerate) eigenvalue  $\frac{1}{2}(1 + \lambda_3^2) = [\nu_2(\Phi)]^2$  is the largest singular value of  $\Omega(\Phi, 2)$  and the corresponding eigenvectors  $|00\rangle$  and  $|11\rangle$  are product states. Thus, Theorem 1 implies that the channel satisfies (3).

## 3. Non-unital maps

We now consider channels similar to those above, but with the image ellipsoid shifted along the longest axis. It suffices to consider  $|\lambda_3| \geq \lambda_1 = \lambda_2 \geq 0$  and  $t_1 = t_2 = 0$ . The same results hold for permutations of 1, 2, 3 and for  $\lambda_1 = \lambda_2 \leq 0$ . However, the analysis in the basis we have chosen to represent  $\Omega(\Phi, 2)$  is simplest when  $|\lambda_{\max}| = |\lambda_3|$ . The matrix representing  $\Omega(\Phi, 2)$  is

$$\frac{1}{2} \begin{pmatrix} 1 + (t_3 + \lambda_3)^2 & 0 & 0 & 0 \\ 0 & 1 + t_3^2 - \lambda_3^2 & 2\lambda_1^2 & 0 \\ 0 & 2\lambda_1^2 & 1 + t_3^2 - \lambda_3^2 & 0 \\ 0 & 0 & 0 & 1 + (t_3 - \lambda_3)^2 \end{pmatrix} \quad (31)$$

which has an “inner” block with eigenvalues  $\frac{1}{2}[1 + t_3^2 - \lambda_3^2 \pm 2\lambda_1^2]$  and a diagonal “outer” block with eigenvalues  $\frac{1}{2}[1 + (t_3 \pm \lambda_3)^2]$  and product eigenvectors. One can verify that the largest singular value is  $\frac{1}{2}[1 + (|t_3| + |\lambda_3|)^2]$ . To see that this equals  $[\nu_2(\Phi)]^2$ , observe that the optimal input state is  $\frac{1}{2}[\mathbb{1} + \frac{t_3}{|t_3|}\sigma_3]$  for which the output state has eigenvalues  $\frac{1}{2}[1 \pm (|t_3| + |\lambda_3|)^2]$ . Thus, we can again use Theorem 1 to conclude that (3) holds.

The methods introduced here are able to handle qubit channels for which the image of the Bloch sphere is an elongated ellipsoid with a symmetry axis, i.e., in the shape of an American football, both when the channel is unital and when it is shifted in the direction of the longest axis. However, it can not handle these channels if the shift is orthogonal to the longest axis, i.e., if  $t_3 = 0$  but  $t_2 \neq 0$  above. When the ellipsoid has a symmetry axis but  $|\lambda_1| = |\lambda_2| \geq |\lambda_3|$  so that it is shaped like a flying saucer, the methods used here can not prove multiplicativity. Even for unital channels, for which multiplicativity has been established [19], neither of the conditions in Theorem 1 holds.

## B. Shifted depolarizing channels

### 1. Shifting and generalizing the depolarizing channel

The unital qubit map with  $\lambda_k = \pm|\lambda_{\max}|$  for all  $k$ , is a special case of the depolarizing channel which has the form  $\mathcal{E}(\gamma) = (1 - x)(\text{Tr}\gamma)\frac{1}{d}\mathbb{1} + x\gamma$ . It is CPT for  $-\frac{1}{3} \leq x \leq 1$ . The non-unital qubit map which takes

$$\begin{aligned} \gamma = \frac{1}{2}[\mathbb{1} + \vec{w} \cdot \vec{\sigma}] &\mapsto \frac{1}{2}[\mathbb{1} + (\vec{t} + \lambda\vec{w}) \cdot \vec{\sigma}] \\ &= (1 - |\vec{t}| - \lambda)\frac{1}{2}\mathbb{1} + |\vec{t}|\frac{1}{2}[\mathbb{1} + \hat{t} \cdot \vec{\sigma}] + \lambda\gamma \end{aligned} \quad (32)$$

can be regarded as a shifted depolarizing channel because it shifts the output toward the point  $\hat{t}$  on the Bloch sphere. By rotating coordinates so that  $\vec{t} = (0, 0, t_3)$ , this is a special case of the qubit maps considered in Section IV A 3 above. It is then natural to define a shifted depolarizing channel in dimension  $d$  by

$$\mathcal{E}(\gamma) = a(\text{Tr}\gamma)\frac{1}{d}\mathbb{1} + b(\text{Tr}\gamma)|\psi\rangle\langle\psi| + c\gamma \quad (33)$$

with the state  $|\psi\rangle$  fixed and  $a + b + c = 1$ . When  $a, b, c$  are positive, this channel is a convex combination of the identity map and two completely noisy channel which maps all states to  $\frac{1}{d}\mathbb{1}$  and to  $b|\psi\rangle\langle\psi|$ , respectively.

We now consider the more general class of channels of the form

$$\mathcal{E}(\gamma) = (1 - c)(\text{Tr}\gamma)\rho + c\gamma \quad (34)$$

where  $\rho$  is a fixed density matrix. For  $\rho = \frac{1}{d}\mathbb{1}$ , this is the usual depolarizing channel; for  $\rho = \frac{1}{a+b}[\frac{a}{d}\mathbb{1} + b|\psi\rangle\langle\psi|]$ , it is the shifted depolarizing channel (33).

When  $c \geq 0$  additivity was proved for the depolarizing channel in  $d$  dimensions using a majorization argument [12] from which multiplicativity immediately follows; for  $-\frac{1}{d^2-1} \leq c \leq 1$ , (which is the range for which the map is CPT) multiplicativity of the depolarizing channel in  $d$  dimensions was proved in [21]. Neither shifted depolarizing channels nor the generalization (34) seem to have been explicitly considered in the literature before. One could obtain a proof of multiplicativity for  $p = 2$  when  $c > 0$  by verifying that the positive element condition in [25] is satisfied. (In fact, these maps satisfy the stronger condition considered in [23].) However, neither of these positive element conditions can be verified when  $c < 0$ . By contrast, the method presented here can establish multiplicativity when  $p = 2$  for all CPT maps of the form (34), including those with  $c < 0$ .

### 2. Convex combinations of the identity and completely noisy maps

It will be useful to write the spectral decomposition of  $\rho$  as  $\rho = \sum_j a_j |j\rangle\langle j|$  with the eigenvalues  $a_j$  in decreasing order. Then, for  $c > 0$ , the state  $\mathcal{E}(|1\rangle\langle 1|)$  majorizes all outputs so that  $[\nu_p(\mathcal{E})]^p = [ca_1 + (1 - c)]^p + c^p \sum_{j>1}^d a_j^p$ .

Since,  $\hat{\mathcal{E}}(B) = c[\text{Tr}B\rho]\mathbb{1} + cB$ , we have

$$\hat{\mathcal{E}}(|j\rangle\langle k|) = (1 - c)\langle k|\rho|j\rangle\mathbb{1} + c|j\rangle\langle k| = (1 - c)\delta_{jk}a_k\mathbb{1} + c|j\rangle\langle k|, \quad (35)$$

and

$$\begin{aligned}\Omega(\mathcal{E}, 2) &= (\widehat{\mathcal{E}} \otimes \widehat{\mathcal{E}})(S) = \sum_{jk} \widehat{\mathcal{E}}(|j\rangle\langle k|) \otimes \widehat{\mathcal{E}}(|k\rangle\langle j|) \\ &= \sum_{jk} \left[ (1-c)^2 \delta_{jk} a_k^2 \mathbb{1} \otimes \mathbb{1} + c(1-c) \delta_{jk} a_k \left( \mathbb{1} \otimes |k\rangle\langle k| + |k\rangle\langle k| \otimes \mathbb{1} \right) + |j\rangle\langle k| \otimes |k\rangle\langle j| \right]\end{aligned}\quad (36)$$

$$= (1-c)^2 (\text{Tr} \rho^2) \mathbb{1} \otimes \mathbb{1} + c(1-c) [\mathbb{1} \otimes \rho + \rho \otimes \mathbb{1}] + c^2 S. \quad (37)$$

From this it is easy to see that  $\Omega(\mathcal{E}, 2)$  has  $d$  product eigenvectors of the form  $|kk\rangle$  with eigenvalues

$$(1-c)^2 (\text{Tr} \rho^2) + 2c(1-c) a_k + c^2 = [(1-c) a_k + c]^2 + (1-c)^2 \sum_{j \neq k} a_j^2, \quad (38)$$

and  $\binom{d}{2}$  blocks of the form  $[(1-c)^2 (\text{Tr} \rho^2) + c(1-c)(a_j + a_k)] \mathbb{1}_2 + c^2 \sigma_x$ , with eigenvalues

$$(1-c)^2 (\text{Tr} \rho^2) + c(1-c)(a_j + a_k) \pm c^2 \quad (39)$$

and entangled eigenvectors  $2^{-1/2}(|jk\rangle \pm |kj\rangle)$ . When  $c > 0$  all eigenvalues are non-negative and the largest singular value is  $[(1-c) a_1 + c]^2 + (1-c)^2 \sum_{j>1} a_j^2 = [\nu_2(\mathcal{E})]^2$  associated with the product eigenvector  $|11\rangle$ . Therefore, one can use Theorem 1, or Corollary 2, to conclude that the channel (34) is multiplicative for  $p = 2$  when  $c > 0$ .

### 3. CPT Maps with a negative contribution from the identity

To analyze the case  $c < 0$ , write  $c = -x$  with  $x = |c| > 0$ , and recall that we assumed that the  $\{a_j\}$  are decreasing. It can still happen that all eigenvalues of  $\Omega(\mathcal{E}, 2)$  are non-negative, in which case the largest singular value is  $[(1+x) a_d - x]^2 + (1+x)^2 \sum_{j<d} a_j^2$  associated with the product eigenvector  $|dd\rangle$ . It turns out that the requirement that  $\mathcal{E}$  be CPT suffices to ensure that the eigenvalues of  $\Omega(\mathcal{E}, 2)$  are non-negative. Therefore, any CPT map of the form (34) is multiplicative for  $p = 2$ .

To see the relevance of the CPT condition, observe that the CP requirement that  $(\mathcal{E} \otimes \mathbb{1}) \left( \sum_{jk} |j\rangle\langle k| \otimes |j\rangle\langle k| \right)$  (which is the Choi matrix) is positive semi-definite holds if and only if  $B = (1+x)\rho - x \begin{pmatrix} 1 & \cdots & 1 \\ \vdots & & \vdots \\ 1 & \cdots & 1 \end{pmatrix}$  is positive semi-definite. Then  $B$  has non-negative diagonal elements, which gives

$$(1+x)a_j - x \geq 0 \quad \Rightarrow \quad x \leq \frac{a_j}{1-a_j} \quad \Rightarrow \quad \frac{x}{1+x} \leq a_j. \quad (40)$$

All  $2 \times 2$  principle minors of  $B$  are non-negative, which implies

$$(1+x)^2 a_1 a_2 - x(1+x)(a_1 + a_2) \geq 0 \quad (41)$$

Now, all eigenvalues of  $\Omega(\mathcal{E}, 2)$  will be positive if  $(1-c)^2 (\text{Tr} \rho^2) + c(1-c)(a_j + a_k) - c^2 \geq 0$  for all  $j, k$ . But the most negative of these is

$$\begin{aligned}(1+x)^2 (\text{Tr} \rho^2) - x(1+x)(a_1 + a_2) - x^2 &\geq (1+x)^2 (a_1^2 + a_2^2) - (1+x)^2 a_1 a_2 - x^2 \\ &= (1+x)^2 \left[ a_1^2 + a_2^2 - a_1 a_2 - \left( \frac{x}{1+x} \right)^2 \right] \\ &\geq (1+x)^2 [a_1^2 + a_2^2 - 2a_1 a_2] = (1+x)^2 (a_1 - a_2)^2 \geq 0,\end{aligned}\quad (42)$$

where the second inequality used (40) with  $j = 1, 2$  to conclude that  $\left( \frac{x}{1+x} \right)^2 \leq a_1 a_2$ .

### C. The Werner-Holevo channel

In our final example, we apply our condition for  $p = 3, 4$  as well as  $p = 2$ . We study the channels  $\mathcal{W}_d$  introduced in [34] to show that multiplicativity does not hold for sufficiently large  $p$ . The channel  $\mathcal{W}_d$  is defined on a  $d$  dimensional



Hilbert space as

$$\mathcal{W}_d(\gamma) \equiv \frac{1}{d-1} [(\text{Tr } \gamma) \mathbb{1}_d - \gamma^T] = \frac{1}{2(d-1)} \sum_{j < k} W_{jk}^\dagger \gamma W_{jk} \quad (43)$$

with  $\mathbb{1}_d$  the identity operator on  $\mathcal{H}$ ,  $\gamma^T$  the matrix transpose with respect to some fixed basis  $\{|i\rangle\}$ , and  $W_{jk}$  the anti-Hermitian operator  $|j\rangle\langle j| - |k\rangle\langle k|$ . (We will often suppress the subscript  $d$  and simply write  $\mathcal{W}$  for  $\mathcal{W}_d$ .) As observed in [34], any pure input state yields an output state  $\mathcal{W}(|\psi\rangle\langle\psi|)$  with eigenvalues  $\frac{1}{(d-1)^2}$  with multiplicity  $d-1$ . This implies

$$\nu_p(\mathcal{W}_d) = (d-1)^{(1-p)/p} \quad (44)$$

Werner and Holevo showed that for  $d=3$  and  $p > 4.79$  this map is not  $\ell_p$  multiplicative, by showing that maximally entangled inputs yield output  $\ell_p$  norm greater than  $(d-1)^{(1-p)/p}$ . For  $d > 3$ , they also showed that multiplicativity fails for sufficiently large  $p$ . Although their results strongly suggest that multiplicativity does hold for smaller  $p$ , they do not preclude the possibility that it fails with inputs that are partially entangled. Our results show that this cannot happen when  $p=2, 3, 4$  and  $d \geq 2^{p-1}$ .

The multiplicativity of  $\mathcal{W}$  for  $p=2$  was established in [25]. The map  $\mathcal{W}$  was also analyzed in [28] and [2]. The former proved additivity of minimal entropy; the latter multiplicativity for all  $1 \leq p \leq 2$ . Here we use Theorem 1 to show that (3) holds for  $p=2$ ; for  $p=3$  when  $d \geq 4$ ; and for  $p=4$  when  $d \geq 8$ .

For  $p=2$  it is straightforward to show that (or see Appendix C)

$$\Omega(\mathcal{W}, 2) = (\mathcal{W} \otimes \mathcal{W})(S) = \frac{1}{(d-1)^2} [(d-2)\mathbb{1} \otimes \mathbb{1} + S]. \quad (45)$$

with  $S$  the SWAP on  $\mathcal{H} \otimes \mathcal{H}$ . The eigenvalues of  $\Omega(\mathcal{W}, 2)$  can be computed from those of  $S$  which has a diagonal block with  $d$  product states  $|jj\rangle$  as eigenvectors with eigenvalue 1, and  $\binom{d}{2}$  blocks of the form  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  with eigenvalues  $+1$  and  $-1$  corresponding to the entangled states  $\frac{1}{\sqrt{2}}(|jk\rangle \pm |kj\rangle)$ . This yields eigenvalues  $\frac{1}{d-1}$  with multiplicity  $\frac{d(d+1)}{2}$  and  $\frac{d-3}{d-1}$  with multiplicity  $\frac{d(d-1)}{2}$ . For  $d \geq 3$ , these are also the singular values of  $\Omega(\mathcal{W}, 2)$ ; for  $d=2$ ,  $\frac{1}{d-1}$  is the only singular value. In both cases  $\|\Omega(\mathcal{W}, 2)\|_\infty = \frac{1}{d-1} = \nu_2(\mathcal{W})$ . Therefore, (21) is satisfied and the result follows from Theorem 1.

For  $p=3, 4$ , the results in Appendix C can be used to conclude that

$$\Omega(\mathcal{W}, 3) = \frac{1}{(d-1)^3} \left[ (d-3)\mathbb{1} + \sum_{a < b} S_{ab} - R_3 \right], \quad (46)$$

$$\Omega(\mathcal{W}, 4) = \frac{1}{(d-1)^4} \left[ (d-4)\mathbb{1} + \sum_{a < b} S_{ab} - \sum_{a < b < c} R_3(a, b, c) + R_4 \right], \quad (47)$$

where the shift  $R_3(a, b, c)$  is defined in Appendix A 2. For  $p=3$ , the multiplicativity condition is satisfied for  $d \geq 4$ ; for  $p=4$ , it is satisfied for  $d \geq 8$ .

Because  $\Omega(\mathcal{W}, p)$  is a linear combination of permutation matrices, it is easy to see that it has a large number of invariant subspaces, giving it a block diagonal structure. This structure is summarized in Table I. In this table,  $i, j, k, \ell$  always denote distinct indices, and, for readability,  $(d-1)^p \xi_{\max}$  is reported in the last three columns. This should be compared to  $(d-1)^p [\nu_p(\mathcal{W})]^p = (d-1)^p$ .

Because  $\Omega(\mathcal{W}, p)$  is a linear combination of matrices for permutation operators, all row and column sums are equal. It follows from (C3) that every row and column sum of  $\Omega(\mathcal{W}, p)$  or, equivalently, every row and column sum of each block of  $\Omega(\mathcal{W}, p)$ , is exactly  $(d-1)^{1-p}$ , which is also the value of  $[\nu_p(\mathcal{W})]^p$ . If a block has only non-negative elements, then it is a multiple of a column stochastic matrix and, hence, the column sum  $(d-1)^{1-p}$  is also the largest singular value of that block. For  $p=3, 4$ , this is true for all blocks whose basis vectors have at most two distinct indices. (To see why this is so, observe that for  $p=3$ , the only negative contribution comes from  $R_3$ , for which  $\langle jjk | R_3 | jkj \rangle = -1$  is the only non-zero element of the row corresponding to  $jjk$ . But  $\langle jjk | \Omega | jkj \rangle \geq \langle jjk | (S_{ac} - R_3) | jkj \rangle = 0$ . Similar results hold for  $|jjkk\rangle$  or  $|jkkk\rangle$  when  $p=4$ .)

Hence, we need to analyze in detail only those blocks with three or more distinct indices. This has been done by explicitly diagonalizing the operators  $\Omega(\mathcal{W}, 3)$  and  $\Omega(\mathcal{W}, 4)$  with the help of Mathematica, confirming the results stated above. (See Appendix C 3 for more details). For  $p=3$ , an analytic argument which does not require determining the eigenvalues of  $\Omega(\mathcal{W}, 3)$  and which confirms the results of Table I is presented in Appendix C 2. In Appendix C 4, some additional structure of the  $24 \times 24$  blocks is described. This leads to the conjecture that  $\mathcal{W}_d$  is  $\ell_p$ -multiplicative when the dimension  $d \geq 2^{p-1}$ .

number of blocks	size	type of vectors	non-neg elements	max sing value $\times (d-1)^p$		
				$d = 3$	$d = 4$	$d \geq 5$
$p = 3$						
$d$	$1 \times 1$	$ kkk\rangle$	yes	2	3	$d-1$
$d(d-1)$	$3 \times 3$	$ jjk\rangle$	yes	2	3	$d-1$
$\binom{d}{3}$	$6 \times 6$	$ ijk\rangle$	no	4	3	$\max\{d-1,  7-d \}$
$p = 4$						
$d$	$1 \times 1$	$ kkkk\rangle$	yes	2	3	$d-1$
$d(d-1)$	$4 \times 4$	$ jjjk\rangle$	yes	2	3	$d-1$
$\binom{d}{2}$	$6 \times 6$	$ jjkk\rangle$	yes	2	3	$d-1$
$\frac{1}{2}d(d-1)(d-2)$	$12 \times 12$	$ ijkk\rangle$	no	2	$\sqrt{13}$	$\max\{d-1, \sqrt{d^2-12d+45}\}$
$\binom{d}{4}$	$24 \times 24$	$ jjk\ell\rangle$	no	12	11	$\max\{d-1,  15-d \}$

TABLE I: Block structure of  $\Omega(\mathcal{W}, p)$ .

## V. CONCLUSION

We have extended the method introduced in [13, 14] to study the maximal  $\ell_p$ -norms of a CPT map when  $p$  is a fixed integer. This yields a sufficient condition for multiplicativity which requires only that one find the singular values of a particular matrix, rather than performing a full optimization. Although the matrix will be  $d^p \times d^p$ , it often has a block structure which makes the problems quite tractable, as shown in several examples. The condition is not necessary, but does allow us to prove new results about multiplicativity in several interesting cases, as well as providing alternative proofs of known results.

## APPENDIX A: SOME OPERATOR PROPERTIES

### 1. Hilbert-Schmidt duality

For a Hilbert space  $\mathcal{H}$  the subspace of operators satisfying  $\text{Tr} A^\dagger A < \infty$  also forms a Hilbert space (the space of Hilbert Schmidt operators) with respect to the inner product

$$\langle A, B \rangle = \text{Tr} A^\dagger B \quad (\text{A1})$$

An operator (sometimes referred to as a “superoperator”)  $\mathcal{E}$  acting on this space has an adjoint which we will denote  $\widehat{\mathcal{E}}$  and which satisfies

$$\text{Tr}[\mathcal{E}(A)]^\dagger B = \text{Tr} A^\dagger \widehat{\mathcal{E}}(B) \quad \forall A, B. \quad (\text{A2})$$

Because  $[\mathcal{E}(A)]^\dagger = \mathcal{E}(A^\dagger)$ , by writing  $C$  for  $A^\dagger$  one easily sees that (A2) is equivalent to the condition

$$\text{Tr}[\mathcal{E}(C)]B = \text{Tr} C \widehat{\mathcal{E}}(B) \quad \forall B, C. \quad (\text{A3})$$

The map  $\widehat{\mathcal{E}}$  is often called the dual of  $\mathcal{E}$  because it is defined by the duality property of the Riesz representation theorem applied to the inner product (A1). When  $\mathcal{E}$  is a CPT map of the form (1), its dual is the unital CP map with the form

$$\widehat{\mathcal{E}}(\gamma) = \sum_k A_k^\dagger B A_k. \quad (\text{A4})$$

One can verify, either directly from (A1) or by using (A4), that the dual of the map  $\mathcal{E}^{\otimes m}$  is given by the  $m$ -fold tensor product of the dual map of  $\mathcal{E}$ , i.e.  $\widehat{\mathcal{E}^{\otimes m}} = (\widehat{\mathcal{E}})^{\otimes m}$ .

## 2. Shift operators

The shift operators defined in (5) are unitary and satisfy  $L_p R_p = \mathbb{1}$  so that  $L_p^\dagger = L_p^{-1} = R_p$ . Moreover, if a vector  $|\Psi\rangle$  in  $\mathcal{H}^{\otimes p}$  has the expansion

$$|\Psi\rangle = \sum_{j_1 j_2 \dots j_p} c_{j_1 j_2 \dots j_p} |\xi_{j_1} \xi_{j_2} \dots \xi_{j_p}\rangle \quad (\text{A5})$$

then

$$L_p |\Psi\rangle = \sum_{j_1 j_2 \dots j_p} c_{j_1 j_2 \dots j_p} |\xi_{j_2} \xi_{j_3} \dots \xi_{j_p} \xi_{j_1}\rangle \quad (\text{A6})$$

$$= \sum_{j_1 j_2 \dots j_p} c_{j_p j_1 \dots j_{p-1}} |\xi_{j_1} \xi_{j_2} \dots \xi_{j_p}\rangle \quad (\text{A7})$$

so that  $L_p$  induces a right shift on the expansion coefficients. From this, it follows that, that  $L_p$  and  $R_p$  induce left and right shifts on all product states, e.g.,

$$L_p |\phi_1, \phi_2, \dots, \phi_p\rangle = |\phi_2, \phi_3, \dots, \phi_p, \phi_1\rangle \quad (\text{A8a})$$

$$R_p |\phi_1, \phi_2, \dots, \phi_p\rangle = |\phi_p, \phi_1, \dots, \phi_{p-1}\rangle \quad (\text{A8b})$$

where  $|\phi_1, \phi_2, \dots, \phi_p\rangle$  denotes  $|\phi_1\rangle \otimes |\phi_2\rangle \otimes \dots \otimes |\phi_p\rangle$ . It also follows from (A8) that the shift operators are independent of the choice of orthonormal basis in (5).

To compute operators associated with the WH-channel, it will be useful to observe that

$$L_p = \sum_{m_1 \dots m_p} |m_2 \dots m_p m_1\rangle \langle m_1 m_2 \dots m_p| \quad (\text{A9a})$$

$$R_p = \sum_{m_1 \dots m_p} |m_p m_1 \dots m_{p-1}\rangle \langle m_1 m_2 \dots m_p| \quad (\text{A9b})$$

where  $|m_j\rangle$  denotes any orthonormal basis of  $\mathcal{H}$ . It will also be useful to introduce some notation for shift operators on a subset of  $\mathcal{H}^{\otimes p}$ . For example, write  $\mathcal{H}^{\otimes 4} = \mathcal{H}_a \otimes \mathcal{H}_b \otimes \mathcal{H}_c \otimes \mathcal{H}_d$ . Then  $L_3(a, b, d)$  denotes the operator which acts as a left shift on  $\mathcal{H}_a \otimes \mathcal{H}_b \otimes \mathcal{H}_d$  and the identity on  $\mathcal{H}_c$ , i.e.,

$$L_3(a, b, d) = \sum_{m_1 \dots m_4} |m_2 m_4 m_3 m_1\rangle \langle m_1 m_2 m_3 m_4|. \quad (\text{A10})$$

The SWAP operators  $L_2(a, b) = R_2(a, b)$  play such a special role that we denote them as  $S_{ab}$ . Using the standard method for writing any permutation as a product of cycles, one can see that any shift can be written as a product of SWAP operators, e.g.  $L_3(a, b, d) = S_{ab} S_{ad}$  and  $L_4(a, b, c, d) = S_{ab} S_{ac} S_{ad}$ .

## 3. Tensor products of shifts

When the underlying Hilbert is itself a tensor product  $\mathcal{H}^{\otimes m}$ , we will let  $\mathbb{L}_p$  denotes the shift operator acting on  $p$  copies of  $\mathcal{H}^{\otimes m}$ , e.g.,  $\mathbb{L}_3|x, y, z\rangle = |y, z, x\rangle$  with  $x, y, z$  denoting vectors in  $\mathcal{H}^{\otimes m}$ . Then,  $\mathbb{L}_p = L_p^{\otimes m} = (L_{mp})^m$ . To avoid notation with double subscripts, we prove this in the case  $p = 3$ . Then

$$\begin{aligned} \mathbb{L}_3|x, y, z\rangle &= \mathbb{L}_3|x_1, x_2, \dots, x_m, y_1, y_2, \dots, y_m, z_1, z_2, \dots, z_m\rangle \\ &= |y_1, y_2, \dots, y_m, z_1, z_2, \dots, z_m, x_1, x_2, \dots, x_m\rangle \\ &= L_3^{\otimes m}|x_1, x_2, \dots, x_m, y_1, y_2, \dots, y_m, z_1, z_2, \dots, z_m\rangle \end{aligned} \quad (\text{A11})$$

where the last line follows by writing

$$L_3^{\otimes m} = (L_3 \otimes \mathbb{1} \otimes \dots \otimes \mathbb{1})(\mathbb{1} \otimes L_3 \otimes \mathbb{1} \otimes \dots \otimes \mathbb{1}) \dots (\mathbb{1} \otimes \mathbb{1} \otimes L_3 \otimes \dots \otimes \mathbb{1} \otimes L_3)$$

and observing that

$$(L_3 \otimes \mathbb{1} \otimes \dots \otimes \mathbb{1})|x_1, x_2, \dots, x_m, y_1, y_2, \dots, y_m, z_1, z_2, \dots, z_m\rangle = |y_1, x_2, \dots, x_m, z_1, y_2, \dots, y_m, x_1, z_2, \dots, z_m\rangle.$$

Note that it is also evident from (A11) that  $L_p^{\otimes m} = (L_{mp})^m$ .

#### 4. An important trace identity

We now show that for any set of operators  $\{B_1, B_2, \dots, B_p\}$  acting on  $\mathcal{H}$ ,

$$\text{Tr}_{\mathcal{H}}[B_1 B_2 B_3 \cdots B_p] = \text{Tr}_{\mathcal{H}^{\otimes p}} [B_1 \otimes B_2 \otimes \cdots \otimes B_p] L_p, \quad (\text{A12})$$

where we have introduced subscripts to emphasize that the trace in the left-hand side of Eq. (A12) is performed on  $\mathcal{H}$ , while the trace in the right-hand side is performed on  $\mathcal{H}^{\otimes p}$ . To verify (A12) observe that

$$\begin{aligned} \text{Tr}[B_1 B_2 \cdots B_p] &= \sum_{\xi_1} \langle \xi_1 | B_1 B_2 \cdots B_{p-1} B_p | \xi_1 \rangle \\ &= \sum_{\xi_1, \dots, \xi_p} \langle \xi_1 | B_1 | \xi_2 \rangle \langle \xi_2 | B_2 | \xi_3 \rangle \cdots \langle \xi_{p-1} | B_{p-1} | \xi_p \rangle \langle \xi_p | B_p | \xi_1 \rangle \\ &= \sum_{\xi_1, \dots, \xi_p} \langle \xi_1, \xi_2, \dots, \xi_p | B_1 \otimes B_2 \otimes \cdots \otimes B_p | \xi_2, \dots, \xi_p, \xi_1 \rangle \\ &= \text{Tr} [B_1 \otimes B_2 \otimes \cdots \otimes B_p] L_p, \end{aligned}$$

where a resolution of the identity operator  $\mathbb{1}$  of  $\mathcal{H}$  was inserted between the products  $B_j B_{j+1}$ .

#### 5. General permutations

Shifts are special cases of permutation operators. Let  $\Pi_p$  denote a permutation of  $\{1, 2, \dots, p\}$ . We will write  $\Pi(j) = k_j$  for the permutation that takes  $j \mapsto k_j$ . For example,  $L_p(j) = j + 1$ . One can then define a permutation operator on  $\mathcal{H}^{\otimes p}$  by

$$\Pi_p |\xi_{j_1} \xi_{j_2} \cdots \xi_{j_p}\rangle = |\xi_{\Pi(j_1)} \xi_{\Pi(j_2)} \cdots \xi_{\Pi(j_p)}\rangle \quad (\text{A13})$$

with  $\{|\xi_j\rangle\}$  an orthonormal basis for  $\mathcal{H}$  as in (5). A permutation of the indices  $\{1, 2, \dots, p\}$  induces a permutation on the  $d^p$  product basis vectors  $\mathcal{H}^{\otimes p}$  via (A13). Although we abuse notation by using the same letter for both, there should be no confusion. The permutation operator on  $\mathcal{H}^{\otimes p}$  is represented by a  $d^p \times d^p$  matrix which has precisely one 1 and  $d^p - 1$  0's in each row and column.

The permutation which takes  $k_1 \mapsto k_2 \mapsto \cdots \mapsto k_q \mapsto k_1$  is called a cycle and written  $P = (k_1, k_2, \dots, k_q)$ , i.e.,  $P(k_j) = (k_{j+1})$  with the understanding that  $P(k_q) = k_1$  and  $\Pi(j) = j$  if  $j$  does not appear as one of the  $k_i$  in the cycle. Any permutation can be written uniquely as a product of *disjoint* cycles, and the length of the disjoint cycles in  $\Pi P \Pi^\dagger$  are the same as those in  $P$ . For example  $(13) L_5 (13) = (14532)$ . If a permutation of  $\{1, 2, \dots, p\}$  has a cycle decomposition with cycles whose length is strictly less than  $p$ , then some subset of  $\{1, 2, \dots, p\}$  is invariant. A permutation  $\Pi_p$  whose shortest cycle is of length  $p$  has no invariant subsets. Permutations satisfying this condition, which is equivalent to  $(\Pi_p)^s(j) \neq j$  for  $s < p$  and  $(\Pi_p)^p(j) = j$  for all  $j$ , are of particular interest.

In fact, when all operators  $B_i = B$  are identical, (A12) can be extended to any permutation  $\Pi_p$  of  $\{1, 2, \dots, p\}$  whose shortest cycle is length  $p$ . One finds

$$\begin{aligned} \text{Tr}_{\mathcal{H}} B^p &= \sum_{\xi_1, \dots, \xi_p} \langle \xi_1 | B | \xi_{\Pi_p(1)} \rangle \langle \xi_{\Pi_p(1)} | B | \xi_{\Pi_p^2(1)} \rangle \cdots \langle \xi_{(\Pi_p)^{p-1}(1)} | B | \xi_1 \rangle \\ &= \sum_{\xi_1, \dots, \xi_p} \langle \xi_1 | B | \xi_{\Pi_p(1)} \rangle \langle \xi_2 | B | \xi_{\Pi_p(2)} \rangle \cdots \langle \xi_p | B | \xi_{\Pi_p(p)} \rangle \\ &= \sum_{\xi_1, \dots, \xi_p} \langle \xi_1, \xi_2, \dots, \xi_p | B \otimes B \otimes \cdots \otimes B | \xi_{k_1}, \xi_{k_2}, \dots, \xi_{k_p} \rangle \\ &= \text{Tr}_{\mathcal{H}^{\otimes p}} [B \otimes B \otimes \cdots \otimes B] \Pi_p = \text{Tr}_{\mathcal{H}^{\otimes p}} B^{\otimes p} \Pi_p, \end{aligned} \quad (\text{A14})$$

To see where the invariance condition is used, consider the permutation (153)(24). Attempting to apply the process above yields

$$\begin{aligned} \text{Tr}_{\mathcal{H}} B^5 &= \sum_{\xi_1, \xi_3, \xi_5} \langle \xi_1 | B | \xi_5 \rangle \langle \xi_5 | B | \xi_3 \rangle \langle \xi_3 | B^3 | \xi_1 \rangle \\ &= \sum_{\xi_1, \xi_3, \xi_5} \langle \xi_1, \xi_5, \xi_3 | B \otimes B \otimes B | \xi_5, \xi_3, \xi_1 \rangle \\ &= \text{Tr}_{\mathcal{H}_a \otimes \mathcal{H}_c \otimes \mathcal{H}_e} [B \otimes B \otimes B^3] L_3 \end{aligned}$$

or  $\text{Tr}_{\mathcal{H}} B^5 = \text{Tr}_{\mathcal{H}^{\otimes 3}} [B \otimes B^3 \otimes B] L_3$  or  $\text{Tr}_{\mathcal{H}} B^5 = \text{Tr}_{\mathcal{H}^{\otimes 2}} [B \otimes B^4] L_2$ .

## 6. Double stochastic matrices

A double stochastic matrix [16] is a matrix with non-negative elements whose row and column sums are all 1, i.e.,  $B$  is double stochastic if and only if  $b_{jk} \geq 0 \ \forall j, k$  and  $\sum_j b_{jk} = \sum_k b_{jk} = 1$ . The vector  $(1, 1, \dots, 1)$  is always an eigenvector with eigenvalue 1. Moreover, all other eigenvalues satisfy  $|\lambda_j| \leq 1$ . A permutation of  $\{1, 2, \dots, p\}$  can be represented by a matrix which has precisely one 1 and  $p-1$  0's in each row and column. This is a special type of double stochastic matrix called a “permutation matrix”. Moreover, a permutation  $\Pi_p$  of  $\{1, 2, \dots, p\}$  has no non-trivial invariant subspaces if and only if its permutation matrix is indecomposable. Note that the corresponding permutation operator on  $\mathcal{H}^p$ , represented by a  $d^p \times d^p$  matrix with precisely one 1 and  $d^p - 1$  0's in each row and column, can have invariant subspaces. In fact, it will be block diagonal.

## APPENDIX B: PROPERTIES OF LINEARIZING OPERATORS $X(\mathcal{E}, p)$

### 1. Kraus operator form of $\Omega(\mathcal{E}, p)$

We first observe that conjugation of a tensor product of operators by a shift operation induces a shift on the tensor product, e.g.,

$$L_p [B_1 \otimes B_2 \otimes \dots \otimes B_p] L_p^{-1} = B_2 \otimes \dots \otimes B_p \otimes B_1. \quad (\text{B1})$$

More generally,

$$\Pi_p [B_1 \otimes B_2 \otimes \dots \otimes B_p] \Pi_p^{-1} = B_{\Pi(1)} \otimes B_{\Pi(2)} \otimes \dots \otimes B_{\Pi(p)}. \quad (\text{B2})$$

To prove (9), one can use (1), and (B1) to see that

$$\begin{aligned} [\widehat{\mathcal{E}}^{\otimes p}(L_p)] R_p &= \left[ \sum_{k_1, \dots, k_p} (A_{k_1}^\dagger \otimes A_{k_2}^\dagger \otimes \dots \otimes A_{k_p}^\dagger) L_p (A_{k_1} \otimes A_{k_2} \otimes \dots \otimes A_{k_p}) \right] L_p^{-1} \\ &= \sum_{k_1, \dots, k_p} (A_{k_1}^\dagger \otimes A_{k_2}^\dagger \otimes \dots \otimes A_{k_p}^\dagger) (A_{k_2} \otimes A_{k_3} \otimes \dots \otimes A_{k_p} \otimes A_{k_1}) \\ &= \sum_{k_1, \dots, k_p} A_{k_1}^\dagger A_{k_2} \otimes A_{k_2}^\dagger A_{k_3} \otimes \dots \otimes A_{k_p}^\dagger A_{k_1} \end{aligned} \quad (\text{B3})$$

which gives the desired result. Moreover, using a similar argument and (B2), one finds

$$\begin{aligned} R_p [\widehat{\mathcal{E}}^{\otimes p}(L_p)] &= R_p \left[ \sum_{k_1, \dots, k_p} (A_{k_1}^\dagger \otimes A_{k_2}^\dagger \otimes \dots \otimes A_{k_p}^\dagger) L_p^{-1} (A_{k_1} \otimes A_{k_2} \otimes \dots \otimes A_{k_p}) \right] \\ &= \sum_{k_1, \dots, k_p} A_{k_p}^\dagger A_{k_1} \otimes A_{k_1}^\dagger A_{k_2} \otimes \dots \otimes A_{k_{p-1}}^\dagger A_{k_p}. \end{aligned} \quad (\text{B4})$$

Then by observing that both (B3) and (B4) involve tensor products of operators of the form  $A_{k_j}^\dagger A_{k_{j+1}}$ , one sees that after a change of variable in the summation indices, e.g,  $k_j \rightarrow k_{j-1}$  in (B4), the two expression are identical. Therefore,  $R_p$  commutes with  $\widehat{\mathcal{E}}^{\otimes p}(L_p)$  and  $\Theta(\mathcal{E}, p) = \Omega(\mathcal{E}, p) R_p = R_p \Omega(\mathcal{E}, p)$ .

### 2. General permutations

Define  $\mathcal{X}(\mathcal{E}, p)$  the set of operators  $X(\mathcal{E}, p)$  of  $\mathcal{H}^{\otimes p}$  that satisfy the property Eq. (4) for all the input states  $\gamma$  of  $\mathcal{H}$ . We have already seen that  $\widehat{\mathcal{E}}^{\otimes p}(L_p)$  is in  $\mathcal{X}(\mathcal{E}, p)$  which implies that it is non-empty. Moreover, the linearity of Eq. (4) with respect to  $X(\mathcal{E}, p)$  implies that whenever  $X(\mathcal{E}, p)$  and  $Y(\mathcal{E}, p)$  are in  $\mathcal{X}(\mathcal{E}, p)$ , then  $aX(\mathcal{E}, p) + (1-a)Y(\mathcal{E}, p)$  is

in  $X(\mathcal{E}, p)$  is also. This true for any real number  $a$  including  $a < 0$  and  $a > 1$ , and even for complex  $a$ . By choosing  $0 < a < 1$ , we can also conclude that  $\mathcal{X}(\mathcal{E}, p)$  is convex; however,  $\mathcal{X}(\mathcal{E}, p)$  is not compact. Because  $\text{Tr}[\mathcal{E}(\gamma)]^p$  is real,

$$\text{Tr} \gamma^{\otimes p} X(\mathcal{E}, p) = \overline{\text{Tr} \gamma^{\otimes p} X(\mathcal{E}, p)} = \text{Tr} [\gamma^{\otimes p} X(\mathcal{E}, p)]^\dagger = \text{Tr} [X(\mathcal{E}, p)]^\dagger \gamma^{\otimes p} = \text{Tr} \gamma^{\otimes p} [X(\mathcal{E}, p)]^\dagger \quad (\text{B5})$$

for all density matrices  $\gamma$ . Therefore, whenever  $X(\mathcal{E}, p)$  is in  $\mathcal{X}(\mathcal{E}, p)$  so are  $[X(\mathcal{E}, p)]^\dagger$  and the self-adjoint operator  $\frac{1}{2}(X(\mathcal{E}, p) + [X(\mathcal{E}, p)]^\dagger)$ .

In view of the discussion in Appendix A 4 we can also conclude that the operator  $\hat{\mathcal{E}}^{\otimes p}(\Pi_p)$  is in  $\mathcal{X}$  whenever  $\Pi_p$  is a permutation whose shortest cycle is length  $p$ . Moreover, a modification of the argument in the preceding section shows that, for these permutations,

$$\hat{\mathcal{E}}^{\otimes p}(\Pi_p) \Pi_p^\dagger = \Pi_p^\dagger \hat{\mathcal{E}}^{\otimes p}(\Pi_p) = \sum_{k_1, \dots, k_p} A_{k_1}^\dagger A_{\Pi(k_1)} \otimes A_{k_2}^\dagger A_{\Pi(k_2)} \otimes \dots \otimes A_{k_p}^\dagger A_{\Pi(k_p)}. \quad (\text{B6})$$

Since  $\Pi_p \gamma^{\otimes p} \Pi_p^\dagger = \Pi_p$  for any permutation,

$$\text{Tr} [\gamma^{\otimes p} (\Pi_p X(\mathcal{E}, p) \Pi_p^\dagger)] = \text{Tr} [(\Pi_p^\dagger \gamma^{\otimes p} \Pi_p) X(\mathcal{E}, p)] = \text{Tr} \gamma^{\otimes p} X(\mathcal{E}, p) = \text{Tr} [\mathcal{E}(\gamma)^p]. \quad (\text{B7})$$

Note that the map  $P_p \mapsto \Pi_p P_p \Pi_p^\dagger$  does not change the cycle structure of  $P_p$ , e.g, if  $P_p$  is a product of a 3-cycle and a disjoint 2-cycle, then so is  $\Pi_p P_p \Pi_p^\dagger$ . Thus,  $\Pi_p L_p \Pi_p^\dagger$  is a permutation whose shortest cycle is length  $p$  irrespective of the cycle structure of  $\Pi_p$ . One can show that  $\Pi_p [\hat{\mathcal{E}}^{\otimes p}(L_p)] \Pi_p^\dagger = \hat{\mathcal{E}}^{\otimes p}(\Pi_p L_p \Pi_p^\dagger)$ , with a similar result when  $L_p$  is replaced by any permutation whose shortest cycle is length  $p$ .

### 3. Linearizing operators for pure inputs

The set  $\mathcal{X}(\mathcal{E}, p)$  is a subset of  $\mathcal{X}_{\text{pure}}(\mathcal{E}, p)$ , the set of operators, which satisfy the property Eq. (4) when  $\gamma = |\psi\rangle\langle\psi|$  is pure. We have already observed that  $\Theta(\mathcal{E}, p) = \Omega(\mathcal{E}, p) R_p$  belongs to  $\mathcal{X}_{\text{pure}}(\mathcal{E}, p)$  but need not belong to  $\mathcal{X}(\mathcal{E}, p)$ . It follows from (B6) that the operators  $\hat{\mathcal{E}}^{\otimes p}(\Pi_p) \Pi_p^\dagger$  are also in  $\mathcal{X}_{\text{pure}}(\mathcal{E}, p)$ . In addition, for any  $X(\mathcal{E}, p) \in \mathcal{X}_{\text{pure}}(\mathcal{E}, p)$  the operators  $X(\mathcal{E}, p) \Pi_p$  and  $\Pi_p X(\mathcal{E}, p)$  are also in  $\mathcal{X}_{\text{pure}}(\mathcal{E}, p)$  for all permutations  $\Pi_p$ . This follows from

$$\begin{aligned} \text{Tr} [\gamma \otimes \gamma \otimes \dots \otimes \gamma] X(\mathcal{E}, p) \Pi_p &= \text{Tr} |\psi\rangle\langle\psi| \otimes |\psi\rangle\langle\psi| \otimes \dots \otimes |\psi\rangle\langle\psi| X(\mathcal{E}, p) \Pi_p \\ &= \text{Tr} \Pi_p |\psi, \dots, \psi\rangle\langle\psi, \dots, \psi| X(\mathcal{E}, p) \\ &= \text{Tr} |\psi, \dots, \psi\rangle\langle\psi, \dots, \psi| X(\mathcal{E}, p) \\ &= \text{Tr} [\gamma \otimes \gamma \otimes \dots \otimes \gamma] X(\mathcal{E}, p) = \text{Tr} \mathcal{E}(\gamma)^p, \end{aligned}$$

whenever  $\gamma = |\psi\rangle\langle\psi|$  is pure.

## APPENDIX C: OPERATORS FOR WERNER-HOLEVO CHANNEL

### 1. General form of $\Omega(\mathcal{W}, p)$

It follows from (43), (A9) and (6) that for the WH channel,

$$\begin{aligned} \Omega(\mathcal{W}, p) &= \sum_{\xi_1 \dots \xi_p} \mathcal{W}(|\xi_2\rangle\langle\xi_1|) \otimes \mathcal{W}(|\xi_3\rangle\langle\xi_2|) \otimes \dots \otimes \mathcal{W}(|\xi_p\rangle\langle\xi_{p-1}|) \mathcal{W}(|\xi_1\rangle\langle\xi_p|) \\ &= \frac{1}{(d-1)^p} \sum_{\xi_1 \dots \xi_p} (\delta_{\xi_2 \xi_1} \mathbb{1} - |\bar{\xi}_1\rangle\langle\bar{\xi}_2|) \otimes (\delta_{\xi_3 \xi_2} \mathbb{1} - |\bar{\xi}_2\rangle\langle\bar{\xi}_3|) \otimes \dots \otimes (\delta_{\xi_1 \xi_p} \mathbb{1} - |\bar{\xi}_p\rangle\langle\bar{\xi}_1|) \\ &= \frac{1}{(d-1)^p} \left[ d \mathbb{1} - \left( \sum_{\xi_1} |\bar{\xi}_1\rangle\langle\bar{\xi}_1| + \sum_{\xi_2} |\bar{\xi}_1\rangle\langle\bar{\xi}_2| + \dots + \sum_{\xi_p} |\bar{\xi}_1\rangle\langle\bar{\xi}_p| \right) \right. \\ &\quad \left. + \sum_{a < b} \left( \sum_{\xi_a \xi_b} |\bar{\xi}_a \bar{\xi}_b\rangle\langle\bar{\xi}_b \bar{\xi}_a| \right) - \dots + (-1)^p \sum_{\xi_1 \dots \xi_p} |\bar{\xi}_1 \bar{\xi}_2 \dots \bar{\xi}_{p-1} \bar{\xi}_p\rangle\langle\bar{\xi}_2 \bar{\xi}_3 \dots \bar{\xi}_p \bar{\xi}_1| \right] \quad (\text{C1}) \\ &= \frac{1}{(d-1)^p} \left[ (d-p) \mathbb{1} + \sum_{a < b} S_{ab} - \sum_{a < b < c} R_3(a, b, c) + \dots + (-1)^p R_p \right] \quad (\text{C2}) \end{aligned}$$

where we have used the notation introduced at the end of Appendix A 2. Note that the orthonormal basis  $\{|\xi_j\rangle\}$  can be chosen real, but even if it is not,  $\{|\bar{\xi}_j\rangle\}$  gives another orthonormal basis for  $\mathcal{H}$  for which the representation (A9) is also valid.

It is useful to compare the structure of (C2) to that of a binomial expansion. The term in square brackets is a sum shift operators  $R_k$  of order  $k = 0, 1, 2, \dots, p$ . For  $k \geq 2$  the number of  $R_k$  is  $\binom{p}{k}$  with coefficient  $(-1)^k$ . In view of (C1), the  $(d-p)\mathbb{1}$  term should be regarded as the sum of a  $k=0$  term  $d\mathbb{1}$  and a  $k=1$  term  $-p\mathbb{1}$ . The coefficient of the  $k=0$  term is anomalous, since it has the value  $d$  rather than 1. This implies that the row and column sums of the matrix representing  $\Omega(\mathcal{W}, p)$  in the orthonormal basis  $\{|\bar{\xi}_{j_1}\bar{\xi}_{j_2}\dots\bar{\xi}_{j_p}\rangle\}$  of  $\mathcal{H}^{\otimes p}$  are

$$\frac{1}{(d-1)^p} \left[ d + \sum_{k=1}^p (-1)^k \binom{p}{k} \right] = \frac{d-1}{(d-1)^p}. \quad (\text{C3})$$

One similarly finds that the sum of the absolute values of elements in any row or column sum is bounded above by

$$\frac{1}{(d-1)^p} \left[ d + \sum_{k=1}^p \binom{p}{k} \right] = \frac{(d-1+2^p)}{(d-1)^p}.$$

## 2. Singular value analysis for $p = 3$

For  $p = 3$ , only the  $6 \times 6$  blocks in  $\Omega(\mathcal{W}, 3)$  of Table I can have non-negative elements and, hence, a singular value greater than  $(d-1)^{1-p}$ . Using an ordered basis whose first three elements are  $\{|ijk\rangle, L_3|ijk\rangle, L_3^2|ijk\rangle\}$  and last three  $S_{ab}\{|ijk\rangle, S_{bc}|ijk\rangle, S_{ac}|ijk\rangle\}$ , one can write each  $6 \times 6$  block as  $(d-1)^{-p}F$  with  $F = (d-3)\mathbb{1}_6 + G$  where

$$G = \begin{pmatrix} -L_3 & V \\ V & -L_3 \end{pmatrix} \text{ and } V = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}. \text{ One then finds}$$

$$\begin{aligned} F^\dagger F &= (d-3)^2 \mathbb{1} + (d-3)[G + G^\dagger] + G^\dagger G \\ &= (d^2 - 5d + 7)\mathbb{1}_6 + \begin{pmatrix} -d+6 & 2d-8 \\ 2d-8 & -d+6 \end{pmatrix} \otimes V. \end{aligned} \quad (\text{C4})$$

Since the eigenvalues of  $V$  are 3, 0, 0, the non-zero eigenvalues of  $F^\dagger F$  are  $d^2 - 5d + 7$  (with 4-fold degeneracy) and  $(d^2 - 5d + 7) + 3[(6-d) \pm (2d-8)]$  or  $(d-7)^2$  and  $(d-1)^2$ . Now  $(d^2 - 5d + 7) \leq (d-1)^2$  when  $d \geq 2$  and  $(d-7)^2 \leq (d-1)^2$  if and only if  $d \geq 4$ . Therefore, the largest singular value is  $d-1$  when  $d \geq 4$ .

## 3. Singular value analysis for $p = 4$

For  $p = 4$ , the singular value analysis was done using Mathematica. From the discussion of Sec. IV C it suffices to consider the blocks associated with two distinct classes of mutually orthogonal subspaces of  $\mathcal{H}^{\otimes 4}$ . The first consists of those subspaces generated by the twelve orthonormal vectors  $\{|ijkk\rangle, |jikk\rangle, \dots, |kjik\rangle\}$  with  $|i\rangle, |j\rangle$  and  $|k\rangle$  distinct orthonormal elements of the canonical basis of  $\mathcal{H}$  defined implicitly in Eq. (43). On each of these subspaces,  $\Omega(\mathcal{W}, 4)$  is described by a real  $12 \times 12$  matrix. The second class consists of those subspaces generated by the twenty four orthonormal vectors  $\{|ijk\ell\rangle, |jik\ell\rangle, \dots, |kjil\rangle\}$  with distinct  $i, j, k, \ell$ . On each of these subspaces,  $\Omega(\mathcal{W}, 4)$  is described by a real  $24 \times 24$  matrix. In both cases, Mathematica is able to produce (analytically) the singular value decomposition of the corresponding matrix. These results are summarized in Table II.

## 4. Odd and even structure for $p \geq 4$

We describe in detail an additional structure of the  $24 \times 24$  block of the operator  $\Omega(\mathcal{W}, 4)$ , which gives some insight into the results reported in Tables I and II. The  $p! \times p!$  block of  $\Omega(\mathcal{W}, p)$  associated with basis vectors  $|j_1 j_1 \dots j_p\rangle$  with distinct  $j_k$  has a similar structure. Moreover, one can show that every block of  $\Omega(\mathcal{W}, p)$  is related to a  $p! \times p!$  block via a partial isometry; more precisely,  $A = XBX^\dagger$  where  $A$  is any block of  $\Omega(\mathcal{W}, p)$ ,  $B$  is one of the  $p! \times p!$  blocks, and  $X$  is a  $\dim(A) \times \dim(B)$  partial isometry satisfying  $XX^\dagger = \mathbb{1}_{\dim(A)}$ . Hence, the singular value analysis of

singular value $\times (d-1)^4$	degeneracy (12 $\times$ 12 blocks)	degeneracy (24 $\times$ 24 blocks)
$\sqrt{d^2 - 12d + 45}$	2	6
$ d-5 $	1	3
$ d-3 $	3	5
$\sqrt{d^2 - 4d + 5}$	4	6
$ d-1 $	2	3
$ d-15 $	0	1

TABLE II: Singular value decomposition of  $\Omega(\mathcal{W}, 4)$  on the twelve dimensional subspace generated by the vectors  $\{|ijkk\rangle, |jjkk\rangle, \dots, |kjik\rangle\}$  and the twenty four dimensional subspace generated by  $\{|ijk\ell\rangle, |jik\ell\rangle, \dots, |kjil\rangle\}$ . In the left column are reported the singular values of  $\Omega(\mathcal{W}, 4)$  while in the central and right column their corresponding degeneracies.

$\Omega(\mathcal{W}, p)$  can be reduced to that of the largest block. Although we did not use this in the analysis above for  $p = 3, 4$ , it allows a considerable simplification in the general case.

When  $p = 4$ ,  $p! = 4! = 24$ . For each fixed  $i < j < k < \ell$ , we can divide the 24 basis vectors into 12 which have the form  $|P_{\text{even}}(ijk\ell)\rangle \equiv P_{\text{even}}|ijk\ell\rangle$  and  $|P_{\text{odd}}(ijk\ell)\rangle \equiv P_{\text{odd}}|ijk\ell\rangle$ , where  $P_{\text{even}}$  and  $P_{\text{odd}}$  represent, respectively, an even and an odd permutation of 4 elements. Now  $\langle ijk\ell|\Pi|ijk\ell\rangle = 0$  unless  $\Pi$  is the identity permutation. Therefore  $\langle P(ijk\ell)|\Pi|\tilde{P}(ijk\ell)\rangle = 0$  unless  $\Pi = P^\dagger \tilde{P}^\dagger$ . Moreover, since the identity is an even permutation

$$\langle P_{\text{even}}(ijk\ell)|\Pi_{\text{odd}}|\tilde{P}_{\text{even}}(ijk\ell)\rangle = \langle P_{\text{odd}}(ijk\ell)|\Pi_{\text{odd}}|\tilde{P}_{\text{odd}}(ijk\ell)\rangle = 0 \quad (\text{C5})$$

$$\langle P_{\text{even}}(ijk\ell)|\Pi_{\text{even}}|\tilde{P}_{\text{odd}}(ijk\ell)\rangle = \langle P_{\text{odd}}(ijk\ell)|\Pi_{\text{even}}|\tilde{P}_{\text{even}}(ijk\ell)\rangle = 0 \quad (\text{C6})$$

Since the permutations  $S_{ab}$  and  $R_4$  are always odd, and  $R_3(a, b, c)$  (which is the only type with a negative coefficient in  $\Omega$ ) is even,  $(d-1)^p \Omega(\mathcal{W}, 4)$  can be written in the form

$$F = (d-4)\mathbb{1} + \begin{pmatrix} -A & C \\ D & -B \end{pmatrix}$$

where  $A, B, C, D$  are  $12 \times 12$  matrices with non-negative elements.

Although not essential, it is convenient to relate the ordering of odd and even permutations. Let  $P_1, P_2, \dots, P_{12}$  denote the even permutations (with  $P_1 = \mathbb{1}$ ) and  $P_{m+12} = S_{ab}P_m$  the odd (for some fixed choice of  $a, b$ ). Let  $|m\rangle = P_m|ijk\ell\rangle$ . Then  $B = S_{ab}AS_{ab}$  with  $S_{ab}$  now the matrix representative of SWAP. Since  $\langle m|\Omega|m'+12\rangle = \langle m|\Omega S_{ab}|m'\rangle = \langle S_{ab}m'|\Omega^\dagger|m\rangle = \langle m'+12|\Omega^\dagger|m\rangle$ , it follows that  $D = C^\dagger$ .  $F$  has the same singular values as

$$\begin{pmatrix} \mathbb{1}_{12} & 0 \\ 0 & -S_{ab} \end{pmatrix} F \begin{pmatrix} \mathbb{1}_{12} & 0 \\ 0 & -S_{ab} \end{pmatrix} = (d-4)\mathbb{1}_{24} - \begin{pmatrix} A & CS_{ab} \\ S_{ab}C^\dagger & A \end{pmatrix}, \quad (\text{C7})$$

and the matrix  $G = \begin{pmatrix} A & CS_{ab} \\ S_{ab}C^\dagger & A \end{pmatrix}$  is now a multiple of a column stochastic matrix. Note that both  $C$  and  $CS_{ab}$  have all row and column sums equal to  $7 = \binom{4}{2} + \binom{4}{4}$ , and that all row and column sums of  $A$  are  $4 = \binom{4}{3}$ . One easily finds two candidates for the largest singular values of  $F$

- The row sum of  $G$  is 11; it is an eigenvalue and the largest singular value of  $G$ , so that  $d-4-11 = d-15$  is an eigenvalue of  $F$ . The corresponding eigenvector of  $F$  is  $\sum_{m=1}^{12} |m\rangle - \sum_{m=13}^{24} |m\rangle = \sum_{\text{even}} P_m|ijk\ell\rangle - \sum_{\text{odd}} P_m|ijk\ell\rangle$ .
- The row sum of  $C - A$  is  $-3$ . It is also an eigenvalue of  $G$  so that  $(d-4)+3 = d-1$  is an eigenvalue of  $F$ . The corresponding eigenvector of  $F$  is  $\sum_{m=1}^{24} |m\rangle = \sum_{\text{even}} P_m|ijk\ell\rangle + \sum_{\text{odd}} P_m|ijk\ell\rangle$ .

(Note that the unitary transformation used in (C7) interchanges these two eigenvectors.) From this we can conclude that the largest singular value of  $F$  is  $\geq \max\{|d-15|, |d-1|\}$ . Since  $|d-15| = 15-d > d-1$  when  $d < 8$ , condition (21) is *not* satisfied when  $d < 8$ . If no other singular value of  $F$  exceeds  $\max\{d-1, |15-d|\}$ , then we can conclude that  $\mathcal{W}$  is multiplicative for  $p = 4$  when  $d \geq 8$ . In fact, this was done using with Mathematica, as described in Appendix C3).

For any  $p$ , the block of  $\Omega(\mathcal{W}_d, p)$  corresponding to basis vectors  $|j_1 j_1 \dots j_p\rangle$  with distinct  $j_k$  always has a similar odd/even structure. (For  $p = 3$ , this is implicit in Appendix C2. However, in that case,  $C = D = V$  has a special



form which we exploited.) One can extend the argument following (C7) to show that  $d - 2^p - 1$  and  $d - 1$  are always eigenvalues of  $F$ . It is then natural to conjecture that the largest singular value of  $\Omega(\mathcal{W}_d, p)$  is  $\max\{(d-1), |d-2^p-1|\}$ . If so, this would imply  $\ell_p$ -multiplicativity for  $\mathcal{W}_d$  when  $d \geq 2^{p-1}$ . If this conjecture is true, then for any integer  $p$  — no matter how large — the channel  $\mathcal{W}_d$  is multiplicative on spaces of sufficiently high dimension, which grows exponentially with  $p$ .

## ACKNOWLEDGMENTS

The contribution of V.G. to this paper was founded by the European Community under contracts IST-SQUIBIT, IST-SQUBIT2, and RTN-Nanoscale Dynamics. V.G. would like to thank P. Zanardi for comments and criticism. V.G. and M.B.R. are grateful to M. D'Ariano for the opportunity of participating to the Quantum Information Processing workshop in Pavia, Italy. The contributions of M.B.R. were partially supported by the National Security Agency (NSA) and Advanced Research and Development Activity (ARDA) under Army Research Office (ARO) contract number DAAD19-02-1-0065, and by the National Science Foundation under Grant DMS-0314228.

- 
- [1] K. M. R. Audenaert and S. L. Braunstein, “On strong superadditivity of the entanglement of formation”, *Commun. Math. Phys.* **246**, 443–452 (2004).
  - [2] R. Alicki and M. Fannes, “Note on multiple additivity of minimal entropy output of extreme  $SU(d)$ -covariant channels” eprint quant-ph/0407033.
  - [3] G. G. Amosov and A. S. Holevo, “On the multiplicativity conjecture for quantum channels”, *Theor. Probab. Appl.* **47**, no.1, 143–146 (2002).
  - [4] G. G. Amosov, A. S. Holevo, and R. F. Werner, “On Some Additivity Problems in Quantum Information Theory”, *Problems in Information Transmission*, **36**, 305 – 313 (2000). eprint math-ph/0003002.
  - [5] C. H. Bennett C. A. Fuchs, and J. A. Smolin, “Entanglement-enhanced classical communication on a noisy quantum channel” in *Quantum Communication, Computing and Measurement* O. Hirota, A.S. Holevo, and C. M. Caves, Eds. (New York, Plenum, 1997), pg. 79.
  - [6] C. H. Bennett and P. W. Shor, “Quantum Information Theory” *IEEE Trans. Inf. Theory* **44**, 2724 (1998).
  - [7] C. Caves and K. Wódkiewicz, “Classical phase-space descriptions of continuous-variable teleportation” eprint quant-ph/0401149.
  - [8] M-D Choi, “Completely Positive Linear Maps on Complex Matrices” *Lin. Alg. Appl.* **10**, 285–290 (1975).
  - [9] B. Davies, *Quantum Theory of Open Systems* (Academic Press, 1976).
  - [10] N. Datta, A. S. Holevo, and Y. Suhov, “A quantum channel with additive minimum output entropy” eprint quant-ph/0403072
  - [11] A. Fujiwara and P. Algoet, “One-to-one parametrization of quantum channels” *Phys. Rev. A* **59**, 3290–3294 (1999).
  - [12] A. Fujiwara and T. Hashizumé, “Additivity of the capacity of depolarizing channels” *Phys. Lett. A*, **299**, 469–475 (2002).
  - [13] V. Giovannetti and S. Lloyd, “Additivity properties of a Gaussian channel” *Phys. Rev. A*, **69**, 062307 (2004).
  - [14] V. Giovannetti, S. Lloyd, L. Maccone, J. H. Shapiro, and B. J. Yen, “Minimum Rényi and Wehrl entropies at the output of bosonic channels” *Phys. Rev. A* in press (2004). eprint quant-ph/0404037.
  - [15] A. S. Holevo, “Some estimates of the information transmitted by quantum communication channel” *Probl. Inf. Trans.* **9**, 177 (1973).
  - [16] R.A. Horn and C.R. Johnson, *Matrix Analysis* (Cambridge University Press, 1985)
  - [17] See also Section 2.1.10 of [29] and Appendix A of [24].
  - [18] C. King, “Maximization of capacity and p-norms for some product channels”, *Journal of Mathematical Physics*, **43**, no. 3, 1247 – 1260 (2002).
  - [19] C. King, “Additivity for unital qubit channels”, *Journal of Mathematical Physics*, **43**, no. 10 4641 – 4653 (2002).
  - [20] C. King, “The capacity of the quantum depolarizing channel”, *IEEE Transactions on Information Theory*, **49**, no. 1 221 – 229, (2003).
  - [21] C. King, “Maximal p-norms of entanglement breaking channels”, *Quantum Information and Computation*, **3**, no. 2, 186 – 190 (2003).
  - [22] C. King, “An application of the Lieb-Thirring inequality in quantum information theory”, to appear in Proceedings of ICMP 2003.
  - [23] C. King, M. Nathanson and M. B. Ruskai, “Multiplicativity results for entrywise positive maps” in preparation.
  - [24] C. King and M. B. Ruskai, “Minimal Entropy of States Emerging from Noisy Quantum Channels”, *IEEE Trans. Info. Theory*, **47**, 192–209 (2001).
  - [25] C. King and M. B. Ruskai, “Comments on multiplicativity of maximal p-norms when  $p = 2$ ” eprint quant-ph/0401026.
  - [26] K. Kraus, *Ann. Phys.* **64**, 311 (1971); K. Kraus, *States, Effects and Operations: Fundamental Notions of Quantum Theory* (Springer, Berlin, 1983).

- [27] K. Matsumoto, T. Shimono and A. Winter, “Remarks on additivity of the Holevo channel capacity and of the entanglement of formation” *Commun. Math. Phys.* **246**, 427–442 (2004).
- [28] K. Matsumoto and F. Yura, “Entanglement cost of antisymmetric states and additivity of capacity of some quantum channels” *J. Phys. A*, **37**, L167 (2004).
- [29] M. A. Nielsen and I. L. Chang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
- [30] A. Rényi, “On measures of entropy and information” pp. 547–561 in *Proc. 4th Berkeley Sympos. Math. Statist. and Prob.*, Vol. I (Univ. California Press, Berkeley, 1961).
- [31] M. B. Ruskai, S. Szarek, E. Werner, “An analysis of completely positive trace-preserving maps  $M_2$ ” *Lin. Alg. Appl.* **347**, 159 (2002).
- [32] A. Serafini, J. Eisert, and M. M. Wolf, “Multiplicativity of maximal output purities of Gaussian channels under Gaussian inputs” eprint quant-ph/0406065.
- [33] P. W. Shor, “Equivalence of Additivity Questions in Quantum Information Theory”, *Commun. Math. Phys.* **246**, 453–472 (2004).
- [34] R. F. Werner and A. S. Holevo, “Counterexample to an additivity conjecture for output purity of quantum channels”, *Jour. Math. Phys.* **43**, no. 9, 4353 – 4357 (2002).
- [35] P. Zanardi and D. A. Lidar, “Purity and State Fidelity of Quantum Channels via Hamiltonians” eprint quant-ph 0403074.