## Statistics of Quantum Sources

Fabio Benatti<sup>1;2</sup>, M ark Fannes<sup>1;3</sup>

Inst. Theor. Phys., K. U. Leuven, Celestijnenlaan 200D, B-3001 Leuven, Belgium
 Dept. Theor. Phys., University of Trieste, Strada Costiera 11, I-34100 Trieste, Italy

<sup>3</sup> Onderzoeksleider FW O

We shall present models of quantum sources with long \words" exhibiting dierent statistics ranging from usual independence to so-called \free independence". This can lead to statistical structures for the uctuations governed by laws other than a Gaussian distribution, such as Wigner's semicircle law.

PACS numbers: 03.67 Hk, 05.45.+b, 02.50.-r

In the eld of quantum chaos, the structure of time correlation functions is of the utmost importance to study relaxation phenomena, to single out the existence of different time scales and to perform the sem i-classical analysis [1]. Correlation functions can also be viewed as a manifestation of an underlying statistical structure that is expected to re ect intrinsic properties of the quantum tim e evolution whether regular or not. It cannot go unnoticed that there is a rem arkable relation between quantum chaos and random matrix theory on one side and, on the other, that the sem icircle distribution for uctuations, typical of random matrix theory, also emerges from the central limit theorem in a probability theory of quantum random variables without any de nite algebraic relations among them [2]. Investigating di erent possibilities becomes even more interesting if one thinks of an evolving quantum system as the sending end of a quantum communication channel, that is as a source of \quantum words" [3]. In such a case, the sender would be interested in the statistics of the source, its entropy per letter and in the distribution law for the uctuations, if any.

In this letter, we will consider the long-time predom inant statistical properties of som e quantum sources that we describe as presenting a sender with a quantum carrier described by operators  $e(t_i)$  at times  $t_1 < t_2 <$ The operator e(t) at tim e t is a xed operator e = e',  $e^2 = 1$ , speci ed at time t = 0 and evolved up to time t according to an underlying quantum evolution. Since we are only interested in the essential features of the time evolution, like regularity or random ness, we do not take into account its detailed structure, but rather resort to a schem atic description. We shall restrict ourselves to discrete tim es t = 1; 2; ... and assum e that the e ect of thedynamics on the source may be described by a so-called \bit-stream " [4,5], that is by a sequence a (1); a (2); ... of zeroes and ones xing the com mutation relations between operators at di erent tim es s;t = 1;2;::::

$$e(s + t) e(s) = (1)^{a(t)} e(s) e(s + t)$$
: (1)

O by iously, these commutation relations strongly depend on the statistical properties of the bit-stream . The sender will use  $\,$  nite sequences of letters from the source thus forming ordered words w (t), where t is a multi-index (t\_1 < t\_2 <  $\,$  \_ n)  $\pm$  Such ordered words are then  $\,$  nite products

$$w(t) := e(t_1)e(t_2) \qquad \text{ne}(t; \qquad (2)$$

with  $t_1 < t_2 < f$ . If the sender has an ensemble of asynchronous sources with the same bit-stream at his disposal, he would construct more general (not time-ordered) words, e.g. w (s) w (t), simply by concatenation of ordered words w (s), w (t) from dierent sources. In particular, it can happen that in the word w (s) w (t) some of the indices  $s_k$  equal some of the indices t. In general, the probabilities for the various words w (s) w (t) w (u) that the sender can construct, are specified by their expectations

$$hw (s) w (t) w (u)i; (3)$$

with respect to a given state of the ensemble of sources. Due to the freedom in constructing words, there are in general no preferred words to single out apart from the identity (all letters being 1). In such a case, a meaningful statistics can only arise from the expectations

$$hw (t)i = 0 ; hli = 1 :$$
 (4)

Notice that, in the usual statistical framework, the expectations in (3) are nothing else than multitime correlation functions. The dynamics during a single time step is given by the shift on the indices of the letters:

$$w(t)$$
 7  $w(t+1) := e(t_1+1)e(t_2+1)$   $e(t_1+1)e(t_2+1)$ 

In spite of their extreme simplicity, the variety of statistics brought about by the expectations in (4) together with the bit-streams is nevertheless noticeable. An extreme case is a set of asynchronous quantum sources giving rise to the so-called \free shift" [6] whose words have no algebraic relations of any sort between their letters e(t) at dierent times, still keeping e = e and

 $e^2=1$ . W ith dynam ics and statistics xed by (5), respectively (4), the sender has no bit-stream structure that he can use to simplify words by commuting letters one over another according to (1).

We shall be interested in the large-time behaviour of time correlation functions of non-commutative dynamical systems. Such a question has a much broader eld of applications than that sketched above and it is therefore better to consider the more general setting of (quantum and classical) dynamical systems of which quantum sources are particular cases.

In quantum mechanics, one usually works with operators X on a Hilbert space H and the Heisenberg evolution generated by some Hamiltonian H

Then, tim e-invariant expectations

$$X$$
7  $hX$ i =  $h$ 7 $X$ j  $i$ ; (7)

are computed by means of a suitable \ground-state"  $H \neq i = 0$ .

In classical mechanics, one has a phase space X , a dynamical (Hamiltonian) ow connecting phase points (q;p) through trajectories (q;pt) and a time-invariant phase density distribution , e.g. the canonical ensemble. It is, however, convenient to adopt an algebraic description in this case too, using complex-valued functions f on X evolving in time according to

$$f(q;p)$$
 7  $f_t(q;p) = f(q_t;p_t)$ : (8)

Expectations are obtained by averaging w.r.t.:

These expectations are time-invariant because is stationary.

In this way, classical and quantum systems can be treated on the same footing. Formally, the only dierence is the algebra considered: commutative in the second one. Nevertheless, this has quite profound consequences on the probabilistic structure of the theory. In fact, one of the hard problems for truly quantum systems is to understand the implications of positivity, i.e. hX  $^{\rm Y}{\rm X}$  i 0, on the structure of the expectations.

The notion of mixing is expressed for both classical and quantum systems by the following request:

$$\lim_{t \to 0} hX Y (t)Z i = hX Z ihY i :$$
 (10)

In the classical case, Z can be commuted over Y (t) so that two observables X and Y su ce: if X = f and Y = g are functions which vanish outside some nite-volume regions F, respectively G, in phase space, (10)

expresses the uniform spreading of G throughout X by the phase space ow. In some cases, m ixing is asking for too m uch and one has to replace the  $\lim_{n \to \infty} i t$  with a time average, in which case the system is called ergodic.

The analogy between classical and quantum systems can be pushed even a bit further by considering the Hilbert space of square summable functions f on the phase space X and (non-commuting) linear operators on it such as jfihfj [7]. The scalar product between functions f and g is computed with respect to the phase space density:

$$\frac{Z}{\text{hf jgi}} := \frac{dqdp}{dqdp} (q;p) \overline{f(q;p)} g(q;p) : (11)$$

Notice that the classical functions act on this Hilbert space simply as multiplication operators: f jg(q;p)i: jf(q;p)g(q;p)i. In this way, we can compute the expectation in (9) as the average of such an f w r.t. the constant wave function 1 on X: hfi = hl fjli. The expectation of a non-classical observable jfihfji is now given by hfihfji = jhl ffjlif. Finally, the dynam ics shifts jfi to jft with jt as in (8).

For quantum systems, a much desired property is that, asymptotically in time, observables commute, namely that

$$\lim_{t} [X;Y(t)] = 0$$
 (12)

as an operator on H . This property is expected to be typical of quantum systems with many freedoms. In the hierarchy of irregular behaviours, next to mixing one and multi-clustering:

$$\lim_{jt_{1}} \lim_{t_{1}, j!} hX^{(1)}(t_{1})X^{(2)}(t_{2}) \qquad {^{(n)}}X(t_{n})i =$$

$$= Y^{n}$$

$$= hX^{(1)}i: \qquad (13)$$

Should operators commute when separated by large tim es, we could make  $t_1 < t_2 < f$  in (13).

In general, one can consider time-asymptotics of correlation functions where consecutive times are well-separated, but times not next to each other may be equal: notice that the words in (3) texactly into this framework. Namely, we shall consider multitime correlation functions of the form:

$$hX^{(1)}(t_1)X^{(2)}(t_2)$$
  $hX^{(n)}(t_n)i;$  (14)

where  $t_1 \in t_2 \in \mathbb{R}$ , but, possibly,  $t_j = t_{j-k}$  when  $k \in 1$ .

Let  $X^{(1)}$ ;  $X^{(2)}$ ; ...;  $X^{(n)}$  be n observables at time t=0. Because of the time evolution, in spite of possible algebraic relations between them, no typical commutation relations might exist between  $X^{(k)}$  and  $X^{(k)}$  (t) for

large t (unless observables separated by large times commute). The only meaningful algebra is what we shall call the \asymptotic free algebra" consisting of (linear commbinations of) words X  $_{i_1}^{(1)}$  X  $_{i_2}^{(2)}$   $_{i_n}^{(n)}$  X. The letters X  $_{i_n}^{(\gamma)}$  are asymptotics, in a sense which will be made clear later on, of time zero observables X  $^{(\gamma)}$ . We will adopt the following rules:

- a) whenever the identity appears it can be dropped;
- b) whenever two consecutive observables  $X_{i_k}^{(k)}$  and  $X_{i_{k+1}}^{(k+1)}$  carry equal subindices  $(i_k=i_{k+1})$ , then they must be considered as the single observable  $X^{(k)}X^{(k+1)}$  pertaining to a same time variable  $t_{i_k}$  and obtained through the corresponding limit.

W ords are multiplied by concatenation without any simpli cation rule between consecutive letters except for a) and b) above.

Expectations on the asymptotic free algebra are computed starting from the multi-time correlation functions:

$$hX_{i_{1}}^{(1)}X_{i_{2}}^{(2)} = \lim_{i_{n}} hX_{i_{1}}^{(n)} = \lim_{i_{1}} hX_{i_{1}}^{(1)} (t_{i_{1}}) X_{i_{2}}^{(2)} (t_{i_{2}}) = \lim_{i_{1} \to \infty} hX_{i_{1}}^{(n)} (t_{i_{1}}) X_{i_{2}}^{(n)} (t_{i_{2}})$$
(15)

where  $_k := t_{i_k}$   $t_{i_{k+1}}$  j. This should be compared with the time  $\lim$  its in the formula (13) for multi-clustering. More in general, one should consider rather ergodic time averages than  $\lim$  its:

These expectations return positive values when used to compute expectations of positive operators and thus the left-hand side member of (16) allows for a consistent probabilistic interpretation.

We want to emphasize that if (12) and (13) hold, as it is generally expected for extended quantum systems, then (15) and (16) can be readily computed. Indeed, one rst has to consider in the denitions (15) or (16) the subsets  $_{\rm j}$  of equal times, i.e. i\_k = i\_ if and only if k and 'belong to the same  $_{\rm j}$ , and then, commuting operators which are separated by large times, one may group together the corresponding observables: of course, one has to keep the original order among the grouped observables that, usually, need not commute. Finally, one gets

$$hX_{i_{1}}^{(1)}X_{i_{2}}^{(2)} \qquad {}^{(n)}_{i_{n}}Xi_{1} = {}^{Y} \quad D \quad Y \qquad {}^{K} \quad E \\ j \quad k2 \quad j \qquad (17)$$

Assuming asymptotic commutativity and multi-time clustering, we now present explicitly a few expectations choosing for notational simplicity X  $^{(1)}$  = A, X  $^{(2)}$  = B

and so on. Remember that the subscripts refer to the times with respect to which the limits in (15) and (16) are computed, so that equal subscripts mean that equal times have been considered:

$$hA_{1}i_{1} = hAi$$

$$hA_{1}B_{2}i_{1} = hAihBi$$

$$hA_{1}B_{2}C_{1}i_{1} = hACihBi$$

$$hA_{1}B_{2}C_{1}D_{2}i_{1} = hACihBDi$$
(18)

$$\frac{1}{N} = \frac{X^{N}}{N} X_{j}; \qquad (19)$$

are distributed according to a Gaussian law [8].

In full generality, we observe that it is easy to construct asym ptotic free algebras for discrete-time dynamical systems and thus to include the models of quantum sources we are considering. In fact, we shall use (15) where limits have to be taken w.r.t. discrete times.

We shall now consider the extreme instance of the quantum source that we referred to as the \free shiff". It is rather obvious that system observables do not commute, even when largely separated in time. We shall presently show that the expectations  $h_{\ i}$  on the asymptotic free algebra of the free shiff exhibit a totally different kind of independence from that expressed in (18), called \freeness". The latter is de ned by the following decoupling scheme [2]:

$$hX_{i_1}^{(1)}X_{i_2}^{(2)}$$
  $i_n^{(n)}Xi_1 = 0$ ; if  $hX^{(')}i = 0$ ; (20)

with 
$$'= 1; :::; n \text{ and } i_1 \in i_2 \in \mathbb{R}$$
. i

In order to prove the assertion, we observe that general centered observables K, i.e. hK i = 0, are obtained by linear combinations of words because of (4). It is therefore necessary and sulcient to consider the time  $\lim_{n \to \infty} \frac{1}{n} = 1$ 

$$\lim_{ \, \dot{t}_{i_k} \ \ \, t_{i_{k+1}} \, j! \, \, 1 } \, h \! \! w^{\, (1)} \, (t_{i_1}) \, w^{\, (2)} \, (t_{i_2}) \qquad {}^{(n)} \! \! w (t_{i_n}) \, \dot{\text{i}} \, : \eqno(21)$$

For large dierences between any two consecutive times, there cannot be simplications due to  $e(t)^2 = 1$ . Therefore, products of words as in (21) will have zero expectation in the limit because of (4). Unlike in the case of standard independence as in (18), freeness leads to a new distribution law for uctuations of asymptotic words, namely to the Wigner semicircle law [2].

It is instructive to compare the expectations in (18) with the ones obtained from freeness (notice that the observables below, like those in (18), need not be centered):

$$\begin{array}{rcl} hA_1i_1 &= hAi\\ hA_1B_2i_1 &= hAihBi\\ hA_1B_2C_1i_1 &= hACihBi\\ hA_1B_2C_1D_2i_1 &= hACihBihDi+hAihBDihCi\\ && hAihBihCihDi \end{array} \eqno(22)$$

If the quantum sources, presented at the beginning of the paper, are governed by generic bit-stream s, their observables will generally not commute when largely separated in time. Therefore, we do not expect the usual independence to hold for the expectations on the asymptotic free algebra. However, neither do we have freeness. Let us consider the simple word w = e and words constructed with alternating products of  $e(t_1)$  and  $e(t_2)$ ,  $t_1 \in t_2$ . The rst, possibly non-zero, expectation of such a word is

he 
$$(t_1)$$
e  $(t_2)$ e  $(t_1)$ e  $(t_2)$ i =  $(1)^{a(jt_2-t_1)}$ : (23)

For irregular bit-stream s, the lim it will not exist and, in order to construct expectations on the asymptotic free algebra, we should use an averaging procedure as in (16). Even if the bit-stream is su ciently irregular to have

$$he_{1}e_{2}e_{1}e_{2}i_{1} = \lim_{T_{1};T_{2}!} \frac{1}{T_{1}T_{2}} \frac{\vec{X}^{1} \quad \vec{X}^{2}}{t_{1}=0 \ t_{2}=0} (1)^{a \ (\hat{J}_{2} \quad t_{1} \ \hat{J})}$$

$$= 0; \qquad (24)$$

nevertheless

$$h(e(t_1)e(t_2))^4 i = 1 = h(e_1e_2)^4 i_1$$
: (25)

Notice that this is the rst non-zero expectation of alternating products.

As yet a dierent model source with simple statistics, we consider a classical ow (q;p) 7  $(q;p_t)$  with mixing properties on phase space, namely:

$$\lim_{t \to 1} \inf jg_t i = \lim_{t \to 1} \int_X dq dp \quad (q;p) \overline{f(q;p)} g(q_t;p_t)$$

$$= \inf i hgi; \qquad (26)$$

where the Hilbert space description for classical systems previously outlined has been used. One can then deduce that

$$\lim_{t \mid 1} hR (f)R (g_t)i = hR (f)ihR (g)i;$$
 (27)

where  $R(f) = f \sinh f$ . The above product structure extends to the set of nite rank matrices and can be used to construct a corresponding asymptotic free algebra with expectations

$$hA_{1}B_{2}$$
  $_{n}iF_{1} =$ 

$$= \lim_{j:t_{i_{k}} t_{i_{k+1}}j!} h1_{j}A_{1}(t_{i_{1}})B_{1}(t_{i_{2}}) \qquad F_{n}(t_{j})i$$

$$= hAihBi \qquad hFi: \qquad (28)$$

Notice that the identity operator 1 is not a nite rank matrix and, in order to construct centered observables K := A hAil, one has to add it to the nite rank operators. Such a dynamical system is not commutative nor asymptotically commutative and therefore, the usual statistical independence (17) and thus a Gaussian distribution of uctuations is not expected to hold. Freeness does not show up either as a property of the asymptotic free algebra. Indeed, considering centered observables K, K, K and K, one can prove that

$$hA_{1}i_{1} = 0$$

$$hA_{1}B_{2}i_{1} = 0$$

$$hA_{1}B_{2}C_{1}i_{1} = hAihBihCi \quad hBihACi$$

$$hA_{1}B_{2}C_{1}D_{2}i_{1} = 0$$
(29)

Notice that the previous way of extending a property of the classical time evolution, in this case phase space mixing, to a \quantum system "was proposed in [7] to provide a counterexample to the claimed incompatibility between chaos and quantum mechanics. Subsequently, a more physical application of these ideas was given in [9]. One wonders whether the decoupling scheme (29) might be a typical signature of true chaotic behaviour in quantum systems.

We conclude by observing that freeness seems to require a total lack of any algebraic structure between observables at dierent times. Multi-clustering for time-ordered correlation functions, such as in (13), is not sufcient to draw any conclusion about correlation functions where equal times appear as in (14). We also notice from the bit-stream example, see (23) and (24), and also from (29) that increasing random behaviours bring us closer to freeness in the sense that more and more asymptotic expectations vanish.

M oreover, we rem ark that, despite their dierent statistical properties, all the previous models of quantum sources share a common feature. Namely, the states on the asymptotic free algebras are invariant w.r.t. to permutations  $(i_1; \dots; i_n)$  7  $((i_1); \dots; (i_n))$  of the lower indices of their arguments:

$$hX_{i_1}^{(1)}X_{i_2}^{(2)} \qquad \stackrel{(n)}{\underset{i_n}{x_n}} i_1 = hX_{(i_1)}^{(1)}X_{(i_2)}^{(2)} \qquad \stackrel{(n)}{\underset{(i_n)}{x_n}} i_1 : (30)$$

It seems thus worthwhile to investigate the general structure of permutation invariant states on asymptotic free algebras.

We warm by thank A. Verbeure for many illuminating suggestions and  $com\ m$  ents.

O ne of the authors (F B .) acknow ledges  $\,$  nancial support from the O nderzoeksfonds K JJ Leuven F/97/60 and the Italian IN F N .

- [L] G. Casati and B. Chirikov, Quantum Chaos (Cambridge University Press, 1995)
- [2] D. V. Voiculescu, K. J. Dykem a, and A. Nica, Free Random Variables (Am erican M athematical Society, Providence, Rhode Island, USA, 1991)
- [3] C.P.W illiams and S.H.C. learwater, Exploration in Quan-

- tum Computing (Springer Telos, New York, Berlin, Heidelberg, 1998)
- [4] G L.Price, Canad.J.Math.39, 492 (1987)
- [5] R.T. Powers, Canad. J.M ath. 40, 86 (1988)
- [6] E.St rm er, Invent.M ath.110,63 (1992)
- [7] M . Berry, True quantum chaos? An instructive example, Proceedings of Yukawa sym posium (Tokyo, 1990)
- [8] D. Goderis, A. Verbeure, and P. Vets, Probab. Th. Rel. Fields 82, 527 (1989)
- [9] R A . Pasm anter, Phys. Rev. A 42, 3622 (1990)