

The Measurement Problem and the Reduction Postulate of Quantum Mechanics

Rodolfo Gambini
 Instituto de Física, Facultad de Ciencias,
 Tristan Narvaja 1674, 11200 Montevideo, Uruguay

It is well known, that the causal Schroedinger evolution of a quantum state is not compatible with the reduction postulate, even when decoherence is taken into account. Here, we analyze the possibility of considering the standard reduction postulate as a good approximation, useful for practical purposes, of a modified version of the postulate, consistent with the Schroedinger evolution, which now is universally valid, and via decoherence with the experimental results. Within this approach, the physical role played by the reduction postulate is as a tool for the computation of relative probabilities and consequently for the determination of the probabilities of consistent histories. From a more philosophical standpoint the reduction of the state is the counterpart in the formalism to the production of events.

It is well known that when two quantum systems temporarily interact, the composite system is generally left, after the interaction, in an "entangled state". That means that the state vector is not just a product of the states vectors corresponding to each system. This raises difficulties if one of the systems is macroscopic, more specifically if it is a measuring apparatus. After the measurement a linear superposition of macroscopically different states of the measuring apparatus is obtained.

It is well known that Bohr's solution to this problem is based on the following assumption: a macroscopic object is not subject to the rules of quantum mechanics but only to classical physics. Even though this rule removes the difficulty, it raises [1] a host of further questions: When is an object big enough to be considered as classical rather than quantum? What laws of physics describe the transition between the quantum world and the classical one? Is it logically consistent to impose classical physics when quantum mechanics find a difficulty?

If one wants to view the quantum theory as the general dynamical theory of all the physical systems, and not just microsystems in the laboratory, difficulties seem to be even harder. One needs to explain why the dynamics expressed by the Schroedinger equation is suddenly rejected in some very special interactions called measurements. This is even more surprising if one takes into account that the same Schroedinger equation is applied to a host of macroscopic objects as long as they are not involved in measurement processes. How could one justify the reduction postulate whose role is to contradict the causal law of evolution and, at the same time, is put outside the reach of the experimental verification?

Decoherence [2], [3], [4] allows to explain the evolution of the density operator of a macroscopic object into an *almost* diagonal matrix, when only collective observables are taken into account.

It can take place, for instance, in a measuring apparatus and destroy any *practical* possibility of observing a quantum superposition of distinct macroscopic events. However if quantum mechanics is exactly correct and the Schroedinger equation rigorously describes the dynamical evolution, the phase coherence is never completely lost. This is the core of the measurement problem that is not completely solved by decoherence: how to combine the observation of an event by an observer O with the fact that the phase is never lost and therefore the information about the other branches is still present, even if not for O . The measurement problem put in a nut shell is the following: The postulate of causal evolution according to Schroedinger and the postulate of reduction of the state cannot hold simultaneously, even when decoherence is taken into account. If one assumes the universal evolution according to the Schroedinger equation, then, the reduction of the density operator from an almost diagonal matrix to a rigorously diagonal one, as it is required by the reduction postulate, cannot occur [5]. But it is not so easy to get rid of the reduction postulate. In fact, if the reduction does not happen, then the events observed by O cannot belong to exactly consistent histories but only satisfy the consistency requirement in some approximation. But if that is the case, what is the counterpart of the production of events in the formalism? What is even more important, if reduction does not occurs when an event is produced it is not even possible to assign probabilities to alternative histories.

Here, I will show that there is a very natural modification of the reduction postulate consistent with the rest of the quantum formalism and with the experimental results that allows to solve the measurement problem. In other words, the new reduction postulate is always consistent with the Schroedinger evolution.

Let us first recall the standard form of the reduction postulate. We consider a system S associated with the Hilbert space $\mathcal{H}$ whose state is given by the density operator ρ , which is a self adjoint positive definite operator with unit trace. Let P_1 be the projector to a certain subset $\mathcal{E}_1$ of the spectrum of an observable operator A . Then, according to standard quantum mechanics, the probability of obtaining one of the eigenvalues of A belonging to the subset $\mathcal{E}_1$ is given by

$$p_1 = \text{Tr}[\rho P_1] \quad (1)$$

and according to the reduction postulate, the corresponding density operator after the measurement is given by

$$\rho_1 = \frac{P_1 \rho P_1}{\text{Tr}[\rho P_1]} \quad (2)$$

Thus if one wants to describe the state of the system after the measurement has taken place, without including the information about the particular outcome of the measurement, it will be given by the density operator

$$\rho_F = \sum_n p_n \rho_n = \sum_n P_n \rho P_n \quad (3)$$

If, for instance the system under consideration is the Zurek's 1982 bit-by-bit model of a spin 1/2 [3] particle coupled with a measuring apparatus, the density operator after the decoherence process is described by

$$\rho = \begin{pmatrix} |a|^2 & z(t) \\ z^*(t) & |b|^2 \end{pmatrix}. \quad (4)$$

where $z(t)$ is a very small quantity. If the reduction of the state takes place, according to Eq 3, it will take the form

$$\rho_F = \begin{pmatrix} |a|^2 & 0 \\ 0 & |b|^2 \end{pmatrix}. \quad (5)$$

Even though the change from ρ to ρ_F would be "fantastically" difficult to detect, this change is a sign of the violation of the causal evolution described by the Schroedinger equation. A fundamental physical law cannot be accepted with the argument that it is very difficult to prove that is false.

We shall now explore the possibility of introducing modifications of the reduction postulate rigorously consistent with the Shroedinger evolution. According to the new postulate the evolution of a state will be always described by the Schroedinger equation, but when an event is observed the information about the density matrix will be actualized. We will therefore be interested in first place in the systems where events are observed. These systems are the macroscopic measurement apparatus coupled with a quantum system. Their states will be described by a partial trace after decoherence has taken place. The density matrix, in the "pointer basis" associated with the macroscopic states, presents after decoherence the well known dumping of the interference terms

$$|\rho_{ij}|/\rho_{ii} \ll 1 \text{ and } |\rho_{ij}|/\rho_{jj} \ll 1 \quad (6)$$

A one parameter family of possible reduced states consistent with Schroedinger can be obtained as follows. If the measurement of the observable A leads to a value lying in the set $\mathcal{E}_1$, the density operator after the measurement may be given by the self-adjoint unit trace matrix

$$\rho_1 = \frac{1}{Tr[\rho P_1]} \{1/2[\rho P_1 + P_1 \rho] - \lambda [\sum_{j \neq 1} |\rho_{1j}|] P_1 + \lambda \sum_{k \neq 1} |\rho_{1k}| P_k\} \quad (7)$$

where λ is an arbitrary real constant. For each value of λ one has an alternative for the reduction postulate. If λ is grater or equal to 1, the density matrix is positive definite. When the outcome of the measurement is unknown, the state of the system after the measurement has taken place is given by

$$\rho_F = \sum_n p_n \rho_n = 1/2 \sum_n (\rho P_n + P_n \rho) = \rho \quad (8)$$

and therefore the reduction process does not imply any abrupt change in the density operator. In other words the reduction of the state and the Scroedinger evolution are now totally compatible. Notice that the form of the state after the measurement differs from the usual expression (2) by interference terms and therefore due to decoherence the modification of the reduction postulate is undetectable in the measurement process.

In order to single out an alternative, the arbitrary constant λ should be fixed by physical considerations. To do that we notice that the reduced state depends explicitly on the pointer basis and therefore is not well suited to describe generic events, it only allows to describe events taking place in a measuring apparatus. A reduction postulate should allow to assign probabilities to any kind of event as the standard postulate does.

If one wants to include microscopic quantum events in the description and assign probabilities to histories, we would not dispose of a preferred pointer basis and therefore the only possible definition of the reduced state is the one obtained for $\lambda = 0$. For this value of λ the reduced density

operator is not always positive definite, but, as we shall prove, it allows to assign probabilities to consistent families of histories and practice retrodiction.

Reduction is the counterpart in the formalism of the production of events. In our approach, reduction can occur at any instant of time provided a "choice" or "branching" among different possibilities occur. That is, if P is the projector corresponding to the event, provided

$$0 < p = \text{Tr}[\rho P] < 1 \quad (9)$$

If this condition is fulfilled, then the property associated with P can be actualized and, in that case, reduction will take place. The density operator will take the form

$$\rho_P = \frac{1}{\text{Tr}[\rho P]} 1/2[\rho P + P\rho] = \frac{1}{\text{Tr}[\rho P]} 1/2[P, \rho]_+ \quad (10)$$

It is now possible to assign probabilities to histories. A family of histories will be consistent if for each set of projectors, of the observables under consideration, on disjoint domains, whose union at a given time, covers the whole spectrum [6–8], and for each union of sets, one can assign a probability p such that

$$0 \leq p = \text{Tr}[P_n(t_n)[P_{n-1}(t_{n-1}), \dots, [P_1(t_1), \rho]_{++}]_+ \leq 1 \quad (11)$$

satisfying the positivity condition. Notice that consistent histories are automatically additive. If $P_i = P^1_i + P^2_i$ then $p = p^1 + p^2$, due to the linearity of the reduction postulate in the projectors. Contrary to what happens with the standard postulate, the main requirement is positivity while additivity is automatically satisfied

Due to decoherence, the consistency condition for macroscopic events would be practically always fulfilled. However, for some observables "fantastically" difficult to observe, events would not occur. For instance if one starts from the state given in Eq[4] and the outcome of the first measurement is $s_z = +1/2$ and one tries to repeat the measurement with a second apparatus rotated with respect to the first an angle of the order of the dumped interference terms, condition (9) could fail. Such an angle (of the order of 2^{-N} , where N is the number of degrees of freedom of the environment) is completely impossible to introduce.

The density operator resulting from a macroscopic measurement will have some negative eigenvalues with absolute values of the order of the interference terms and therefore it could be called quasi-positive definite. Due to the unitarity of the Schroedinger evolution it is immediate to check that the quasi-positivity of the density operator resulting from a macroscopic measurement is conserved by the evolution.

To conclude, we have proposed a modification of the reduction postulate. Due to decoherence, this postulate practically coincides with the usual one, leading after a measurement process to a state which differs from the standard projected state by terms proportional to the interference components in the decohered matrix, and insures an evolution rigorously consistent with the Schroedinger equation. It allows to assign probabilities to histories and consequently to speak of quantum events. The reduction of the state in a macroscopic system corresponds to an actualization of the information available for the observers of the event [9] while the evolution of the Universe is always described by the Schroedinger equation. The new postulate involves the use of non positive definite density matrices as intermediate steps in the calculation of probabilities for consistent histories. As we have stressed these matrices are never obtained as the outcome of a macroscopic measurement. In this approach the physical role played by the reduction postulate

is as a tool for the computation of relative probabilities and consequently for the determination of the probabilities of consistent histories. From a more philosophical standpoint the reduction of the state is the counterpart in the formalism to the production of events.

- [1] R. Omnès, *The Interpretation of Quantum Mechanics*, Princeton Series in Physics, Princeton University Press, Princeton, New Jersey (1994).
- [2] K.Hepp and E.H. Lieb, *Helv. Phys. Acta* **46**,573(1973).
- [3] W.H.Zurek, *Phys.Rev.* **D 26**, 1862,(1982).
- [4] E. Joos and H. D. Zeh, *Z.Phys.* **59**,223 (1985).
- [5] B. d’Espagnat, *Found.Phys.* **20**,1147 (1990)
- [6] R.B.Griffiths, *J. Stat. Phys.***36**,219 (1984).
- [7] R. Omnès, *J. Stat. Phys.***53**,893 (1988).
- [8] M. Gell-Mann and J. B. Hartle, in *Complexity, Entropy and the physics of Information*, W. H. Zurek ed., Santa Fe Institute Studies in the Science of Complexity, No.8, Addison-Wesley, Redwood City, California (1990).
- [9] C. Rovelli, *International Journal of Theoretical Physics*,**35**,1637 (1996)