

Statistical Mechanics of a Trapped Bose–Einstein Condensate ^{1 2}

Luca Salasnich

Istituto Nazionale per la Fisica della Materia, Unità di Milano,
Dipartimento di Fisica, Università di Milano,
Via Celoria 16, 20133 Milano, Italy
Istituto Nazionale di Fisica Nucleare, Sezione di Padova,
Via Marzolo 8, 35131 Padova, Italy
Dipartimento di Matematica Pura ed Applicata,
Università di Padova, Via Belzoni 7, 35131 Padova, Italy

Bose–Einstein condensation (BEC) in a gas has now been achieved [1]. Alkali atoms (^{87}Rb , ^{23}Na and ^7Li) have been cooled to the point of condensation (temperature of 100 nK) using laser cooling and trapping, followed by magnetic trapping and evaporative cooling. This important experimental result has also renewed the interest on theoretical studies of BEC [2]. In this contribution we discuss the statistical mechanics of the trapped BEC at zero temperature [3]. In particular, we study the stability of the condensate by using a variational method with local and non–local interaction between the particles [4].

The quantum Hamiltonian operator of a gas of N identical bosonic particles of mass m and without spin in a trap is given by

$$\hat{H} = \sum_{i=1}^N \left(-\frac{\hbar^2}{2m} \nabla_i^2 + V_0(\mathbf{r}_i) \right) + \frac{1}{2} \sum_{ij=1}^N V(\mathbf{r}_i, \mathbf{r}_j), \quad (1)$$

where $V_0(\mathbf{r})$ is the trapping potential and $V(\mathbf{r}, \mathbf{r}')$ is the interaction potential [3]. Let $\Psi_\alpha(\mathbf{r})$ be the single particle wave function, where α represents a complete set of quantum numbers. The totally symmetric many–particle wave function with quantum numbers $\alpha_1, \dots, \alpha_N$ results

$$\Phi_{\alpha_1 \dots \alpha_N}(\mathbf{r}_1, \dots, \mathbf{r}_N) = \sqrt{\frac{N_{\alpha_1}! \dots N_{\alpha_N}!}{N!}} \sum_{\hat{P}} \Psi_{\hat{P}\alpha_1}(\mathbf{r}_1) \dots \Psi_{\hat{P}\alpha_N}(\mathbf{r}_N), \quad (2)$$

where \hat{P} is the permutation operator and N_{α_i} is the number of particles in the state α_i . The system is a Bose–Einstein condensate if all the N particles are in the same quantum state α . Thus, for $\alpha = \alpha_1$, we have $N_{\alpha_1} = N$, $N_{\alpha_2} = \dots = N_{\alpha_N} = 0$ and the many–body wave function reads

$$\Phi_\alpha(\mathbf{r}_1, \dots, \mathbf{r}_N) = \Psi_\alpha(\mathbf{r}_1) \dots \Psi_\alpha(\mathbf{r}_N). \quad (3)$$

¹Meccanica Statistica di un Condensato di Bose–Einstein Intrappolato.

²Presented to SIMAI98, IV National Conference of the Società Italiana di Matematica Applicata ed Industriale, Giardini Naxos, Messina, June 1998.

We observe that the mean energy per particle, which is given by

$$\epsilon = \langle \Phi_\alpha | \frac{\hat{H}}{N} | \Phi_\alpha \rangle = \frac{1}{N} \int d^3\mathbf{r}_1 \dots d^3\mathbf{r}_N \Phi_\alpha^*(\mathbf{r}_1, \dots, \mathbf{r}_N) \hat{H} \Phi_\alpha(\mathbf{r}_1, \dots, \mathbf{r}_N) , \quad (4)$$

can be written as

$$\epsilon[\Psi_\alpha] = \int d^3\mathbf{r} \frac{\hbar^2}{2m} |\nabla \Psi_\alpha(\mathbf{r})|^2 + V_0(\mathbf{r}) |\Psi_\alpha(\mathbf{r})|^2 + \frac{(N-1)}{2} \int d^3\mathbf{r} d^3\mathbf{r}' |\Psi_\alpha(\mathbf{r})|^2 V(\mathbf{r}, \mathbf{r}') |\Psi_\alpha(\mathbf{r}')|^2 , \quad (5)$$

where ϵ is a functional of the wave function Ψ_α . By considering a weakly interacting gas, such that $V(\mathbf{r}, \mathbf{r}') = B\delta^3(\mathbf{r} - \mathbf{r}')$, where $B = 4\pi\hbar^2 a_s/m$ is the scattering amplitude, the mean energy per particle results

$$\epsilon[\Psi] = \int d^3\mathbf{r} \frac{\hbar^2}{2m} |\nabla \Psi(\mathbf{r})|^2 + V_0(\mathbf{r}) |\Psi(\mathbf{r})|^2 + \frac{BN}{2} |\Psi(\mathbf{r})|^4 , \quad (6)$$

where we have omitted the quantum number α and we have written N instead of $N-1$ because we consider a large number of particles. This is the so-called Gross-Pitaevskii functional [5]. We observe that the scattering length a_s is supposed to be positive for ^{87}Rb and ^{23}Na , but negative for ^7Li . It means that for ^{87}Rb and ^{23}Na the interatomic interaction is repulsive while for ^7Li the atom-atom interaction is effectively attractive [1].

To find the minimum of the energy functional $\epsilon[\Psi]$ we impose that the first order variation of the energy functional is zero

$$\delta_\Psi \epsilon[\Psi] = 0 \quad (7)$$

with the constraint

$$\int d^3\mathbf{r} |\Psi(\mathbf{r})|^2 = 1 . \quad (8)$$

In this way we get the following Euler-Lagrange equation

$$\left[-\frac{\hbar^2}{2m} \nabla^2 + V_0(\mathbf{r}) + BN |\Psi(\mathbf{r})|^2 \right] \Psi(\mathbf{r}) = \mu \Psi(\mathbf{r}) , \quad (9)$$

where μ (Lagrange multiplier) is the chemical potential. This equation (Gross-Pitaevskii equation) has the form of a nonlinear stationary Schrödinger equation [5].

We analyze the Gross-Pitaevskii functional by using a variational technique. The external potential of the trap is well approximated by an harmonic oscillator $V_0(\mathbf{r}) = (m\omega^2/2)r^2$, where the frequency ω is proportional to the amplitude of the trapping magnetic field. We choose a Gaussian trial wave function for the BEC with a free parameter σ , which is the standard deviation of the Gaussian, i.e. the mean radius of the condensate

$$\Psi(\mathbf{r}) = \frac{1}{\pi^{3/4} \sigma^{3/2}} \exp\left(-\frac{r^2}{2\sigma^2}\right) . \quad (10)$$

The resulting energy function is

$$\epsilon(\sigma) = \frac{3}{2} \left(\frac{\hbar^2}{2m} \right) \frac{1}{\sigma^2} + \frac{3}{2} \left(\frac{m\omega^2}{2} \right) \sigma^2 + \frac{BN}{2} \frac{1}{(2\pi)^{3/2} \sigma^3} . \quad (11)$$

We study when the first derivative of $\epsilon(\sigma)$ is zero and we obtain the formula

$$N = \frac{2(2\pi)^{3/2}}{B} \left[\left(\frac{m\omega^2}{2} \right) \sigma^5 - \left(\frac{\hbar^2}{2m} \right) \sigma \right]. \quad (12)$$

For $B > 0$ there is only one solution, which is a minimum of the energy functional (stable solution), instead for $B < 0$ there are two solutions: one is a maximum (unstable solution) and the other a minimum (metastable solution) of the energy functional. In fact, for $B < 0$ the absolute minimum of the energy function is $\epsilon = \infty$ at $\sigma = 0$. It follows that for $B > 0$, when $N = 0$ we have $\sigma = \sqrt{\hbar/(m\omega)}$ and $\sigma \rightarrow \infty$ for $N \rightarrow \infty$. For $B < 0$ the mean radius σ decreases by increasing the number of bosons N to a minimum radius σ^{min} , with a maximum number N^{max} of bosons. For greater values of N the condensate becomes unstable. It is not difficult to obtain this critical number of bosons. We put $\tilde{B} = -B$ and get

$$0 = \frac{dN}{d\sigma} = \frac{2(2\pi)^{3/2}}{\tilde{B}} \left[\left(\frac{\hbar^2}{2m} \right) - 5 \left(\frac{m\omega^2}{2} \right) \sigma^4 \right], \quad (13)$$

from which we have the minimum radius $\sigma^{min} = 5^{-1/4} \sqrt{\hbar/(m\omega)}$ and the maximum number of bosons

$$N^{max} = \frac{8}{5^{5/4}} \frac{(2\pi)^{3/2}}{\tilde{B}} \frac{\hbar^2}{2m} \sqrt{\frac{\hbar}{m\omega}}. \quad (14)$$

Thus we obtain an analytical formula of the maximum number N^{max} of bosons for which the condensate, with attractive interaction, is metastable.

We observe that in general we can not neglect the non-zero range dependence of the interaction potential. Let us now consider a more realistic non-local interaction between the particles

$$V(\mathbf{r}, \mathbf{r}') = B\delta^3(\mathbf{r} - \mathbf{r}') - A \frac{\exp(-\Gamma|\mathbf{r} - \mathbf{r}'|)}{|\mathbf{r} - \mathbf{r}'|}, \quad (15)$$

where the constants A , B and Γ are chosen to reproduce phenomenologically the experimental data [4]. By using again the trial Gaussian wave function and studying when the first derivative of the energy function $\epsilon(\sigma)$ is zero we get the following formula

$$N = \frac{\left(\frac{m\omega^2}{2} \right) \sigma^4 - \left(\frac{\hbar^2}{2m} \right)}{\frac{B}{2(2\pi)^{3/2}} \sigma^{-1} + \frac{1}{3} \frac{A\Gamma^2}{\sqrt{2\pi}} \sigma^3 - \frac{A}{6} \sqrt{\frac{2}{\pi}} \sigma - \frac{A\Gamma^3}{6} \sigma^4 \exp\left(\frac{\sigma^2 \Gamma^2}{2}\right) \text{erfc}(\sigma \Gamma \sqrt{2})}, \quad (16)$$

where $\text{erfc}(x) = 1 - \text{erf}(x)$ is the complementary error function. For $B > 0$ there is only one solution, which is a minimum (stable solution), instead for $B < 0$ there are three solutions: one is a maximum (unstable solution) and the other two are minima (metastable solutions). Thus, we find that with a non-local interaction for $B < 0$ there is a new metastable branch at high density.

To summarize, we study the stability of a gas of weakly interacting bosons in a harmonic trap with local and non-local interaction. To minimize the energy functional, we choose a Gaussian trial wave function, where its standard deviation represents the mean radius of the condensate. In the case of repulsive interaction the mean radius of the condensate grows by increasing the number of bosons. In the case of attractive interaction the mean radius decreases by increasing the number of bosons until it becomes unstable.

Our variational technique can be applied also to systems with different shapes of the trapping potential. In the future it will be interesting to investigate the time evolution of the condensate with exotic initial conditions.

This work has been supported by INFM under the Research Advanced Project (PRA) on "Bose–Einstein Condensation". The author is grateful to A. Parola and L. Reatto for stimulating discussions and thanks V.R. Manfredi and G. Maino for their interest on the subject of this contribution.

- [1] M.H. Anderson, et al., *Science* **269**, 189 (1995); K.B. Davis, et al., *Phys. Rev. Lett.* **75**, 3969 (1995); C.C. Bradley, et al., *Phys. Rev. Lett.* **75**, 1687 (1995).
- [2] F. Dalfovo and S. Stringari, *Phys. Rev. A* **53**, 2477 (1996); C.C. Bradley, C.A. Sackett, and R.G. Hulet, *Phys. Rev. Lett.* **78**, 985 (1997).
- [3] A.L. Fetter and J.D. Walecka, *Quantum Theory of Many-Particle Systems* (McGraw–Hill, New York, 1971);
- [4] A. Parola, L. Salasnich and L. Reatto, to appear in *Phys. Rev. A*.
- [5] E.P. Gross, *Nuovo Cimento* **20**, 454 (1961); L.P. Pitaevskii, *Sov. Phys. JETP* **13**, 451 (1961).