DARBOUX-TYPE TRANSFORMATIONS AND HYPERELLIPTIC CURVES

FRITZ GESZTESY AND HELGE HOLDEN

ABSTRACT. We systematically study Darboux-type transformations for the KdV and AKNS hierarchies and provide a complete account of their effects on hyperelliptic curves associated with algebro-geometric solutions of these hierarchies.

1. INTRODUCTION

Since the early days of completely integrable evolution equations in the late sixties, Darboux-type transformations (cf., e.g., [42] and the references therein) played an important role and turned out to be an integral part of (auto-)Bäcklund transformations between soliton equations. The canonical example in this context is Miura's transformation [49] between the Korteweg-de Vries (KdV) hierarchy and modified Korteweg-de Vries (mKdV) hierarchy. As described in (2.35)–(2.42), Miura's transformation, disregarding the time variable (i.e., considering the stationary case), is effected by the classical Crum-Darboux transformation, which in turn is based on factorizing the second-order Lax (one-dimensional Schrödinger) operator

$$L = -\frac{d^2}{dx^2} + V \tag{1.1}$$

for the KdV hierarchy into a product of first-order differential expressions plus a shift,

$$L = AA^{+} + z_{0}, \quad A = \frac{d}{dx} + \phi, \quad A^{+} = -\frac{d}{dx} + \phi, \quad V = \phi^{2} + \phi_{x} + z_{0}.$$
(1.2)

Assuming V to satisfy one of the stationary KdV equations and reversing the order of the two factors A and A^+ produces a new Lax operator \widetilde{L} ,

$$\widetilde{L} = A^+ A + z_0 = -\frac{d^2}{dx^2} + \widetilde{V}, \quad \widetilde{V} = \phi^2 - \phi_x + z_0,$$
 (1.3)

whose potential \widetilde{V} is a new solution of one of the equations in the stationary KdV hierarchy. In short, the transformation

$$V \mapsto V$$
 (1.4)

¹⁹⁹¹ Mathematics Subject Classification. Primary 35Q53, 35Q55, 58F07; Secondary 35Q51, 35Q58.

 $Key\ words\ and\ phrases.$ Darboux transformations, hyperelliptic curves, KdV hierarchy, AKNS hierarchy.

Supported in part by the Research Council of Norway under grant 107510/410 and the University of Missouri Research Board grant RB-97-086.

represents a Darboux transformation, or equivalently, an auto-Bäcklund transformation of the stationary KdV hierarchy. Incidentally, $\pm \phi$ in (1.2), (1.3) represent solutions of one of the stationary equations of the mKdV hierarchy and hence

$$V \mapsto \phi \mapsto -\phi \mapsto V \tag{1.5}$$

represents a Bäcklund transformation from the (stationary) KdV to the mKdV hierarchy $(V \mapsto \phi)$ as well as auto-Bäcklund transformations for the KdV $(V \mapsto \tilde{V})$ and mKdV hierarchy $(\phi \mapsto -\phi)$. However, while ϕ and $-\phi$ satisfy the identical equation(s) within the stationary mKdV hierarchy, and hence

$$D = \begin{pmatrix} 0 & A^+ \\ A & 0 \end{pmatrix} \mapsto \widetilde{D} = \begin{pmatrix} 0 & -A \\ -A^+ & 0 \end{pmatrix}$$
(1.6)

represents an isospectral deformation of D, V and \tilde{V} in general do not satisfy the same stationary equation of the KdV hierarchy, that is,

$$L \mapsto \widetilde{L},$$
 (1.7)

in general, is not an isospectral deformation of L. More precisely, each solution V of (one of) the *n*th stationary KdV equations is associated with a (possibly singular) hyperelliptic curve \mathcal{K}_n of the type

$$\mathcal{K}_n: y^2 = \prod_{m=0}^{2n} (z - E_m), \quad \{E_m\}_{m=0,\dots,2n} \subset \mathbb{C}.$$
 (1.8)

Similarly, \widetilde{V} corresponds to a curve $\widetilde{\mathcal{K}}_{\tilde{n}}$ of the type

$$\widetilde{\mathcal{K}}_{\tilde{n}}: y^2 = \prod_{m=0}^{2n} (z - \widetilde{E}_m), \quad \{\widetilde{E}_m\}_{m=0,\dots,2\tilde{n}} \subset \mathbb{C}$$
(1.9)

and hence V and \widetilde{V} (resp. L and \widetilde{L}) are isospectral if and only if $\mathcal{K}_n = \widetilde{\mathcal{K}}_{\tilde{n}}$ (i.e., $\{E_m\}_{m=0,\ldots,2n} = \{\widetilde{E}_m\}_{m=0,\ldots,2\tilde{n}}, n = \tilde{n}\}.$

The principal aim of the paper is to re-examine the relation between \mathcal{K}_n and $\mathcal{\tilde{K}}_{\tilde{n}}$, depending on various choices of the function ϕ in (1.2), and to provide a complete, yet elementary solution of this problem. Historically, the first attempts to link \mathcal{K}_n and $\mathcal{\tilde{K}}_{\tilde{n}}$ were made by Drach [14], [15], [16]. The first solution of this problem was obtained by Ehlers and Knörrer in 1982 on the basis of purely algebro-geometric techniques. An elementary but quite elaborate approach was recently developed by Ohmiya [52] (based on two additional papers [51], [53]). Our proof of Theorem 2.3 appears to be the only elementary and short one at this point.

It seems worthwhile pointing out that the curves (1.8) and (1.9) may of course be singular, that is, some (or even all) of the E_m 's may coincide. In fact, the class of rational KdV solutions as discussed by Adler and Moser [2] (see also [50], [53]) arises in precisely this manner (with all $E_m = 0$). Similarly, the class of soliton solutions and more generally, solitons relative to an algebro-geometric background potential, as described, for instance, in [11], [12], [17], [33], [35], [36], [43], [47], [48], App. A, [55] results in *n* pairs of coinciding E_m 's, that is, $\{E_m\}_{m=0,...,2n} =$ $\{E_0, E_1 = E_2, \ldots, E_{2n-1} = E_{2n}\}$ for an appropriate enumeration of the E_m 's.

Finally, a quick description of the content of this paper. Section 2 is devoted to the KdV case and starts by summarizing the basic formalism for the KdV hierarchy and its algebro-geometric stationary solutions. Subsequently, we review Darboux transformations $V \mapsto \tilde{V}$ and finally derive the precise connection between \mathcal{K}_n and $\tilde{\mathcal{K}}_{\tilde{n}}$ in Theorem 2.3. Section 3 then presents the analogous results for the AKNS hierarchy. Besides of being an important hierarchy of soliton evolution equations, the AKNS hierarchy allows us to study hyperelliptic curves without a branch point at infinity as opposed to the KdV case, which necessarily leads to hyperelliptic curves branched at infinity. The corresponding Theorem 3.3, detailing the connection between \mathcal{K}_n and $\tilde{\mathcal{K}}_{\tilde{n}}$ in the AKNS case, to the best of our knowledge, appears to be without precedent.

2. The stationary KDV hierarchy

In this section we study Bäcklund transformations of the KdV hierarchy. Our principal aim is to prove a result by Ehlers and Knörrer [19] on hyperelliptic curves and Darboux transformations by entirely elementary means and at the same time offer a more detailed treatment (cf. Theorem 2.3).

We start by introducing a polynomial recursion formalism following Al'ber [3], [4] (see also [13], Ch. 12, [26], [27]) presented in detail in [32], [36] (see also [29], [30], [37]).

Suppose $V : \mathbb{C} \to \mathbb{C}_{\infty}$, with $\mathbb{C}_{\infty} = \mathbb{C} \cup \{\infty\}$, is meromorphic and consider the Schrödinger operator

$$L = -\frac{d^2}{dx^2} + V(x), \quad x \in \mathbb{C}.$$
(2.1)

Introducing $\{f_j\}_{j\in\mathbb{N}_0}$, with $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, recursively by

$$f_0 = 1, \quad f_{j,x} = -\frac{1}{4}f_{j-1,xxx} + Vf_{j-1,x} + \frac{1}{2}V_x f_{j-1}, \quad j \in \mathbb{N},$$
 (2.2)

one finds explicitly,

$$f_0 = 1,$$

$$f_1 = \frac{1}{2}V + c_1,$$

$$f_2 = -\frac{1}{8}V_{xx} + \frac{3}{8}V^2 + c_1\frac{1}{2}V + c_2, \quad \text{etc.},$$
(2.3)

where $\{c_\ell\}_{\ell\in\mathbb{N}}\subset\mathbb{C}$ denote integration constants. The f_j are well-known to be differential polynomials in V (see, e.g., [20]). Given V and f_j one defines differential expressions P_{2n+1} of order 2n + 1,

$$P_{2n+1} = \sum_{j=0}^{n} \left(f_{n-j}(x) \frac{d}{dx} - \frac{1}{2} f_{n-j,x}(x) \right) L^{j}, \quad n \in \mathbb{N}_{0},$$
(2.4)

and one verifies

$$[P_{2n+1}, L] = 2f_{n+1,x}, \quad n \in \mathbb{N}_0, \tag{2.5}$$

with $[\cdot, \cdot]$ the commutator. The stationary KdV hierarchy is then defined in terms of the stationary Lax relations

$$[P_{2n+1}, L] = 2f_{n+1,x} = 0, \quad n \in \mathbb{N}_0.$$
(2.6)

Explicitly, one finds

$$n = 0: \quad V_x = 0,$$

$$n = 1: \quad \frac{1}{4}V_{xxx} - \frac{3}{2}VV_x - c_1V_x = 0, \quad \text{etc.}$$
(2.7)

V(x) is called an *algebro-geometric* KdV *potential* if it satisfies one (and hence infinitely many) of the equations in the stationary KdV hierarchy (2.6).

Introducing the fundamental polynomial $F_n(z, x)$ of degree n in z,

$$F_n(z,x) = \sum_{j=0}^n f_{n-j}(x) z^j,$$
(2.8)

equations (2.2) and (2.6), that is, $f_{n+1,x}(x) = 0, x \in \mathbb{C}$, imply

$$F_{n,xxx} - 4(V-z)F_{n,x} - 2V_xF_n = 0.$$
(2.9)

Multiplying (2.9) by F_n and integrating yields

$$\frac{1}{2}F_{n,xx}(z,x)F_n(z,x) - \frac{1}{4}F_{n,x}(z,x)^2 - (V(x)-z)F_n(z,x)^2 = R_{2n+1}(z), \quad (2.10)$$

where the integration constant $R_{2n+1}(z)$ is a monic polynomial in z of degree 2n+1, and hence of the form

$$R_{2n+1}(z) = \prod_{m=0}^{2n} (z - E_m), \quad \{E_m\}_{m=0,\dots,2n} \subset \mathbb{C}.$$
 (2.11)

Introducing the algebraic eigenspace,

$$\ker(L-z) = \{\psi \colon \mathbb{C} \mapsto \mathbb{C}_{\infty} \text{ meromorphic } | (L-z)\psi = 0\}, \quad z \in \mathbb{C}$$
(2.12)

one verifies

$$P_{2n+1}\Big|_{\ker(L-z)} = \left(F_n(z,x)\frac{d}{dx} - \frac{1}{2}F_{n,x}(z,x)\right)\Big|_{\ker(L-z)}.$$
(2.13)

Moreover, a celebrated result by Burchnall–Chaundy [6], [7], [8] (see also [25], [54], [55], [57], [59]) then yields

$$P_{2n+1}^2 + R_{2n+1}(L) = 0. (2.14)$$

Equation (2.14) naturally leads to the hyperelliptic curve $\dot{\mathcal{K}}_n$ defined by

$$\dot{\mathcal{K}}_n: \mathcal{F}_n(z,y) = y^2 - R_{2n+1}(z) = 0, \quad R_{2n+1}(z) = \prod_{m=0}^{2n} (z - E_m).$$
 (2.15)

The one-point compactification of $\dot{\mathcal{K}}_n$ by joining P_{∞} , the point at infinity, is then denoted by \mathcal{K}_n . A general point $P \in \mathcal{K}_n \setminus \{P_{\infty}\}$ will be denoted by P = (z, y), where

$$\mathcal{F}_n(z,y) = 0. \tag{2.16}$$

Moreover, we define the involution * on \mathcal{K}_n by

$$*: \mathcal{K}_n \mapsto \mathcal{K}_n, \quad P = (z, y) \mapsto P^* = (z, -y), \quad P^*_{\infty} = P_{\infty}.$$
(2.17)

Introducing the polynomial H_{n+1} of degree n+1 in z,

$$H_{n+1}(z,x) = \frac{1}{2}F_{n,xx}(z,x) + (z - V(x))F_n(z,x), \qquad (2.18)$$

(2.10) implies

$$R_{2n+1}(z) + \frac{1}{4}F_{n,x}(z,x)^2 = F_n(z,x)H_{n+1}(z,x)$$
(2.19)

and we may define the following fundamental meromorphic function $\phi(P, x)$ on \mathcal{K}_n ,

$$\phi(P,x) = \frac{iy(P) + \frac{1}{2}F_{n,x}(z,x)}{F_n(z,x)}$$
(2.20a)

$$= \frac{-H_{n+1}(z,x)}{iy(P) - \frac{1}{2}F_{n,x}(z,x)}, \quad P = (z,y) \in \mathcal{K}_n, \ x \in \mathbb{C},$$
(2.20b)

where y(P) denotes the meromorphic function on \mathcal{K}_n obtained upon solving $y^2 =$ $R_{2n+1}(z)$ with P = (z, y).

Introducing the stationary Baker–Akhiezer function $\psi(P, x, x_0)$ on $\mathcal{K}_n \setminus \{P_\infty\}$ by

$$\psi(P, x, x_0) = \exp\left(\int_{x_0}^x dx' \,\phi(P, x')\right), \quad P = (z, y) \in \mathcal{K}_n \setminus \{P_\infty\}, \, x, x_0 \in \mathbb{C}, \quad (2.21)$$

choosing a smooth non-selfintersecting path from x_0 to x which avoids singularities of $\phi(P, x)$, one easily verifies from (2.9), (2.10), (2.13), (2.20), and (2.21) the following result.

Lemma 2.1. (see, e.g., [30], [32]) Suppose $f_{n+1,x}(x) = 0$ and let $P = (z, y) \in$ $\mathcal{K}_n \setminus \{P_\infty\}, x, x_0 \in \mathbb{C}$. Then $\phi(P, x)$ satisfies the Riccati-type equation

$$\phi_x(P,x) + \phi(P,x)^2 = V(x) - z, \qquad (2.22)$$

and

$$\phi(P, x)\phi(P^*, x) = H_{n+1}(z, x)/F_n(z, x), \qquad (2.23)$$

$$\phi(P,x) + \phi(P^*,x) = F_{n,x}(z,x)/F_n(z,x), \qquad (2.24)$$

$$\phi(P, x) - \phi(P^*, x) = 2iy(P)/F_n(z, x), \qquad (2.25)$$

$$\phi(P,x) = iz^{-n}y(P) + O(|z|^{-1/2}) \text{ as } P = (z,y) \to P_{\infty}.$$
 (2.26)

 $\psi(P, x, x_0)$ satisfies

$$(L - z(P))\psi(P, \cdot, x_0) = 0, \quad (P_{n+1} - iy(P))\psi(P, \cdot, x_0) = 0,$$
 (2.27)

and

$$\psi(P, x, x_0)\psi(P^*, x, x_0) = F_n(z, x)/F_n(z, x_0), \qquad (2.28)$$

$$\psi_x(P, x, x_0)\psi_x(P^*, x, x_0) = H_{n+1}(z, x)/F_n(z, x_0),$$

$$\psi(P, x, x_0)\psi_x(P^*, x, x_0) + \psi(P^*, x, x_0)\psi_x(P, x, x_0)$$
(2.29)

$$P, x, x_0)\psi_x(P^*, x, x_0) + \psi(P^*, x, x_0)\psi_x(P, x, x_0)$$

$$= F_{n,x}(z,x)/F_n(z,x_0), \qquad (2.30)$$

$$W(\psi(P, \cdot, x_0), \psi(P^*, \cdot, x_0)) = -2iy(P)/F_n(z, x_0),$$
(2.31)

$$\psi(P, x, x_0) = \exp\left(iz^{-n}y(P)(x - x_0)\right)(1 + O(|z|^{-1/2})) \text{ as } P = (z, y) \to P_{\infty}.$$
(2.32)

(Here W(f,g)(x) = f(x)g'(x) - f'(x)g(x) denotes the Wronskian of f and g.) In addition, we introduce the formal diagonal Green's function G(P, x, x) associated with the differential expression L by

$$G(P, x, x) = \frac{\psi(P, x, x_0)\psi(P^*, x, x_0)}{W(\psi(P, \cdot, x_0), \psi(P^*, \cdot, x_0))}$$
(2.33a)

$$=\frac{iF_n(z,x)}{2y(P)}, \quad P=(z,y)\in\mathcal{K}_n\setminus\{P_\infty\}, \ x\in\mathbb{C},$$
(2.33b)

using $\phi = \psi_x/\psi$, (2.25), and (2.28). Equations (2.10) and (2.33b) then yield the universal equation

$$-2G''(P, x, x)G(P, x, x) + G'(P, x, x)^{2} + 4(V(x) - z)G(P, x, x)^{2} = 1,$$

$$P = (z, y) \in \mathcal{K}_{n} \setminus \{P_{\infty}\}.$$
 (2.34)

These observations complete our review of the stationary KdV hierarchy. The timedependent KdV hierarchy could readily be defined at this point but since it is not essential for our purpose we resist the temptation to do so (see, e.g., [30], [32] for the time-dependent formalism).

Next we turn to the Darboux transformations in connection with $L = -d^2/dx^2 + V$. Define

$$\psi(P, x, x_0, \sigma) = \begin{cases} \frac{1}{2}(1+\sigma)\psi(P, x, x_0) + \frac{1}{2}(1-\sigma)\psi(P^*, x, x_0) & \text{for } \sigma \in \mathbb{C}, \\ \psi(P, x, x_0) - \psi(P^*, x, x_0) & \text{for } \sigma = \infty, \\ P \in \mathcal{K}_n \setminus \{P_\infty\}, \quad (2.35) \end{cases}$$

pick $Q_0 = (z_0, y_0) \in \mathcal{K}_n \setminus \{P_\infty\}$, and introduce the differential expressions

$$A_{\sigma}(Q_0) = \frac{d}{dx} + \phi(Q_0, x, \sigma), \quad A_{\sigma}^+(Q_0) = -\frac{d}{dx} + \phi(Q_0, x, \sigma), \quad \sigma \in \mathbb{C}_{\infty}, \quad (2.36)$$

where

$$\phi(P, x, \sigma) = \psi_x(P, x, x_0, \sigma) / \psi(P, x, x_0, \sigma), \quad P \in \mathcal{K}_n \setminus \{P_\infty\}, \, \sigma \in \mathbb{C}_\infty.$$
(2.37)

One verifies (cf. (2.22))

$$L = A_{\sigma}(Q_0)A_{\sigma}^+(Q_0) + z_0 = -\frac{d^2}{dx^2} + V, \qquad (2.38)$$

with

$$V(x) = \phi(Q_0, x, \sigma)^2 + \phi_x(Q_0, x, \sigma) + z_0, \qquad (2.39)$$

independent of the choice of $\sigma \in \mathbb{C}_{\infty}$. Interchanging the order of the differential expressions $A_{\sigma}(Q_0)$ and $A_{\sigma}^+(Q_0)$ in (2.38) then yields

$$\widetilde{L}_{\sigma}(Q_0) = A_{\sigma}^+(Q_0)A_{\sigma}(Q_0) + z_0 = -\frac{d^2}{dx^2} + \widetilde{V}_{\sigma}(x, Q_0), \qquad (2.40)$$

with

$$\widetilde{V}_{\sigma}(x,Q_0) = \phi(Q_0,x,\sigma)^2 - \phi_x(Q_0,x,\sigma) + z_0$$

= $V(x) - 2(\ln(\psi(Q_0,x,x_0,\sigma)))_{xx}, \quad \sigma \in \mathbb{C}_{\infty}.$ (2.41)

The transformation

$$V(x) \mapsto V_{\sigma}(x, Q_0), \quad Q_0 \in \mathcal{K}_n \setminus \{P_\infty\}, \, \sigma \in \mathbb{C}_\infty$$
 (2.42)

is usually called the Darboux transformation (also Crum-Darboux transformation or single commutation method) and goes back at least to Jacobi [40] and Darboux [10]. While we only aim at its properties from an algebraic point of view, its analytic properties in connection with spectral deformations (isospectral and non-isospectral ones) have received enormous attention in the context of spectral theory (especially, regarding the insertion of eigenvalues into spectral gaps), inverse spectral theory, and Bäcklund tranformations for the (time-dependent) KdV hierarchy. A complete bibliography in this context being impossible, we just refer to [1], [9], [11], [12], [18], Ch. 4, [19], [21], [23], [24], [28], [33], [34], [35], [36], [44], [45], [46], [56], [58] and the extensive literature therein. From a historical point of view it is very interesting to note that Drach [14], [15], [16] in his 1919 studies of Darboux transformations (being a student of Darboux) not only introduced a set of nonlinear differential equations for V, which today can be identified with the stationary KdV hierarchy, but also studied the effect of Darboux transformations on the underlying hyperelliptic curve. As a consequence, he seems to have been the

 $\mathbf{6}$

first to explicitly establish the connection between integrable systems and spectral theory. For modern treatments of this connection see, for instance, [5], Ch. 3 and [30].

Before analyzing the Darboux tranformation (2.42) in some detail, we briefly return to the formal diagonal Green's function G(P, x, x) in (2.33). In view of the possibility of linear combinations as displayed in the definition of $\psi(P, x, x_0, \sigma)$ in (2.35), and our lack of natural boundary conditions (resp. L^2 -conditions) in connection with the (possibly singular) differential expression L, our definition of G(P, x, x) in (2.33) appears to be highly arbitrary. However, the following considerations will show that our choice in (2.33) is the unique one that yields a bounded Green's function as $P \to P_{\infty}$ (for $x \in \mathbb{C} \setminus \{x_0\}$ fixed). In fact,

$$G(P, x, x) = \frac{iz^n}{2y(P)} + O(|z|^{-1}) \text{ as } P = (z, y) \to P_{\infty}.$$
 (2.43)

Introducing

$$H(P, x, x, x_0, \sigma_1, \sigma_2) = \frac{\psi(P, x, x_0, \sigma_1)\psi(P, x, x_0, \sigma_2)}{W(\psi(P, \cdot, x_0, \sigma_1), \psi(P, \cdot, x_0, \sigma_2))}, \\ \sigma_1, \sigma_2 \in \mathbb{C}_{\infty}, \, \sigma_1 \neq \sigma_2, \quad (2.44)$$

one infers from Lemma 2.1, (2.32), and (2.35)

$$H(P, x, x, x_0, \sigma_1, \sigma_2) = \frac{i(1+\sigma_1)(1+\sigma_2)z^n}{4(\sigma_1-\sigma_2)y(P)} \exp(2iz^{-n}y(P)(x-x_0))(1+O(|z|^{-1/2})) + \frac{i(1-\sigma_1)(1-\sigma_2)z^n}{4(\sigma_1-\sigma_2)y(P)} \exp(-2iz^{-n}y(P)(x-x_0))(1+O(|z|^{-1/2})) + \frac{i(1-\sigma_1\sigma_2)z^n}{2(\sigma_1-\sigma_2)y(P)} + O(|z|^{-1/2}) \text{ as } P = (z,y) \to P_{\infty}$$

for $\sigma_1, \sigma_2 \in \mathbb{C}$, $\sigma_1 \neq \sigma_2$, and similarly if $\sigma_1 = \infty$, $\sigma_2 \in \mathbb{C}$, or $\sigma_1 \in \mathbb{C}$, $\sigma_2 = \infty$. Thus, $H(P, x, x, x_0, \sigma_1, \sigma_2)$ is uniformly bounded for P in a neighborhood of P_{∞} and $x \in \mathbb{C} \setminus \{x_0\}$ fixed, if and only if

$$\sigma_1 = -\sigma_2 \in \{-1, 1\}. \tag{2.46}$$

Since $\psi(P, x, x_0, -\sigma) = \psi(P^*, x, x_0, \sigma)$, this shows that our choice for G(P, x, x) in (2.32) yields the unique diagonal Green's function of L which is bounded for P in a neighborhood of P_{∞} for fixed $x \in \mathbb{C} \setminus \{x_0\}$. In addition, $H(P, x, x, x_0, \sigma_1, \sigma_2)$ is independent of x_0 if and only if (2.46) holds.

Next, assuming that $\psi \in \ker(L-z)$,

$$L\psi(z) = z\psi(z), \qquad (2.47)$$

one infers $A^+_{\sigma}(Q_0)\psi(z) \in \ker(\widetilde{L}_{\sigma}(Q_0)-z),$

$$\widetilde{L}_{\sigma}(Q_0)(A_{\sigma}^+(Q_0)\psi(z)) = zA_{\sigma}^+(Q_0)\psi(z), \qquad (2.48)$$

and

$$W(A_{\sigma}^{+}(Q_{0})\psi_{1}(z), A_{\sigma}^{+}(Q_{0})\psi_{2}(z)) = (z - z_{0})W(\psi_{1}(z), \psi_{2}(z)),$$

$$\psi_{1}(z), \psi_{2}(z) \in \ker(L - z). \quad (2.49)$$

Since

GESZTESY AND HOLDEN

$$(A_{\sigma}^{+}(Q_{0})\psi(P,\cdot,x_{0}))(x) = (\phi(Q_{0},x,\sigma) - \phi(P,x))\psi(P,x,x_{0}),$$

$$P \in \mathcal{K}_{n} \setminus \{Q_{0},P_{\infty}\}, \quad (2.50)$$

we define

$$\psi_{\sigma}(P, x, x_0, Q_0) = (A_{\sigma}^+(Q_0)\psi(P, \cdot, x_0))(x)$$

= $(\phi(Q_0, x, \sigma) - \phi(P, x))\psi(P, x, x_0),$ (2.51)
 $P \in \mathcal{K}_n \setminus \{Q_0, P_\infty\}, \sigma \in \mathbb{C}_{\infty}.$

Then

$$(\widetilde{L}_{\sigma}(Q_0)\widetilde{\psi}_{\sigma}(P,\cdot,x_0,Q_0))(x) = z\widetilde{\psi}_{\sigma}(P,x,x_0,Q_0),$$
$$P = (z,y) \in \mathcal{K}_n \setminus \{Q_0, P_\infty\}, \quad (2.52)$$

and we define in analogy to (2.33a) the diagonal Green's function $\widetilde{G}_{\sigma}(P, x, x, Q_0)$ of $\widetilde{L}_{\sigma}(Q_0)$ by

$$\widetilde{G}_{\sigma}(P, x, x, Q_0) = \frac{\widetilde{\psi}_{\sigma}(P, x, x_0, Q_0) \widetilde{\psi}_{\sigma}(P^*, x, x_0, Q_0)}{W(\widetilde{\psi}_{\sigma}(P, \cdot, x_0, Q_0), \widetilde{\psi}_{\sigma}(P^*, \cdot, x_0, Q_0))},$$

$$P = (z, y) \in \mathcal{K}_n \setminus \{Q_0, P_\infty\}.$$
(2.53)

Lemma 2.2. Assume $f_{n+1,x}(x) = 0$ and let $Q_0 = (z_0, y_0) \in \mathcal{K}_n \setminus \{P_\infty\}, P = (z, y) \in \mathcal{K}_n \setminus \{Q_0, P_\infty\}, \sigma \in \mathbb{C}_\infty$. Then the diagonal Green's function $\widetilde{G}_{\sigma}(P, x, x, Q_0)$ in (2.34) explicitly reads

$$\widetilde{G}_{\sigma}(P, x, x, Q_0) = \frac{H_{n+1}(z, x) + \phi(Q_0, x, \sigma)^2 F_n(z, x) - \phi(Q_0, x, \sigma) F_{n,x}(z, x)}{-2i(z - z_0)y(P)}$$
(2.54a)

$$=\frac{(\phi(P,x)-\phi(Q_0,x,\sigma))(\phi(P^*,x)-\phi(Q_0,x,\sigma))F_n(z,x)}{-2i(z-z_0)y(P)} \quad (2.54b)$$

$$=\frac{i\widetilde{F}_{\sigma,\tilde{n}}(z,x)}{2\tilde{y}(P)},$$
(2.54c)

where $\tilde{y}(P)$ denotes the meromorphic solution on $\widetilde{\mathcal{K}}_{\sigma,\tilde{n}}(Q_0)$ obtained upon solving $y^2 = \widetilde{R}_{\sigma,2\tilde{n}+1}(z)$, P = (z,y) for some polynomial $\widetilde{R}_{\sigma,2\tilde{n}+1}(z)$ of degree $2\tilde{n} + 1 \in \mathbb{N}_0$ (cf. (2.15) and (2.58)) and $\widetilde{F}_{\sigma,\tilde{n}}(z,x)$ denotes a polynomial with respect to z of degree \tilde{n} , with $0 \leq \tilde{n} \leq n + 1$. In particular, the Darboux transformation (2.42), $V(x) \mapsto \widetilde{V}_{\sigma}(x,Q_0)$ maps the class of algebro-geometric KdV potentials into itself.

Proof. (2.54a) and (2.54b) follow upon use of $\phi(P, x) = \psi_x(P, x, x_0)/\psi(P, x, x_0)$, (2.23), (2.24), (2.28)–(2.31), and (2.49). Since the numerator in (2.54a) is a polynomial in z, and

$$\widetilde{G}_{\sigma}(P, x, x, Q_0) = \frac{iz^n}{2y(P)} + O(|z|^{-1}) \text{ as } P = (z, y) \to P_{\infty}$$
 (2.55)

again by (2.54a), one concludes (2.54c) and $0 \leq \tilde{n} \leq n+1$. By inspection, $\tilde{F}_{\sigma,\tilde{n}}(z,x)$ satisfies equation (2.10) with V(x) replaced by $\tilde{V}_{\sigma}(x,Q_0)$, n by \tilde{n} , and $R_{2n+1}(z)$ by $\tilde{R}_{\sigma,2\tilde{n}+1}(z)$. As a consequence, V(x) being an algebro-geometric KdV potential implies that $\tilde{V}_{\sigma}(x,Q_0)$ is one as well.

The following theorem, the principal result of this section, will clarify the dependence of $\tilde{n} = \tilde{n}(n, Q_0, \sigma)$ on its variables. This result was originally derived using an entirely different algebro-geometric approach by Ehlers and Knörrer [19].

Theorem 2.3. Suppose $f_{n+1,x}(x) = 0$ and let $Q_0 = (z_0, y_0) \in \mathcal{K}_n \setminus \{P_\infty\}, n \in \mathbb{N}_0, \sigma \in \mathbb{C}_\infty$, and $\tilde{n} = \tilde{n}(n, Q_0, \sigma)$ as in (2.54c). Then

$$\tilde{n}(n,Q_{0},\sigma) = \begin{cases} n+1 & \text{for } \sigma \in \mathbb{C}_{\infty} \setminus \{-1,1\} \text{ and } y_{0} \neq 0, \\ n+1 & \text{for } \sigma = \infty \text{ and } y_{0} = 0, \\ n & \text{for } \sigma \in \{-1,1\} \text{ and } y_{0} \neq 0, \\ n & \text{for } \sigma \in \mathbb{C}, y_{0} = 0, \text{ and } R_{2n+1,z}(z_{0}) \neq 0, \\ n-1 & \text{for } \sigma \in \mathbb{C}, y_{0} = 0, \text{ and } R_{2n+1,z}(z_{0}) = 0, n \in \mathbb{N}, \end{cases}$$

$$(2.56)$$

and hence the hyperelliptic curve $\tilde{\mathcal{K}}_{\sigma,\tilde{n}}(Q_0)$ associated with $\tilde{V}_{\sigma}(x,Q_0)$ is of the type

$$\widetilde{\mathcal{K}}_{\sigma,\tilde{n}}(Q_0) \colon \widetilde{\mathcal{F}}_{\sigma,\tilde{n}}(z,y,Q_0) = y^2 - \widetilde{R}_{\sigma,2\tilde{n}+1}(z,Q_0) = 0, \qquad (2.57)$$

with

$$\widetilde{R}_{\sigma,2\widetilde{n}+1}(z,Q_0) = \begin{cases}
(z-z_0)^2 R_{2n+1}(z) & \text{for } \sigma \in \mathbb{C}_{\infty} \setminus \{-1,1\} \text{ and } y_0 \neq 0, \\
(z-z_0)^2 R_{2n+1}(z) & \text{for } \sigma = \infty \text{ and } y_0 = 0, \\
R_{2n+1}(z) & \text{for } \sigma \in \{-1,1\} \text{ and } y_0 \neq 0, \\
R_{2n+1}(z) & \text{for } \sigma \in \mathbb{C}, y_0 = 0, \text{ and } R_{2n+1,z}(z_0) \neq 0, \\
(z-z_0)^{-2} R_{2n+1}(z) & \text{for } \sigma \in \mathbb{C}, y_0 = 0, \text{ and } R_{2n+1,z}(z_0) = 0, n \in \mathbb{N}.
\end{cases}$$
(2.58)

Here

$$R_{2n+1}(z) = \prod_{m=0}^{2n} (z - E_m).$$
(2.59)

Proof. Our starting point will be (2.54b) and a careful case distinction taking into account whether or not Q_0 is a branch point, and distinguishing the cases $\sigma \in \mathbb{C} \setminus \{-1, 1\}, \sigma \in \{-1, 1\}, \text{ and } \sigma = \infty$.

Case (i). $\sigma \in \mathbb{C}_{\infty} \setminus \{-1, 1\}$ and $y_0 \neq 0$: One computes from (2.35) and (2.37),

$$\phi(Q_0, x, \sigma) = \begin{cases} \frac{(1+\sigma)\psi_x(Q_0, x, x_0) + (1-\sigma)\psi_x(Q_0^*, x, x_0)}{(1+\sigma)\psi(Q_0, x, x_0) + (1-\sigma)\psi(Q_0^*, x, x_0)} & \text{for } \sigma \in \mathbb{C} \setminus \{-1, 1\}, \\ \frac{\psi_x(Q_0, x, x_0) - \psi_x(Q_0^*, x, x_0)}{\psi(Q_0, x, x_0) - \psi(Q_0^*, x, x_0)} & \text{for } \sigma = \infty, \end{cases}$$
(2.60)

and upon comparison with $\phi(Q_0, x) \neq \phi(Q_0^*, x)$,

$$\phi(Q_0, x) = \frac{\psi_x(Q_0, x, x_0)}{\psi(Q_0, x, x_0)}, \quad \phi(Q_0^*, x) = \frac{\psi_x(Q_0^*, x, x_0)}{\psi(Q_0^*, x, x_0)}, \tag{2.61}$$

one concludes that no cancellations can occur in (2.54b), proving $\tilde{n}(n, Q_0, \sigma) = n+1$ and the first statement in (2.58).

Case (ii). $\sigma = \infty$ and $y_0 = 0$: Combining (2.20a), (2.21), (2.25), and (2.35) one computes

$$\begin{split} \phi(Q_0, x, \infty) &= \lim_{P \to Q_0} \phi(P, x, \infty) \\ &= \lim_{P \to Q_0} \left(\frac{\phi(P, x) \exp\left(\int_{x_0}^x dx' \, \phi(P, x')\right) - \phi(P^*, x) \exp\left(\int_{x_0}^x dx' \, \phi(P^*, x')\right)}{\exp\left(\int_{x_0}^x dx' \, \phi(P, x')\right) - \exp\left(\int_{x_0}^x dx' \, \phi(P^*, x')\right)} \right) \end{split}$$

$$= \phi(Q_0, x)$$

$$+ \lim_{P \to Q_0} \left(\frac{\phi(P, x) - \phi(P^*, x)}{\exp\left(\int_{x_0}^x dx' \, \phi(P, x')\right) - \exp\left(\int_{x_0}^x dx' \, \phi(P^*, x')\right)} \times \exp\left(\int_{x_0}^x dx' \, \phi(P^*, x')\right) \right)$$

$$= \phi(Q_0, x) + \exp\left(\int_{x_0}^x dx' \, \phi(Q_0, x')\right) \times \lim_{P \to Q_0} \left(\frac{2iy(P)/F_n(z, x)}{\exp\left(iy(P)\int_{x_0}^x \frac{dx'}{F_n(z, x')}\right) - \exp\left(-iy(P)\int_{x_0}^x \frac{dx'}{F_n(z, x')}\right)} \times \exp\left(-\frac{1}{2}\int_{x_0}^x dx' \frac{F_{n,x}(z, x')}{F_n(z, x')}\right) \right)$$

$$= \phi(Q_0, x)$$

$$= \phi(Q_0, x) + \psi(Q_0, x, x_0) \frac{1}{F_n(z_0, x)\psi(Q_0, x, x_0)} \lim_{P \to Q_0} \left(\frac{2iy(P)}{2iy(P) \int_{x_0}^x \frac{dx'}{F_n(z, x')} + O(y(P)^2)} \right)$$

$$= \phi(Q_0, x) + \left(F_n(z_0, x) \int_{x_0}^x \frac{dx'}{F_n(z, x')} \right)^{-1}, \quad x \in \mathbb{C} \setminus \{x_0\},$$

$$(2.62)$$

using $\lim_{P \to Q_0} y(P) = y(Q_0) = y_0 = 0$. From

$$\phi(Q_0, x) = \frac{1}{2} \frac{F_{n,x}(z_0, x)}{F_n(z_0, x)}$$
(2.63)

one concludes again that no cancellations can occur in (2.54b). Thus $\tilde{n}(n, Q_0, \infty) = n + 1$ and the second statement in (2.58) holds.

The remainder of the proof requires a more refined argument, the basis of which will be derived next. First, replacing V(x) by $V(x) - z_0$, we may assume without loss of generality that $z_0 = 0$ in the following. Moreover, from this point on we exclude the trivial case V(x) = 0, $x \in \mathbb{C}$ (the case $V(x) = E_0$ will be studied in Example 2.7 below). Writing

$$y(z)^{2} = R_{2n+1}(z) \underset{z \to 0}{=} y_{0}^{2} + \tilde{y}_{1}z + \tilde{y}_{2}z^{2} + O(z^{3}), \qquad (2.64)$$

a comparison of the powers z^0 and z^1 in (2.10) yields

$$2f_{n,xx}f_n = f_{n,x}^2 + 4Vf_n^2 + 4y_0^2 \tag{2.65}$$

and

$$f_{n-1,xx}f_n + f_{n,xx}f_{n-1} - f_{n,x}f_{n-1,x} - 4Vf_nf_{n-1} + 2Vf_n^2 - 2\tilde{y}_1 = 0.$$
(2.66)

Inserting (2.65) into (2.66), a little algebra proves the basic identity

$$f_n^2 (f_{n-1}/f_n)_{xx} + f_{n,x} f_n (f_{n-1}/f_n)_x + 2V f_n^2 + 4y_0^2 (f_{n-1}/f_n) - 2\tilde{y}_1 = 0.$$
(2.67)

Case (iii). $\sigma \in \{-1, 1\}$ and $y_0 \neq 0$: Then (2.35) yields

$$\phi(Q_0, x, 1) = \phi(Q_0, x), \quad \phi(Q_0, x, -1) = \phi(Q_0^*, x), \tag{2.68}$$

with $\phi(Q_0, x) \neq \phi(Q_0^*, x)$ since $y_0 \neq 0$. In this case there is a cancellation in (2.54b). For instance, choosing $\sigma = 1$ one computes from (2.8) and (2.20a),

$$\phi(P, x) - \phi(Q_0, x, 1) = \phi(P, x) - \phi(Q_0, x)$$

$$=_{P \to Q_0} \frac{i(y(P) - y_0)}{f_n(x)} - iy_0 \frac{f_{n-1}(x)}{f_n(x)^2} z + \frac{1}{2} \left(\frac{f_{n-1}(x)}{f_n(x)}\right)_x z + O(z^2)$$

$$=_{P \to Q_0} c_1(x) z + O(z^2)$$
(2.69)

since

$$y(P) - y_0 =_{P \to Q_0} y_1 z + O(z^2), \quad \tilde{y}_1 = 2y_0 y_1.$$
 (2.70)

It remains to show that $c_1(x)$ does not vanish identically in $x \in \mathbb{C}$. Arguing by contradiction we assume

$$0 = c_1(x) = \frac{iy_1}{f_n(x)} - \frac{iy_0}{f_n(x)} \frac{f_{n-1}(x)}{f_n(x)} + \frac{1}{2} \left(\frac{f_{n-1}(x)}{f_n(x)}\right)_x, \quad x \in \mathbb{C}.$$
 (2.71)

Differentiating (2.71) with respect to x and inserting the ensuing expression for $(f_{n-1}/f_n)_{xx}$ and the one for $(f_{n-1}/f_n)_x$ from (2.71) into (2.67) then results in the contradiction

$$0 = 2V(x)f_n(x)^2, \quad x \in \mathbb{C}.$$
 (2.72)

Moreover, since

$$\phi(P^*, x) - \phi(Q_0, x, x, 1) = \phi(P^*, x) - \phi(Q_0, x) \underset{P \to Q_0}{=} -\frac{2iy_0}{f_n(x)} + O(z), \quad (2.73)$$

one concludes that precisely one factor of z cancels in (2.54b). Hence $\tilde{n}(n, Q_0, 1) = n$ and the third relation in (2.58) holds. The case $\sigma = -1$ in treated analogously.

Case (iv). $\sigma \in \mathbb{C}$, $y_0 = 0$, and $R_{2n+1,z}(0) \neq 0$: Taking into account that $\phi(Q_0, x, \sigma) = \phi(Q_0, x)$ (using (2.37) and $Q_0 = Q_0^*$) is independent of $\sigma \in \mathbb{C}$, (2.8) and (2.20a) yield

$$\left(\phi(P,x) - \phi(Q_0,x)\right) \left(\phi(P^*,x) - \phi(Q_0,x)\right) = \frac{y_1^2 z}{f_n(x)^2} + O(z^2)$$
(2.74)

since

$$y(P) =_{P \to Q_0} y_1 z^{1/2} + O(z^{3/2}), \quad y_1 = \left(\prod_{E_m \neq 0} E_m\right)^{1/2}.$$
 (2.75)

Thus we infer again that precisely one factor of z cancels in (2.54b). Hence $\tilde{n}(n, Q_0, \sigma) = n$ and the fourth relation in (2.58) is proved.

Case (v). $\sigma \in \mathbb{C}$, $y_0 = \tilde{y}_1 = 0$, and $\tilde{y}_2 \neq 0$ (cf. (2.64)): One calculates as in (2.74),

$$\phi(P,x) - \phi(Q_0,x) \Big) \Big(\phi(P^*,x) - \phi(Q_0,x) \Big) = \sum_{P \to Q_0} \left(\frac{y_1^2}{f_n(x)^2} + \frac{1}{4} \left(\left(\frac{f_{n-1}(x)}{f_n(x)} \right)_x \right)^2 \right) z^2 + O(z^3) = \sum_{P \to Q_0} c_2(x) z^2 + O(z^3)$$
(2.76)

since

(

$$y(P) =_{P \to Q_0} y_1 z + O(z^2), \quad y_1 = \left(\prod_{E_m \neq 0} E_m\right)^{1/2}.$$
 (2.77)

Next we show that $c_2(x)$ does not vanish identically in $x \in \mathbb{C}$. Arguing again by contradiction we suppose that

$$0 = c_2(x) = \frac{y_1^2}{f_n(x)^2} + \frac{1}{4} \left(\left(\frac{f_{n-1}(x)}{f_n(x)} \right)_x \right)^2, \quad x \in \mathbb{C}.$$
 (2.78)

Thus

$$\left(\frac{f_{n-1}}{f_n}\right)_x = \frac{C}{f_n} \tag{2.79}$$

for some constant $C \in \mathbb{C}$. Insertion of (2.79) and its x-derivative into (2.67) then again yields the contradiction

$$0 = 2V(x)f_n(x)^2, \quad x \in \mathbb{C}.$$
 (2.80)

Hence $\tilde{n}(n, Q_0, \sigma) = n - 1$ and the last relation in (2.58) holds in this case. Case (vi). $\sigma \in \mathbb{C}$, $y_0 = \tilde{y}_1 = \tilde{y}_2 = 0$ (cf. (2.64)): As in (2.76) one obtains

$$\left(\phi(P,x) - \phi(Q_0,x)\right) \left(\phi(P^*,x) - \phi(Q_0,x)\right) = \frac{1}{P \to Q_0} \frac{1}{4} \left(\left(\frac{f_{n-1}(x)}{f_n(x)}\right)_x \right)^2 z^2 + O(z^3)$$
(2.81)

since

$$y(P) =_{P \to Q_0} O(z^{3/2}).$$
 (2.82)

The remainder of the proof of case (vi) is now a special case of case (v) (with $y_1 = C = 0$) and one concludes again that $\tilde{n}(n, Q_0, \sigma) = n - 1$.

We can summarize the previous theorem in the following table.

		$\sigma \in \mathbb{C} \backslash \{-1, 1\}$	$\sigma \in \{-1, 1\}$	$\sigma = \infty$
$y_0 \neq 0$		n+1	n	
$y_0 = 0$	$\tilde{y}_1 \neq 0$	n		n+1
	$\tilde{y}_1 = 0$	n-1		

Table 2.4. The table shows the value of the arithmetic genus \tilde{n} associated with the Darboux transformation. Here $R_{2n+1}(z) = y_0^2 + \tilde{y}_1(z-z_0) + O((z-z_0)^2)$ as $z \to z_0$.

These results show, in particular, that Darboux transformations do not change the local structure of the original curve $y^2 = R_{2n+1}(z)$, except, of course, near the point Q_0 .

Remark 2.5. Theorem 2.3 was first derived by purely algebro-geometric means by Ehlers and Knörrer [19] in 1982. An elementary but rather lengthy derivation of Theorem 2.3 (focusing on the case when $\tilde{n}(n,\sigma) = n-1$) was provided by Ohmiya [52] in 1997 (based on two other papers [51], [53]). The current proof seems to be the only elementary and short one available at this point. As mentioned in the paragraph following (2.42), Drach [14], [15], [16] appears to have been the first to study particular aspects (the case $\tilde{n} = n + 1$) of Theorem 2.3 around 1919. Moreover, it seems worthwhile to point out that the case $\sigma = \infty$ and $y_0 = 0$, which leads to $\tilde{n}(n, Q_0, \sigma) = n + 1$, necessarily constructs an algebro-geometric KdV potential $\tilde{V}_{\infty}(x, Q_0)$ singular at $x = x_0$ (cf. (2.62)). In fact, the class of rational algebro-geometric solutions constructed by Adler and Moser [2] arises exactly in this manner.

Remark 2.6. The results described in Theorem 2.3 are not confined to hyperelliptic curves \mathcal{K}_n of finite (arithmetic) genus n. In fact, upon shifting the emphasis from $F_n(z, x)$ to G(P, x, x), the results in Theorem 2.3 extend to certain classes of transcendental hyperelliptic curves of infinite (arithmetic) genus \mathcal{K}_∞ (including those associated with periodic potentials V). More details will be presented elsewhere.

We conclude with the following elementary illustration.

Example 2.7. The case n = 0.

$$y(P)^{2} = R_{1}(z) = z - E_{0},$$

$$F_{0}(z, x) = 1, \quad H_{1}(z, x) = z - E_{0}, \quad V(x) = E_{0},$$

$$\phi(P, x) = iy(P), \quad \psi(P, x, x_{0}) = \exp(iy(P)(x - x_{0})),$$

$$G(P, x, x) = \frac{i}{2y(P)},$$

$$\phi(P, x, \sigma) = \begin{cases} iy(P)\frac{(1+\sigma)\exp(iy(P)(x-x_{0})) - (1-\sigma)\exp(-iy(P)(x-x_{0}))}{(1+\sigma)\exp(iy(P)(x-x_{0})) + (1-\sigma)\exp(-iy(P)(x-x_{0}))}, & \sigma \in \mathbb{C}, \\ iy(P)\frac{\exp(iy(P)(x-x_{0})) - \exp(-iy(P)(x-x_{0}))}{\exp(iy(P)(x-x_{0})) + \exp(-iy(P)(x-x_{0}))}, & \sigma = \infty, \end{cases}$$

$$\phi((E_{0}, 0), x, \sigma) = \begin{cases} 0, & \sigma \in \mathbb{C}, \\ (x - x_{0})^{-1}, & \sigma = \infty. \end{cases}$$
(2.83)

More generally, the case $V(x) = E_0$ can be associated with any curve $y(P)^2 = R_{2n+1}(z)$ since $f_{n,x}(x) = 0$, $n \in \mathbb{N}_0$ in this special case.

3. The stationary AKNS hierarchy

This section is devoted to Darboux (gauge) transformations for the AKNS hierarchy. In particular, we derive the KdV analogs of Section 2 and hence determine the effect of gauge transformations on hyperelliptic AKNS curves in the spirit of our approach to the KdV result of Ehlers and Knörrer [19].

We start by introducing a polynomial recursion for the AKNS hierarchy following the derivation in [31]. Suppose $p, q: \mathbb{C} \to \mathbb{C}_{\infty}$ are meromorphic and introduce the Dirac-type differential expression

$$D = \begin{pmatrix} \frac{d}{dx} & -q\\ p & -\frac{d}{dx} \end{pmatrix}, \quad x \in \mathbb{C}.$$
(3.1)

Introducing $\{f_j(x)\}_{j\in\mathbb{N}_0}, \{g_j(x)\}_{j\in\mathbb{N}_0}$, and $\{h_j(x)\}_{j\in\mathbb{N}_0}$ recursively by

$$f_{0}(x) = -iq(x), \quad g_{0}(x) = 1, \quad h_{0}(x) = ip(x),$$

$$f_{j}(x) = \frac{i}{2}f_{j-1,x}(x) - iq(x)g_{j}(x), \quad j \in \mathbb{N},$$

$$g_{j,x}(x) = p(x)f_{j-1}(x) + q(x)h_{j-1}(x), \quad j \in \mathbb{N},$$

$$h_{j}(x) = -\frac{i}{2}h_{j-1,x}(x) + ip(x)g_{j}(x), \quad j \in \mathbb{N}.$$
(3.2)

Explicitly, one computes

$$f_0 = -iq, \quad f_1 = \frac{1}{2}q_x + c_1(-iq),$$

GESZTESY AND HOLDEN

$$f_{2} = \frac{i}{4}q_{xx} - \frac{i}{2}pq^{2} + c_{1}\frac{1}{2}q_{x} + c_{2}(-iq),$$

$$g_{0} = 1, \quad g_{1} = c_{1}, \quad g_{2} = \frac{1}{2}pq + c_{2},$$

$$g_{3} = -\frac{i}{4}(p_{x}q - pq_{x}) + c_{1}\frac{1}{2}pq + c_{3},$$

$$h_{0} = ip, \quad h_{1} = \frac{1}{2}p_{x} + c_{1}ip,$$

$$h_{2} = -\frac{i}{4}p_{xx} + \frac{i}{2}p^{2}q + c_{1}\frac{1}{2}p_{x} + c_{2}ip, \quad \text{etc.},$$
(3.3)

where $\{c_j\}_{j\in\mathbb{N}_0} \subset \mathbb{C}$ are integration constants. The coefficients f_j , g_j , h_j are wellknown to be differential polynomials in p and q (see, e.g., [30]). Given p, q and f_j , g_j , and h_j one defines the matrix-valued differential expression of order n + 1,

$$E_{n+1} = i \sum_{j=0}^{n+1} \begin{pmatrix} -g_{n+1-j} & f_{n-j} \\ -h_{n-j} & g_{n+1-j} \end{pmatrix} D^j, \quad n \in \mathbb{N}_0, \quad f_{-1} = h_{-1} = 0,$$
(3.4)

and verifies

$$[E_{n+1}, D] = \begin{pmatrix} 0 & -2if_{n+1} \\ 2ih_{n+1} & 0 \end{pmatrix}, \quad n \in \mathbb{N}_0.$$
(3.5)

The stationary AKNS hierarchy is then defined by the stationary Lax relations

 $[E_{n+1}, D] = 0, \quad n \in \mathbb{N}, \text{ that is, } f_{n+1} = h_{n+1} = 0, \quad n \in \mathbb{N}.$ (3.6)

Explicitly, one finds

$$n = 0: \quad \begin{pmatrix} -p_x + c_1(-2ip) \\ -q_x + c_1(2iq) \end{pmatrix} = 0,$$

$$n = 1: \quad \begin{pmatrix} \frac{i}{2}p_{xx} - ip^2q + c_1(-p_x) + c_2(-2ip) \\ -\frac{i}{2}q_{xx} + ipq^2 + c_1(-q_x) + c_2(2iq) \end{pmatrix} = 0, \text{ etc.}$$
(3.7)

(p(x), q(x)) are called *algebro-geometric* AKNS *potentials* if they satisfy one (and hence infinitely many) of the equations of the stationary AKNS hierarchy (3.6).

Introducing the polynomials $F_n(z, x)$, $G_n(z, x)$, and $H_n(z, x)$ with respect to z,

$$F_{n}(z,x) = \sum_{j=0}^{n} f_{n-j}(x)z^{j},$$

$$G_{n+1}(z,x) = \sum_{j=0}^{n+1} g_{n+1-j}(x)z^{j},$$

$$H_{n}(z,x) = \sum_{j=0}^{n} h_{n-j}(x)z^{j},$$
(3.8)

equations (3.2) and (3.6), that is, $f_{n+1}(x) = h_{n+1}(x) = 0, x \in \mathbb{C}$, imply

$$F_{n,x} = -2izF_n + 2qG_{n+1}, \quad G_{n+1,x} = pF_n + qH_n, \quad H_{n,x} = 2izH_n + 2pG_{n+1}.$$
(3.9)

Equations (3.9) yield

$$G_{n+1}(z,x)^2 - F_n(z,x)H_n(z,x) = R_{2n+2}(z), \qquad (3.10)$$

where $R_{2n+2}(z)$ is a monic polynomial in z of degree 2n+2 and hence of the form 2n+1

$$R_{2n+2}(z) = \prod_{m=0}^{2n+1} (z - E_m), \quad \{E_m\}_{m=0,\dots,2n+1} \subset \mathbb{C}.$$
 (3.11)

Moreover, (3.9) and (3.10) imply

$$F_{n}(z,x)F_{n,xx}(z,x) - \frac{q_{x}(x)}{q(x)}F_{n}(z,x)F_{n,x}(z,x) - \frac{1}{2}F_{n,x}(z,x)^{2} + \left(2z^{2} - 2iz\frac{q_{x}(x)}{q(x)} - 2p(x)q(x)\right)F_{n}(z,x)^{2} = -2q(x)^{2}R_{2n+2}(z), \qquad (3.12)$$
$$H_{n}(z,x)H_{n,xx}(z,x) - \frac{p_{x}(x)}{p(x)}H_{n}(z,x)H_{n,x}(z,x) - \frac{1}{2}H_{n,x}(z,x)^{2} - \left(2z^{2} - \frac{p_{x}(x)}{p(x)} - \frac{p_{x}(x)}{p(x)}\right)F_{n}(z,x)H_{n,x}(z,x) - \frac{1}{2}H_{n,x}(z,x)^{2}$$

$$+\left(2z^{2}+2iz\frac{p_{x}(x)}{p(x)}-2p(x)q(x)\right)H_{n}(z,x)^{2}=-2p(x)^{2}R_{2n+2}(z).$$
(3.13)

Introducing the algebraic eigenspace

$$\ker(D-z) = \left\{ \Psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} : \mathbb{C} \mapsto (\mathbb{C}_{\infty})^2 \text{ meromorphic } \mid (D-z)\Psi = 0 \right\}, \quad z \in \mathbb{C},$$
(3.14)

one verifies

$$E_{n+1}\Big|_{\ker(D-z)} = i \begin{pmatrix} -G_{n+1}(z,x) & F_n(z,x) \\ -H_n(z,x) & G_{n+1}(z,x) \end{pmatrix} \Big|_{\ker(D-z)}.$$
 (3.15)

Moreover, the analog of the Burchnall–Chaundy polynomial for the KdV case in Section 2 now reads

$$E_{n+1}^2 + R_{2n+2}(D) = 0. (3.16)$$

Equation (3.16) naturally leads to the hyperelliptic curve $\dot{\mathcal{K}}_n$ defined by

$$\dot{\mathcal{K}}_n: \mathcal{F}_n(z,y) = y^2 - R_{2n+2}(z) = 0, \quad R_{2n+2}(z) = \prod_{m=0}^{2n+1} (z - E_m), \ z \in \mathbb{C}.$$
 (3.17)

The compactification of $\dot{\mathcal{K}}_n$, by joining $\{P_{\infty_-}, P_{\infty_+}\}$, the points at infinity, is then denoted \mathcal{K}_n . As in Section 2 we denote points $P \in \mathcal{K}_n \setminus \{P_{\infty_-}, P_{\infty_+}\}$ by P = (z, y), where $\mathcal{F}_n(z, y) = 0$. Moreover the involution * on \mathcal{K}_n is defined by

*:
$$\mathcal{K}_n \mapsto \mathcal{K}_n, \quad P = (z, y) \mapsto P^* = (z, -y), \quad P^*_{\infty_{\mp}} = P_{\infty_{\pm}}.$$
 (3.18)

Because of (3.10) we may define the following fundamental meromorphic function $\phi(P,x)$ on \mathcal{K}_n

$$\phi(P,x) = \frac{y(P) + G_{n+1}(z,x)}{F_n(z,x)}$$
(3.19a)

$$=\frac{-H_n(z,x)}{y(P)-G_{n+1}(z,x)}, \quad P=(z,y)\in\mathcal{K}_n, x\in\mathbb{C},$$
(3.19b)

where y(P) denotes the meromorphic function on \mathcal{K}_n obtained upon solving $y^2 = R_{2n+2}(z)$ denoting P = (z, y). The associated Baker–Akhiezer vector $\Psi(P, x, x_0)$ on $\mathcal{K}_n \setminus \{P_{\infty_-}, P_{\infty_+}\}$ is then defined by

$$\psi_{1}(P, x, x_{0}) = \exp\left(\int_{x_{0}}^{x} dx' \left(-iz + q(x')\phi(P, x')\right)\right),$$

$$\psi_{2}(P, x, x_{0}) = \phi(P, x)\psi_{1}(P, x, x_{0}),$$

$$\Psi(P, x, x_{0}) = \begin{pmatrix}\psi_{1}(P, x, x_{0})\\\psi_{2}(P, x, x_{0})\end{pmatrix}, \quad P = (z, y) \in \mathcal{K}_{n} \setminus \{P_{\infty_{-}}, P_{\infty_{+}}\}, x, x_{0} \in \mathbb{C}, \quad (3.20)$$

choosing a small non-selfintersecting path from x_0 to x avoiding singularities of q(x) and $\phi(P, x)$. The analog of Lemma 2.1 then reads as follows.

Lemma 3.1. (see, e.g., [30], [31]) Suppose $f_{n+1} = h_{n+1} = 0$ and let $P = (z, y) \in \mathcal{K}_n \setminus \{P_{\infty_-}, P_{\infty_+}\}, x, x_0 \in \mathbb{C}$. Then $\phi(P, x)$ satisfies the Riccati-type equation

$$\phi_x(P,x) + q(x)\phi(P,x)^2 - 2iz\phi(P,x) = p(x), \qquad (3.21)$$

and

$$\phi(P, x)\phi(P^*, x) = H_n(z, x)/F_n(z, x), \tag{3.22}$$

$$\phi(P, x) + \phi(P^*, x) = 2G_{n+1}(z, x) / F_n(z, x), \qquad (3.23)$$

$$\phi(P, x) - \phi(P^*, x) = 2y(P)/F_n(z, x). \tag{3.24}$$

Moreover, if $p(x)q(x) \neq 0$,

$$\phi(P, x) = O(z^{\pm 1}) \text{ as } P = (z, x) \to P_{\infty_{\pm}}.$$
 (3.25)

 $\Psi(P, x, x_0)$ satisfies

$$(D - z(P))\Psi(P, \cdot, x_0) = 0, \quad (E_{n+1} - iy(P))\Psi(P, \cdot, x_0) = 0, \quad (3.26)$$

and

$$\psi_1(P, x, x_0)\psi_1(P^*, x, x_0) = F_n(z, x)/F_n(z, x_0), \qquad (3.27)$$

$$\psi_2(P, x, x_0)\psi_2(P^*, x, x_0) = H_n(z, x)/F_n(z, x_0), \qquad (3.28)$$

$$\psi_1(P, x, x_0)\psi_2(P^*, x, x_0) + \psi_1(P^*, x, x_0)\psi_2(P, x, x_0)$$

$$= 2G_{n+1}(z,x)/F_n(z,x_0), \qquad (3.29)$$

$$W(\Psi(P, \cdot, x_0), \Psi(P^*, \cdot, x_0)) = -2y(P)/F_n(z, x_0).$$
(3.30)

Moreover, if $p(x)q(x) \neq 0$, $q(x_0) \neq 0$, then

$$\psi_1(P, x, x_0) = \exp(z^{-n}y(P)(x - x_0))(1 + O(z^{-1})),$$

$$\psi_2(P, x, x_0) = \exp(z^{-n}y(P)(x - x_0))O(z^{\mp 1}) \text{ as } P = (z, y) \to P_{\infty_{\pm}}.$$
 (3.31)

(Here $W(F,G) = f_1g_2 - f_2g_1$ denotes the Wronskian of $F = \begin{pmatrix} f_1 \\ f_2 \end{pmatrix}$ and $G = \begin{pmatrix} g_1 \\ g_2 \end{pmatrix}$.) Since the diagonal elements $G_{\ell,\ell}(P,x,x')$, $\ell = 1,2$ of the 2 × 2 Green's matrix G(P,x,x'), $x \neq x'$ associated with D are discontinuous as $x \to x'$ in contrast to the off-diagonal elements $G_{\ell,\ell'}(P,x,x')$, $\ell \neq \ell'$, $\ell,\ell' = 1,2$, we consider

$$G_{1,2}(P,x,x) = -i\frac{\psi_1(P,x,x_0)\psi_1(P^*,x,x_0)}{W(\Psi(P,\cdot,x_0),\Psi(P^*,\cdot,x_0))}$$
(3.32a)

$$=\frac{iF_n(z,x)}{2y(P)}, \quad P=(z,y)\in\mathcal{K}_n\setminus\{P_{\infty_-},P_{\infty_+}\}, x\in\mathbb{C},$$
(3.32b)

using $\phi = \psi_2/\psi_1$, (3.24), and (3.27). Equations (3.12) and (3.32b) then yield the universal equation

$$2G_{1,2}''(P,x,x)G_{1,2}(P,x,x) - 2\frac{q_x(x)}{q(x)}G_{1,2}'(P,x,x)G_{1,2}(P,x,x) - G_{1,2}'(P,x,x)^2 + 2\left(2z^2 - 2iz\frac{q_x(x)}{q(x)} - 2p(x)q(x)\right)G_{1,2}(P,x,x)^2 = q(x)^2,$$

$$P = (z,y) \in \mathcal{K}_n \setminus \{P_{\infty_-}, P_{\infty_+}\}.$$
(3.33)

Similarly, we find

$$G_{2,1}(P,x,x) = -i\frac{\psi_2(P,x,x_0)\psi_2(P^*,x,x_0)}{W(\Psi(P,\cdot,x_0),\Psi(P^*,\cdot,x_0))}$$
(3.34a)

$$=\frac{iH_n(z,x)}{2y(P)}, \quad P=(z,y)\in\mathcal{K}_n\setminus\{P_{\infty_-},P_{\infty_+}\}, x\in\mathbb{C},$$
(3.34b)

and

$$G_{2,1}^{\prime\prime}(P,x,x)G_{2,1}(P,x,x) - 2\frac{p_x(x)}{p(x)}G_{2,1}^{\prime}(P,x,x)G_{2,1}(P,x,x) - G_{2,1}^{\prime}(P,x,x)^2 + 2\left(2z^2 + 2iz\frac{p_x(x)}{p(x)} - 2p(x)q(x)\right)G_{2,1}(P,x,x)^2 = p(x)^2,$$

$$P = (z,y) \in \mathcal{K}_n \setminus \{P_{\infty_-}, P_{\infty_+}\}, x \in \mathbb{C}. \quad (3.35)$$

This completes our review of the stationary AKNS hierarchy. The corresponding time-dependent AKNS hierarchy can now readily be introduced, see, for instance, [30], [31].

Next we turn to gauge (i.e., Darboux-type) transformations in connection with $D = i \begin{pmatrix} d/dx & -q \\ p & -d/dx \end{pmatrix}$. Introducing

$$U(z,x) = \begin{pmatrix} -iz & -q(x) \\ p(x) & iz \end{pmatrix},$$
(3.36)

the equation

$$D\Psi(z,x) = z\Psi(z,x), \quad \Psi(z,x) = \begin{pmatrix} \psi_1(z,x)\\ \psi_2(z,x) \end{pmatrix}$$
(3.37)

is equivalent to

$$\Psi_x(z,x) = U(z,x)\Psi(z,x).$$
(3.38)

The formal gauge transformation

$$\Psi(z,x) \mapsto \widetilde{\Psi}(z,x) = \Gamma(z,x)\Psi(z,x), \tag{3.39}$$

$$U(z,x)) \mapsto \widetilde{U}(z,x) = \begin{pmatrix} -iz & -\widetilde{q}(x)\\ \widetilde{p}(x) & iz \end{pmatrix}$$
$$= \Gamma(z,x)U(z,x)\Gamma(z,x)^{-1} + \Gamma_x(z,x)\Gamma(z,x)^{-1}, \qquad (3.40)$$

with $\Gamma(z, x) \ge 2 \times 2$ matrix to be chosen later, then implies

$$\widetilde{\Psi}_x(z,x) = \widetilde{U}(z,x)\widetilde{\Psi}(z,x).$$
(3.41)

Hence,

$$\widetilde{D}\widetilde{\Psi}(z,x) = z\widetilde{\Psi}(z,x), \quad \widetilde{\Psi}(z,x) = \begin{pmatrix} \widetilde{\psi}_1(z,x)\\ \widetilde{\psi}_2(z,x) \end{pmatrix}, \quad (3.42)$$

with

$$\widetilde{D} = i \begin{pmatrix} d/dx & -\widetilde{q} \\ \widetilde{p} & -d/dx \end{pmatrix}.$$
(3.43)

Next, introduce

$$\Psi(P, x, x_0, \sigma) = \begin{cases} \frac{1}{2}(1+\sigma)\Psi(P, x, x_0) + \frac{1}{2}(1-\sigma)\Psi(P^*, x, x_0) & \text{for } \sigma \in \mathbb{C}, \\ \Psi(P, x, x_0) - \Psi(P^*, x, x_0) & \text{for } \sigma = \infty, \end{cases}$$

$$P = (z, y) \in \mathcal{K}_n \setminus \{P_{\infty_-}, P_{\infty_+}\}, \quad (3.44)$$

pick $Q_0 = (z_0, y_0) \in \mathcal{K}_n \setminus \{P_{\infty_-}, P_{\infty_+}\}$, and define

$$\Gamma(Q_0, x, \sigma) = \begin{pmatrix} z - z_0 - \frac{i}{2}q(x)\phi(Q_0, x, \sigma) & \frac{i}{2}q(x) \\ \frac{i}{2}\phi(Q_0, x, \sigma) & -\frac{i}{2} \end{pmatrix}.$$
 (3.45)

Here $\Psi(P, x, x_0)$ is defined in (3.19) and

$$\phi(P, x, \sigma) = \psi_2(P, x, x_0, \sigma) / \psi_1(P, x, x_0, \sigma),$$

$$P = (z, y) \in \mathcal{K}_n \setminus \{P_{\infty_-}, P_{\infty_+}\}, \ \sigma \in \mathbb{C}_\infty.$$
(3.46)

According to (3.40) one then obtains

$$\widetilde{D}_{\sigma}(Q_0) = i \begin{pmatrix} d/dx & -\widetilde{q}_{\sigma}(Q_0) \\ \widetilde{p}_{\sigma}(Q_0) & -d/dx \end{pmatrix}, \quad \sigma \in \mathbb{C}_{\infty},$$
(3.47a)

$$\tilde{p}_{\sigma}(x, Q_0) = \phi(Q_0, x, \sigma), \qquad (3.47b)$$

$$\tilde{q}_{\sigma}(x,Q_0) = -2iz_0 q(x) - q_x(x) + \phi(Q_0, x, \sigma)q(x)^2, \qquad (3.47c)$$

$$Q_0 = (z_0, y_0) \in \mathcal{K}_n \setminus \{P_{\infty_-}, P_{\infty_+}\}, \ \sigma \in \mathbb{C}_\infty,$$
(3.47d)

utilizing the fact that $\phi(P, x, \sigma)$ satisfies the Riccati-type equation (3.21) for all $\sigma \in \mathbb{C}_{\infty}$. The gauge transformation (or equivalently, Darboux transformation)

$$(p(x), q(x)) \mapsto (\tilde{p}_{\sigma}(x), \tilde{q}_{\sigma}(x))$$
 (3.48)

can be inferred from the results in [41] (with a bit of additional work). Adding solitons (i.e., inserting eigenvalues into the spectrum of D) and its effect on the Baker-Akhiezer vector Ψ has also been studied in [22], [24].

We also mention that as in Section 2 (cf. (2.43)–(2.44)), our choice of the offdiagonal elements of the Green's matrix of D in (3.32) yields the only bounded Green's matrix for P near $P_{\infty_{\pm}}$ and generic $x \in \mathbb{C}$ when comparing with more general linear combinations $\Psi(P, x, x_0, \sigma)$ in (3.44) as opposed to $\Psi(P, x, x_0)$ or $\Psi(P^*, x, x_0)$. Introducing

$$\Psi_{\sigma}(P, x, x_0, Q_0) = \Gamma(Q_0, x, \sigma) \Psi(P, x, x_0),$$

$$P = (z, y) \in \mathcal{K}_n \setminus \{P_{\infty_-}, P_{\infty_+}\}, \ \sigma \in \mathbb{C}_{\infty}, \quad (3.49)$$

where $\Psi(P, \cdot, x_0) \in \ker(D-z)$ and $\Gamma(Q_0, x, \sigma)$ is defined in (3.45), one infers

$$\widetilde{D}_{\sigma}(Q_0)\widetilde{\Psi}_{\sigma}(P, x, x_0, Q_0) = z\widetilde{\Psi}_{\sigma}(P, x, x_0, Q_0).$$
(3.50)

Moreover,

$$W(\tilde{\Psi}_{\sigma,1}(P,\cdot,x_0,Q_0),\tilde{\Psi}_{\sigma,2}(P,\cdot,x_0,Q_0)) = -\frac{i}{2}(z-z_0)W(\Psi_1(P,\cdot,x_0),\Psi_2(P,\cdot,x_0)), \qquad (3.51)$$

where

$$\widetilde{\Psi}_{\sigma,j}(P, x, x_0, Q_0) = \Gamma(Q_0, x, \sigma) \Psi_j(P, x, x_0), \Psi_j(P, \cdot, x_0) \in \ker(D-z), \ j = 1, 2.$$
(3.52)

Given these facts we define in analogy to (3.32a) and (3.34a) the off-diagonal Green's matrix elements associated with $\widetilde{D}_{\sigma}(Q_0)$ by

$$\widetilde{G}_{\sigma,1,2}(P,x,x,Q_0) = -i \frac{\widetilde{\psi}_{\sigma,1}(P,x,x_0,Q_0)\widetilde{\psi}_{\sigma,1}(P^*,x,x_0,Q_0)}{W(\widetilde{\Psi}_{\sigma}(P,\cdot,x_0,Q_0),\widetilde{\Psi}_{\sigma}(P^*,\cdot,x_0,Q_0))},$$
(3.53)

DARBOUX-TYPE TRANSFORMATIONS

$$\widetilde{G}_{\sigma,2,1}(P,x,x,Q_0) = -i \frac{\widetilde{\psi}_{\sigma,2}(P,x,x_0,Q_0)\widetilde{\psi}_{\sigma,2}(P^*,x,x_0,Q_0)}{W(\widetilde{\Psi}_{\sigma}(P,\cdot,x_0,Q_0),\widetilde{\Psi}_{\sigma}(P^*,\cdot,x_0,Q_0))},$$

$$P = (z,y) \in \mathcal{K}_n \setminus \{Q_0, P_{\infty_-}, P_{\infty_+}\}, \ \sigma \in \mathbb{C}_{\infty}.$$
(3.54)

Lemma 3.2. Assume $f_{n+1} = h_{n+1} = 0$ and let $Q_0 = (z_0, y_0) \in \mathcal{K}_n \setminus \{P_{\infty_-}, P_{\infty_+}\}, P = (z, y) \in \mathcal{K}_n \setminus \{Q_0, P_{\infty_-}, P_{\infty_+}\}, \sigma \in \mathbb{C}_{\infty}$. Then the off-diagonal elements of the Green's matrix of $\widetilde{D}_{\sigma}(Q_0)$ read

$$\widetilde{G}_{\sigma,1,2}(P,x,x,Q_0) = i \left(\left(i(z-z_0)q(x) + \frac{1}{2}q(x)^2\phi(Q_0,x,\sigma) \right) G_{n+1}(z,x) - \frac{1}{4}q(x)^2 H_n(z,x) + \left((z-z_0)^2 - i(z-z_0)q(x)\phi(Q_0,x,\sigma) - \frac{1}{4}q(x)^2\phi(Q_0,x,\sigma)^2 \right) F_n(z,x) \right) \times \\
\times \left(- (z-z_0)y(P) \right)^{-1} \tag{3.55a}$$

$$= i \left((z - z_0) + \frac{i}{2} q(x) (\phi(P, x) - \phi(Q_0, x, \sigma)) \right) \times \\ \times \left((z - z_0) + \frac{i}{2} q(x) (\phi(P^*, x) - \phi(Q_0, x, \sigma)) \right) F_n(z, x) (-(z - z_0) y(P))^{-1}$$
(3.55b)

$$=\frac{iF_{\sigma,\tilde{n}}(z,x)}{2\tilde{y}(P)}$$
(3.55c)

and

=

$$\widetilde{G}_{\sigma,2,1}(P, x, x, Q_0) = \frac{i}{4} \big(\phi(Q_0, x, \sigma)^2 F_n(z, x) - 2\phi(Q_0, x, \sigma) G_{n+1}(z, x) + H_n(z, x) \big) \times \\ \times \big((z - z_0) y(P) \big)^{-1}$$

$$= \frac{i}{4} \big(\phi(P, x) - \phi(Q_0, x, \sigma) \big) \big(\phi(P^*, x) - \phi(Q_0, x, \sigma) \big) F_n(z, x) \big((z - z_0) y(P) \big)^{-1}$$

$$(3.56b)$$

$$=\frac{iH_{\sigma,\tilde{n}}(z,x)}{2\tilde{y}(P)},\tag{3.56c}$$

where $\tilde{y}(P)$ denotes the meromorphic solution on $\tilde{\mathcal{K}}_{\sigma,\tilde{n}}(Q_0)$ obtained upon solving $y^2 = \tilde{R}_{\sigma,2\tilde{n}+2}(z), P = (z,y)$ for some polynomial $\tilde{R}_{\sigma,2\tilde{n}+2}(z)$ of degree $2\tilde{n}+2$ and $\tilde{F}_{\sigma,\tilde{n}}(z,x)$ and $\tilde{H}_{\sigma,\tilde{n}}(z,x)$ denote polynomials in z of degree \tilde{n} , with $0 \leq \tilde{n} \leq n+1$. In particular, the Darboux transformation (3.48), $(p(x), q(x)) \mapsto (\tilde{p}_{\sigma}(x), \tilde{q}_{\sigma}(x))$ maps the class of algebro-geometric AKNS potentials into itself.

Proof. We present the argument for $\widetilde{G}_{\sigma,1,2}$ only, the case $\widetilde{G}_{\sigma,2,1}$ follows similarly. As in Lemma 3.2, $\phi(P, x) = \psi_2(P, x, x_0)/\psi_1(P, x, x_0)$, (3.22), (3.23), (3.27)–(3.30), and (3.51) imply equations (3.55a) and (3.55b). Since the numerator in (3.55a) is a polynomial in z, and

$$\widetilde{G}_{\sigma,1,2}(P,x,x,Q_0) = \frac{\widetilde{q}(x)z^n}{2y(P)} + O(|z|^{-2}) \text{ as } P = (z,y) \to P_{\infty_{\pm}}$$
(3.57)

by (3.55a), one infers (3.55c) and $0 \leq \tilde{n} \leq n+1$. Again one verifies that $\widetilde{F}_{\sigma,\tilde{n}}(z,x)$ satisfies equation (3.12) with (p(x), q(x)) replaced by $(\tilde{p}_{\sigma}(x), \tilde{q}_{\sigma}(x)), n$ by \tilde{n} , and $R_{2n+2}(z)$ by $\widetilde{R}_{\sigma,2\tilde{n}+2}(z)$, proving that the Darboux transformation (3.48) leaves the class of algebro-geometric AKNS potentials invariant.

The following theorem, the principal result of this section and in complete analogy to Theorem 2.3, will clarify the dependence of $\tilde{n} = \tilde{n}(n, Q_0, \sigma)$ on its variables.

Theorem 3.3. Suppose $f_{n+1} = h_{n+1} = 0$, let $Q_0 = (z_0, y_0) \in \mathcal{K}_n \setminus \{P_{\infty_-}, P_{\infty_+}\}$, $n \in \mathbb{N}_0, \sigma \in \mathbb{C}_\infty$, and $\tilde{n} = \tilde{n}(n, Q_0, \sigma)$ as in (3.55c). Then

$$\tilde{n}(n,Q_{0},\sigma) = \begin{cases} n+1 & \text{for } \sigma \in \mathbb{C}_{\infty} \setminus \{-1,1\} \text{ and } y_{0} \neq 0, \\ n+1 & \text{for } \sigma = \infty \text{ and } y_{0} = 0, \\ n & \text{for } \sigma \in \{-1,1\} \text{ and } y_{0} \neq 0, \\ n & \text{for } \sigma \in \mathbb{C}, y_{0} = 0, \text{ and } R_{2n+2,z}(z_{0}) \neq 0, \\ n-1 & \text{for } \sigma \in \mathbb{C}, y_{0} = 0, \text{ and } R_{2n+2,z}(z_{0}) = 0, n \in \mathbb{N}, \end{cases}$$

$$(3.58)$$

and hence the hyperelliptic curve $\dot{\widetilde{\mathcal{K}}}_{\sigma,\widetilde{n}}(Q_0)$ associated with $(\widetilde{p}_{\sigma}(x,Q_0),\widetilde{q}_{\sigma}(x,Q_0))$ is of the type

$$\widetilde{\mathcal{K}}_{\sigma,\tilde{n}} \colon \widetilde{\mathcal{F}}_{\sigma,\tilde{n}}(z,y,Q_0) = y^2 - \widetilde{R}_{\sigma,2\tilde{n}+2}(z,Q_0) = 0,$$
(3.59)

with

()

$$\widetilde{R}_{\sigma,2\widetilde{n}+2}(z,Q_{0}) = \begin{cases}
(z-z_{0})^{2}R_{2n+2}(z) & \text{for } \sigma \in \mathbb{C}_{\infty} \setminus \{-1,1\} \text{ and } y_{0} \neq 0, \\
(z-z_{0})^{2}R_{2n+2}(z) & \text{for } \sigma = \infty \text{ and } y_{0} = 0, \\
R_{2n+2}(z) & \text{for } \sigma \in \{-1,1\} \text{ and } y_{0} \neq 0, \\
R_{2n+2}(z) & \text{for } \sigma \in \mathbb{C}, y_{0} = 0, \text{ and } R_{2n+2,z}(z_{0}) \neq 0, \\
(z-z_{0})^{-2}R_{2n+2}(z) & \text{for } \sigma \in \mathbb{C}, y_{0} = 0, \text{ and } R_{2n+2,z}(z_{0}) = 0, n \in \mathbb{N}.
\end{cases}$$
(3.60)

Here

$$R_{2n+2}(z) = \prod_{m=0}^{2n+1} (z - E_m).$$
(3.61)

Proof. The following arguments closely parallel those in the proof of Theorem 2.3. Again our starting point will be (3.55b) and (3.56b) and a careful case distinction between $\sigma \in \mathbb{C}_{\infty} \setminus \{-1, 1\}, \sigma \in \{-1, 1\}, \sigma = \infty$, and whether or not Q_0 is a branch point.

Case (i). $\sigma \in \mathbb{C}_{\infty} \setminus \{-1, 1\}$ and $y_0 \neq 0$: One calculates using (3.44) and (3.46),

$$\phi(Q_0, x, \sigma) = \begin{cases} \frac{(1+\sigma)\psi_2(Q_0, x, x_0) + (1-\sigma)\psi_2(Q_0^*, x, x_0)}{(1+\sigma)\psi_1(Q_0, x, x_0) + (1-\sigma)\psi_1(Q_0^*, x, x_0)} & \text{for } \sigma \in \mathbb{C} \setminus \{-1, 1\}, \\ \frac{\psi_2(Q_0, x, x_0) - \psi_2(Q_0^*, x, x_0)}{\psi_1(Q_0, x, x_0) - \psi_1(Q_0^*, x, x_0)} & \text{for } \sigma = \infty, \end{cases}$$
(3.62)

and since

$$\phi(Q_0, x) = \frac{\psi_1(Q_0, x, x_0)}{\psi_2(Q_0, x, x_0)} \neq \phi(Q_0^*, x) = \frac{\psi_1(Q_0^*, x, x_0)}{\psi_2(Q_0^*, x, x_0)},$$
(3.63)

one concludes that no cancellations can occur in (3.55b) or (3.56b) and hence $\tilde{n}(n, Q_0, \sigma) = n + 1.$

Case (ii). $\sigma = \infty$ and $y_0 = 0$: Then (3.19a), (3.20), (3.24) and (3.44) imply $\phi(Q_0, x, \infty) = \lim_{P \to Q_0} \phi(P, x, \infty)$ $= \lim_{P \to Q_0} \left(\left(\phi(P, x) \exp\left(\int_{x_0}^x dx' \left(-iz + q(x')\phi(P^*, x') \right) \right) \right) \times \left(\exp\left(\int_{x_0}^x dx' \left(-iz + q(x')\phi(P^*, x') \right) \right) \right) \right) \times \left(\exp\left(\int_{x_0}^x dx' \left(-iz + q(x')\phi(P^*, x') \right) \right) \right) \right)$ $= \phi(Q_0, x)$ $+ \lim_{P \to Q_0} \left(\frac{\phi(P, x) - \phi(P^*, x)}{\exp\left(\int_{x_0}^x dx' q(x')\phi(P, x')\right) - \exp\left(\int_{x_0}^x dx' q(x')\phi(P^*, x')\right) \right) \right)$ $= \phi(Q_0, x) + \exp\left(\int_{x_0}^x dx' q(x')\phi(Q_0, x')\right) \times \left(\exp\left(\int_{x_0}^x dx' q(x')\phi(Q_0, x')\right) \right) \right)$ $= \phi(Q_0, x) + \exp\left(\int_{x_0}^x dx' q(x')\phi(Q_0, x')\right) \times \left(\exp\left(\int_{x_0}^x dx' q(x')\phi(Q_0, x')\right) \right) \right)$

$$\times \exp\left(\int_{x_0}^x dx' \, \frac{q(x')G_{n+1}(z,x')}{F_n(z,x')}\right)\right)^{-1} \frac{2y(P)}{F_n(z,x)}$$

$$= \phi(Q_0,x) + \left(F_n(z_0,x)\int_{x_0}^x dx' \, \frac{q(x')}{F_n(z,x')}\right)^{-1}, \quad x \in \mathbb{C} \setminus \{x_0\}, \tag{3.64}$$

using $\lim_{P \to Q_0} y(P) = y(Q_0) = y_0 = 0$. Since by (3.9)

$$\phi(Q_0, x) = \frac{G_{n+1,x}(z_0, x)}{F_n(z_0, x)} = \frac{1}{q(x)} \left(\frac{F_{n,x}(z_0, x)}{2F_n(z_0, x)} + iz_0\right),\tag{3.65}$$

one again concludes that no cancellation occurs in (3.55b) or (3.56b), and hence $\tilde{n}(n, Q_0, \infty) = n + 1$.

The rest of the proof relies on some additional arguments to be discussed next. First, we will assume without loss of generality that $z_0 = 0$. This can be achieved by noticing that

$$D = i \begin{pmatrix} d/dx & -q(x) \\ p(x) & -d/dx \end{pmatrix} \text{ and } D_a = i \begin{pmatrix} d/dx & -q(x)e^{-2ax} \\ p(x)e^{2ax} & -d/dx \end{pmatrix}$$
(3.66)

are related by

$$UD_a U^{-1} = D - iaI, \quad U = \begin{pmatrix} e^{2ax} & 0\\ 0 & e^{-2ax} \end{pmatrix}.$$
 (3.67)

Moreover, we will exclude the trivial case where p(x) and q(x) are proportional to e^{-2ic_1x} and e^{2ic_1x} , $x \in \mathbb{C}$, respectively (cf. Example 3.4 below). We write

$$y(z)^{2} = R_{2n+2}(z) \underset{z \to 0}{=} y_{0}^{2} + \tilde{y}_{1}z + \tilde{y}_{2}z^{2} + O(z^{3}).$$
(3.68)

Case (iii). $\sigma \in \{-1, 1\}$ and $y_0 \neq 0$: Then (3.44) yields

$$\phi(Q_0, x, 1) = \phi(Q_0, x), \quad \phi(Q_0, x, -1) = \phi(Q_0^*, x), \tag{3.69}$$

with $\phi(Q_0, x) \neq \phi(Q_0^*, x)$ since $y_0 \neq 0$. In this case there is a cancellation in (3.55b) and (3.56b). Choosing $\sigma = 1$ one computes from (3.8) and (3.19a),

$$\phi(P^*, x) - \phi(Q_0, x, 1) = \phi(P^*, x) - \phi(Q_0, x) = -\frac{2y_0}{f_n(x)} + O(z).$$
(3.70)

Furthermore,

$$\begin{split} \phi(P,x) &- \phi(Q_0,x,1) = \phi(P,x) - \phi(Q_0,x) \\ &= \frac{y(P) - y_0}{f_n(x)} - \left(y_0 \frac{f_{n-1}(x)}{f_n(x)^2} - \frac{g_n(x)}{f_n(x)} + \frac{g_{n+1}(x)f_{n-1}(x)}{f_n(x)^2}\right) z + O(z^2) \\ &= \frac{y_0(x)}{f_n(x)} - \frac{y_0 f_{n-1}(x)}{f_n(x)^2} + \frac{g_n(x)}{f_n(x)} - \frac{g_{n+1}(x)f_{n-1}(x)}{f_n(x)^2}\right) z + O(z^2) \\ &= \frac{z_0}{P \to Q_0} c_1(x) z + O(z^2) \end{split}$$
(3.71)

since

$$y(P) - y_0 = \underset{P \to Q_0}{=} y_1 z + O(z^2), \quad \tilde{y}_1 = 2y_0 y_1.$$
 (3.72)

Similarly, we find

$$z + \frac{i}{2}q(x)\big(\phi(P,x) - \phi(Q_0,x,1)\big) =_{P \to Q_0} \left(1 + \frac{i}{2}q(x)c_1(x)\right)z + O(z^2).$$
(3.73)

It remains to show that c_1 does not vanish identically. We assume temporarily that $G_{\sigma,1,2}$ and $G_{\sigma,2,1}$ have cancellations of the same order as $z \to 0$. Arguing by contradiction we suppose that c_1 vanishes identically. But (3.71) and (3.73) then show that $\tilde{G}_{\sigma,1,2}$ and $\tilde{G}_{\sigma,2,1}$ would have cancellations of different order, which is a contradiction. We conclude that precisely one factor of z cancels in (3.55b) and (3.56b), and hence $\tilde{n}(n, Q_0, 1) = n$. The case $\sigma = -1$ is treated analogously. It remains to show that $\widetilde{G}_{\sigma,1,2}$ and $\widetilde{G}_{\sigma,2,1}$ necessarily have cancellations of the same order as $z \to 0$. A comparison of $\widetilde{D}_{\sigma} = i \begin{pmatrix} d/dx & -\tilde{q}_{\sigma} \\ \tilde{p}_{\sigma} & -d/dx \end{pmatrix}$ and its formal adjoint $\widetilde{D}_{\sigma}^* = i \begin{pmatrix} \frac{d}{dx} & -\overline{\widetilde{p}_{\sigma}} \\ \overline{\widetilde{q}_{\sigma}} & -d/dx \end{pmatrix}$ yields the replacement of $(\widetilde{p}_{\sigma}, \widetilde{q}_{\sigma})$ by $(\overline{\widetilde{q}_{\sigma}}, \overline{\widetilde{p}_{\sigma}})$ and hence the corresponding replacements of $(\widetilde{F}_{\sigma,\tilde{n}}(z,x),\widetilde{G}_{\sigma,\tilde{n}+1}(z,x),\widetilde{H}_{\sigma,\tilde{n}}(z,x))$ by $\left(\overline{\widetilde{F}_{\sigma,\tilde{n}}(\overline{z},x)},\overline{\widetilde{G}_{\sigma,\tilde{n}+1}(\overline{z},x)},\overline{\widetilde{H}_{\sigma,\tilde{n}}(\overline{z},x)}\right)$ (cf. (3.2) and (3.8)) and $\widetilde{R}_{\sigma,2\tilde{n}+2}(z)$ by $\widetilde{R}_{\sigma,2\tilde{n}+2}(\overline{z})$ (cf. the notation employed in Lemma 3.2). This fact has two consequences: Firstly, from relation (3.10) we infer that the corresponding algebraic curves associated with D_{σ} and D_{σ}^* have complex conjugate branch points, that is, if $\{\widetilde{E}_{\sigma,m}\}_{m=0,\ldots,2\tilde{n}+1}$ corresponds to \widetilde{D}_{σ} , then $\{\overline{\widetilde{E}_{\sigma,m}}\}_{m=0,\ldots,2\tilde{n}+1}$ \widetilde{D}_{σ}^* , where $\widetilde{R}_{\sigma,2\tilde{n}+2}(z) = \prod_{m=0}^{2\tilde{n}+1} (z - \widetilde{E}_{\sigma,m})$. Secondly, we infer corresponds to $\widetilde{G}_{\sigma,2,1}(P,x,x) = \overline{\widetilde{G}_{\sigma,2,1}^*(\overline{P},x,x)},$ (3.74)

where P(z, y), $\overline{P} = (\overline{z}, \overline{y})$, and $\widetilde{G}^*_{\sigma}(P, x, x)$ denotes the Green's matrix associated with \widetilde{D}^*_{σ} . This shows that any cancellations in $\widetilde{G}_{\sigma,1,2}$ and $\widetilde{G}_{\sigma,2,1}$ as $z \to 0$ are necessarily of identical order.

Case (iv). $\sigma \in \mathbb{C}$, $y_0 = 0$, and $R_{2n+1,z}(0) \neq 0$: Using $\phi(Q_0, x, \sigma) = \phi(Q_0, x)$ for all $\sigma \in \mathbb{C}$, (3.8) and (3.19a) yield

$$\left(z + \frac{i}{2}q(x)\left(\phi(P, x) - \phi(Q_0, x)\right)\right) \left(z + \frac{i}{2}q(x)\left(\phi(P^*, x) - \phi(Q_0, x)\right)\right)$$

= $\sum_{P \to Q_0} \frac{q(x)^2 y_1^2}{4f_n(x)^2} z + O(z^2)$ (3.75)

since

$$y(P) =_{P \to Q_0} y_1 z^{1/2} + O(z^{3/2}), \quad y_1 = \left(\prod_{E_m \neq 0} E_m\right)^{1/2}.$$
 (3.76)

Thus again precisely one factor of z cancels in (3.55b) (similarly, one factor cancels in (3.56b)), and hence $\tilde{n}(n, Q_0, \sigma) = n$.

Case (v). $\sigma \in \mathbb{C}$, $y_0 = \tilde{y}_1 = 0$, and $\tilde{y}_2 \neq 0$ (cf. (3.68)): One computes as in (3.75),

$$\left(z + \frac{i}{2}q(x)\left(\phi(P, x) - \phi(Q_0, x)\right)\right) \left(z + \frac{i}{2}q(x)\left(\phi(P^*, x) - \phi(Q_0, x)\right)\right) \\ = \frac{1}{P \to Q_0} \left(\frac{q(x)^2 y_1^2}{4f_n(x)^2} + \left(1 + \frac{i}{2}q(x)\left(\frac{g_n(x)}{f_n(x)} - \frac{g_{n+1}(x)f_{n-1}(x)}{f_n(x)^2}\right)\right)^2\right) z^2 + O(z^3) \\ = \frac{1}{P \to Q_0} c_2(x)z^2 + O(z^3) \tag{3.77}$$

since

$$y(P) =_{P \to Q_0} y_1 z + O(z^2), \quad y_1 = \left(\prod_{E_m \neq 0} E_m\right)^{1/2}.$$
 (3.78)

Similarly, we find

$$\begin{pmatrix} \phi(P,x) - \phi(Q_0,x) \end{pmatrix} \begin{pmatrix} \phi(P^*,x) - \phi(Q_0,x) \end{pmatrix} = \\ P \to Q_0 - \left(\frac{y_1^2}{f_n(x)^2} - \left(\frac{g_n(x)}{f_n(x)} - \frac{g_{n+1}(x)f_{n-1}(x)}{f_n(x)^2} \right)^2 \right) z^2 + O(z^3) = \\ P \to Q_0 c_3(x) z^2 + O(z^3).$$
(3.79)

We see that both $c_2(x)$ and $c_3(x)$ cannot vanish simultaneously, and hence precisely a factor z^2 cancels in (3.55b) and (3.56b). Thus $\tilde{n}(n, Q_0, \sigma) = n - 1$.

Case (vi). $\sigma \in \mathbb{C}$, $y_0 = \tilde{y}_1 = \tilde{y}_2 = 0$ (cf. (3.68)): In analogy to (3.77) and (3.79) one obtains

$$\left(z + \frac{i}{2}q(x)\left(\phi(P, x) - \phi(Q_0, x)\right)\right) \left(z + \frac{i}{2}q(x)\left(\phi(P^*, x) - \phi(Q_0, x)\right)\right)$$

= $\left(1 + \frac{i}{2}q(x)\left(\frac{g_n(x)}{f_n(x)} - \frac{g_{n+1}f_{n-1}}{f_n^2}\right)\right)^2 z^2 + O(z^3)$ (3.80)

and

$$\left(\phi(P,x) - \phi(Q_0,x) \right) \left(\phi(P^*,x) - \phi(Q_0,x) \right)$$

$$= \frac{g_n(x)}{f_n(x)} - \frac{g_{n+1}f_{n-1}}{f_n^2} \right)^2 z^2 + O(z^3)$$
(3.81)

respectively, since

$$y(P) =_{P \to Q_0} O(z^{3/2}).$$
 (3.82)

Thus this case subordinates to case (v) resulting again in $\tilde{n}(n, Q_0, \sigma) = n - 1$.

While Lemma 3.2 has first been noted in [39] (see also [38]), Theorem 3.3 appears to be new. We also emphasize that Table 2.4 and Remark 2.6 apply in the present AKNS context.

We conclude this section with the elementary genus zero example.

Example 3.4. The case n = 0.

$$\begin{split} y(P)^2 &= R_2(z) = (z - E_0)(z - E_1), \\ c_1 &= -(E_0 + E_1)/2, \\ p(x) &= p(x_0)e^{-2ic_1(x - x_0)}, \quad q(x) = q(x_0)e^{2ic_1(x - x_0)}, \\ p(x)q(x) &= (E_0 - E_1)^2/4, \\ F_0(z, x) &= -iq(x), \quad G_1(z, x) = z + c_1, \quad H_0(z, x) = ip(x), \\ \phi(P, x) &= \frac{y(P) + z + c_1}{-iq(x)} = \frac{ip(x)}{y(P) - z - c_1}, \\ \psi_1 &= e^{i(y(P) + c_1)(x - x_0)}, \quad \psi_2 = \frac{y(P) + z + c_1}{-iq(x_0)}e^{i(y(P) - c_1)(x - x_0)}, \\ \phi(P, x, \sigma) &= \frac{i}{q(x)} \times \\ \times \begin{cases} \frac{(1 + \sigma)(y(P) + z + c_1)\exp(iy(P)(x - x_0)) + (1 - \sigma)(-y(P) + z + c_1)\exp(-iy(P)(x - x_0)))}{(1 + \sigma)\exp(iy(P)(x - x_0)) + (1 - \sigma)\exp(-iy(P)(x - x_0)))}, \quad \sigma \in \mathbb{C}, \\ \frac{(y(P) + z + c_1)\exp(iy(P)(x - x_0)) + (1 - \sigma)\exp(-iy(P)(x - x_0))}{\exp(iy(P)(x - x_0)) + \exp(-iy(P)(x - x_0))}, \quad \sigma = \infty, \end{cases} \\ \phi((E_j, 0), x, \sigma) &= \frac{1}{2q(x)} \begin{cases} i(E_j + c_1), \qquad \sigma \in \mathbb{C}, \\ i(E_j + c_1) + (x - x_0)^{-1}, \qquad \sigma = \infty, \end{cases} \end{cases}$$
(3.83)

Acknowledgments. We are indebted to Rudi Weikard for discussions on Darbouxtype transformations for AKNS systems.

References

- M. Adler and P. van Moerbeke, Birkhoff strata, Bäcklund transformations, and regularization of isospectral operators, Adv. Math. 108, 140–204 (1994).
- [2] M. Adler and J. Moser, On a class of polynomials connected with the Korteweg-de Vries equation, Comm. Math. Phys. 61, 1–30 (1978).
- [3] S. I. Al'ber, Investigation of equations of Korteweg-de Vries type by the method of recurrence relations, J. London Math. Soc. 19, 467–480 (1979) (Russian).
- [4] S. I. Al'ber, On stationary problems for equations of Korteweg-de Vries type, Commun. Pure Appl. Math. 34, 259–272 (1981).
- [5] E. D. Belokolos, A. I. Bobenko, V. Z. Enol'skii, A.R. Its, and V. B. Matveev, Algebro-Geometric Approach to Nonlinear Integrable Equations, Springer, Berlin, 1994.
- [6] J. L. Burchnall and T. W. Chaundy, Commutative ordinary differential operators, Proc. London Math. Soc. (2), 21, 420–440 (1923).
- [7] J. L. Burchnall and T. W. Chaundy, Commutative ordinary differential operators, Proc. Roy. Soc. London A118, 557–583 (1928).
- [8] J. L. Burchnall and T. W. Chaundy, Commutative ordinary differential operators II. The identity Pⁿ = Q^m, Proc. Roy. Soc. London A134, 471–485 (1932).

- [9] M. M. Crum, Associated Sturm-Liouville systems, Quart. J. Math. Oxford Ser. (2) 6, 121–127 (1955).
- [10] G. Darboux, Sur une proposition relative aux équations linéaire, C. R. Acad. Sci. Paris Sér. I Math. 94, 1456–1459 (1882).
- [11] P. A. Deift, Applications of a commutation formula, Duke Math. J. 45, 267–310 (1978).
- [12] P. Deift and E. Trubowitz, *Inverse scattering on the line*, Comm. Pure Appl. Math. 32, 121–251 (1979).
- [13] L. A. Dickey, Soliton Equations and Hamiltonian Systems, World Scientific, Singapore, 1991.
- [14] J. Drach, Sur les groupes complexes de rationalité et sur l'intégration par quadratures, C. R. Acad. Sci. Paris 167 (1918), 743–746.
- [15] J. Drach, Détermination des cas de réduction de l'équation différentielle d²y/dx² = [φ(x) + h]y, C. R. Acad. Sci. Paris 168 (1919), 47–50.
- [16] J. Drach, Sur l'intégration par quadratures de l'équation $d^2y/dx^2 = [\phi(x) + h]y$, C. R. Acad. Sci. Paris **168** (1919), 337–340.
- [17] B. A. Dubrovin, V. B. Matveev, and S. P. Novikov, Non-linear equations of Korteweg-de Vries type, finite-zone linear operators, and Abelian varieties, Russian Math. Surv. 31:1, 59–146 (1976).
- [18] M. S. P. Eastham and H. Kalf, Schrödinger-Type Operators with Continuous Spectra, Pitman, Boston, 1982.
- [19] F. Ehlers and H. Knörrer, An algebro-geometric interpretation of the Bäcklund transformation for the Korteweg-de Vries equation, Comment. Math. Helv. 57, 1–10 (1982).
- [20] G. Eilenberger, Solitons, Springer, Berlin, 1983.
- [21] N. M. Ercolani and H. Flaschka, The geometry of the Hill equation and of the Neumann system, Phil. Trans. Roy. Soc. London A 315, 405–422 (1985).
- [22] H. Flaschka, Relations between infinite-dimensional and finite-dimensional isospectral equations, in Non-linear Integrable Systems – Classical Theory and Quantum Theory, M. Sato (ed.), World Scientific, Singapore, 1983, pp. 219–240.
- [23] H. Flaschka and D. W. McLaughlin, Some comments on Bäcklund transformations, canonical transformations, and the inverse scattering method, in Bäcklund Transformations, the Inverse Scattering Method, Solitons, and their Applications, R. M. Miura (ed.), Lecture Notes in Math. 515, Springer, Berlin, 1976, pp. 252–295.
- [24] H. Flaschka and A. C. Newell, Multiphase similarity solutions of integrable evolution equations, Physica 3D, 203–221 (1981).
- [25] L. Gatto and S. Greco, Algebraic curves and differential equations: an introduction, The Curves Seminar at Queen's, Vol. VIII (ed. by A. V. Geramita), Queen's Papers Pure Appl. Math. 88, Queen's Univ., Kingston, Ontario, Canada, 1991, B1–B69.
- [26] I. M. Gel'fand and L. A. Dikii, Asymptotic behaviour of the resolvent of Sturm-Liouville equations and the algebra of the Korteweg-de Vries equations, Russian Math. Surv. 30:5, 77-113 (1975).
- [27] I. M. Gel'fand and L. A. Dikii, Integrable nonlinear equations and the Liouville theorem, Funct. Anal. Appl. 13, 6–15 (1979).
- [28] F. Gesztesy, A complete spectral characterization of the double commutation method, J. Funct. Anal. 117, 401–446 (1993).
- [29] F. Gesztesy and H. Holden, Dubrovin equations and integrable systems on hyperelliptic curves, preprint, 1998.
- [30] F. Gesztesy and H. Holden, Hierarchies of Soliton Equations and Their Algebro-Geometric Solutions, monograph in preparation.
- [31] F. Gesztesy and R. Ratneseelan, An alternative approach to algebro-geometric solutions of the AKNS hierarchy, Rev. Math. Phys. 10, 345–391 (1998).
- [32] F. Gesztesy, R. Ratnaseelan, and G. Teschl, The KdV hierarchy and associated trace formulas, in Recent Developments in Operator Theory and its Applications. Operator Theory: Advances and Applications, 87, I. Gohberg, P. Lancaster, and P. N. Shivakumar (eds.), Birkhäuser, Basel, 1996, pp. 125–163.
- [33] F. Gesztesy, W. Schweiger, and B. Simon, Commutation methods applied to the mKdVequation, Trans. Amer. Math. Soc. 324, 465–525 (1991).
- [34] F. Gesztesy, B. Simon and G. Teschl, Spectral deformation of one-dimensional Schrödinger operators, J. Analyse Math. 70, 267–324 (1996).

GESZTESY AND HOLDEN

- [35] F. Gesztesy and R. Svirsky, (m)KdV solitons on the background of quasi-periodic finite-gap solutions, Mem. Amer. Math. Soc. 118, no 563, (1995).
- [36] F. Gesztesy and R. Weikard, Spectral deformations and soliton equations, in Differential Equations with Applications to Mathematical Physics, W. F. Ames, E. M. Harrell II, J. V. Herod (eds.), Academic Press, Boston, 1993, pp. 101–139.
- [37] F. Gesztesy and R. Weikard, Picard potentials and Hill's equation on a torus, Acta Math. 176, 73–107 (1996).
- [38] F. Gesztesy and R. Weikard, Elliptic algebro-geometric solutions of the KdV and AKNS hierarchies – an analytic approach, Bull. Amer. Math. Soc. 35, 271–317 (1998).
- [39] F. Gesztesy and R. Weikard, A characterization of all elliptic algebro-geometric solutions of the AKNS hierarchy, Acta Math. 181, 63–108(1998).
- [40] C. G. T. Jacobi, Über eine neue Methode zur Integration der hyperelliptischen Differentialgleichungen und über die rationale Form ihrer vollständigen algebraischen Integralgleichungen, J. Reine Angew. Math. 32, 220–226 (1846).
- [41] B. G. Konopelchenko and C. Rogers, Bäcklund and reciprocal transformations: Gauge connections, in Nonlinear Equations in the Applied Sciences, W. F. Ames and C. Rogers (eds.), Academic Press, Boston, 1992, pp. 317–362.
- [42] V. B. Matveev and M. A. Salle, Darboux Transformations and Solitons, Springer, Berlin, 1991.
- [43] H. P. McKean, Theta functions, solitons, and singular curves, in Partial Differential Equations and Geometry, C. I. Byrnes (ed.), Marcel Dekker, New York, 1979, pp. 237–254.
- [44] H. P. McKean, Variation on a theme of Jacobi, Commun. Pure Appl. Math. 38, 669–678 (1985).
- [45] H. P. McKean, Geometry of KdV(1): Addition and the unimodular spectral classes, Rev. Mat. Iberoamericana 2, 235–261 (1986).
- [46] H. P. McKean, Geometry of KdV(2): Three examples, J. Statist. Phys. 46, 1115–1143 (1987).
- [47] D. McRae and R. Weikard, Theta functions on singular hyperelliptic surfaces, preprint, 1997.
- [48] P. D. Miller, N. M. Ercolani, I. M. Krichever, and C. D. Levermore, *Finite genus solutions to the Ablowitz-Ladik equations*, Commun. Pure Appl. Math. 48, 1369–1440 (1995).
- [49] R. M. Miura, Korteweg-de Vries equation and generalizations. I. A remarkable explicit nonlinear transformation, J. Math. Phys. 9, 1202–1204 (1968).
- [50] M. Ohmiya, On the Darboux transformation of the second order differential operator of Fuchsian type on the Riemann sphere, Osaka J. Math., 25, 607–632 (1988).
- [51] M. Ohmiya, KdV polynomials and Λ-operator, Osaka J. Math., 32, 409-430 (1995).
- [52] M. Ohmiya, Spectrum of Darboux transformation of differential operator, preprint, Tokushima University, 1997.
- [53] M. Ohmiya and Y. P. Mishev, Darboux transformation and Λ-operator, J. Math. Tokushima Univ., 27, 1–15 (1993).
- [54] E. Previato, Seventy years of spectral curves: 1923–1993, in Integrable Systems and Quantum Groups, M. Francaviglia and S. Greco (eds.), Lecture Notes in Math., Vol. 1620, Springer, Berlin, 1996, pp. 419–481.
- [55] E. Previato, Burchnall-Chaundy bundles, in Algebraic Geometry, P. E. Newstead (ed.), Marcel Dekker, New York, 1998, pp. 377–383.
- [56] U.-W. Schmincke, On Schrödinger's factorization method for Sturm-Liouville operators, Proc. Roy. Soc. Edinburgh Sect. A, 80, 67–84 (1978).
- [57] G. Segal and G. Wilson, Loop groups and equations of KdV type, Publ. Math. IHES 61, 5–65 (1985).
- [58] A. P. Veselov and A. B. Shabat, Dressing chains and the spectral theory of the Schrödinger operator, Funct. Appl. 27, 81–96 (1993).
- [59] G. Wilson, Algebraic curves and soliton equations, in Geometry Today, E. Arbarello, C. Procesi, and E. Strickland (eds.), Birkhäuser, Boston, 1985, pp. 303–329.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA *E-mail address*: fritz@math.missouri.edu *URL*: http://www.math.missouri.edu/people/fgesztesy.html

Department of Mathematical Sciences, Norwegian University of Science and Technology, N–7034 Trondheim, Norway

E-mail address: holden@math.ntnu.no URL: http://www.math.ntnu.no/~holden/