arXiv:1312.6331v1 [math.AC] 22 Dec 2013

ON MODULAR COMPUTATION OF GRÖBNER BASES WITH INTEGER COEFFICIENTS

S. Yu. Orevkov

§1. Introduction. Let $A = \mathbb{Z}[X]$, $X = (x_1, \dots, x_n)$. We consider A as a subset of $\mathbb{Q}[X]$, so, if I is an ideal of A, then $\mathbb{Q}I$ is the ideal of $\mathbb{Q}[X]$ generated by I. Given a ring R, we denote the natural mapping $A \to A \otimes R = R[X]$ by ι_R. This note is devoted to the following algorithmic problem (see [1] for a definition and properties of Gröbner bases of ideals in polynomial rings over $\mathbb{Z}$).

Problem $(\mathcal{P})$**.** *Suppose that we have an infinite sequence $f_1, f_2, \dots$ of elements of A (a blackbox producing them one by one). Let $I = (f_1, f_2, \dots)$ be the ideal generated by all the f_i's. Compute the Gröbner base of I under the assumption that the Gröbner base of $\mathbb{Q}I$ and the Gröbner bases of $\iota_{\mathbb{Z}/m\mathbb{Z}}(I)$, $m \in \mathbb{Z}$, are known.*

This problem arises in [5], [6] (see §2 for more details). Our solution is based on a computation of the exponent of the torsion part of A/I (see [3]), and Main Lemma in §3 below.

§2. Motivation. More precisely, the algorithmic problem appeared in [5], [6] is the following. Let F be a finite rank free A-module, M_0 a submodule of F (given by a finite generating set G_0), $\tau \in \mathrm{Hom}_A(F, A)$, and $\rho_1, \dots, \rho_p \in \mathrm{End}_A(F)$. Compute (a Gröbner base of) $\tau(M)$ where M is the minimal submodule of F such that $M_0 \subset M$ and $\rho_i(M) \subset M$ for any $i = 1, \dots, p$. Theoretically, this problem has an evident solution, namely, we could compute one by one the Gröbner bases of the modules M_j defined recursively by $M_{j+1} = \sum_{i=1}^{p} \rho_i(M_j)$ till the moment when $M_{j+1} = M_j$. We set at this moment $M = M_j$ and $I = \tau(M)$.

However, if the rank of F is very big, then the computation of Gröbner bases of M_j's may be so long that it cannot be performed in practice whereas the Gröbner bases of $(\mathbb{Z}/m\mathbb{Z})[X]$-modules $M_j \otimes (\mathbb{Z}/m\mathbb{Z})$ can be computed much faster. When we know them, we can compute the Gröbner bases of $\mathbb{Q}[X]$-modules $M_j \otimes \mathbb{Q}$ (see [2] and §5 below). Therefore we can compute the modules $M \otimes (\mathbb{Z}/m\mathbb{Z})$ and $M \otimes \mathbb{Q}$, and hence the ideals $\iota_{\mathbb{Z}/m\mathbb{Z}}(I)$ and $\mathbb{Q}I$. Also we can successively compute the generators of I of the form $\tau\rho_{i_1}\rho_{i_2}\dots\rho_{i_k}(g)$, $g \in G_0$. So, the problem $(\mathcal{P})$ naturally arises.

§3. Solution to Problem $(\mathcal{P})$**.** Let the notation be as in Introduction.

Main Lemma. *Let I and J be ideals in A such that $J \subset I$. Let m be the exponent of the torsion part of the abelian group A/J, i. e., m is the minimal positive integer such that $m(\mathbb{Q}J \cap A) \subset J$. Let $m_1, \dots, m_t$ be pairwise coprime integers whose product is equal to m. Suppose that*

(i) $\mathbb{Q}J = \mathbb{Q}I$, *i. e., I and J generate the same ideal in $\mathbb{Q}[X]$,*
(ii) $\iota_{\mathbb{Z}/m_i\mathbb{Z}}(J) = \iota_{\mathbb{Z}/m_i\mathbb{Z}}(I)$, $i = 1, \dots, t$;

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

Then $I = J$.

Proof. Let $c_i = m/m_i = \prod_{j\neq i} m_j$, $i = 1, \dots, t$. Since m_i's are pairwise coprime, there exist $b_1, \dots, b_t \in \mathbb{Z}$ such that

$$b_1c_1 + \cdots + b_tc_t = 1 \tag{1}$$

We have to show that $I \subset J$. Let $f \in I$. By Condition (ii), for any $i = 1, \dots, t$, we have $f = f_i + m_ih_i$ with $f_i \in J$ and $h_i \in A$. By Condition (i), we have $kf \in J$ for some $k \in \mathbb{Z}$, hence, for any i, we have $km_ih_i = kf - kf_i \in J$, i. e., h_i represents an element of the torsion part of the group A/J. By definition of m, this implies $mh_i \in J$, hence $c_if = c_if_i + c_im_ih_i = c_if_i + mh_i \in J$. Combined with (1), this yields

$$f = b_1(c_1f) + \cdots + b_t(c_tf) \in J \quad \square$$

Thus we obtain the following solution to Problem $(\mathcal{P})$. We fix any term order in A. We try one by one all ideals J of the form $(f_1, \dots, f_k)$, $k = 1, 2, \dots$ and for each of them do the following:

(1) Compute the reduced Gröbner bases of the ideals $\mathbb{Q}I$ and $\mathbb{Q}J$ of the ring $\mathbb{Q}[X]$. If they do not coincide, then we pass to the next J.
(2) Compute m as in Main Lemma using one of the algorithms given in [3] (see also §4). Let $m = p_1^{\alpha_1} \dots p_t^{\alpha_t}$ be the factorization of m into prime factors and let $m_i = p_i^{\alpha_i}$.
(3) For each $i = 1, \dots, t$, compute the reduced Gröbner bases of $\iota_{\mathbb{Z}/m_i\mathbb{Z}}(I)$ and $\iota_{\mathbb{Z}/m_i\mathbb{Z}}(J)$. If they do not coincide, then we pass to the next J. If they do coincide, then we compute the Gröbner base of J and stop the computation.

§4. Computation of m. Several algorithms are discussed in [3] to compute m (defined in Main Lemma). For the reader's convenience, we reproduce here the algorithm which (we quote [3]) 'seems better suited for practical computations'. It consists in the following steps:

(1) Fix any term ordering and compute a Gröbner base G of J. Let s be the least common multiple of the leading coefficients of the elements of G. Then $\mathbb{Q}J \cap A = \mathbb{Z}[1/s]J \cap A$.
(2) Let Y be a new indeterminate. Then $\mathbb{Z}[1/s]J \cap A = (I, sY - 1) \cap A$. A Gröbner base G_s of this ideal can be computed using an elimination term ordering.
(3) Compute m such that $mg \in J$ for any $g \in G_s$.

§5. On the modular computation of Gröbner bases over $\mathbb{Q}$. In our notation, Elizabeth Arnold's theorem can be formulated as follows (similar theorem holds also for modules).

Theorem. [2; Th. 7.1]. *Let I be a* **homogeneous** *ideal in $\mathbb{Z}[X]$ and $G \subset \mathbb{Z}[X]$. Let $\mathbb{Q}I$ be the ideal in $\mathbb{Q}[X]$ generated by I. Let p be a prime number. We set $I_p = \iota_{\mathbb{F}_p}(I)$, $G_p = \iota_{\mathbb{F}_p}(G)$. Suppose that:*

(1) *G_p is a Gröbner base of I_p,*
(2) *G is a Gröbner base of the ideal $\langle G\rangle_{\mathbb{Q}}$ that it generates in $\mathbb{Q}[X]$,*
(3) *$\mathbb{Q}I \subset \langle G\rangle_{\mathbb{Q}}$,*
(4) *$\mathrm{LM}(G_p) = \mathrm{LM}(G)$ where "LM" denotes the set of leading monomials.*

Then G is a Gröbner base of $\mathbb{Q}I$.

It is claimed in [4; Th. 2.4] that the same statement holds for non-homogeneous ideals. However, it is wrong as one can see in the simplest example $I = (px + 1)$, $G = \{1\}$.

When the ideal is not homogeneous, it can be homogenized by adding a new variable and after that one can apply E. Arnold's theorem. For instance, if we homogenize the above counter-example, i. e., if we set $I = (px + y)$, $G = \{y\}$, then the condition (3) no longer holds true, so, the contradiction disappears.

Acknowledgment. I am grateful to Vladimir Gerdt for useful discussions.

References

1. W. W. Adams, P. Loustaunau, *An Introduction to Gröbner Bases*, Graduate Studies in Mathematics, vol. 3, A.M.S., Providence, RI, 1994.
2. E. A. Arnold, *Modular algorithms for computing Gröbner bases*, J. of Symbolic Computations **35** (2003), 403–419.
3. M. Aschenbrenner, *Algorithms for computing saturations of ideals in finitely generated commutative rings.* Appendix to: *Automorphisms mapping a point into a subvariety*, J. of Algebraic Geom. **20** (2011), 785–794 (by B. Poonen).
4. V. Idrees, G. Pfister, S. Steidel, *Parallelization of modular algorithms*, J. of Symbolic Computations **46** (2011), 672–684.
5. S. Yu. Orevkov, *Markov trace on the Funar algebra*, arXiv:1206.0765.
6. S. Yu. Orevkov, *Cubic Hecke algebras and invariants of transversal links*, Doklady Math. (to appear); arXiv:1307.5862.

Steklov Math. Institute, Moscow

IMT, Université Paul Sabatier (Toulouse-3)
E-mail address: `orevkov@math.ups-tlse.fr`