arXiv:1902.09534v1 [math.CV] 24 Feb 2019

ON GENERALIZED $p-$VALENT NON-BAZILEVIČ FUNCTIONS OF ORDER $\alpha + i\beta$

A.A. AMOURAH, A. G. ALAMOUSH, AND M. DARUS

Abstract. In this paper, we introduce a subclass $N_{p,\mu}^{n}(\alpha, \beta, A, B)$ of p-valent non-Bazilevič functions of order $\alpha + i\beta$. Some subordination relations and the inequality properties of $p-$valent functions are discussed. The results presented here generalize and improve some known results.

1. Introduction and preliminaries

Let $\mathcal{A}_p$ denote the class of functions f of the form:

$$f(z) = z^p + \sum_{k=n}^{\infty} a_{k+p} z^{k+p} \quad (p, n \in \mathbb{N} = \{1, 2, 3, ...\}), \tag{1.1}$$

which are *analytic* and $p-$*valent* in the open unit disc $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$. If $f(z)$ and $g(z)$ are analytic in $\mathbb{U}$, we say that $f(z)$ is *subordinate* to $g(z)$, and we write:

$$f \prec g \text{ in } \mathbb{U} \text{ or } f(z) \prec g(z),\ z \in \mathbb{U}, \tag{1.2}$$

if there exists a Schwarz function $w(z)$, which is analytic in $\mathbb{U}$ with

$$|w(0)| = 0 \text{ and } |w(z)| < 1,\ z \in \mathbb{U},$$

such that

$$f(z) = g(w(z)),\ z \in \mathbb{U}.$$

Furthermore, if the function $g(z)$ is *univalent* in $\mathbb{U}$, then we have the following equivalence, see Miller & Mocanu ([3], [4]),

$$f(z) \prec g(z)\ (z \in \mathbb{U}) \Leftrightarrow f(0) = g(0) \text{ and } f(\mathbb{U}) \subset g(\mathbb{U}).$$

We define a subclass of $\mathcal{A}_p$ as follows:

Definition 1.1. Let $N_{p,\mu}^{n}(\alpha, \beta, A, B)$ denote the class of functions $f(z) \in \mathcal{A}_p$ satisfying the inequality:

$$\left\{(1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\right\} \prec \frac{1+Az}{1+Bz},\ (z \in \mathbb{U}), \tag{1.3}$$

where $\mu \in \mathbb{C}$, $\alpha \geq 0$, $\beta \in \mathbb{R}$, $-1 \leq B \leq 1$, $A \neq B$, $p \in \mathbb{N}$ and $A \in \mathbb{R}$. All the powers in (1.3) are principal values.

2000 *Mathematics Subject Classification.* 30C45.

Key words and phrases. Analytic functions, non-Bazilevič functions, differential subordination.

We say that the function $f(z)$ in this class is p-valent non-Bazilevič functions of type $\alpha + i\beta$.

Definition 1.2. Let $f \in N_{p,\mu}^n(\alpha, \beta, \rho)$ if and only if $f(z) \in \mathcal{A}_p$ and it satisfies

$$\operatorname{Re}\left\{(1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\right\} > \rho, \ (z \in \mathbb{U}), \tag{1.4}$$

where $\mu \in \mathbb{C}$, $\alpha \geq 0$, $\beta \in \mathbb{R}$, $p \in \mathbb{N}$ and $0 \leq \rho < p$.

Special Cases:

(1) When $p = 1$, then $N_{1,\mu}^n(\alpha, \beta, A, B)$ is the class studied by AlAmoush and Darus [6].

(2) When $p = 1$, $\beta = 0$, then $N_{1,\mu}^n(\alpha, 0, A, B)$ is the class studied by Wang et al [1].

(3) When $p = 1$, $\beta = 0$, $\mu = -1$, $A = 1$ and $B = -1$, then $N_{\mu}^n(\alpha)$ is the class studied by Obradovic [10].

(4) If $p = 1$, $\beta = 0$, $\mu = B = -1$ and $A = 1 - 2\rho$, then the class $N_{1,-1}^n(\alpha, 0, 1-2\rho, -1)$ reduces to the class of non-Bazilevič functions of order ρ $(0 \leq \rho < 1)$. The Fekete-Szegö problem of the class $N_{1,-1}^n(\alpha, 0, 1-2\rho, -1)$ were considered by Tuneski and Darus [2].

We will need the following lemmas in the next section.

Lemma 1.3. [7] *Let the function $h(z)$ be analytic and convex in $\mathbb{U}$ with $h(0) = 1$. Suppose also that the function $\Phi(z)$ given by*

$$\Phi(z) = 1 + c_n z^n + c_{n+1} z^{n+1} + ...$$

is analytic in $\mathbb{U}$. If

$$\Phi(z) + \frac{1}{\gamma} z\Phi'(z) \prec h(z) \ (z \in \mathbb{U}, \operatorname{Re}\gamma \geq 0, \gamma \neq 0), \tag{1.5}$$

then

$$\Phi(z) \prec \Psi(z) = \frac{\gamma}{n} z^{-\gamma/n} \int_0^z t^{(\gamma/n)-1} h(t)dt \prec h(z) \ (z \in \mathbb{U}),$$

and $\Psi(z)$ is the best dominant for the differential subordination (1.5).

Lemma 1.4. [8] *Let $-1 \leq B_1 \leq B_2 < A_2 < A_1 \leq 1$, then*

$$\frac{1 + A_2 z}{1 + B_2 z} \prec \frac{1 + A_1 z}{1 + B_1 z}.$$

Lemma 1.5. [9] *Let $\Phi(z)$ be analytic and convex in $\mathbb{U}$, $f(z) \in \mathcal{A}_p$, $g(z) \in \mathcal{A}_p$.*

If $f(z) \prec \Phi(z)$, $g(z) \prec \Phi(z)$, $0 \leq \mu \leq 1$ then

$$\mu f(z) + (1-\mu) g(z) \prec \Phi(z).$$

Lemma 1.6. [11] *Let $q(z)$ be a convex univalent function in $\mathbb{U}$ and let $\sigma \in \mathbb{C}$, $\eta \in \mathbb{C}^* = \mathbb{C} \backslash \{0\}$ with*

$$\operatorname{Re}\left\{1 + \frac{zq''(z)}{q'(z)}\right\} > \max\left\{0, -\operatorname{Re}\left(\frac{\sigma}{\eta}\right)\right\}.$$

If the function $\Phi(z)$ is analytic in $\mathbb{U}$ and

$$\sigma\Phi(z) + \eta z\Phi^{'}(z) \prec \sigma q(z) + \eta z q^{'}(z),$$

then, $\Phi(z) \prec q(z)$ and $q(z)$ is the best dominant.

Lemma 1.7. [12] *Let $q(z)$ be a convex univalent in $\mathbb{U}$ and $\eta \in \mathbb{C}$. Further, assume that $\mathrm{Re}\{\overline{\eta}\} > 0$. If $\Phi(z) \in H[q(0), 1] \cap Q$, and $\Phi(z) + \eta z\Phi^{'}(z)$ is univalent in $\mathbb{U}$. Then*

$$q(z) + \eta z q^{'}(z) \prec \Phi(z) + \eta z\Phi^{'}(z),$$

signifies that $q(z) \prec \Phi(z)$ and $q(z)$ are the best subordinat.

We employ techniques similar to these used earlier by Yousef et al. [13], Amourah et al. ([14], [15]), AlAmoush and Darus [16] and Al-Hawary et al. [13].

In the present paper, we shall obtain results concerning the subordination relations and inequality properties of the class $N^{n}_{p,\mu}(\alpha, \beta, A, B)$. The results obtained generalize the related works of some authors.

2. Main Result

Theorem 2.1. *Let $\mu \in \mathbb{C}$, $\alpha \geq 0$, $\beta \in \mathbb{R}$, $-1 \leq B \leq 1$, $A \neq B$, $\alpha + i\beta \neq 0$, $p \in \mathbb{N}$ and $A \in \mathbb{R}$. If $f(z) \in N^{n}_{p,\mu}(\alpha, \beta, A, B)$, Then*

$$\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \prec \frac{p(\alpha+i\beta)}{\mu n}\int_0^1 \frac{1+Azu}{1+Bzu}u^{\frac{p(\alpha+i\beta)}{\mu n}-1}du \prec \frac{1+Az}{1+Bz}. \tag{2.1}$$

Proof. Let

$$\Phi(z) = \left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}. \tag{2.2}$$

Then $\Phi(z)$ is analytic in $\mathbb{U}$ with $\Phi(0) = 1$. Taking logarithmic differentiation of (2.2) in both sides, we obtain

$$p(\alpha+i\beta) - (\alpha+i\beta)\frac{zf(z)^{'}}{f(z)} = \frac{z\Phi^{'}(z)}{\Phi(z)}.$$

In the above equation, we have

$$1 - \frac{zf'(z)}{pf(z)} = \frac{1}{p(\alpha+i\beta)}\frac{z\Phi^{'}(z)}{\Phi(z)}.$$

From this we can easily deduce that

$$\left\{(1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\right\}. \tag{2.3}$$

On a class of $p-$valent non-Bazilevič functions

$$= \Phi(z) + \frac{\mu z\Phi^{'}(z)}{p(\alpha+i\beta)} \prec \frac{1+Az}{1+Bz}. \tag{2.4}$$

Now, by Lemma 1.3 for $\gamma = \frac{p(\alpha+i\beta)}{\mu}$, we deduce that

$$\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \prec q(z) = \frac{p(\alpha+i\beta)}{\mu n} z^{-\frac{p(\alpha+i\beta)}{\mu n}} \int_0^z t^{\frac{p(\alpha+i\beta)}{\mu n}-1} (\frac{1+At}{1+Bt}) dt.$$

Putting $t = zu \Rightarrow dt = zdu$. Then we have the above equation with

$$= \frac{p(\alpha+i\beta)}{\mu n} \int_0^1 \frac{1+Azu}{1+Bzu} u^{\frac{p(\alpha+i\beta)}{\mu n}-1} du \prec \frac{1+Az}{1+Bz},$$

and the proof is complete. □

Corollary 2.2. *Let $\mu \in \mathbb{C}$, $\alpha \geq 0$, $\beta \in \mathbb{R}$, $\alpha + i\beta \neq 0$, $p \in \mathbb{N}$ and $\rho \neq 0$. If $f(z) \in \mathcal{A}_p$ satisfies*

$$(1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \prec \frac{1+(1-2\rho)z}{1-z}, \ (z \in \mathbb{U}),$$

then

$$\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \prec \frac{p(\alpha+i\beta)}{\mu n} \int_0^1 \frac{1+(1-2\rho)zu}{1-zu} u^{\frac{p(\alpha+i\beta)}{\mu n}-1} du, \ (z \in \mathbb{U}),$$

or equivalent to

$$\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \prec \rho + \frac{p(\alpha+i\beta)(1-\rho)}{\mu n} \int_0^1 \frac{1+zu}{1-zu} u^{\frac{p(\alpha+i\beta)}{\mu n}-1} du, \ (z \in \mathbb{U}).$$

Corollary 2.3. *Let $\mu \in \mathbb{C}$, $\alpha \geq 0$, $\beta \in \mathbb{R}$, $\alpha + i\beta \neq 0$, $p \in \mathbb{N}$ and $\operatorname{Re}\{\mu\} \geq 0$, then*

$$N^n_{p,\mu}(\alpha,\beta,A,B) \subset N^n_{p,0}(\alpha,\beta,A,B).$$

Theorem 2.4. *Let $0 \leq \mu_1 \leq \mu_2$, $\alpha \geq 0$, $\beta \in \mathbb{R}$, $p \in \mathbb{N}$, $\alpha + i\beta \neq 0$ and $-1 \leq B_1 \leq B_2 < A_2 \leq A_1 \leq 1$, then*

$$N^n_{p,\mu_2}(\alpha,\beta,A_2,B_2) \subset N^n_{p,\mu_1}(\alpha,\beta,A_1,B_1). \tag{2.5}$$

Proof. Suppose that $f(z) \in N^n_{p,\mu_2}(\alpha,\beta,A_2,B_2)$ we have $f(z) \in \mathcal{A}_p$ and

$$\left\{(1+\mu_2)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu_2\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\right\} \prec \frac{1+A_2z}{1+B_2z}, \ (z \in \mathbb{U}).$$

Since $-1 \leq B_1 \leq B_2 < A_2 \leq A_1 \leq 1$, therefore it follows from Lemma 1.4 that

$$\left\{(1+\mu_2)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu_2\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\right\} \prec \frac{1+A_1z}{1+B_1z}, \ (z \in \mathbb{U}), \tag{2.6}$$

that is $f(z) \in N^n_{p,\mu_2}(\alpha,\beta,A_1,B_1)$. So Theorem 2.4 is proved when $\mu_1 = \mu_2 \geq 0$.

When $\mu_2 > \mu_1 \geq 0$, then we can see from Corollary 2.3 that $f(z) \in N^n_{p,0}(\alpha,\beta,A_1,B_1)$, then

$$\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \prec \frac{1+A_1z}{1+B_1z}. \tag{2.7}$$

But

$$\left\{(1+\mu_1)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}-\mu_1\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\right\}$$
$$=(1-\frac{\mu_1}{\mu_2})\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}+\frac{\mu_1}{\mu_2}\left\{\begin{array}{c}(1+\mu_1)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\\ -\mu_1\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\end{array}\right\}.$$

It is obvious that $\frac{1+A_1z}{1+B_1z}$ is analytic and convex in $\mathbb{U}$. Sowe obtain fromLemma 1.5 and differential subordinations (2.6) and (2.7) that

$$\left\{(1+\mu_1)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}-\mu_1\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\right\}\frac{1+A_1z}{1+B_1z},$$

that is, $f(z)\in N^n_{p,\mu_1}(\alpha,\beta,A_1,B_1)$. Thuswe have

$$N^n_{p,\mu_2}(\alpha,\beta,A_2,B_2)\subset N^n_{p,\mu_1}(\alpha,\beta,A_1,B_1).$$

□

Corollary 2.5. *Let* $0\le\mu_1\le\mu_2$, $0\le\rho_1\le\rho_2$, $\alpha\ge0$, $\beta\in\mathbb{R}$, $p\in\mathbb{N}$ *and* $\alpha+i\beta\neq0$ *then*

$$N^n_{p,\mu_2}(\alpha,\beta,\rho_2)\subset N^n_{p,\mu_1}(\alpha,\beta,\rho_1).$$

Theorem 2.6. *Let* $\mu\in\mathbb{C}$, $\alpha\ge0$, $\beta\in\mathbb{R}$, $\mu+i\beta\neq0$, $p\in\mathbb{N}$, $-1\le B\le1$, $A\neq B$ *and* $A\in\mathbb{R}$. *If* $f(z)\in N^n_{p,\mu}(\alpha,\beta,A,B)$, *then*

$$\inf_{z\in\mathbb{U}}\operatorname{Re}\left\{\frac{p(\alpha+i\beta)}{\mu n}\int_0^1\frac{1+Azu}{1+Bzu}u^{\frac{p(\alpha+i\beta)}{\mu n}-1}du\right\}$$
$$<\operatorname{Re}\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}<\sup_{z\in\mathbb{U}}\operatorname{Re}\left\{\frac{p(\alpha+i\beta)}{\mu n}\int_0^1\frac{1+Azu}{1+Bzu}u^{\frac{p(\alpha+i\beta)}{\mu n}-1}du\right\}.$$

Proof. Suppose that $f(z)\in N^n_{p,\mu}(\alpha,\beta,A,B)$, then from Theorem 2.1 we know that

$$\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\prec\frac{p(\alpha+i\beta)}{\mu n}\int_0^1\frac{1+Azu}{1+Bzu}u^{\frac{p(\alpha+i\beta)}{\mu n}-1}du. \tag{2.8}$$

Therefore, from the definition of the subordination, we have

$$\operatorname{Re}\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}>\inf_{z\in\mathbb{U}}\operatorname{Re}\left\{\frac{p(\alpha+i\beta)}{\mu n}\int_0^1\frac{1+Azu}{1+Bzu}u^{\frac{p(\alpha+i\beta)}{\mu n}-1}du\right\},$$

$$\operatorname{Re}\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}<\sup_{z\in\mathbb{U}}\operatorname{Re}\left\{\frac{p(\alpha+i\beta)}{\mu n}\int_0^1\frac{1+Azu}{1+Bzu}u^{\frac{p(\alpha+i\beta)}{\mu n}-1}du\right\}.$$

□

Corollary 2.7. *Let $\mu \in \mathbb{C}$, $\alpha \geq 0$, $\beta \in \mathbb{R}$, $\mu + i\beta \neq 0$, $p \in \mathbb{N}$ and $\rho < 1$. If $f(z) \in N^n_{p,\mu}(\alpha, \beta, 1-2\rho, -1)$, then*

$$\begin{aligned}
&\rho + (1-\rho) \inf_{z\in\mathbb{U}} \operatorname{Re}\left\{ \frac{p(\alpha+i\beta)}{\mu n} \int_0^1 \frac{1+zu}{1-zu} u^{\frac{p(\alpha+i\beta)}{\mu n}-1} du \right\} \\
&< \operatorname{Re}\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \\
&< \rho + (1-\rho) \sup_{z\in\mathbb{U}} \operatorname{Re}\left\{ \frac{p(\alpha+i\beta)}{\mu n} \int_0^1 \frac{1+zu}{1-zu} u^{\frac{p(\alpha+i\beta)}{\mu n}-1} du \right\}. \qquad (2.9)
\end{aligned}$$

Corollary 2.8. *Let $\mu \in \mathbb{C}$, $\alpha \geq 0$, $\beta \in \mathbb{R}$, $\mu + i\beta \neq 0$, $p \in \mathbb{N}$ and $\rho > 1$. If $f(z) \in N^n_{p,\mu}(\alpha, \beta, 1-2\rho, -1)$, then*

$$\operatorname{Re}\left\{ (1+\mu_1)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu_1 \left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \right\} < \rho, \ z \in \mathbb{U},$$

then

$$\begin{aligned}
&\rho + (1-\rho) \sup_{z\in\mathbb{U}} \operatorname{Re}\left\{ \frac{p(\alpha+i\beta)}{\mu n} \int_0^1 \frac{1+zu}{1-zu} u^{\frac{p(\alpha+i\beta)}{\mu n}-1} du \right\} \\
&< \operatorname{Re}\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \\
&< \rho + (1-\rho) \inf_{z\in\mathbb{U}} \operatorname{Re}\left\{ \frac{p(\alpha+i\beta)}{\mu n} \int_0^1 \frac{1+zu}{1-zu} u^{\frac{p(\alpha+i\beta)}{\mu n}-1} du \right\}.
\end{aligned}$$

Next, several new differential suborination results of non-Bazilevič class of order $\alpha + i\beta$ are established.

3. Suborination for $p-$Valent Non-Bazilevič Class of Order $\alpha + i\beta$

By employing lemma 1.6, the following result is given.

Theorem 3.1. *Let q be univalent in $\mathbb{U}$, $\mu \in \mathbb{C}^*$, $\alpha \geq 0$, $\beta \in \mathbb{R}$ and $\alpha + i\beta \neq 0$. Suppose that q satisfies*

$$\operatorname{Re}\left\{ 1 + \frac{zq''(z)}{q'(z)} \right\} > \max\left\{ 0, -\operatorname{Re}\left\{ \frac{p(\alpha+i\beta)}{\mu} \right\} \right\}. \qquad (3.1)$$

If $f \in \mathcal{A}_p$ satisfies the subordination

$$\left\{ (1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu \left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \right\} \prec q(z) + \mu \frac{zq'(z)}{p(\alpha+i\beta)}, \qquad (3.2)$$

then

$$\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \prec q(z) \qquad (3.3)$$

and $q(z)$ is the best dominant.

Proof. Define the function $\Phi(z)$ by

$$\Phi(z) = \left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}. \tag{3.4}$$

Having (3.4) differentiated logarithmically in connection with z, we have

$$\frac{z\Phi^{'}(z)}{\Phi(z)} = p(\alpha+i\beta)\left(1-\frac{zf(z)^{'}}{pf(z)}\right), \tag{3.5}$$

which, with respect to hypothesis (3.2) of Theorem 3.1, the following subordination is obtained:

$$\Phi(z) + \frac{\mu z\Phi^{'}(z)}{p(\alpha+i\beta)} \prec q(z) + \mu\frac{zq^{'}(z)}{p(\alpha+i\beta)}. \tag{3.6}$$

Theorem 3.1 assertion is now followed by the use of Lemma 1.6 with $\eta = \frac{\mu}{p(\alpha+i\beta)}$ and $\sigma = 1$. □

Remark 3.2. For the choice $q(z) = \frac{1+Az}{1+Bz}$ in Theorem 3.1, the following the corollary is obtained.

Corollary 3.3. *Let $\mu \in \mathbb{C}^*$, $-1 \leq B < A \leq 1$, and*

$$\mathrm{Re}\left\{\frac{1-Bz}{1+Bz}\right\} > \max\left\{0, -\mathrm{Re}\left\{\frac{p(\alpha+i\beta)}{\mu}\right\}\right\}, \ (z \in \mathbb{U}). \tag{3.7}$$

If $f \in \mathcal{A}_p$, and

$$\left\{(1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\right\} \prec \frac{\mu(A-B)z}{p(\alpha+i\beta)(1+Bz)^2} + \frac{1+Az}{1+Bz}, \tag{3.8}$$

then

$$\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \prec \frac{1+Az}{1+Bz} \tag{3.9}$$

and $\frac{1+Az}{1+Bz}$ is the best dominant.

Remark 3.4. For the choice $q(z) = \frac{1+z}{1-z}$ in Theorem 3.1, the following the corollary is obtained.

Corollary 3.5. *Let $\mu \in \mathbb{C}^*$, and*

$$\mathrm{Re}\left\{\frac{1+z}{1-z}\right\} > \max\left\{0, -\mathrm{Re}\left\{\frac{p(\alpha+i\beta)}{\mu}\right\}\right\}, \ (z \in \mathbb{U}). \tag{3.10}$$

If $f \in \mathcal{A}_p$, and

$$\left\{(1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\right\} \prec \frac{2\mu z}{p(\alpha+i\beta)(1-z)^2} + \frac{1+z}{1-z}, \tag{3.11}$$

then

$$\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \prec \frac{1+z}{1-z} \tag{3.12}$$

and $\frac{1+Az}{1+Bz}$ is the best dominant.

4. Superordination for $p-$Valent Non-Bazilevič Class of Order $\alpha+i\beta$

Theorem 4.1. *Let q be convex univalent in $\mathbb{U}$, $\mu \in \mathbb{C}$, $\alpha \geq 0$, $\beta \in \mathbb{R}$ and $\alpha + i\beta \neq 0$. Suppose that q satisfies*

$$\operatorname{Re}\{\mu\} > 0 \tag{4.1}$$

and $\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \in H[q(0),1] \cap Q$. Let

$$(1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \tag{4.2}$$

be univalent in $\mathbb{U}$, If

$$q(z) + \mu\frac{zq'(z)}{p(\alpha+i\beta)} \prec (1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}, \tag{4.3}$$

then

$$q(z) \prec \left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \tag{4.4}$$

and $q(z)$ is the best subordinant.

Proof. Define the function $\Phi(z)$ by

$$\Phi(z) = \left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}. \tag{4.5}$$

Then based on a computation, it indicates that

$$\Phi(z) + \frac{\mu z\Phi'(z)}{p(\alpha+i\beta)} \prec (1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}. \tag{4.6}$$

Theorem 4.1 follows from Lemma 1.7. □

Remark 4.2. Taking $q(z) = \frac{1+Az}{1+Bz}$ in Theorem 4.1, the following the corollary is obtained.

Corollary 4.3. *Let $-1 \leq B < A \leq 1$. Let q be convex univalent in $\mathbb{U}$. Suppose that q satisfies $\operatorname{Re}\{\mu\} > 0$, and $\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \in H[q(0),1] \cap Q$. Let*

$$(1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \tag{4.7}$$

be univalent in $\mathbb{U}$, If

$$\frac{\mu(A-B)z}{p(\alpha+i\beta)(1+Bz)^2} + \frac{1+Az}{1+Bz} \prec \left\{(1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}\right\}, \tag{4.8}$$

then

$$\frac{1+Az}{1+Bz} \prec \left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \tag{4.9}$$

and $\frac{1+Az}{1+Bz}$ is the best subordinant.

5. Sandwich Results for $p-$Valent Non-Bazilevič Class of Order $\alpha + i\beta$

Combining the differential subordination and supordination results, the sandwich results are highlighted as follows.

Theorem 5.1. *Let q_1 be convex univalent and let q_2 be univalent in $\mathbb{U}$, $\mu \in \mathbb{C}$, $\alpha \geq 0$, $\beta \in \mathbb{R}$ and $\alpha + i\beta \neq 0$. Suppose q_1 satisfies (4.1) and q_2 satisfies (3.1). If $0 \neq \left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \in H[q(0), 1] \cap Q$, $(1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta}$ is univalent in $\mathbb{U}$, and if $f \in \mathcal{A}_p$ satisfies*

$$q_1(z) + \mu\frac{zq_1'(z)}{p(\alpha+i\beta)} \prec (1+\mu)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} - \mu\left(\frac{zf'(z)}{pf(z)}\right)\left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \prec q_2(z) + \mu\frac{zq_2'(z)}{p(\alpha+i\beta)}, \tag{5.1}$$

then

$$q_1(z) \prec \left(\frac{z^p}{f(z)}\right)^{\alpha+i\beta} \prec q_2(z) \tag{5.2}$$

and $q_1(z)$ and $q_2(z)$ are respectively, the best subordinant and best dominant.

Proof. Simultaneously applying the techniques of the proof of Theorem 3.1 and Theorem 4.1. □

Acknowledgement: The work here is supported by FRGSTOPDOWN/2013/ST06/UKM/01/1.

References

[1] Z. Wang, C. Gao & M. Liao, *On certain generalized class of non-Bazilevic functions*, Acta Mathematica. Academiae Paedagogicae Nyıregyháziensis. New Series 21(2005), 147-154.

[2] N. Tuneski & M. Darus, *Fekete-Szegö functional for non-Bazilevic functions*, Acta Mathematica. Academiae Paedagogicae Nyıregyháziensis. New Series 18(2002), 63-65.

[3] S. S. Miller & P. T. Mocanu, *Differential subordinations and univalent functions*, The Michigan Mathematical Journal 28. 2(1981), 157-172.

[4] S. S. Miller & P. T. Mocanu, *Differential Subordination: Theory and Applications, Ser,* Monogr. Textbooks Pure Appl. Math 225(2000).

[5] J. L. Liu & K. I. Noor, *Some properties of Noor integral operator*, Journal of Natural Geometry 21. 1/2(2002), 81-90.

[6] A. G. Alamoush & M. Darus, *On Certain Class of Non-Bazilevič Functions of Order $\alpha + i\beta$ Defined by a Differential Subordination*, International Journal of Differential Equations 2014(2014).

[7] S. S. Miller & P. T. Mocanu, *Differential subordinations and univalent functions*, The Michigan Mathematical Journal 28. 2(1981), 157-172.

[8] L. Mingsheng, *On a Subclass of $p-$valent Close-to-convex Functions of Order β and Type α*, Journal of Mathematical Study, 30. 1(1997), 102–104.

[9] L. Mingsheng, *On certain class of analytic functions defined by differential subordination*, Acta Mathematica Scientia 22. 3(2002), 388-392.

[10] M. Obradovic, *A class of univalent functions*, Hokkaido Mathematical Journal 27. 2(1998), 329-335.

[11] T. N. Shanmugam, V. Ravichandran & S. Sivasubramanian, *Differential sandwich theorems for some subclasses of analytic functions*, J. Austr. Math. Anal. Appl 3. 1(2006), 1-11.

[12] S. S. Miller & P. T. Mocanu, *Subordinants of differential superordinations*, Complex variables 48 .10(2003), 815-826.

[13] F. Yousef, A. A. Amourah, and M. Darus, *Differential sandwich theorems for p-valent functions associated with a certain generalized differential operator and integral operator*, Italian Journal of Pure and Applied Mathematics 36 (2016), 543-556.

[14] A. A. Amourah, F. Yousef, T. Al-Hawary and M. Darus, *A certain fractional derivative operator for p-valent functions and new class of analytic functions with negative coefficients*, Far East Journal of Mathematical Sciences, 99. (1)(2016), 75-87.
[15] A. A. Amourah, F. Yousef, T. Al-Hawary and M. Darus, *On a Class of p- Valent non-Bazilevič Functions of Order* $\mu + i\beta$, International Journal of Mathematical Analysis, 10. (15)(2016), 701-710.
[16] A. G. Alamoush and M. Darus, *Subordination results for a certain subclass of non-bazilevic analytic functions defined by linear operator*, italian journal of pure and applied mathematics, 34 (2015), 375-388.
[17] T. Al-Hawary, A. A. Amourah, and M. Darus, *Differential sandwich theorems for p-valent functions associated with two generalized differential operator and integral operator*, International Information Institute (Tokyo). Information, 17. (8)(2014), 3559.

A.A. AMOURAH: Department of Mathematics, Faculty of Science and Technology, Irbid National University, Irbid, Jordan.

E-mail address: alaammour@yahoo.com.

A. G. AlAmoush: School of Mathematical Sciences Faculty of Science and Technology Universiti Kebangsaan Malaysia Bangi 43600 Selangor D. Ehsan, Malaysia.

E-mail address: adnan-omoush@yahoo.com.

M. DARUS : School of Mathematical Sciences Faculty of Science and Technology Universiti Kebangsaan Malaysia Bangi 43600 Selangor D. Ehsan, Malaysia.

E-mail address: maslina@ukm.edu.my.