arXiv:1906.06771v2 [math.RA] 16 Aug 2019

3-LIE BIALGEBRAS AND 3-PRE-LIE ALGEBRAS INDUCED BY INVOLUTIVE DERIVATIONS

RUIPU BAI, SHUAI HOU, AND CHUANGCHUANG KANG

Abstract. In this paper, we study the structure of 3-Lie algebras with involutive derivations. We prove that if A is an m-dimensional 3-Lie algebra with an involutive derivation D, then there exists a compatible 3-pre-Lie algebra $(A, \{,,\}_D)$ such that A is the sub-adjacent 3-Lie algebra, and there is a local cocycle 3-Lie bialgebraic structure on the $2m$-dimensional semi-direct product 3-Lie algebra $A \ltimes_{ad^*} A^*$, which is associated to the adjoint representation (A, ad). By means of involutive derivations, the skew-symmetric solution of the 3-Lie classical Yang-Baxter equation in the 3-Lie algebra $A \ltimes_{ad^*} A^*$, a class of 3-pre-Lie algebras, and eight and ten dimensional local cocycle 3-Lie bialgebras are constructed.

[1]

1. Introduction

In recent years, quantum 3-Lie algebras [1], 3-Lie bialgebras [2, 3], local cocycle 3-Lie bialgebras, 3-pre Lie algebras and 3-Lie classical Yang-Baxter equation [4] are provided. It is well known that Lie bialgebra is the algebraic structure corresponding to a Poisson-Lie group and the classical structure of a quantized universal enveloping algebra [6, 7], and it has a coboundary theory, which leads to the construction of Lie bialgebras from solutions of the classical Yang-Baxter equation. For a 3-Lie algebra A, and $r = \sum\limits_i x_i \otimes y_i \in A \otimes A$, the equation

$$[[r, r, r]] = 0$$

is called the 3-Lie classical Yang-Baxter equation (3-Lie CYBE) [4], where

$$\begin{aligned}[[r, r, r]] &:= [r_{12}, r_{13}, r_{14}] + [r_{12}, r_{23}, r_{24}] + [r_{13}, r_{23}, r_{34}] + [r_{14}, r_{24}, r_{34}] \\ &= \sum_{i,j,k} \big([x_i, x_j, x_k] \otimes y_i \otimes y_j \otimes y_k + x_i \otimes [y_i, x_j, x_k] \otimes y_j \otimes y_k \\ &\quad + x_i \otimes x_j \otimes [y_i, y_j, x_k] \otimes y_k + x_i \otimes x_j \otimes x_k \otimes [y_i, y_j, y_k]\big).\end{aligned}$$

But from the solutions of 3-Lie classical Yang-Baxter equation, the 3-Lie bialgebra which was introduced in [2], can not be constructed. For excavating the ideal connections between the solutions of 3-Lie classical Yang-Baxter equation and 3-Lie bialgebraic structures, authors in [4], introduced a new 3-Lie coproduct $\triangle$ on a 3-Lie algebra, and the pair $(A, \triangle)$ is called a local cocycle 3-Lie bialgebra, where $\triangle = \triangle_1 + \triangle_2 + \triangle_3 \colon A \to A^{\wedge 3}$ satisfies that the dual $(A^*, \triangle^*)$ is a 3-Lie algebra, and

Δ_1 is a 1-cocycle associated to the representation $(A \otimes A \otimes A, ad \otimes 1 \otimes 1)$;

Δ_2 is a 1-cocycle associated to the representation $(A \otimes A \otimes A, 1 \otimes ad \otimes 1)$;

2010 *Mathematics Subject Classification.* 17B05, 17D99.

Key words and phrases. 3-Lie algebra; involutive derivation; local cocycle 3-Lie bialgebra; 3-Lie Yang-Baxter equation.

[1] Corresponding author: Ruipu Bai, E-mail: bairuipu@hbu.edu.cn.

Δ_3 is a 1-cocycle associated to the representation $(A \otimes A \otimes A, 1 \otimes 1 \otimes ad)$.

For a skew-symmetric $r = \sum\limits_i x_i \otimes y_i \in A \otimes A$, if r is a solution of the 3-Lie classical Yang-Baxter equation, then $(A, \triangle)$ is a local cocycle 3-Lie bialgebra, where for all $x \in A$,

$$\triangle_1(x) = \sum_{ij} [x, x_i, x_j] \otimes y_j \otimes y_i; \triangle_2(x) = \sum_{ij} y_i \otimes [x, x_i, x_j] \otimes y_j; \triangle_3(x) = \sum_{ij} y_j \otimes y_i \otimes [x, x_i, x_j].$$

The question that how to get solutions of 3-Lie classical Yang-Baxter equation and how to construct local cocycle 3-Lie bialgebras is a hard task. Motivated by the object, we study the structure of involutive derivations on 3-Lie algebras. And in terms of involutive derivations, solutions of 3-Lie classical Yang-Baxter equation, a class of eight and ten dimensional local cocycle 3-Lie bialgebras, and a class of 3-pre-Lie algebras are constructed.

In the following we assume that all algebras are over an algebraically closed field F with characteristic zero, if $\{x_1, \cdots, x_m\}$ is a basis of A, then $\{x_1^*, \cdots, x_m^*\}$ is the basis of the dual space A^*, where $\langle x_i^*, x_j \rangle = \delta_{ij}$, $1 \leq i, j \leq m$, and Z is the set of integers.

2. Involutive derivations and compatible 3-pre Lie algebras

A **3-Lie algebra** $(A, [,,])$ over a field F is a vector space A with a linear multiplication $[,,] : A^{\wedge 3} \to A$ satisfying that for all $x_1, x_2, x_3, x_4, x_5 \in A$,

$$[x_1, x_2, [x_3, x_4, x_5]] = [[x_1, x_2, x_3], x_4, x_5] + [x_3, [x_1, x_2, x_4], x_5] + [x_3, x_4, [x_1, x_2, x_5]],$$

which is usually called *the generalized Jacobi identity*, or *Filippov identity*.

The subalgebra generated by $[x_1, \cdots, x_n], \forall x_1, \cdots, x_n \in A$, is called the derived algebra of A, and is denoted by A^1.

A derivation D is an endomorphism of A satisfying,

$$D[x_1, x_2, x_3] = [Dx_1, x_2, x_3] + [x_1, Dx_2, x_3] + [x_1, x_2, Dx_3], \forall x_1, x_2, x_3 \in A. \tag{1}$$

Further, if D satisfies $D^2 = I_d$ (identity), then D is called an **involutive derivation** on A, and A has the decomposition

$$A = A_1 \dot{+} A_{-1}, \tag{2}$$

where $A_1 = \{v \in A \mid Dv = v\}$ and $A_{-1} = \{v \in A \mid Dv = -v\}$ which are abelian subalgebras.

$Der(A)$ denotes the derivation algebra of A. For all $x_1, x_2 \in A$, the left multiplications $ad(x_1, x_2)$: $A \to A$, $ad(x_1, x_2)(x) = [x_1, x_2, x]$, $\forall x \in A$, are derivations which are called inner derivations, and $ad(A)$ denotes all the inner derivations.

A representation of A (or **an** A**-module**) is a pair (V, ρ), where V is a vector space, $\rho : A \wedge A \to End(V)$ is a linear mapping such that for all $x_i \in A, 1 \leq i \leq 4$,

$$\begin{aligned}[\rho(x_1, x_2), \rho(x_3, x_4)] &= \rho(x_1, x_2)\rho(x_3, x_4) - \rho(x_3, x_4)\rho(x_1, x_2) \\ &= \rho([x_1, x_2, x_3], x_4) - \rho([x_1, x_2, x_4], x_3), \\ \rho([x_1, x_2, x_3], x_4) &= \rho(x_1, x_2)\rho(x_3, x_4) + \rho(x_2, x_3)\rho(x_1, x_4) + \rho(x_3, x_1)\rho(x_2, x_4).\end{aligned}$$

A linear mapping $T : V \to A$ is called **an** $\wp$**-operator** associated to an A-module (V, ρ) if T satisfies

$$[Tu, Tv, Tw] = T(\rho(Tu, Tv)w + \rho(Tv, Tw)u + \rho(Tw, Tu)v), \ \forall u, v, w \in V. \tag{3}$$

(A, ad) is called **the adjoint representation** of A. The dual representation of (A, ad) is denoted by (A^*, ad^*), where A^* is the dual space of A, and $ad^* : A^{\wedge^2} \to EndA^*$, for all $x_1, x_2, x \in A$

and $\xi \in A^*$,

$$\langle ad^*(x_1, x_2)\xi, x\rangle = -\langle \xi, ad(x_1, x_2)x\rangle. \tag{4}$$

There is an equivalent description of an A-module (V, ρ), that is, $(B, [,,]_\rho)$ is a 3-Lie algebra, where $B = A\dot{+}V$, for all $x_1, x_2, x_3 \in A, v \in V$,

$$[x_1, x_2, x_3]_\rho = [x_1, x_2, x_3],\ [x_1, x_2, v]_\rho = \rho(x_1, x_2)v,\ [A, V, V] = [V, V, V] = 0.$$

For any 3-Lie algebra A, we have **the semi-direct product 3-Lie algebra** (B, μ), where $B = A\dot{+}A^*$, $[,,]_* : (A\dot{+}A^*)^{\wedge 3} \to A\dot{+}A^*$, for all $x_i \in A, \xi_i \in A^*, 1 \le i \le 3$,

$$\mu(x_1 + \xi_1, x_2 + \xi_2, x_3 + \xi_3) = [x_1, x_2, x_3] + ad^*(x_1, x_2)\xi_3 + ad^*(x_2, x_3)\xi_1 + ad^*(x_3, x_1)\xi_2, \tag{5}$$

which is denoted by $A \ltimes_{ad^*} A^*$.

Theorem 2.1. *Let $(A, [\cdot,\cdot,\cdot])$ be a 3-Lie algebra with an involutive derivation D. Then D is an $\wp$-operator of A associated to the adjoint representation (A, ad), and D satisfies*

$$[Dx, Dy, Dz] = D([Dx, Dy, z] + [Dy, Dz, x] + [Dz, Dx, y]),\ \forall x, y, z \in A. \tag{6}$$

Proof. By Eq (1), for all $x, y, z \in A$,

$D(ad(Dx, Dy)z + ad(Dy, Dz)x + ad(Dz, Dx)y)$

$= D([Dx, Dy, z] + [Dy, Dz, x] + [Dz, Dx, y])$

$= D([Dx, Dy, D^2z] + [D^2x, Dy, Dz] + [Dx, D^2y, Dzy]) = [Dx, Dy, Dz].$

The proof is completed. □

A 3-pre-Lie algebra $(A, \{,,\})$ [4] is a vector space A with a 3-ary linear mapping $\{\cdot,\cdot,\cdot\}: A \otimes A \otimes A \to A$, satisfying $\forall x_1, x_2, x_3 \in A$,

$$\{x_1, x_2, x_3\} = -\{x_2, x_1, x_3\}, \tag{7}$$

$$\begin{aligned}\{x_1, x_2, \{x_3, x_4, x_5\}\} = &\{\{x_1, x_2, x_3\}_c, x_4, x_5\} + \{x_3, \{x_1, x_2, x_4\}_c, x_5\}\\ &+\{x_3, x_4, \{x_1, x_2, x_5\}\},\end{aligned} \tag{8}$$

$$\begin{aligned}\{\{x_1, x_2, x_3\}_c, x_4, x_5\} = &\{x_1, x_2, \{x_3, x_4, x_5\}\} + \{x_2, x_3, \{x_1, x_4, x_5\}\}\\ &+\{x_3, x_1, \{x_2, x_4, x_5\}\},\end{aligned} \tag{9}$$

$$\{x_1, x_2, x_3\}_c = \{x_1, x_2, x_3\} + \{x_2, x_3, x_1\} + \{x_3, x_1, x_2\}. \tag{10}$$

Therefore, if $(A, \{,,\})$ is a 3-pre-Lie algebra, then $(A, \{,,\}_c)$ is a 3-Lie algebra, which is called **the sub-adjacent 3-Lie algebra of the 3-pre-Lie algebra** $(A, \{,,\})$, and $(A, \{,,\})$ is called **the compatible 3-pre-Lie algebra of the 3-Lie algebra** $(A, \{,,\}_c)$.

Theorem 2.2. *Let $(A, [\cdot,\cdot,\cdot])$ be a 3-Lie algebra, $D \in Der(A)$ be an involutive derivation. Then $(A, \{\cdot,\cdot,\cdot\}_D)$ is a 3-pre-Lie algebra, where*

$$\{x, y, z\}_D = [Dx, Dy, z]. \tag{11}$$

Furthermore,

$$\{x,y,z\}_D = \begin{cases} 0,\ x,y,z \in A_1, \text{or } x,y,z \in A_{-1}, \\ [x,y,z],\ x,y \in A_1, z \in A_{-1}, \\ -[x,y,z],\ x \in A_1, y \in A_{-1}, z \in A_{-1}, \\ [x,y,z],\ x,y \in A_{-1}, z \in A_1, \\ -[x,y,z],\ x \in A_1, y \in A_{-1}, z \in A_1. \end{cases} \tag{12}$$

And $(A, \{\cdot,\cdot,\cdot\}_D)$ *is called* **the 3-pre-Lie algebra associated with the involutive derivation** D.

Proof. By Theorem 2.1, D is an $\wp$-operator associate to the adjoint representation (A, ad), and for all $x_i \in A, 1 \leq i \leq 5$,

$[[Dx_1, Dx_2, x_3] + [Dx_1, x_2, Dx_3] + [x_1, Dx_2, Dx_3], Dx_4, x_5] = -[[x_1, x_2, x_3], Dx_4, x_5]$,

$[D[Dx_1, Dx_2, Dx_3], Dx_4, x_5] = -[D[x_1, x_2, x_3], Dx_4, x_5]$.

By Eq (11), $\{x_1, x_2, x_3\}_D = [Dx_1, Dx_2, x_3] = -\{x_2, x_1, x_3\}_D$, Eq (7) holds. Since

$$[Dx_3, Dx_4, [Dx_1, Dx_2, x_5]] = [Dx_1, Dx_2, [Dx_3, Dx_4, x_5]] - [[Dx_1, Dx_2, Dx_3], Dx_4, x_5,] + [Dx_3, [Dx_1, Dx_2, Dx_4], x_5],$$

$$\{\{x_1, x_2, x_3\}_{Dc}, x_4, x_5\}_D + \{x_3, \{x_1, x_2, x_4\}_{Dc}, x_5\}_D + \{x_3, x_4, \{x_1, x_2, x_5\}_D\}_D$$

$$= 2([([Dx_1, x_2, x_3] + [x_1, Dx_2, x_3] + [x_1, x_2, Dx_3]), Dx_4, x_5]$$

$$+[Dx_3, ([Dx_1, x_2, x_4] + [x_1, Dx_2, x_4] + [x_1, x_2, Dx_4]), x_5]$$

$$+[[Dx_1, Dx_2, Dx_3], Dx_4, x_5] + [Dx_3, [Dx_1, Dx_2, Dx_4], x_5]) + [Dx_1, Dx_2, [Dx_3, Dx_4, x_5]]$$

$$= 2([D[x_1, x_2, x_3], Dx_4, x_5] + [[Dx_1, Dx_2, Dx_3], Dx_4, x_5]$$

$$+[D[x_3, D[x_1, x_2, x_4], x_5] + [Dx_3, [Dx_1, Dx_2, Dx_4], x_5]) + [Dx_1, Dx_2, [Dx_3, Dx_4, x_5]]$$

$= [Dx_1, Dx_2, [Dx_3, Dx_4, x_5]]$, we get Eq (8).

Therefore, A is a 3-pre-Lie algebra in the multiplication (11). Eq (12) follows from Eqs (2) and (11), and a direct computation. □

Theorem 2.3. *Let* $(A, [\cdot,\cdot,\cdot])$ *be a 3-Lie algebra, D be an involutive derivation on A. Then D is an algebra isomorphism from the sub-adjacent 3-Lie algebra* $(A, \{\cdot,\cdot,\cdot\}_{Dc})$ *of the 3-pre-Lie algebra* $(A, \{\cdot,\cdot,\cdot\}_D)$ *to the 3-Lie algebra* $(A, [\cdot,\cdot,\cdot])$*, and*

$$\{x,y,z\}_{Dc} = \{x,y,z\}_D + \{y,z,x\}_D + \{z,x,y\}_D = D[Dx, Dy, Dz],\ \forall x,y,z \in A. \tag{13}$$

Furthermore,

$$\{x,y,z\}_{Dc} = \begin{cases} 0,\ x,y,z \in A_1, \text{or } x,y,z \in A_{-1}, \\ -[x,y,z],\ x,y \in A_1, z \in A_{-1}, \\ -[x,y,z],\ x,y \in A_{-1}, z \in A_1. \end{cases} \tag{14}$$

Proof. By Eq (12), the sub-adjacent 3-Lie algebra $(A, \{\cdot,\cdot,\cdot\}_{Dc})$ with the multiplication

$$\{x,y,z\}_{Dc} = \{x,y,z\}_D + \{y,z,x\}_D + \{z,x,y\}_D = [Dx, Dy, z] + [Dy, Dz, x] + [Dz, Dx, y] = D[Dx, Dy, Dz].$$

It follows Eq (13). Since for all $x,y,z \in A$,

$D(\{x, y, z\}_{Dc}) = D(D[Dx, Dy, Dz] = D^2[Dx, Dy, Dz] = [Dx, Dy, Dz]$,
D is an algebra isomorphism. Thanks to Eqs (13) and (2), Eq(14) holds. □

Theorem 2.4. *Let $(A, [\cdot, \cdot, \cdot])$ be a 3-Lie algebra with an involutive derivation D. Then there exists a compatible 3-pre Lie algebra $(A, \{, , \}_A)$, where*

$$\{x, y, z\}_A = D[x, y, Dz]. \tag{15}$$

Proof. By Eq (15), for all $x_i \in A, 1 \leq i \leq 5$,
$\{x_1, x_2, x_3\}_A = D[x_1, x_2, Dx_3] = -D[x_2, x_1, Dx_3] = -\{x_2, x_1, x_3\}_A$,

$\{\{x_1, x_2, \{x_3, x_4, x_5\}_A\}_A = D[x_1, x_2, D^2[x_3, x_4, Dx_5]] = D[x_1, x_2, [x_3, x_4, Dx_5]]$,
Eq (7) holds. And

$$[[Dx_1, Dx_2, x_3] + [Dx_1, x_2, Dx_3] + [x_1, Dx_2, Dx_3], x_4, Dx_5] = -[[x_1, x_2, x_3], x_4, Dx_5],$$

$\{\{x_1, x_2, x_3\}_{Ac}, x_4, x_5\}_A + \{x_3, \{x_1, x_2, x_4\}_{Ac}, x_5\}_A + \{x_3, x_4, \{x_1, x_2, x_5\}_A\}_A$
$= D([D([x_1, x_2, Dx_3] + [x_2, x_3, Dx_1] + [x_3, x_1, Dx_2]), x_4, Dx_5]$
$+[x_3, D([x_1, x_2, Dx_4] + [x_2, x_4, Dx_1] + [x_4, x_1, Dx_2]), Dx_5]+[x_3, x_4, [x_1, x_2, Dx_5]])$
$= D(2[[Dx_1, x_2, Dx_3], x_4, Dx_5] + 2[[x_1, Dx_2, Dx_3], x_4, Dx_5] + 2[[Dx_1, Dx_2, x_3], x_4, Dx_5]$
$+2[x_3, [Dx_1, Dx_2, x_4], Dx_5] + 2[x_3, [x_1, Dx_2, Dx_4], Dx_5] + 2[x_3, [Dx_1, x_2, Dx_4], Dx_5])$
$+3[[x_1, x_2, x_3], x_4, Dx_5] + 3[x_3, [x_1, x_2, x_4], Dx_5] + 3[x_3, x_4, [x_1, x_2, Dx_5]])$
$= D([x_1, x_2, [x_3, x_4, Dx_5]] + [[x_1, x_2, x_3], x_4, Dx_5] + [x_3, [x_1, x_2, x_4], Dx_5]$
$+[D[Dx_1, Dx_2, Dx_3], x_4, Dx_5] + [x_3, D[Dx_1, Dx_2, Dx_4], Dx_5])$
$= D[x_1, x_2, [x_3, x_4, Dx_5]] = \{x_1, x_2, \{x_3, x_4.x_5\}_A\}_A$,
it follows Eq (8). Similar discussion to the above, we get Eq (9). Thanks to

$$\{x_1, x_2, x_3\}_A = D([x_1, x_2, Dx_3] + [x_2, x_3, Dx_1] + [x_3, x_1, Dx_2]),$$

$(A, \{, , \}_A)$ is the compatible 3-pre-Lie algebra of $(A, [, ,])$. □

Theorem 2.5. *Let A be an m-dimensional 3-Lie algebra. If $m = 4$, then there are compatible 3-pre Lie algebras. If $m = 5$, $\dim A^1 \leq 3$, or, $\dim A^1 = 4$ and $Z(A) \neq 0$, then there are compatible 3-pre-Lie algebras.*

Proof. The result follows from the straightforward checking of the existence of involutive derivations by the classification theorems in [5] and Theorem 2.4, we omit the computation process. □

3. Local cocycle 3-Lie bialgebras induced by involutive derivations

In paper [2], the concept of 3-Lie bialgebra was introduced, but it has not close relationship with 3-Lie classical Yang-Baxter equation. Authors in [4], gave another 3-Lie bialgebraic structures on a 3-Lie algebra, it is called the local cocycle 3-Lie bialgebra. It has close relationship with 3-pre-Lie algebras, Manin triples, matched pairs of 3-Lie algebras and 3-Lie classical Yang-Baxter equation. So, in this section, we study local cocycle 3-Lie bialgebras induced by involutive derivations.

Definition 3.1. [4] *A **local cocycle** 3-**Lie bialgebra** is a triple* $(A, [,,], \Delta)$*, where* $(A, [,,])$ *is a* 3*-Lie algebra and* $\Delta = \Delta_1 + \Delta_2 + \Delta_3 : A \to A \otimes A \otimes A$ *is a linear mapping such that* (A^*, Δ^*) *is a 3-Lie algebra, and*

- Δ_1 *is a* 1*-cocycle associated to the representataion* $(A \otimes A \otimes A, ad \otimes 1 \otimes 1)$*,*
- Δ_2 *is a* 1*-cocycle associated to the representataion* $(A \otimes A \otimes A, 1 \otimes ad \otimes 1)$*,*
- Δ_3 *is a* 1*-cocycle associated to the representataion* $(A \otimes A \otimes A, 1 \otimes 1 \otimes ad)$*,*

where $\Delta^* : A^* \otimes A^* \otimes A^* \to A^*$ *is the dual mapping of* Δ*, that is, for all* $\alpha, \beta, \gamma \in A^*$ *and* $x \in A$*,*

$$\langle \Delta^*(\alpha, \beta, \gamma), x \rangle = \langle \alpha \otimes \beta \otimes \gamma, \Delta x \rangle.$$

Let A be a 3-Lie algebra, $r = \sum\limits_{i=1}^{m} x_i \otimes y_i \in A \otimes A$, $p, q \in \mathbb{Z}_{>0}$, and $1 \leq p \neq q \leq m$. Define an inclusion $\cdot_{pq} : \otimes^2 A \longrightarrow \otimes^n A$ by sending $r = \sum\limits_i x_i \otimes y_i \in A \otimes A$ to

$$r_{pq} := \sum_i z_{i1} \otimes \cdots \otimes z_{in}, \quad \text{where } z_{ij} = \begin{cases} x_i, & j = p, \\ y_i, & j = q, \\ 1, & j \neq p, q, \end{cases} \quad 1 \leq p, q, i, j \leq n, \tag{16}$$

that is, r_{pq} puts x_i at the p-th position, y_i at the q-th position and 1 elsewhere in the n-tensor, where 1 is a symbol playing a similar role of unit.

Define an operator $\phi_{pq} : A^{\otimes m} \to A^{\otimes m}$, $\forall \sum x_1 \otimes x_2 \otimes \cdots \otimes x_p \otimes \cdots \otimes x_q \otimes \cdots \otimes x_m \in A^{\otimes m}$,

$$\phi_{pq}(x_1 \otimes x_2 \otimes \cdots \otimes x_p \otimes \cdots \otimes x_q \otimes \cdots \otimes x_m) = x_1 \otimes x_2 \otimes \cdots \otimes x_q \otimes \cdots \otimes x_p \otimes \otimes x_m,$$

that is, ϕ_{pq} exchanges the position of x_p with x_q, where $m \geq 2$.

Definition 3.2. [4] *Let* $(A, [\cdot, \cdot, \cdot])$ *be a* 3*-Lie algebra and* $r = \sum_i x_i \otimes y_i \in A \otimes A$*, denote*

$$\begin{aligned} [[r, r, r]] :\equiv \sum_{i,j,k} \big(& [x_i, x_j, x_k] \otimes y_i \otimes y_j \otimes y_k + x_i \otimes [y_i, x_j, x_k] \otimes y_j \otimes y_k \\ & + x_i \otimes x_j \otimes [y_i, y_j, x_k] \otimes y_k + x_i \otimes x_j \otimes x_k \otimes [y_i, y_j, y_k] \big). \end{aligned} \tag{17}$$

The equation

$$[[r, r, r]] = 0 \tag{18}$$

is called the 3**-Lie classical Yang-Baxter equation** *in the 3-Lie algebra A, and simply denoted by* **CYBE**.

Let $D : A \to A$ be a linear mapping. Define the tensor $\overline{D} \in A^* \otimes A$, for all $x \in A, \xi \in A^*$,

$$\overline{D}(x, \xi) = \langle \xi, Dx \rangle. \tag{19}$$

Theorem 3.3. *Let* $(A, [\cdot, \cdot, \cdot])$ *be a* 3*-Lie algebra with a basis* $\{x_1, \cdots, x_n\}$*,* $\{x_1^*, \cdots, x_n^*\}$ *be the dual basis of* A^**, and* D *be an involutive derivation on* A*. Then*

$$r = \overline{D} - \sigma_{12} \overline{D} \tag{20}$$

is a skew-symmetric solution of **CYBE** *in the semi-direct product* 3*-Lie algebra* $A \ltimes_{ad^*} A^*$*, and*

$$\overline{D} = \sum_{i=1}^{n} x_i^* \otimes Dx_i, \ r = \sum_{i=1}^{n} x_i^* \otimes Dx_i - \sum_{i=1}^{n} Dx_i \otimes x_i^* \in A^* \otimes A. \tag{21}$$

Proof. By Theorem 2.1, D is an $\wp$-operator associative to the adjoint representation (A, ad). By Theorem 3.19 in [4], $r = \overline{D} - \sigma_{12}\overline{D}$ is a skew-symmetric solution of **CYBE** in the semi-direct product 3-Lie algebra $A \ltimes_{ad^*} A^*$. Thanks to $D \in Der(A)$ and $D^2 = I_d$, we get Eq (21). The proof is completed. □

Theorem 3.4. *Let A be a 3-Lie algebra, D be an involutive derivation on A, $\overline{D}$ and r be defined as Eq* (21) *and Eq* (20)*. Then r induces a local cocycle 3-Lie bialgebra $(A \ltimes_{ad^*} A^*, \Delta)$, where for all $x \in A \oplus A^*$,*

$$
\begin{cases}
\Delta(x) = \Delta_1(x) + \Delta_2(x) + \Delta_3(x), \\
\Delta_1(x) = \sum_{i,j=1}^{n} \mu(x, x_i^*, -Dx_j) \otimes x_j^* \otimes Dx_i \\
\qquad\quad + \sum_{i,j=1}^{n} \mu(x, -Dx_i, x_j^*) \otimes Dx_j \otimes x_i^* \\
\qquad\quad + \sum_{i,j=1}^{n} \mu(x, Dx_i, Dx_j) \otimes x_j^* \otimes x_i^*, \\
\Delta_2(x) = \phi_{13}\phi_{12}\Delta_1(x), \\
\Delta_3(x) = \phi_{12}\phi_{13}\Delta_1(x).
\end{cases} \tag{22}
$$

Proof. Suppose $\dim A = n$, and the decomposition of A associated to D is

$$A = A_1 + A_{-1},$$

and $\{x_1, ..., x_s, x_{s+1}, ...x_n\}$ is a basis of A, $x_1, ..., x_s \in A_1$, $x_{s+1}, ...x_n \in A_{-1}$. By Theorem 3.3, and Eq (21), the tensor

$$r = \sum_{i=1}^{s} (x_i^* \otimes x_i - x_i \otimes x_i^*) - \sum_{i=s+1}^{n} (x_i^* \otimes x_i - x_i \otimes x_i^*) \tag{23}$$

is a solution of **CYBE** in the semi-direct product 3-Lie algebra $A \ltimes_{ad^*} A^*$.

For convenience, let $r = \sum_{i=1}^{2n} u_i \otimes v_i$. By [4], r determines a local cocycle 3-Lie bialgebra $(A \ltimes_{ad^*} A^*, \Delta)$, where for all $x \in A \ltimes_{ad^*} A^*$,

$\Delta_1(x) = \sum_{i,j} \mu(x, u_i, u_j) \otimes v_j \otimes v_i$,
$\Delta_2(x) = \phi_{12}\Delta_1(x) = \sum_{i,j} v_i \otimes \mu(x, u_i, u_j) \otimes v_j$,

$\Delta_3(x) = \phi_{12}\phi_{13}\Delta_1(x) = \sum_{i,j} v_j \otimes v_i \otimes \mu(x, u_i, u_j)$. Thanks to Eq (23),

$$u_i = \begin{cases} x_i^* & 1 \le i \le n, \\ -D(x_{i-n}) & n+1 \le i \le 2n, \end{cases} \quad u_j = \begin{cases} x_j^* & 1 \le j \le n, \\ -D(x_{j-n}) & n+1 \le j \le 2n, \end{cases}$$

$$v_i = \begin{cases} y_i^* & 1 \le i \le n, \\ D(x_i) & n+1 \le i \le 2n, \end{cases}, \quad v_j = \begin{cases} D(x_j) & 1 \le j \le n, \\ x_{j-n}^* & n+1 \le j \le 2n. \end{cases}$$

Therefore,

$$
\begin{aligned}
\Delta_1(x) &= \sum_{i=1}^{n}\sum_{j=1}^{n} \mu(x, u_i, u_j) \otimes v_j \otimes v_i + \sum_{i=1}^{n}\sum_{j=n+1}^{2n} \mu(x, u_i, u_j) \otimes v_j \otimes v_i \\
&\quad + \sum_{i=n+1}^{2n}\sum_{j=1}^{n} \mu(x, u_i, u_j) \otimes v_j \otimes v_i + \sum_{i=n+1}^{2n}\sum_{j=n+1}^{2n} \mu(x, u_i, u_j) \otimes v_j \otimes v_i \\
&= \sum_{i=1}^{n}\sum_{j=1}^{n} \mu(x, x_i^*, x_j^*) \otimes Dx_j \otimes Dx_i + \sum_{i=1}^{n}\sum_{j=n+1}^{2n} \mu(x, x_i^*, -Dx_{j-n}) \otimes x_{j-n}^* \otimes Dx_i \\
&\quad + \sum_{i=n+1}^{2n}\sum_{j=1}^{n} \mu(x, -Dx_{i-n}, x_j^*) \otimes Dx_j \otimes x_{i-n}^* \\
&\quad + \sum_{i=n+1}^{2n}\sum_{j=n+1}^{2n} \mu(x, -Dx_{i-n}, -Dx_{j-n}) \otimes x_{j-n}^* \otimes x_{i-n}^* \\
&= \sum_{i,j} \mu(x, x_i^*, -Dx_j) \otimes x_j^* \otimes Dx_i + \sum_{i,j} \mu(x, -Dx_i, x_j^*) \otimes Dx_j \otimes x_i^*
\end{aligned}
$$

$+\sum_{i,j}\mu(x,Dx_i,Dx_j)\otimes x_j^*\otimes x_i^*$. The proof is completed. □

Corollary 3.5. *Let $(A,[\cdot,\cdot,\cdot])$ be an n-dimensional 3-Lie algebra with an involutive derivation D, $\{x_1,\cdots,x_s,x_{s+1},\cdots,x_n\}$ be a basis of A and $x_i\in A_1, 1\leq i\leq s, x_j\in A_{-1}, s+1\leq j\leq n$. Then $(A\ltimes_{ad^*}A^*,\mu,\Delta)$ is a local cocycle 3-Lie bialgebra, where $\Delta=\Delta_1+\Delta_2+\Delta_3$, $\Delta_2(x)=\phi_{13}\phi_{12}\Delta_1(x), \Delta_3(x)=\phi_{12}\phi_{13}\Delta_1(x)$, and*

$$
\begin{aligned}
\Delta_1(x) &= \sum_{i=1}^{s}\sum_{j=1}^{s}\mu(x,x_i^*,-x_j)\otimes x_j^*\otimes x_i+\sum_{i=1}^{s}\sum_{j=s+1}^{n}\mu(x,x_i^*,x_j)\otimes x_j^*\otimes x_i\\
&+\sum_{i=s+1}^{n}\sum_{j=1}^{s}\mu(x,x_i^*,-x_j)\otimes x_j^*\otimes(-x_i)+\sum_{i=s+1}^{n}\sum_{j=s+1}^{n}\mu(x,x_i^*,x_j)\otimes x_j^*\otimes(-x_i)\\
&+\sum_{i=1}^{s}\sum_{j=1}^{s}\mu(x,-x_i,x_j^*)\otimes x_j\otimes x_i^*+\sum_{i=1}^{s}\sum_{j=s+1}^{n}\mu(x,-x_i,x_j^*)\otimes(-x_j)^*\otimes x_i^*\\
&+\sum_{i=s+1}^{n}\sum_{j=1}^{s}\mu(x,x_i,x_j^*)\otimes x_j\otimes x_i^*+\sum_{i=s+1}^{n}\sum_{j=s+1}^{n}\mu(x,x_i,x_j^*)\otimes(-x_j)\otimes x_i^*\\
&+\sum_{i=1}^{s}\sum_{j=1}^{s}\mu(x,x_i,x_j)\otimes x_j^*\otimes x_i^*+\sum_{i=1}^{s}\sum_{j=s+1}^{n}\mu(x,x_i,-x_j)\otimes x_j^*\otimes x_i^*\\
&+\sum_{i=s+1}^{n}\sum_{j=1}^{s}\mu(x,-x_i,x_j)\otimes x_j^*\otimes x_i^*+\sum_{i=s+1}^{n}\sum_{j=s+1}^{n}\mu(x,x_i,x_j)\otimes x_j^*\otimes x_i^*.
\end{aligned}
\tag{24}
$$

Proof. The result follows from Theorem 3.4, direcrly. □

4. Eight and Ten dimensional 3-Lie bialgebras

In this section, we construct 8 and 10-dimensional local cocycle 3-Lie bialgebras by involutive derivations. We need the classification theorems of 4 and 5-dimensional 3-Lie algebras in [5].

Lemma 4.1. [5] *Let $(A,[,,])$ be a 4-dimensional non-abelian 3-Lie algebra with a basis $\{x_1,x_2,x_3,x_4\}$. Then up to isomorphisms, A is one and only one of the following possibilities*

(b_1) $[x_2,x_3,x_4]=x_1$; (b_2) $[x_1,x_2,x_3]=x_1$; (c_3) $\begin{cases}[x_1,x_3,x_4]=x_1,\\ [x_2,x_3,x_4]=x_2;\end{cases}$

(c_1) $\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_1,x_3,x_4]=x_2;\end{cases}$ $(c_2)\begin{cases}[x_2,x_3,x_4]=\alpha x_1+x_2,\\ [x_1,x_3,x_4]=e_2,\alpha\in F \text{ and } \alpha\neq 0;\end{cases}$

(d_1) $\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_1,x_3,x_4]=x_2,\\ [x_1,x_2,x_3]=x_3;\end{cases}$ (e_1) $\begin{cases}[x_2,x_3,x_4]=-x_2,\\ [x_1,x_3,x_4]=x_1,\\ [x_1,x_2,x_3]=x_3,\\ [x_1,x_2,x_4]=-x_4.\end{cases}$

Lemma 4.2. [5] *Let $(A,[\cdot,\cdot,\cdot])$ be a 5-dimensional non-abelian 3-Lie algebra with a basis $\{x_1,x_2,x_3,x_4,x_5\}$. Then up to isomorphisms, A is one and only one of the following possibilities*

$(b_1)\ [x_2, x_3, x_4] = x_1$; $(b_2)\ [x_1, x_2, x_3] = x_1$,

$$(c_1)\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_3,x_4,x_5]=x_2;\end{cases}\quad (c_2)\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_2,x_4,x_5]=x_2,\\ [x_1,x_4,x_5]=x_1;\end{cases}\quad (c_3)\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_1,x_3,x_4]=x_2;\end{cases}$$

$$(c_4)\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_1,x_3,x_4]=x_2,\\ [x_2,x_4,x_5]=x_2,\\ [x_1,x_4,x_5]=x_1;\end{cases}\quad (c_5)\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_2,x_4,x_5]=\alpha x_1+x_2;\end{cases}$$

$$(c_6)\begin{cases}[x_2,x_3,x_4]=\alpha x_1+x_2,\\ [x_1,x_3,x_4]=x_2,\\ [x_2,x_4,x_5]=x_2,\\ [x_1,x_4,x_5]=x_1;\end{cases}\quad (c_7)\begin{cases}[x_1,x_3,x_4]=x_1,\\ [x_2,x_3,x_4]=x_2;\end{cases}\quad \alpha\in F \text{ and } \alpha\neq 0.$$

$$(d_1)\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_2,x_4,x_5]=-x_2,\\ [x_3,x_4,x_5]=x_3;\end{cases}\quad (d_2)\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_3,x_4,x_5]=\alpha x_2+x_3;\end{cases}$$

$$(d_2)\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_3,x_4,x_5]=\alpha x_2+x_3;\end{cases}\quad (d_3)\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_3,x_4,x_5]=x_3,\\ [x_2,x_4,x_5]=x_2,\\ [x_1,x_4,x_5]=2x_1;\end{cases}$$

$$(d_4)\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_1,x_3,x_4]=x_2,\\ [x_1,x_2,x_4]=x_3;\end{cases}\quad (d_5)\begin{cases}[x_1,x_4,x_5]=x_1,\\ [x_2,x_4,x_5]=x_3,\\ [x_3,x_4,x_5]=\beta x_2+(1+\beta)x_3;\end{cases}$$

$$(d_6)\begin{cases}[x_1,x_4,x_5]=x_1,\\ [x_2,x_4,x_5]=x_2,\\ [x_3,x_4,x_5]=x_3;\end{cases}\quad (d_7)\begin{cases}[x_1,x_4,x_5]=x_2,\\ [x_2,x_4,x_5]=x_3,\\ [x_3,x_4,x_5]=sx_1+tx_2+ux_3;\end{cases}$$

$\beta, s, t, u \in F$, $\beta s \neq 0$.

$$(e_1)\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_3,x_4,x_5]=x_2,\\ [x_2,x_4,x_5]=x_3,\\ [x_2,x_3,x_5]=x_4;\end{cases}\quad (e_2)\begin{cases}[x_2,x_3,x_4]=x_1,\\ [x_1,x_3,x_4]=x_2,\\ [x_1,x_2,x_4]=x_3,\\ [x_1,x_2,x_3]=x_4.\end{cases}$$

Theorem 4.3. *Let* $(A, [\cdot,\cdot,\cdot])$ *be a 4-dimensional* 3*-Lie algebra with a basis* $\{x_1, x_2, x_3, x_4\}$*. Then we have* 8*-dimensional local cocycle* 3*-Lie bialgebras* $(A \ltimes_{ad^*} A^*, \mu_i, \Delta^i)$*,* $1 \leq i \leq 7$*, where*

$$(25)\qquad \begin{cases}\mu_1(x_2,x_3,x_4)=x_1,\\ \mu_1(x_2,x_3,x_1^*)=-x_4^*,\\ \mu_1(x_2,x_4,x_1^*)=x_3^*,\\ \mu_1(x_3,x_4,x_1^*)=-x_2^*.\end{cases}\qquad \begin{cases}\Delta^1(x_1^*)=x_2^*\wedge x_4^*\wedge x_3^*,\\ \Delta^1(x_2)=x_1\wedge x_3^*\wedge x_4^*,\\ \Delta^1(x_3)=x_1\wedge x_4^*\wedge x_2^*,\\ \Delta^1(x_4)=x_1\wedge x_2^*\wedge x_3^*,\\ \Delta^1(x_2^*)=\Delta^1(x_3^*)=\Delta^1(x_4^*)=\Delta^1(x_1)=0.\end{cases}$$

$$
\begin{cases}
\mu_2(x_1, x_2, x_3) = x_1, \\
\mu_2(x_1, x_2, x_1^*) = -x_3^*, \\
\mu_2(x_2, x_3, x_1^*) = -x_1^*, \\
\mu_2(x_1, x_3, x_1^*) = x_2^*.
\end{cases}
\quad
\begin{cases}
\Delta^2(x_1^*) = x_1^* \wedge x_3^* \wedge x_2^*, \\
\Delta^2(x_1) = x_1 \wedge x_2^* \wedge x_3^*, \\
\Delta^2(x_2) = x_1 \wedge x_3^* \wedge x_1, \\
\Delta^2(x_3) = x_1 \wedge x_1^* \wedge x_2^*, \\
\Delta^2(x_4) = \Delta^2(x_2^*) = \Delta^2(x_3^*) = \Delta^2(x_4^*) = 0.
\end{cases} \tag{26}
$$

$$
\begin{cases}
\mu_3(x_2, x_3, x_4) = x_1, \\
\mu_3(x_1, x_3, x_4) = x_2, \\
\mu_3(x_2, x_3, x_1^*) = -x_4^*, \\
\mu_3(x_3, x_4, x_1^*) = -x_2^*, \\
\mu_3(x_2, x_4, x_1^*) = x_3^*, \\
\mu_3(x_1, x_3, x_2^*) = -x_4^*, \\
\mu_3(x_1, x_4, x_2^*) = x_3^*, \\
\mu_3(x_3, x_4, x_2^*) = -x_1^*.
\end{cases}
\quad
\begin{cases}
\Delta^3(x_1^*) = x_2^* \wedge x_4^* \wedge x_3^*, \\
\Delta^3(x_2^*) = x_1^* \wedge x_4^* \wedge x_3^*, \\
\Delta^3(x_1) = x_2 \wedge x_3^* \wedge x_4^*, \\
\Delta^3(x_2) = x_1 \wedge x_3^* \wedge x_4^*, \\
\Delta^3(x_3) = x_1 \wedge x_4^* \wedge x_2^* + x_2 \wedge x_4^* \wedge x_1^*, \\
\Delta^3(x_4) = x_1 \wedge x_2^* \wedge x_3^* + x_1^* \wedge x_3^* \wedge x_2, \\
\Delta^3(x_3^*) = \Delta^3(x_4^*) = 0.
\end{cases} \tag{27}
$$

$$
\begin{cases}
\mu_4(x_2, x_3, x_4) = \alpha x_1 + x_2, \\
\mu_4(x_1, x_3, x_4) = x_2, \\
\mu_4(x_2, x_3, x_1^*) = -\alpha x_4^*, \\
\mu_4(x_2, x_3, x_2^*) = -x_4^*, \\
\mu_4(x_2, x_4, x_1^*) = \alpha x_3^*, \\
\mu_4(x_2, x_4, x_2^*) = x_3^*, \\
\mu_4(x_3, x_4, x_1^*) = -\alpha x_2^*, \\
\mu_4(x_3, x_4, x_2^*) = -x_1^* - x_2^*, \\
\mu_4(x_1, x_3, x_2^*) = -x_4^*, \\
\mu_4(x_1, x_4, x_2^*) = x_3^*.
\end{cases}
\quad
\begin{cases}
\Delta^4(x_1^*) = \alpha x_3^* \wedge x_2^* \wedge x_4^* \\
\Delta^4(x_2^*) = x_1^* \wedge x_4^* \wedge x_3^* + x_2^* \wedge x_4^* \wedge x_3^*, \\
\Delta^4(x_1) = x_2 \wedge x_3^* \wedge x_4^*, \\
\Delta^4(x_2) = \alpha x_1 \wedge x_3^* \wedge x_4^* + x_3^* \wedge x_4^* \wedge x_2, \\
\Delta^4(x_3) = \alpha x_1 \wedge x_4^* \wedge x_2^* + x_2 \wedge x_4^* \wedge x_2^* \\
\qquad\quad + x_1^* \wedge x_2 \wedge x_4^*, \\
\Delta^4(x_4) = \alpha x_1 \wedge x_2^* \wedge x_3^* + x_2 \wedge x_2^* \wedge x_3^* \\
\qquad\quad + x_1^* \wedge x_3^* \wedge x_2, \\
\Delta^4(x_3^*) = \Delta^4(x_4^*) = 0.
\end{cases} \tag{28}
$$

(29)
$$\begin{cases} \mu_6(x_2,x_3,x_4)=x_1,\\ \mu_6(x_1,x_3,x_4)=x_2,\\ \mu_6(x_1,x_2,x_4)=x_3,\\ \mu_6(x_2,x_3,x_1^*)=-x_4^*,\\ \mu_6(x_3,x_4,x_1^*)=-x_2^*,\\ \mu_6(x_2,x_4,x_1^*)=x_3^*,\\ \mu_6(x_1,x_3,x_2^*)=-x_4^*,\\ \mu_6(x_1,x_4,x_2^*)=x_3^*,\\ \mu_6(x_3,x_4,x_2^*)=-x_1^*,\\ \mu_6(x_1,x_2,x_3^*)=-x_4^*,\\ \mu_6(x_1,x_4,x_3^*)=x_2^*,\\ \mu_6(x_2,x_4,x_3^*)=-x_1^*. \end{cases} \quad \begin{cases} \Delta^6(x_1^*)=x_2^*\wedge x_4^*\wedge x_3^*,\\ \Delta^6(x_2^*)=x_1^*\wedge x_4^*\wedge x_3^*,\\ \Delta^6(x_3^*)=x_1^*\wedge x_4^*\wedge x_2^*\\ \Delta^6(x_1)=x_2\wedge x_3^*\wedge x_4^*+x_2^*\wedge x_4^*\wedge x_3,\\ \Delta^6(x_2)=x_1\wedge x_3^*\wedge x_4^*+x_4^*\wedge x_1^*\wedge x_3,\\ \Delta^6(x_3)=x_4^*\wedge x_2^*\wedge x_1+x_4^*\wedge x_1^*\wedge x_2,\\ \Delta^6(x_4)=x_1^*\wedge x_3^*\wedge x_2+x_1^*\wedge x_2^*\wedge x_3,\\ \qquad\quad +x_1\wedge x_2^*\wedge x_3^*,\\ \Delta^6(x_4^*)=0. \end{cases}$$

(30)
$$\begin{cases} \mu_7(x_2,x_3,x_4)=-x_2,\\ \mu_7(x_1,x_3,x_4)=x_1,\\ \mu_7(x_1,x_2,x_3)=x_3,\\ \mu_7(x_1,x_2,x_4)=-x_4,\\ \mu_7(x_2,x_3,x_2^*)=x_4^*,\\ \mu_7(x_3,x_4,x_2^*)=x_2^*,\\ \mu_7(x_2,x_4,x_2^*)=-x_3^*,\\ \mu_7(x_1,x_3,x_1^*)=-x_4^*,\\ \mu_7(x_3,x_4,x_1^*)=-x_1^*,\\ \mu_7(x_1,x_4,x_1^*)=x_3^*,\\ \mu_7(x_1,x_2,x_3^*)=-x_3^*,\\ \mu_7(x_1,x_3,x_3^*)=x_2^*,\\ \mu_7(x_2,x_3,x_3^*)=-x_1^*,\\ \mu_7(x_1,x_2,x_4^*)=x_4^*,\\ \mu_7(x_1,x_4,x_4^*)=-x_2^*,\\ \mu_7(x_2,x_4,x_4^*)=x_1^*. \end{cases} \quad \begin{cases} \Delta^7(x_1^*)=x_1^*\wedge x_4^*\wedge x_3^*,\\ \Delta^7(x_2^*)=x_2^*\wedge x_3^*\wedge x_4^*,\\ \Delta^7(x_3^*)=x_1^*\wedge x_3^*\wedge x_2^*,\\ \Delta^7(x_4^*)=x_1^*\wedge x_2^*\wedge x_4^*,\\ \Delta^7(x_1)=x_1\wedge x_3^*\wedge x_4^*+x_2^*\wedge x_3^*\wedge x_3\\ \qquad\quad +x_4^*\wedge x_2^*\wedge x_4,\\ \Delta^7(x_2)=x_2\wedge x_4^*\wedge x_3^*+x_3^*\wedge x_1^*\wedge x_3\\ \qquad\quad +x_1^*\wedge x_4^*\wedge x_4,\\ \Delta^7(x_3)=x_4^*\wedge x_1^*\wedge x_1+x_2^*\wedge x_4^*\wedge x_2\\ \qquad\quad +x_1^*\wedge x_2^*\wedge x_3,\\ \Delta^7(x_4)=x_3^*\wedge x_2^*\wedge x_2+x_1^*\wedge x_3^*\wedge x_1\\ \qquad\quad +x_4\wedge x_2^*\wedge x_1^*,\\ \Delta^7(x_4^*)=0. \end{cases}$$

$$
(31)\qquad
\begin{cases}
\mu_5(x_1,x_3,x_4)=x_1,\\
\mu_5(x_2,x_3,x_4)=x_2,\\
\mu_5(x_1,x_3,x_1^*)=-x_4^*,\\
\mu_5(x_1,x_4,x_1^*)=x_3^*,\\
\mu_5(x_3,x_4,x_1^*)=-x_1^*,\\
\mu_5(x_2,x_3,x_2^*)=-x_4^*,\\
\mu_5(x_2,x_4,x_2^*)=x_3^*,\\
\mu_5(x_3,x_4,x_2^*)=-x_2^*.
\end{cases}
\qquad
\begin{cases}
\Delta^5(x_1^*)=x_1^*\wedge x_4^*\wedge x_3^*,\\
\Delta^5(x_2^*)=x_2^*\wedge x_4^*\wedge x_3^*,\\
\Delta^5(x_1)=x_1\wedge x_3^*\wedge x_4^*,\\
\Delta^5(x_2)=x_2\wedge x_3^*\wedge x_4^*,\\
\Delta^5(x_3)=x_1\wedge x_4^*\wedge x_1^*+x_2\wedge x_4^*\wedge x_2^*,\\
\Delta^5(x_4)=x_1\wedge x_1^*\wedge x_3^*+x_2\wedge x_2^*\wedge x_3^*,\\
\Delta^5(x_3^*)=\Delta^5(x_4^*)=0.
\end{cases}
$$

Proof. By Lemma 4.1, and Eq (5), if A is the case of (b_1), then there is an involutive derivation D on A such that $x_1, x_2, x_3 \in A_1$ and $x_4 \in A_{-1}$. Then the multiplication of the semi-direct product 3-Lie algebra $(A \ltimes_{ad^*} A^*, \mu_1)$ in the basis $\{x_1, \cdots, x_4, x_1^*, \cdots, x_4^*\}$ is as follows

$\mu_1(x_2,x_3,x_4)=x_1$, $\mu_1(x_2,x_3,x_1^*)=ad^*(x_2,x_3)(x_1^*)=-x_4^*$,

$\mu_1(x_2,x_4,x_1^*)=ad^*(x_2,x_4)(x_1^*)=-x_3^*$, $\mu_1(x_3,x_4,x_1^*)=ad^*(x_3,x_4)(x_1^*)=-x_2^*$.

Thanks to Theorem 3.3 and Theorem 3.4, $(A \ltimes_{ad^*} A^*, \mu_1, \Delta^1)$ is a local cocycle 3-Lie bialgebra, where

$$
\begin{aligned}
\Delta_1^1(x_1^*) &= [x_1^*,x_2,x_3]\otimes x_3^*\otimes x_2^*+[x_1^*,x_3,x_2]\otimes x_2^*\otimes x_3^*+[x_1^*,x_2,-x_4]\otimes x_4^*\otimes x_2^*\\
&\quad+[x_1^*,x_3,-x_4]\otimes x_4^*\otimes x_3^*+[x_1^*,-x_4,x_2]\otimes x_2^*\otimes x_4^*+[x_1^*,-x_4,x_3]\otimes x_3^*\otimes x_4^*\\
&= -x_4^*\otimes x_3^*\otimes x_2^*+x_4^*\otimes x_2^*\otimes x_3^*-x_3^*\otimes x_4^*\otimes x_2^*\\
&\quad+x_2^*\otimes x_4^*\otimes x_3^*+x_3^*\otimes x_2^*\otimes x_4^*-x_2^*\otimes x_3^*\otimes x_4^*,\\
\Delta_2^1(x_1^*) &= \phi_{13}\phi_{12}\Delta_1^1(x_1^*),\ \Delta_3^1(x_1^*)=\phi_{12}\phi_{13}\Delta_1^1(x_1^*),\\
\Delta^1(x_1^*) &= \Delta_1^1(x_1^*)+\Delta_2^1(x_1^*)+\Delta_3^1(x_1^*)\\
&= -x_4^*\otimes x_3^*\otimes x_2^*-x_4^*\otimes x_2^*\otimes x_3^*+x_2^*\otimes x_4^*\otimes x_3^*\\
&\quad-x_2^*\otimes x_3^*\otimes x_4^*+x_3^*\otimes x_2^*\otimes x_4^*-x_3^*\otimes x_4^*\otimes x_2^*=x_2^*\wedge x_4^*\wedge x_3^*.
\end{aligned}
$$

Similarly, we have $\Delta^1(x_2)=x_1\wedge x_3^*\wedge x_4^*$, $\Delta^1(x_3)=x_1\wedge x_4^*\wedge x_2^*$, $\Delta^1(x_4)=x_1\wedge x_2^*\wedge x_3^*$, and $\Delta^1(x_2^*)=\Delta^1(x_3^*)=\Delta^1(x_4^*)=\Delta^1(x_1)=0$. It follows Eq (25). Eq (26) follows from the completely similar discussion.

For the case $\dim A^1=2$, we get local cocycle 3-Lie bialgebras $(A\ltimes_{ad^*}A^*,\mu_i,\Delta^i))$, $i=3,4,5$. If $\dim A^1=3$, then we get local cocycle 3-Lie bialgebras $(A\ltimes_{ad^*}A^*,\mu_i,\Delta^i))$, $i=5,6$. If $\dim A^1=4$, then we get local cocycle 3-Lie bialgebras $(A\ltimes_{ad^*}A^*,\mu_7,\Delta^7))$. We omit the proving process since it is similar to the case $\dim A^1=1$. □

Theorem 4.4. *Let A be a 5-dimensional 3-Lie algebra with a basis $\{x_1,x_2,x_3,x_4,x_5\}$ and satisfy $\dim A^1\le 3$, or, $\dim A^1=4$ and $Z(A)\neq 0$. Then we have 10-dimensional local cocycle 3-Lie bialgebras $(A\ltimes_{ad^*}A^*,\psi_j,\widetilde{\Delta}^j)$, $1\le j\le 17$, where $(A\ltimes_{ad^*}A^*,\psi_j)$ are semi-direct product 3-Lie*

algebras, and

$$
\begin{cases} \psi_1(x_2,x_3,x_4)=x_1, \\ \psi_1(x_2,x_3,x_1^*)=-x_4^*, \\ \psi_1(x_3,x_4,x_1^*)=-x_2^*, \\ \psi_1(x_2,x_4,x_1^*)=x_3^*. \end{cases} \quad \begin{cases} \widetilde{\Delta}^1(x_1^*)=x_2^*\wedge x_4^*\wedge x_3^*, \\ \widetilde{\Delta}^1(x_2)=x_1\wedge x_3^*\wedge x_4^*, \\ \widetilde{\Delta}^1(x_3)=x_1\wedge x_4^*\wedge x_2^*, \\ \widetilde{\Delta}^1(x_4)=x_1\wedge x_2^*\wedge x_3^*, \\ \widetilde{\Delta}^1(x_2^*)=\widetilde{\Delta}^1(x_5)=\widetilde{\Delta}^1(x_3^*)=0, \\ \widetilde{\Delta}^1(x_4^*)=\widetilde{\Delta}^1(x_5^*)=\widetilde{\Delta}^1(x_1)=0. \end{cases} \tag{32}
$$

$$
\begin{cases} \psi_2(x_2,x_3,x_4)=x_1, \\ \psi_2(x_3,x_4,x_5)=x_2, \\ \psi_2(x_2,x_3,x_1^*)=-x_4^*, \\ \psi_2(x_2,x_4,x_1^*)=x_3^*, \\ \psi_2(x_3,x_4,x_1^*)=-x_2^*, \\ \psi_2(x_3,x_4,x_2^*)=-x_5^*, \\ \psi_2(x_4,x_5,x_2^*)=-x_3^*, \\ \psi_2(x_3,x_5,x_2^*)=x_4^*. \end{cases} \quad \begin{cases} \widetilde{\Delta}^2(x_1^*)=x_2^*\wedge x_4^*\wedge x_3^*, \\ \widetilde{\Delta}^2(x_2^*)=x_3^*\wedge x_5^*\wedge x_4^*, \\ \widetilde{\Delta}^2(x_2)=x_1\wedge x_3^*\wedge x_4^*, \\ \widetilde{\Delta}^2(x_3)=x_1\wedge x_4^*\wedge x_2^*+x_2\wedge x_4^*\wedge x_5^*, \\ \widetilde{\Delta}^2(x_4)=x_1\wedge x_2^*\wedge x_3^*+x_5^*\wedge x_3^*\wedge x_2, \\ \widetilde{\Delta}^2(x_5)=x_2\wedge x_3^*\wedge x_4^*, \\ \widetilde{\Delta}^2(x_3^*)=\widetilde{\Delta}^2(x_4^*)=\widetilde{\Delta}^2(x_5^*)=\widetilde{\Delta}^2(x_1)=0. \end{cases} \tag{33}
$$

$$
\begin{cases} \psi_3(x_2,x_3,x_4)=x_1, \\ \psi_3(x_2,x_4,x_5)=x_2, \\ \psi_3(x_1,x_4,x_5)=x_1, \\ \psi_3(x_2,x_3,x_1^*)=-x_4^*, \\ \psi_3(x_2,x_4,x_1^*)=x_3^*, \\ \psi_3(x_3,x_4,x_1^*)=-x_2^*, \\ \psi_3(x_2,x_4,x_2^*)=-x_5^*, \\ \psi_3(x_2,x_5,x_2^*)=x_4^*, \\ \psi_3(x_4,x_5,x_2^*)=-x_2^*, \\ \psi_3(x_1,x_4,x_1^*)=-x_5^*, \\ \psi_3(x_4,x_5,x_1^*)=-x_1^*, \\ \psi_3(x_1,x_5,x_1^*)=x_4^*. \end{cases} \quad \begin{cases} \widetilde{\Delta}^3(x_1^*)=x_1^*\wedge x_5^*\wedge x_4^*+x_2^*\wedge x_4^*\wedge x_3^*, \\ \widetilde{\Delta}^3(x_2^*)=x_2^*\wedge x_5^*\wedge x_4^*, \\ \widetilde{\Delta}^3(x_1)=x_1\wedge x_4^*\wedge x_5^*, \\ \widetilde{\Delta}^3(x_2)=x_1\wedge x_3^*\wedge x_4^*+x_2\wedge x_4^*\wedge x_5^*, \\ \widetilde{\Delta}^3(x_3)=x_1\wedge x_4^*\wedge x_2^*, \\ \widetilde{\Delta}^3(x_4)=x_1\wedge x_2^*\wedge x_3^*+x_2\wedge x_5^*\wedge x_2^* \\ \qquad\qquad +x_5^*\wedge x_1^*\wedge x_1, \\ \widetilde{\Delta}^3(x_5)=x_2^*\wedge x_4^*\wedge x_2+x_1^*\wedge x_4^*\wedge x_1, \\ \widetilde{\Delta}^3(x_3^*)=\widetilde{\Delta}^3(x_4^*)=\widetilde{\Delta}^3(x_5^*)=0. \end{cases} \tag{34}
$$

(35)
$$
\begin{cases}
\psi_4(x_2,x_3,x_4)=\alpha x_1+x_2,\\
\psi_4(x_1,x_3,x_4)=x_2,\\
\psi_4(x_2,x_4,x_5)=x_2,\\
\psi_4(x_1,x_4,x_5)=x_1,\\
\psi_4(x_2,x_3,x_1^*)=-\alpha x_4^*,\\
\psi_4(x_2,x_4,x_1^*)=\alpha x_3^*,\\
\psi_4(x_3,x_4,x_1^*)=-\alpha x_2^*,\\
\psi_4(x_2,x_3,x_2^*)=-x_4^*,\\
\psi_4(x_2,x_4,x_2^*)=x_3^*,\\
\psi_4(x_3,x_4,x_2^*)=-x_2^*,\\
\psi_4(x_1,x_3,x_2^*)=-x_4^*,\\
\psi_4(x_1,x_4,x_2^*)=x_3^*,\\
\psi_4(x_3,x_4,x_2^*)=-x_1^*,\\
\psi_4(x_2,x_4,x_2^*)=-x_5^*,\\
\psi_4(x_4,x_5,x_2^*)=-x_2^*,\\
\psi_4(x_2,x_5,x_2^*)=x_4^*,\\
\psi_4(x_1,x_4,x_1^*)=-x_5^*,\\
\psi_4(x_4,x_5,x_1^*)=-x_1^*,\\
\psi_4(x_1,x_5,x_1^*)=x_4^*.
\end{cases}
\qquad
\begin{cases}
\widetilde{\Delta}^4(x_1^*)=x_1^*\wedge x_5^*\wedge x_4^*+\alpha x_2^*\wedge x_4^*\wedge x_3^*,\\
\widetilde{\Delta}^4(x_2^*)=x_2^*\wedge x_4^*\wedge x_3^*+x_1^*\wedge x_4^*\wedge x_3^*\\
\qquad\qquad +x_2^*\wedge x_5^*\wedge x_4^*,\\
\widetilde{\Delta}^4(x_1)=x_2\wedge x_3^*\wedge x_4^*+x_1\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^4(x_2)=\alpha x_1\wedge x_3^*\wedge x_4^*+x_2\wedge x_3^*\wedge x_4^*\\
\qquad\qquad +x_2\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^4(x_3)=\alpha x_1\wedge x_4^*\wedge x_2^*+x_2\wedge x_4^*\wedge x_2^*\\
\qquad\qquad +x_2\wedge x_4^*\wedge x_1^*,\\
\widetilde{\Delta}^4(x_4)=\alpha x_1\wedge x_2^*\wedge x_3^*+x_2\wedge x_2^*\wedge x_3^*\\
\qquad\qquad +x_1^*\wedge x_3^*\wedge x_2,+x_5^*\wedge x_2^*\wedge x_2\\
\qquad\qquad +x_5^*\wedge x_1^*\wedge x_1,\\
\widetilde{\Delta}^4(x_5)=x_2^*\wedge x_4^*\wedge x_2+x_1^*\wedge x_4^*\wedge x_1,\\
\widetilde{\Delta}^4(x_3^*)=\widetilde{\Delta}^4(x_4^*)=\widetilde{\Delta}^4(x_5^*)=0.
\end{cases}
$$

(36)
$$
\begin{cases}
\psi_5(x_2,x_3,x_4)=\alpha x_1+x_2,\\
\psi_5(x_1,x_3,x_4)=x_2,\\
\psi_5(x_2,x_3,x_1^*)=-\alpha x_4^*,\\
\psi_5(x_3,x_4,x_1^*)=-\alpha x_2^*,\\
\psi_5(x_2,x_4,x_1^*)=\alpha x_3^*,\\
\psi_5(x_2,x_3,x_2^*)=-x_4^*,\\
\psi_5(x_2,x_4,x_2^*)=x_3^*,\\
\psi_5(x_3,x_4,x_2^*)=-x_2^*-x_1^*,\\
\psi_5(x_1,x_3,x_2^*)=-x_4^*,\\
\psi_5(x_1,x_4,x_2^*)=x_3^*.
\end{cases}
\qquad
\begin{cases}
\widetilde{\Delta}^5(x_1^*)=\alpha x_2^*\wedge x_4^*\wedge x_3^*,\\
\widetilde{\Delta}^5(x_2^*)=x_1^*\wedge x_4^*\wedge x_3^*+x_2^*\wedge x_4^*\wedge x_3^*,\\
\widetilde{\Delta}^5(x_1)=\alpha x_2\wedge x_3^*\wedge x_4^*,\\
\widetilde{\Delta}^5(x_2)=x_1\wedge x_3^*\wedge x_4^*+x_2\wedge x_3^*\wedge x_4^*,\\
\widetilde{\Delta}^5(x_3)=x_1^*\wedge x_2\wedge x_4^*+x_2\wedge x_4^*\wedge x_2^*\\
\qquad\qquad +\alpha x_1\wedge x_4^*\wedge x_2^*,\\
\widetilde{\Delta}^5(x_4)=x_1^*\wedge x_3^*\wedge x_2+\alpha x_1\wedge x_2^*\wedge x_3^*\\
\qquad\qquad +x_2\wedge x_2^*\wedge x_3^*,\\
\widetilde{\Delta}^5(x_3^*)=\widetilde{\Delta}^5(x_4^*)=0,\\
\widetilde{\Delta}^5(x_5^*)=\widetilde{\Delta}^5(x_5)=0.
\end{cases}
$$

(37)
$$\begin{cases} \psi_6(x_2,x_3,x_4)=x_1,\\ \psi_6(x_1,x_3,x_4)=x_2,\\ \psi_6(x_2,x_4,x_5)=x_2,\\ \psi_6(x_1,x_4,x_5)=x_1,\\ \psi_6(x_2,x_3,x_1^*)=-x_4^*,\\ \psi_6(x_2,x_4,x_1^*)=x_3^*,\\ \psi_6(x_3,x_4,x_1^*)=-x_2^*,\\ \psi_6(x_1,x_3,x_2^*)=-x_4^*,\\ \psi_6(x_3,x_4,x_2^*)=-x_1^*,\\ \psi_6(x_1,x_4,x_2^*)=x_3^*,\\ \psi_6(x_2,x_4,x_2^*)=-x_5^*,\\ \psi_6(x_4,x_5,x_2^*)=-x_2^*,\\ \psi_6(x_2,x_5,x_2^*)=x_4^*,\\ \psi_6(x_1,x_4,x_1^*)=-x_5^*,\\ \psi_6(x_4,x_5,x_1^*)=-x_1^*,\\ \psi_6(x_1,x_5,x_2^*)=x_4^*. \end{cases} \quad \begin{cases} \widetilde{\Delta}^6(x_1^*)=x_2^*\wedge x_4^*\wedge x_3^*+x_1^*\wedge x_5^*\wedge x_4^*,\\ \widetilde{\Delta}^6(x_2^*)=x_1^*\wedge x_4^*\wedge x_3^*+x_2^*\wedge x_5^*\wedge x_4^*,\\ \widetilde{\Delta}^6(x_1)=x_1\wedge x_4^*\wedge x_5^*+x_2\wedge x_3^*\wedge x_4^*,\\ \widetilde{\Delta}^6(x_2)=x_1\wedge x_3^*\wedge x_4^*+x_2\wedge x_4^*\wedge x_5^*,\\ \widetilde{\Delta}^6(x_3)=x_1\wedge x_4^*\wedge x_2^*+x_2\wedge x_4^*\wedge x_1^*,\\ \widetilde{\Delta}^6(x_4)=x_1\wedge x_2^*\wedge x_3^*+x_2\wedge x_1^*\wedge x_3^*,\\ \qquad +x_5^*\wedge x_2^*\wedge x_2+x_5^*\wedge x_1^*\wedge x_1,\\ \widetilde{\Delta}^6(x_5)=x_2^*\wedge x_4^*\wedge x_2+x_1^*\wedge x_4^*\wedge x_1,\\ \widetilde{\Delta}^6(x_3^*)=\widetilde{\Delta}^6(x_4^*)=\widetilde{\Delta}^6(x_5^*)=0. \end{cases}$$

(38)
$$\begin{cases} \psi_7(x_2,x_3,x_4)=x_1,\\ \psi_7(x_2,x_4,x_5)=x_2,\\ \psi_7(x_3,x_4,x_5)=x_3,\\ \psi_7(x_2,x_3,x_1^*)=-x_4^*,\\ \psi_7(x_2,x_4,x_1^*)=x_3^*,\\ \psi_7(x_3,x_4,x_1^*)=-x_2^*,\\ \psi_7(x_2,x_4,x_2^*)=-x_5^*,\\ \psi_7(x_4,x_5,x_2^*)=-x_2^*,\\ \psi_7(x_2,x_5,x_2^*)=x_4^*,\\ \psi_7(x_3,x_4,x_3^*)=-x_5^*,\\ \psi_7(x_4,x_5,x_3^*)=-x_3^*,\\ \psi_7(x_3,x_5,x_3^*)=x_4^*. \end{cases} \quad \begin{cases} \widetilde{\Delta}^7(x_1^*)=x_2^*\wedge x_4^*\wedge x_3^*,\\ \widetilde{\Delta}^7(x_2^*)=x_2^*\wedge x_5^*\wedge x_4^*,\\ \widetilde{\Delta}^7(x_3^*)=x_3^*\wedge x_5^*\wedge x_4^*,\\ \widetilde{\Delta}^7(x_2)=x_1\wedge x_3^*\wedge x_4^*+x_2\wedge x_4^*\wedge x_5^*,\\ \widetilde{\Delta}^7(x_3)=x_1\wedge x_4^*\wedge x_2^*+x_3\wedge x_4^*\wedge x_5^*,\\ \widetilde{\Delta}^7(x_4)=x_1\wedge x_2^*\wedge x_3^*+x_2\wedge x_5^*\wedge x_2^*\\ \qquad +x_3\wedge x_5^*\wedge x_3^*,\\ \widetilde{\Delta}^7(x_5)=x_2\wedge x_2^*\wedge x_4^*+x_3\wedge x_3^*\wedge x_4^*,\\ \widetilde{\Delta}^7(x_4^*)=\widetilde{\Delta}^7(x_5^*)=\widetilde{\Delta}^7(x_1)=0. \end{cases}$$

(39)
$$\begin{cases} \psi_8(x_2,x_3,x_4)=x_2,\\ \psi_8(x_1,x_3,x_4)=x_1,\\ \psi_8(x_1,x_3,x_1^*)=-x_4^*,\\ \psi_8(x_1,x_4,x_1^*)=x_3^*,\\ \psi_8(x_3,x_4,x_1^*)=-x_1^*,\\ \psi_8(x_2,x_3,x_2^*)=-x_4^*,\\ \psi_8(x_2,x_4,x_2^*)=x_3^*,\\ \psi_8(x_3,x_4,x_2^*)=-x_2^*. \end{cases} \quad \begin{cases} \widetilde{\Delta}^8(x_1^*)=x_1^*\wedge x_4^*\wedge x_3^*,\\ \widetilde{\Delta}^8(x_1)=x_1\wedge x_3^*\wedge x_4^*,\\ \widetilde{\Delta}^8(x_2)=x_2\wedge x_3^*\wedge x_4^*,\\ \widetilde{\Delta}^8(x_3)=x_1\wedge x_4^*\wedge x_1^*+x_2\wedge x_4^*\wedge x_2^*,\\ \widetilde{\Delta}^8(x_4)=x_1\wedge x_1^*\wedge x_3^*+x_2\wedge x_2^*\wedge x_3^*,\\ \widetilde{\Delta}^8(x_2^*)=x_2^*\wedge x_4^*\wedge x_3^*,\\ \widetilde{\Delta}^8(x_3^*)=\widetilde{\Delta}^8(x_4^*)=0,\\ \widetilde{\Delta}^8(x_5^*)=\widetilde{\Delta}^8(x_5)=0. \end{cases}$$

(40)
$$\begin{cases} \psi_9(x_2,x_3,x_4)=x_1,\\ \psi_9(x_3,x_4,x_5)=x_3+\alpha x_2,\\ \psi_9(x_2,x_4,x_5)=x_3,\\ \psi_9(x_1,x_4,x_5)=x_1,\\ \psi_9(x_2,x_3,x_1^*)=-x_4^*,\\ \psi_9(x_2,x_4,x_1^*)=x_3^*,\\ \psi_9(x_3,x_4,x_1^*)=-x_2^*,\\ \psi_9(x_3,x_4,x_3^*)=-x_5^*,\\ \psi_9(x_4,x_5,x_3^*)=-x_3^*,\\ \psi_9(x_3,x_5,x_3^*)=x_4^*,\\ \psi_9(x_3,x_4,x_2^*)=-\alpha x_5^*,\\ \psi_9(x_4,x_5,x_2^*)=-\alpha x_3^*,\\ \psi_9(x_3,x_5,x_2^*)=\alpha x_4^*,\\ \psi_9(x_2,x_4,x_3^*)=-x_5^*,\\ \psi_9(x_4,x_5,x_3^*)=-x_2^*,\\ \psi_9(x_2,x_5,x_3^*)=x_4^*,\\ \psi_9(x_1,x_4,x_1^*)=-x_5^*,\\ \psi_9(x_4,x_5,x_1^*)=-x_1^*,\\ \psi_9(x_1,x_5,x_1^*)=x_4^*. \end{cases} \quad \begin{cases} \widetilde{\Delta}^9(x_1^*)=x_3^*\wedge x_2^*\wedge x_4^*+x_4^*\wedge x_1^*\wedge x_5^*,\\ \widetilde{\Delta}^9(x_2^*)=\alpha x_4^*\wedge x_3^*\wedge x_5^*,\\ \widetilde{\Delta}^9(x_3^*)=x_4^*\wedge x_3^*\wedge x_5^*+x_4^*\wedge x_2^*\wedge x_5^*,\\ \widetilde{\Delta}^9(x_1)=x_1\wedge x_4^*\wedge x_5^*,\\ \widetilde{\Delta}^9(x_2)=x_1\wedge x_3^*\wedge x_4^*+x_3\wedge x_4^*\wedge x_5^*,\\ \widetilde{\Delta}^9(x_3)=x_1\wedge x_4^*\wedge x_2^*+\alpha x_2\wedge x_4^*\wedge x_5^*\\ \qquad +x_3\wedge x_4^*\wedge x_5^*,\\ \widetilde{\Delta}^9(x_4)=x_1\wedge x_2^*\wedge x_3^*+x_3\wedge x_5^*\wedge x_3^*\\ \qquad +\alpha x_2\wedge x_5^*\wedge x_3^*+x_3\wedge x_5^*\wedge x_2^*\\ \qquad +x_1\wedge x_5^*\wedge x_1^*,\\ \widetilde{\Delta}^9(x_5)=x_3^*\wedge x_4^*\wedge x_3+\alpha x_3^*\wedge x_4^*\wedge x_2\\ \qquad +x_2^*\wedge x_4^*\wedge x_3+x_1^*\wedge x_4^*\wedge x_1,\\ \widetilde{\Delta}^9(x_4^*)=\widetilde{\Delta}^9(x_5^*)=0. \end{cases}$$

(41)
$$\begin{cases}
\psi_{10}(x_2,x_3,x_4)=x_1,\\
\psi_{10}(x_3,x_4,x_5)=x_3,\\
\psi_{10}(x_2,x_4,x_5)=x_2,\\
\psi_{10}(x_1,x_4,x_5)=2x_2,\\
\psi_{10}(x_2,x_3,x_1^*)=-x_4^*,\\
\psi_{10}(x_3,x_4,x_1^*)=-x_2^*,\\
\psi_{10}(x_2,x_4,x_1^*)=x_3^*,\\
\psi_{10}(x_3,x_4,x_3^*)=-x_5^*,\\
\psi_{10}(x_4,x_5,x_3^*)=-x_3^*,\\
\psi_{10}(x_3,x_5,x_3^*)=x_4^*,\\
\psi_{10}(x_2,x_4,x_2^*)=-x_5^*,\\
\psi_{10}(x_4,x_5,x_2^*)=-x_2^*,\\
\psi_{10}(x_2,x_5,x_2^*)=x_4^*,\\
\psi_{10}(x_1,x_4,x_2^*)=-2x_5^*,\\
\psi_{10}(x_4,x_5,x_2^*)=-2x_1^*,\\
\psi_{10}(x_1,x_5,x_2^*)=2x_4^*.
\end{cases}
\qquad
\begin{cases}
\widetilde{\Delta}^{10}(x_1^*)=x_3^*\wedge x_2^*\wedge x_4^*,\\
\widetilde{\Delta}^{10}(x_2^*)=x_4^*\wedge x_2^*\wedge x_5^*+2x_4^*\wedge x_1^*\wedge x_5^*,\\
\widetilde{\Delta}^{10}(x_3^*)=x_4^*\wedge x_3^*\wedge x_5^*,\\
\widetilde{\Delta}^{10}(x_1)=2x_2\wedge x_4^*\wedge x_5^*\\
\widetilde{\Delta}^{10}(x_2)=x_1\wedge x_3^*\wedge x_4^*+x_2\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^{10}(x_3)=x_1\wedge x_4^*\wedge x_2^*+x_3\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^{10}(x_4)=x_1\wedge x_2^*\wedge x_3^*+x_3\wedge x_5^*\wedge x_3^*,\\
\qquad\qquad +x_2\wedge x_5^*\wedge x_2^*+2x_2\wedge x_5^*\wedge x_1^*\\
\widetilde{\Delta}^{10}(x_5)=x_3\wedge x_3^*\wedge x_4^*+x_2\wedge x_2^*\wedge x_4^*\\
\qquad\qquad +2x_2\wedge x_1^*\wedge x_4^*,\\
\widetilde{\Delta}^{10}(x_4^*)=\widetilde{\Delta}^{10}(x_5^*)=0.
\end{cases}$$

(42)
$$\begin{cases}
\psi_{11}(x_2,x_3,x_4)=x_1,\\
\psi_{11}(x_1,x_3,x_4)=x_2,\\
\psi_{11}(x_1,x_2,x_4)=x_3,\\
\psi_{11}(x_2,x_3,x_1^*)=-x_4^*,\\
\psi_{11}(x_3,x_4,x_1^*)=-x_2^*,\\
\psi_{11}(x_2,x_4,x_1^*)=x_3^*,\\
\psi_{11}(x_1,x_3,x_2^*)=-x_4^*,\\
\psi_{11}(x_3,x_4,x_2^*)=-x_1^*,\\
\psi_{11}(x_1,x_4,x_2^*)=x_3^*,\\
\psi_{11}(x_1,x_4,x_3^*)=x_2^*,\\
\psi_{11}(x_1,x_2,x_3^*)=-x_4^*,\\
\psi_{11}(x_2,x_4,x_3^*)=-x_1^*.
\end{cases}
\qquad
\begin{cases}
\widetilde{\Delta}^{11}(x_1^*)=x_3^*\wedge x_2^*\wedge x_4^*,\\
\widetilde{\Delta}^{11}(x_2^*)=x_3^*\wedge x_1^*\wedge x_4^*,\\
\widetilde{\Delta}^{11}(x_3^*)=x_2^*\wedge x_1^*\wedge x_4^*,\\
\widetilde{\Delta}^{11}(x_1)=x_2\wedge x_3^*\wedge x_4^*+x_3\wedge x_2^*\wedge x_4^*,\\
\widetilde{\Delta}^{11}(x_2)=x_1\wedge x_3^*\wedge x_4^*+x_3\wedge x_4^*\wedge x_1^*,\\
\widetilde{\Delta}^{11}(x_3)=x_1\wedge x_4^*\wedge x_2^*+x_2\wedge x_4^*\wedge x_1^*,\\
\widetilde{\Delta}^{11}(x_4)=x_1\wedge x_2^*\wedge x_3^*+x_2\wedge x_1^*\wedge x_3^*\\
\qquad\qquad +x_3\wedge x_1^*\wedge x_2^*,\\
\widetilde{\Delta}^{11}(x_4^*)=\widetilde{\Delta}^{11}(x_5^*)=0,\\
\widetilde{\Delta}^{11}(x_5)=0.
\end{cases}$$

(43)
$$\begin{cases}
\psi_{12}(x_1,x_2,x_5)=x_1,\\
\psi_{12}(x_2,x_4,x_5)=x_3,\\
\psi_{12}(x_3,x_4,x_5)=\beta x_2\\
\qquad\qquad +(1+\beta)x_3,\\
\psi_{12}(x_1,x_4,x_1^*)=-x_5^*,\\
\psi_{12}(x_4,x_5,x_1^*)=-x_1^*,\\
\psi_{12}(x_1,x_5,x_1^*)=x_4^*,\\
\psi_{12}(x_2,x_4,x_3^*)=-x_5^*,\\
\psi_{12}(x_4,x_5,x_3^*)=-x_2^*,\\
\psi_{12}(x_2,x_5,x_3^*)=x_4^*,\\
\psi_{12}(x_3,x_4,x_2^*)=-\beta x_5^*,\\
\psi_{12}(x_4,x_5,x_2^*)=-\beta x_3^*,\\
\psi_{12}(x_3,x_5,x_2^*)=\beta x_4^*,\\
\psi_{12}(x_3,x_4,x_3^*)=-(1+\beta)x_5^*,\\
\psi_{12}(x_4,x_5,x_3^*)=-(1+\beta)x_3^*,\\
\psi_{12}(x_3,x_5,x_3^*)=(1+\beta)x_4^*.
\end{cases}
\quad
\begin{cases}
\widetilde{\Delta}^{12}(x_1^*)=x_4^*\wedge x_1^*\wedge x_5^*,\\
\widetilde{\Delta}^{12}(x_2^*)=\beta x_4^*\wedge x_3^*\wedge x_5^*,\\
\widetilde{\Delta}^{12}(x_3^*)=x_4^*\wedge x_2^*\wedge x_5^*\\
\qquad\qquad +(1+\beta)x_4^*\wedge x_3^*\wedge x_5^*,\\
\widetilde{\Delta}^{12}(x_1)=x_1\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^{12}(x_2)=x_3\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^{12}(x_3)=\beta x_2\wedge x_4^*\wedge x_5^*\\
\qquad\qquad +(1+\beta)x_3\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^{12}(x_4)=x_1\wedge x_5^*\wedge x_1^*+x_3\wedge x_5^*\wedge x_2^*\\
\qquad\qquad +\beta x_2\wedge x_5^*\wedge x_3^*\\
\qquad\qquad +(1+\beta)x_3\wedge x_5^*\wedge x_3^*,\\
\widetilde{\Delta}^{12}(x_5)=x_1\wedge x_1^*\wedge x_4^*+x_3\wedge x_2^*\wedge x_4^*\\
\qquad\qquad +\beta x_2\wedge x_3^*\wedge x_4^*\\
\qquad\qquad +(1+\beta)x_3\wedge x_3^*\wedge x_4^*,\\
\widetilde{\Delta}^{12}(x_4^*)=\widetilde{\Delta}^{12}(x_5^*)=0.
\end{cases}$$

(44)
$$\begin{cases}
\psi_{13}(x_1,x_4,x_5)=x_1,\\
\psi_{13}(x_2,x_4,x_5)=x_2,\\
\psi_{13}(x_3,x_4,x_5)=x_3,\\
\psi_{13}(x_1,x_4,x_1^*)=-x_5^*,\\
\psi_{13}(x_4,x_5,x_1^*)=-x_1^*,\\
\psi_{13}(x_1,x_5,x_1^*)=x_4^*,\\
\psi_{13}(x_2,x_4,x_2^*)=-x_5^*,\\
\psi_{13}(x_4,x_5,x_2^*)=-x_2^*,\\
\psi_{13}(x_2,x_5,x_2^*)=x_4^*,\\
\psi_{13}(x_3,x_4,x_3^*)=-x_5^*,\\
\psi_{13}(x_4,x_5,x_3^*)=-x_3^*,\\
\psi_{13}(x_3,x_5,x_3^*)=x_4^*.
\end{cases}
\quad
\begin{cases}
\widetilde{\Delta}^{13}(x_1^*)=x_4^*\wedge x_1^*\wedge x_5^*,\\
\widetilde{\Delta}^{13}(x_2^*)=x_4^*\wedge x_2^*\wedge x_5^*,\\
\widetilde{\Delta}^{13}(x_3^*)=x_4^*\wedge x_3^*\wedge x_5^*,\\
\widetilde{\Delta}^{13}(x_1)=x_1\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^{13}(x_2)=x_2\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^{13}(x_3)=x_3\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^{13}(x_4)=x_5^*\wedge x_1^*\wedge x_1+x_5^*\wedge x_2^*\wedge x_2\\
\qquad\qquad +x_5^*\wedge x_3^*\wedge x_3,\\
\widetilde{\Delta}^{13}(x_5)=x_1^*\wedge x_4^*\wedge x_1+x_2^*\wedge x_4^*\wedge x_2\\
\qquad\qquad +x_3^*\wedge x_4^*\wedge x_3,\\
\widetilde{\Delta}^{13}(x_4^*)=\widetilde{\Delta}^{13}(x_5^*)=0.
\end{cases}$$

(45)
$$\begin{cases}
\psi_{14}(x_1,x_4,x_5)=x_2,\\
\psi_{14}(x_2,x_4,x_5)=x_3,\\
\psi_{14}(x_3,x_4,x_5)=sx_1\\
+tx_2+ux_3,\\
\psi_{14}(x_1,x_4,x_2^*)=-x_5^*,\\
\psi_{14}(x_4,x_5,x_2^*)=-x_1^*,\\
\psi_{14}(x_1,x_5,x_2^*)=x_4^*,\\
\psi_{14}(x_2,x_4,x_3^*)=-x_5^*,\\
\psi_{14}(x_4,x_5,x_3^*)=-x_2^*,\\
\psi_{14}(x_2,x_5,x_3^*)=x_4^*,\\
\psi_{14}(x_3,x_4,x_1^*)=-sx_5^*,\\
\psi_{14}(x_4,x_5,x_1^*)=-sx_3^*,\\
\psi_{14}(x_3,x_5,x_1^*)=sx_4^*,\\
\psi_{14}(x_3,x_4,x_2^*)=-tx_5^*,\\
\psi_{14}(x_4,x_5,x_2^*)=-tx_3^*,\\
\psi_{14}(x_3,x_5,x_2^*)=tx_4^*,\\
\psi_{14}(x_3,x_4,x_3^*)=-ux_5^*,\\
\psi_{14}(x_4,x_5,x_3^*)=-ux_3^*,\\
\psi_{14}(x_3,x_5,x_3^*)=ux_4^*,
\end{cases}
\qquad
\begin{cases}
\widetilde{\Delta}^{14}(x_1^*)=sx_4^*\wedge x_3^*\wedge x_5^*,\\
\widetilde{\Delta}^{14}(x_2^*)=x_4^*\wedge x_1^*\wedge x_5^*+tx_4^*\wedge x_3^*\wedge x_5^*,\\
\widetilde{\Delta}^{14}(x_3^*)=x_4^*\wedge x_2^*\wedge x_5^*+ux_4^*\wedge x_3^*\wedge x_5^*,\\
\widetilde{\Delta}^{14}(x_1)=x_2\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^{14}(x_2)=x_3\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^{14}(x_3)=sx_1\wedge x_4^*\wedge x_5^*+tx_2\wedge x_4^*\wedge x_5^*\\
\qquad\qquad +ux_3\wedge x_4^*\wedge x_5^*,\\
\widetilde{\Delta}^{14}(x_4)=x_2\wedge x_5^*\wedge x_1^*+x_3\wedge x_5^*\wedge x_2^*\\
\qquad\qquad +sx_1\wedge x_5^*\wedge x_3^*+tx_2\wedge x_5^*\wedge x_3^*\\
\qquad\qquad +ux_3\wedge x_5^*\wedge x_3^*,\\
\widetilde{\Delta}^{14}(x_5)=x_2\wedge x_1^*\wedge x_4^*+x_3\wedge x_2^*\wedge x_4^*\\
\qquad\qquad +sx_1\wedge x_3^*\wedge x_4^*+tx_2\wedge x_3^*\wedge x_4^*\\
\qquad\qquad +ux_3\wedge x_3^*\wedge x_4^*,\\
\widetilde{\Delta}^{14}(x_4^*)=\widetilde{\Delta}^{14}(x_5^*)=0.
\end{cases}$$

(46)
$$\begin{cases}
\psi_{15}(x_2,x_3,x_4)=x_1,\\
\psi_{15}(x_1,x_3,x_4)=x_2,\\
\psi_{15}(x_2,x_3,x_1^*)=-x_4^*,\\
\psi_{15}(x_3,x_4,x_1^*)=-x_2^*,\\
\psi_{15}(x_2,x_4,x_1^*)=x_3^*,\\
\psi_{15}(x_1,x_3,x_2^*)=-x_4^*,\\
\psi_{15}(x_3,x_4,x_2^*)=-x_1^*,\\
\psi_{15}(x_1,x_4,x_2^*)=x_3^*.
\end{cases}
\qquad
\begin{cases}
\widetilde{\Delta}^{15}(x_1^*)=x_2^*\wedge x_4^*\wedge x_3^*,\\
\widetilde{\Delta}^{15}(x_2^*)=x_1^*\wedge x_4^*\wedge x_3^*,\\
\widetilde{\Delta}^{15}(x_1)=x_2\wedge x_3^*\wedge x_4^*,\\
\widetilde{\Delta}^{15}(x_2)=x_1\wedge x_3^*\wedge x_4^*,\\
\widetilde{\Delta}^{15}(x_3)=x_1\wedge x_4^*\wedge x_2^*+x_1^*\wedge x_2\wedge x_4^*,\\
\widetilde{\Delta}^{15}(x_4)=x_1\wedge x_2^*\wedge x_3^*+x_1^*\wedge x_3^*\wedge x_2,\\
\widetilde{\Delta}^{15}(x_3^*)=\widetilde{\Delta}^{15}(x_4^*)=0,\\
\widetilde{\Delta}^{15}(x_5)=\widetilde{\Delta}^{15}(x_5^*)=0.
\end{cases}$$

$$
\begin{cases}
\psi_{16}(x_2,x_3,x_4)=-x_2,\\
\psi_{16}(x_1,x_3,x_4)=x_1,\\
\psi_{16}(x_1,x_2,x_3)=x_3,\\
\psi_{16}(x_1,x_2,x_4)=-x_4,\\
\psi_{16}(x_2,x_3,x_2^*)=x_4^*,\\
\psi_{16}(x_3,x_4,x_2^*)=x_2^*,\\
\psi_{16}(x_2,x_4,x_2^*)=-x_3^*,\\
\psi_{16}(x_1,x_3,x_1^*)=-x_4^*,\\
\psi_{16}(x_3,x_4,x_1^*)=-x_1^*,\\
\psi_{16}(x_1,x_4,x_1^*)=x_3^*,\\
\psi_{16}(x_1,x_2,x_3^*)=-x_3^*,\\
\psi_{16}(x_1,x_3,x_3^*)=x_2^*,\\
\psi_{16}(x_2,x_3,x_3^*)=-x_1^*,\\
\psi_{16}(x_1,x_2,x_4^*)=x_4^*,\\
\psi_{16}(x_1,x_4,x_4^*)=-x_2^*,\\
\psi_{16}(x_2,x_4,x_4^*)=x_1^*.
\end{cases}
\qquad
\begin{cases}
\widetilde{\Delta}^{16}(x_1^*)=x_1^*\wedge x_4^*\wedge x_3^*,\\
\widetilde{\Delta}^{16}(x_2^*)=x_2^*\wedge x_3^*\wedge x_4^*,\\
\widetilde{\Delta}^{16}(x_3^*)=x_1^*\wedge x_3^*\wedge x_2^*,\\
\widetilde{\Delta}^{16}(x_4^*)=x_1^*\wedge x_2^*\wedge x_4^*,\\
\widetilde{\Delta}^{16}(x_1)=x_1\wedge x_3^*\wedge x_4^*+x_2^*\wedge x_3^*\wedge x_3\\
\qquad\qquad +x_4^*\wedge x_2^*\wedge x_4,\\
\widetilde{\Delta}^{16}(x_2)=x_3\wedge x_3^*\wedge x_1^*+x_4^*\wedge x_3^*\wedge x_2\\
\qquad\qquad +x_4\wedge x_1^*\wedge x_4^*,\\
\widetilde{\Delta}^{16}x_3)=x_1^*\wedge x_2^*\wedge x_3+x_2^*\wedge x_4^*\wedge x_2\\
\qquad\qquad +x_4^*\wedge x_1^*\wedge x_1,\\
\widetilde{\Delta}^{16}(x_4)=x_1^*\wedge x_4\wedge x_2^*+x_1\wedge x_1^*\wedge x_3^*\\
\qquad\qquad +x_2\wedge x_3^*\wedge x_2^*,\\
\widetilde{\Delta}^{16}(x_5^*)=\widetilde{\Delta}^{16}(x_5)=0.
\end{cases}
\tag{47}
$$

$$
\begin{cases}
\psi_{17}(x_1,x_2,x_3)=x_1,\\
\psi_{17}(x_1,x_2,x_1^*)=-x_3^*,\\
\psi_{17}(x_2,x_3,x_1^*)=-x_1^*,\\
\psi_{17}(x_1,x_3,x_1^*)=x_2^*.
\end{cases}
\qquad
\begin{cases}
\widetilde{\Delta}^{17}(x_1^*)=x_1^*\wedge x_3^*\wedge x_2^*,\\
\widetilde{\Delta}^{17}(x_1)=x_1\wedge x_2^*\wedge x_3^*,\\
\widetilde{\Delta}^{17}(x_2)=x_3^*\wedge x_1^*\wedge x_1,\\
\widetilde{\Delta}^{17}(x_3)=x_1^*\wedge x_2^*\wedge x_1,\\
\widetilde{\Delta}^{17}(x_2^*)=\widetilde{\Delta}^{17}(x_3^*)=\widetilde{\Delta}^{17}(x_4^*)=0,\\
\widetilde{\Delta}^{17}(x_5^*)=\widetilde{\Delta}^{17}(x_4)=\widetilde{\Delta}^{17}(x_5)=0.
\end{cases}
\tag{48}
$$

Proof. Apply Lemma 4.1, and Eq(5), and Theorem 3.4 and Theorem 3.3. □

Acknowledgements. The first named author was supported in part by the Natural Science Foundation (11371245) and the Natural Science Foundation of Hebei Province (A2018201126).

References

[1] R. Bai, L. Lin and W. Guo, q-Deformation of 3-Lie algebras, *Algebra Colloquium,* 2017, 24:3: 519-540

[2] R. Bai, Y. Cheng and J. Li, 3-Lie bialgebras, *Acta Mathematica Scientia*, 2014, 34B(2):513-522

[3] R. Bai, W. Guo, L. Lin and Y. Zhang, n-Lie bialgebras, *Linear and Multilinear Algebra*, 2018, 66(2): 382-397.

[4] C. Bai, L. Guo and Y. Sheng, Bialgebra classical yang-baxter equation and manin triples for 3-Lie algebras, *J. Phys. A, Math. Theor.,* 2016, 42, 035207

[5] R. Bai, G. Song and Y. Zhang, On classification of n-Lie algebras, *Front. Math. China*, 2011, 6(4): 581-606

[6] V. Chari and A. Pressley, A guide to quantum groups. Cambridge University Press, Cambridge (1994). 2, 3

[7] V. Drinfeld, Hamiltonian structure on the Lie groups, Lie bialgebras and the geometric sense of the classical Yang-Baxter equations. Soviet Math. Dokl. 1983, 27: 68-71.

College of Mathematics and Information Science, Hebei University, Key Laboratory of Machine Learning and Computational, Intelligence of Hebei Province, Baoding 071002, P.R. China
E-mail address: bairuipu@hbu.edu.cn

College of Mathematics and Information Science, Hebei University, Baoding 071002, China
E-mail address: hshuaisun@163.com

College of Mathematics and Information Science, Hebei University, Baoding 071002, China
E-mail address: kangchuang2016@163.com