arXiv:2306.02973v1 [math.AP] 5 Jun 2023

SIGN CHANGING BUBBLE TOWER SOLUTIONS TO A SLIGHTLY SUBCRITICAL ELLIPTIC PROBLEM WITH NON-POWER NONLINEARITY

SHENGBING DENG AND FANG YU

Abstract. We study the following elliptic problem involving slightly subcritical non-power nonlinearity

$$\begin{cases} -\Delta u = \frac{|u|^{2^*-2}u}{[\ln(e+|u|)]^{\varepsilon}} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded smooth domain in $\mathbb{R}^n$, $n \geq 3$, $2^* = \frac{2n}{n-2}$ is the critical Sobolev exponent, $\varepsilon > 0$ is a small parameter. By the finite dimensional Lyapunov-Schmidt reduction method, we construct a sign changing bubble tower solution with the shape of a tower of bubbles as ε goes to zero.

1. Introduction

In this paper, we consider the following elliptic problem involving slightly subcritical non-power nonlinearity

$$\begin{cases} -\Delta u = \frac{|u|^{2^*-2}u}{[\ln(e+|u|)]^{\varepsilon}} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \tag{1.1}$$

where Ω is a bounded smooth domain in $\mathbb{R}^n$, $n \geq 2$, $2^* = \frac{2n}{n-2}$ is the critical Sobolev exponent for the embedding $H_0^1(\Omega) \hookrightarrow L^{2^*}(\Omega)$, $\varepsilon > 0$ is a small parameter.

The main feature of problem (1.1) is the non-power type nonlinearity, which is first proposed by Castro and Pardo [12], they proved the existence of a priori L^∞ bounds for positive solutions of Laplacian problem involving the nonlinearity $f(u) = \frac{u^{\frac{n+2}{n-2}}}{\ln(2+u)^\alpha}$ with $\alpha > \frac{2}{n-2}$. Then, Mavinga and Pardo [36] obtained a priori estimates for positive classical solutions to the following Hamiltonian elliptic system

$$\begin{cases} -\Delta u = \frac{v^p}{[\ln(e+v)]^\alpha} & \text{in } \Omega, \\ -\Delta v = \frac{u^q}{[\ln(e+u)]^\beta} & \text{in } \Omega, \\ u = v = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded convex domain with smooth boundary in $\mathbb{R}^n$ for $n > 2$, $1 < p, q < \infty$ and $\alpha, \beta > 0$, $\frac{1}{p+1} + \frac{1}{q+1} = \frac{n-2}{n}$. For more results of non-power nonlinearity, we refer to [13, 22, 39] for slightly subcritical problem and [26] for supercritical problem.

On the one hand, problem (1.1) is related to the following slightly subcritical elliptic problem

$$\begin{cases} -\Delta u = |u|^{2^*-2-\varepsilon}u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases} \tag{1.2}$$

2020 *Mathematics Subject Classification.* 35B33; 35B40; 35J15.

Key words and phrases. non-power nonlinearity; sign changing bubble tower solutions; Lyapunov-Schmidt reduction.

When $\varepsilon = 0$, Pohozaev [41] proved that the non existence of nontrivial solution if Ω is a star-sharped domain. When Ω is a annulus, Kazdan and Warner [33] obtained the existence of a positive radial solution. Bahri and Coron [1] studied a positive solution provided that Ω has nontrivial topology. For the existence of sign changing solution, there are few results. When $\Omega = \mathbb{R}^n$, Ding [27] showed the infinitely many sign changing solutions by Ljusternik-Schnirlman category theory. In specific case like torii, Hebey and Vaugon [32] investigated the existence and multiple sign changing solutions. The existence and multiplicity of sign changing solutions are also treated in some contractible domains with an involution symmetry by Clapp and Weth [19].

When ε is a positive parameter, problem (1.2) has a positive least energy solution u_ε, that is, u_ε is a solution for the variational problem

$$\inf\Big\{\|u\|^2 = \int_\Omega |\nabla u|^2 dx : u \in H_0^1(\Omega), \int_\Omega |u|^{2^*-\varepsilon} dx = 1\Big\}.$$

The blow-up phenomenon for positive and sign changing solutions to (1.2) has been studied extensively. When ε goes to 0, Rey [42] and Han [31] studied that the solution to (1.2) blows up and concentrates at a critical point of Robin function. Moreover, Flucher and Wei [28] proved that the concentration point is the minimum point of the Robin function. Furthermore, if ξ^* is a stable critical point of Robin function, then (1.2) has a positive solution which blows up at ξ^*, this result is obtained in [37,42]. For the multiple concentration points, Rey [43] showed that the two blow up and concentration points (ξ_1^*, ξ_2^*), which is a critical point of a function involving Robin function and Green's function. If the domain is convex, Grossi and Takahashi in [30] proved that (1.2) does not admit any positive solution blowing up more than two points. The positive solution to (1.2) concentrate simultaneously at different points $\xi_1, \cdots, \xi_k \in \Omega$, $k \geq 2$, has been established in [2,37]. If any ξ_i, $i = 1, \cdots, k$, is a simple blow up point, Li [34] characterized the form of solution u_ε near each blow up point ξ_i as

$$u_\varepsilon(x) \sim \frac{\mu_i \sqrt{\varepsilon}}{(\mu_i^2 \varepsilon + |x - \xi_i|^2)^{\frac{n-2}{2}}}, \quad \text{with } \mu_i > 0.$$

On the other hand, the existence of one sign changing solution to (1.2) is first proved in [5,11], and multiple sign changing solutions with their nodal properties are treated in [6,7] for $\varepsilon \in (0, \frac{4}{n-2})$. Moreover, they proved that (1.2) has a least energy nodal solution with two nodal domains. Ben Ayed et al. [8] obtained that the low energy sign-changing solutions blow up at two points, and the energy converges to the value $2S^{\frac{n}{2}}$, where S is the Sobolev constant for the embedding $H_0^1(\Omega)$ into $L^{\frac{2n}{n-2}}(\Omega)$. Bartsch et al. [4] considered that (1.2) has k pairs of sign changing solutions $\pm u_\varepsilon^{(i)}$, $i = 1, \cdots, k$, which satisfies that $u_\varepsilon^{(i)}$ blows up positively at a point $\xi_1^{(i)} \in \Omega$ and $-u_\varepsilon^{(i)}$ blows up negatively at a point $\xi_2^{(i)} \in \Omega$ with $\xi_1^{(i)} \neq \xi_2^{(i)}$. Bartsch et al. [3] proved a sign changing four-bubble solution with two positive and two negative blow-up points provided that Ω is convex and satisfies some symmetry conditions.

In contrast to the result of positive and sign changing solutions to (1.2), there are some papers of bubble tower. If Ω is a smooth bounded domain in $\mathbb{R}^n$ symmetric with respect to $x_1, \cdots, x_n$ and contains the origin, Pistoia and Weth [40] constructed a sign changing bubble tower solution u_ε concentrating at the center of symmetry of Ω. The same consequence in any bounded smooth domain is considered in [38], and they removed the assumption on non-degeneracy of critical point of Robin's function. If the domain has holes like $\Omega \setminus B(a, \varepsilon) \cup B(b, \varepsilon)$ with center at point a, b and radius $\varepsilon > 0$, Ge et al. [29] constructed sign changing solutions blowing up both at a and b. For any other bubble tower results of elliptic problem, see [14,15,17,20,24] and references therein. In particular, we refer to the papers [21,23] for fractional and biharmonic operators involving almost critical Soblev exponent.

Recently, by Lyapunov-Schmidt reduction method, Clapp et al. [9,18] constructed solutions to problem (1.1). Before stating the results, let us introduce some definitions and notations. For $\xi \in \Omega$ and $\mu > 0$, let

$$U(x) = \alpha_n \frac{1}{(1+|x|^2)^{\frac{n-2}{2}}}, \quad U_{\mu,\xi}(x) = \frac{\alpha_n \mu^{\frac{n-2}{2}}}{(\mu^2 + |x-\xi|^2)^{\frac{n-2}{2}}}, \quad \text{with } \alpha_n = (n(n-2))^{\frac{n-2}{4}}, \tag{1.3}$$

which are the only solutions of the equation

$$-\Delta u = u^{2^*-1}, \quad u > 0 \quad \text{in} \quad \mathbb{R}^n. \tag{1.4}$$

Let us denote by $G(x,y)$ the Green's function of $-\Delta$ in Ω with Dirichlet boundary condition, and by $H(x,y)$ its regular part, so that

$$H(x,y) = \frac{1}{(n-2)|\partial B|}\Big(\frac{1}{|x-y|^{n-2}} - G(x,y)\Big), \quad \text{for every} \quad x, y \in \Omega,$$

where $|\partial B|$ denotes the surface area of the unit sphere in $\mathbb{R}^n$. The Robin function is defined as $\varphi(x) = H(x,x)$ for every $x \in \Omega$. Let ξ_* is a non-degenerate critical point of Robin function, Clapp et al. [18] constructed a single bubble solution of the form

$$u_\varepsilon = U_{\mu_\varepsilon,\xi_\varepsilon} + \phi_\varepsilon,$$

with $\mu_\varepsilon\Big(\frac{|\ln\varepsilon|}{\varepsilon}\Big)^{\frac{1}{n-2}} \to d > 0$, $\xi_\varepsilon \to \xi_*$, $\phi_\varepsilon \in H_0^1(\Omega)$ such that $\int_\Omega |\nabla\phi_\varepsilon|^2 dx = O\Big(\frac{\varepsilon}{|\ln\varepsilon|}\Big)$ as $\varepsilon \to 0$. Liu et al. [35] established a solution concentrating at the origin point for a critical Hénon problem with non-power type. Ben Ayed et al. [9] obtained positive as well as sign changing solutions concentrating at several points, which involving Robin function and Green's function.

In present paper, motivated by several results [9, 18, 38, 40], we construct a solution with the shape of a tower of sign changing bubbles to problem (1.1) by finite Lyapunov-Schmidt dimensional reduction procedure.

Our result can be stated as follows.

Theorem 1.1. *Assume that $n \geq 3$, for any integer $k \geq 1$, there exists $\varepsilon_0 > 0$ such that for every $\varepsilon \in (0, \varepsilon_0)$, there are some points $\xi_{i_\varepsilon} \in \Omega$ and positive constants d_{i_ε} for $i = 1, \cdots k$, problem (1.1) has a solution u_ε of the form,*

$$u_\varepsilon(x) = \alpha_n \sum_{i=1}^{k} (-1)^i \left(\frac{d_{i_\varepsilon} \big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{2i-1}{n-2}}}{\Big(d_{i_\varepsilon}\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{2i-1}{n-2}}\Big)^2 + |x-\xi_{i_\varepsilon}|^2} \right)^{\frac{n-2}{2}} + \Theta_\varepsilon(x),$$

where $\|\Theta_\varepsilon\| \to 0$ as $\varepsilon \to 0$, $\varphi(\xi_{i_\varepsilon}) \to \min\limits_{z\in\Omega} \varphi(z)$ and $d_{i_\varepsilon} \to d_i > 0$ for $i = 1, \cdots k$.

Observe that in the above construction, the solutions behaves like a superposition of bubbles of different blow up orders centered at around the minimum point of the Robin function, thus, it is called bubble tower solutions. It was first studied by del Pino et al. [25] for a slightly supercritical Brezis-Nirenberg problem in a ball, and this type solutions has been constructed in many problems, see [16, 21, 23, 29, 38, 40] and references therein.

The paper is organized as follows. In Section 2, we give the scheme of proof for Theorem 1.1. We show the finite dimensional reduction process in Section 3. Proposition 2.2 is proved in Section 4. Finally, there are some estimates in the Appendix. We will use $C > 0$ to denote various positive constants.

2. Scheme of the proof

In this section, let us give the sketch proof of Theorem 1.1. We first introduce some notations. The Sobolev space $H_0^1(\Omega)$ is endowed with inner product $\langle\cdot,\cdot\rangle$ defined by

$$\langle u, v\rangle = \int_\Omega \nabla u \nabla v dx,$$

for all u, $v \in H_0^1(\Omega)$, and $L^p(\Omega)$ is the Lebesgue space with the norm $|u|_q = \Big(\int_\Omega |u|^q dx\Big)^{\frac{1}{q}}$, for $1 < q < \infty$.

Let $i^* : L^{\frac{2n}{n+2}}(\Omega) \hookrightarrow H_0^1(\Omega)$ be the adjoint operator of the embedding $i : H_0^1(\Omega) \hookrightarrow L^{\frac{2n}{n-2}}(\Omega)$, that is, for $v \in L^{\frac{2n}{n+2}}(\Omega)$, $u = i^*(v)$ if and only if

$$-\Delta u = v \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega.$$

Then, it holds

$$\|i^*(v)\| \le c|u|_{\frac{2n}{n+2}}, \tag{2.1}$$

for some constant $c > 0$ depending only on Ω and n. Using these definitions and notations, problem (1.1) is equivalent to the following equation

$$u = i^*[f_\varepsilon(u)], \quad u \in H_0^1(\Omega),$$

where $f_\varepsilon(u) = \frac{|u|^{2^*-2}u}{[\ln(e+|u|)]^\varepsilon}$.

In order to describe the shape of the solutions to problem (1.1), we give an integer number k, and define the positive parameters μ_i as

$$\mu_i = \Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{2i-1}{n-2}} d_i, \quad \text{with} \quad d_i > 0, \quad i = 1, \cdots, k. \tag{2.2}$$

Let ξ be a point in Ω, $\xi_i \in \Omega$, $i = 1, \cdots, k$, is given by

$$\xi_i = \xi + \mu_i \sigma_i, \quad \text{for some points } \sigma_i \in \mathbb{R}^n, \tag{2.3}$$

where $\sigma_k = 0$. We will assume the following bounds on the parameters and points appearing in (2.2) and (2.3): given $\eta > 0$ small,

$$\operatorname{dist}(\xi, \partial\Omega) > \eta, \quad \eta < d_i < \frac{1}{\eta}, \quad |\sigma_i| \le \frac{1}{\eta}, \quad i = 1, \cdots, k. \tag{2.4}$$

It is an immediate observation that

$$\mu_1 = \Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{1}{n-2}} d_1 \quad \text{and} \quad \frac{\mu_{i+1}}{\mu_i} = \Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{2}{n-2}} \frac{d_{i+1}}{d_i}.$$

We denote by $PU_{\mu,\xi}$ the projection onto $H_0^1(\Omega)$ of $U_{\mu,\xi}$, that is

$$\begin{cases} -\Delta PU_{\mu,\xi} = -\Delta U_{\mu,\xi} & \text{in } \Omega, \\ PU_{\mu,\xi} = 0 & \text{on } \partial\Omega. \end{cases}$$

Let $k \ge 1$, the approximate solutions are given by

$$u(x) = V(x) + \phi(x), \quad V(x) = V_{\bar d, \bar\sigma, \xi}(x) = \sum_{i=1}^k (-1)^i PU_{\mu_i,\xi_i}(x), \tag{2.5}$$

where

$$\bar d = (d_1, \cdots, d_k) \in \mathbb{R}_+^k, \ \bar\sigma = (\sigma_1, \cdots, \sigma_k) \in (\mathbb{R}_+^n)^k. \tag{2.6}$$

The term ϕ is small in some sense.

Let us describe ϕ. As it is shown in [10], any solution of

$$-\Delta\psi = f_0^{'}(U_{\mu,\xi})\psi \quad \text{in } \mathbb{R}^n, \tag{2.7}$$

can be expressed as a linear combination of

$$\psi^0(y) = \frac{(n-2)\alpha_n}{2}\frac{|y|^2-1}{(1+|y|^2)^{\frac{n}{2}}}, \quad \psi^h(y) = (n-2)\alpha_n\frac{y_h}{(1+|y|^2)^{\frac{n}{2}}}, \quad \text{for } h = 1,\cdots,n. \tag{2.8}$$

Moreover, we set

$$\begin{aligned}
\psi^0_{\mu,\xi}(x) &= \mu^{-\frac{n-2}{2}}\psi^0(\frac{x-\xi}{\mu}) = \frac{n-2}{2}\alpha_n\mu^{\frac{n-2}{2}}\frac{|x-\xi|^2-\mu^2}{(\mu^2+|x-\xi|^2)^{\frac{n}{2}}},\\
\psi^h_{\mu,\xi}(x) &= \mu^{-\frac{n-2}{2}}\psi^h(\frac{x-\xi}{\mu}) = (n-2)\alpha_n\mu^{\frac{n}{2}}\frac{x_h-\xi_h}{(\mu^2+|x-\xi|^2)^{\frac{n}{2}}}, \text{ for } h = 1,\cdots,n,
\end{aligned} \tag{2.9}$$

then

$$\psi^0_{\mu,\xi}(x) = \mu\frac{\partial U_{\mu,\xi}}{\partial\mu}, \quad \psi^h_{\mu,\xi}(x) = \mu\frac{\partial U_{\mu,\xi}}{\partial\xi_h}. \tag{2.10}$$

We denote that $P\psi^h_{\mu,\xi}$ is the projection of $\psi^h_{\mu,\xi}$, $h = 0,\cdots,n$, and define the subspace of $H^1_0(\Omega)$

$$\begin{aligned}
E_{\mu,\xi} &= \text{span}\left\{P\psi^h_{\mu,\xi} : h = 0,1,\cdots,n,\ i = 1,\cdots,k\right\},\\
E^{\perp}_{\mu,\xi} &= \left\{\phi\in H^1_0(\Omega) : \langle\phi, P\psi^h_{\mu,\xi}\rangle = 0 : h = 0,1,\cdots,n,\ i = 1,\cdots,k\right\}.
\end{aligned}$$

Let

$$\Pi_{\mu,\xi} : H^1_0(\Omega)\to E_{\mu,\xi} \quad \text{and} \quad \Pi^{\perp}_{\mu,\xi} : H^1_0(\Omega)\to E^{\perp}_{\mu,\xi},$$

be the corresponding projections. To solve (1.1), it is equivalent to solve the couple of following equations

$$\Pi^{\perp}_{\mu,\xi}\Big(V+\phi-i^*[f_\varepsilon(V+\phi]\Big) = 0, \tag{2.11}$$

and

$$\Pi_{\mu,\xi}\Big(V+\phi-i^*[f_\varepsilon(V+\phi)]\Big) = 0. \tag{2.12}$$

We solve equation (2.11) in the following result, whose proof can be found in Section 3.

Proposition 2.1. *There exists $\varepsilon_0 > 0$ such that for any $\xi\in\Omega$, $\bar{d}\in\mathbb{R}^k_+$, $\bar{\sigma}\in(\mathbb{R}^n_+)^k$ satisfying (2.4), for $\varepsilon\in(0,\varepsilon_0)$, there is a unique function $\phi\in E^{\perp}_{\mu,\xi}$ which solves (2.11). Moreover*

$$\|\phi\| = \begin{cases} O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big) & \text{if } 3\le n\le 6,\\ O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n+2}{2(n-2)}}\Big) & \text{if } n\ge 7. \end{cases} \tag{2.13}$$

From Proposition 2.1, there is a unique $\phi\in E^{\perp}_{\mu,\xi}$ such that (2.11) holds, it means that there are some constants $c_{il}s$ $(i = 1,\cdots,k,\ l = 0,\cdots,n)$ such that

$$V+\phi-i^*[f_\varepsilon(V+\phi)] = \sum_{i=1}^{k}\sum_{l=0}^{n}c_{il}P\psi^l_{\mu_i,\xi_i}, \tag{2.14}$$

which equals to solve equation (2.12), that is, the following result is valid, whose proof is postponed to Section 4.

Proposition 2.2. *The following facts hold.*
Part a. If $(\bar{d}_\varepsilon, \bar{\sigma}_\varepsilon, \xi_\varepsilon)$ *satisfies*

$$\Big\langle V+\phi-i^*[f_\varepsilon(V+\phi)], P\psi^h_{\mu_{j_\varepsilon},\xi_{j_\varepsilon}}\Big\rangle = 0, \quad \text{for } h=0,\cdots,n, \tag{2.15}$$

where $j=1,\cdots,k$. *Then* $V+\phi$ *is a solution of problem (1.1).*
Part b. For $\xi\in\Omega$, $\bar{d}=(d_1,\cdots,d_k)\in\mathbb{R}^k_+$, $\bar{\sigma}=(\sigma_1,\cdots,\sigma_k)\in(\mathbb{R}^n_+)^k$, *there holds*

$$\Big\langle V+\phi-i^*[f_\varepsilon(V+\phi)], P\psi^h_{\mu_j,\xi_j}\Big\rangle = \begin{cases} \frac{\varepsilon}{|\ln\varepsilon|^2}G^\varepsilon_0(\bar{d},\bar{\sigma},\xi) - \frac{2k^2}{(n-2)^2}a_4\varepsilon\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big| & \text{for } h=0,\\ \frac{\varepsilon}{|\ln\varepsilon|^2}G^\varepsilon_h(\bar{d},\bar{\sigma},\xi) & \text{for } h=1,\cdots,n,\end{cases}$$

where $j=1,\cdots,k$, *and* $G^\varepsilon=(G^\varepsilon_0,G^\varepsilon_h)$ *is given by*

$$\begin{cases} G^\varepsilon_0(\bar{d},\bar{\sigma},\xi)=\alpha_n a_1 d_1^{n-2}\varphi(\xi)+a_3\sum\limits_{i=1}^{k-1}\Big(\frac{d_{i+1}}{d_i}\Big)^{\frac{n-2}{2}}g(\sigma_i)-a_4\sum\limits_{i=1}^{k}\frac{2}{2i-1}|\ln d_i|+o(1),\\ G^\varepsilon_h(d_1,\xi)=\frac{\alpha_n}{2}a_2\partial_{\xi_h}\varphi(\xi)d_1^{n-1} \quad \text{for } h=1,\cdots,n,\end{cases}$$

with $G^\varepsilon_0:[0,+\infty]\times[0,+\infty]\times\Omega\to\mathbb{R}\times\mathbb{R}\times\mathbb{R}^n$, $G^\varepsilon_h:[0,+\infty]\times\Omega\to\mathbb{R}\times\mathbb{R}^n$ *and*

$$\begin{aligned} a_1&=(2^*-1)\int_{\mathbb{R}^n}U^{2^*-2}(y)\psi^0(y)dy,\\ a_2&=\int_{\mathbb{R}^n}U^{2^*-1}(y)dy,\\ a_3&=\frac{n-2}{2}\alpha_n^{p+1},\\ a_4&=\int_{\mathbb{R}^n}\Big|\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}\ln\Big(\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}\Big)\psi^0(y)\Big|dy>0,\\ g(\sigma)&=\int_{\mathbb{R}^n}\frac{y^{2-n}}{(1+|y-\sigma|^2)^{\frac{n+2}{2}}}dy.\end{aligned}$$

From Propositions 2.1 and 2.2, we view that $V+\phi$ is the solution to problem (1.1) if there are $d_\varepsilon>0$, $\sigma_\varepsilon>0$ and $\xi_\varepsilon\in\Omega$ such that $c_{il}(d_\varepsilon,\sigma_\varepsilon,\xi_\varepsilon)$ are zero when ε small enough.

The sequel of this section is devoted to the proof of the main result.

Proof of Theorem 1.1. By Proposition 2.2, equation (2.15) is equivalent to

$$\begin{cases}\alpha_n a_1 d_1^{n-2}\varphi(\xi)+a_3\sum\limits_{i=1}^{k-1}\Big(\frac{d_{i+1}}{d_i}\Big)^{\frac{n-2}{2}}g(\sigma_i)-a_4\sum\limits_{i=1}^{k}\frac{2}{2i-1}|\ln d_i|=o(1),\\ \frac{\alpha_n}{2}a_2\partial_{\xi_h}\varphi(\xi)d_1^{n-1}=0,\end{cases} \tag{2.16}$$

for $h=1,\cdots,n$. We note that $G^\varepsilon\to G$ uniformly on compact set of $[0,+\infty]\times[0,+\infty]\times\Omega$, and the vector functional $G(\bar{d},\bar{\sigma},\xi)=\Big(G_0(\bar{d},\bar{\sigma},\xi),G_h(d_1,\xi)\Big)$ is the principal part of defined by

$$\begin{aligned} G_0(\bar{d},\bar{\sigma},\xi)&=\alpha_n a_1 d_1^{n-2}\varphi(\xi)+a_3\sum_{i=1}^{k-1}\Big(\frac{d_{i+1}}{d_i}\Big)^{\frac{n-2}{2}}g(\sigma_i)-a_4\sum_{i=1}^{k}\frac{2}{2i-1}|\ln d_i|,\\ G_h(d_1,\xi)&=\frac{\alpha_n}{2}a_2\partial_{\xi_h}\varphi(\xi)d_1^{n-2}, \quad \text{for } h=1,\cdots,n.\end{aligned}$$

Let us set $s_1=d_1$, $s_i=\frac{d_i}{d_{i-1}}$, $i=2,\cdots,k$, then in the new variables $\bar{s}=(s_1,\cdots,s_k)$, $G_h(d_1,\xi)$ and $G_0(\bar{d},\bar{\sigma},\xi)$ can be rewrite as

$$\bar{G}_h(s_1,\xi)=\frac{\alpha_n}{2}a_2\partial_{\xi_h}\varphi(\xi)s_1^{n-2}, \quad \text{for} \quad h=1,\cdots,n,$$

$$\bar{G}_0(\bar{s},\bar{\sigma},\xi) =\alpha_n a_1 s_1^{n-2}\varphi(\xi) + a_3\sum_{i=2}^{k} s_i^{\frac{n-2}{2}} g(\sigma_i) - a_4\sum_{i=1}^{k}\frac{2}{2i-1}|\ln s_i|.$$

We denote $\bar{G} = \Big(\bar{G}_0(\bar{s},\bar{\sigma},\xi), \bar{G}_h(s_1,\xi)\Big)$. Let $\xi^0 \in \Omega$ be a strict minimum point of Robin function φ, which is the zero point of function $\bar{G}_h$ for $h = 1,\cdots,n$. Observe that $\sigma_i = 0$ is a strict minimum point of g. On the other hand, when s_i is close to 0, the function $\bar{G}_0$ tends to $-\infty$, and $\bar{G}_0 > 0$ as $s_i > 0$ large enough, thus, by intermediate value theorem, there exists a $\bar{s}_0$ such that $\bar{G}(\bar{s}_0, 0, \xi^0) = 0$. Moreover, $(\bar{s}_0, 0, \xi^0)$ is an isolated zero of $\bar{G}$ whose Brouwer degree is not zero. Therefore, if ε is small enough, (2.16) has a solution $(\bar{s}_\varepsilon, \bar{\sigma}_\varepsilon, \xi_\varepsilon)$ near $(\bar{s}_0, 0, \xi_0)$. We conclude that the right hand side of (2.14) is zero, i.e.,

$$\sum_{i=1}^{k}\sum_{l=0}^{n} c_{il}\Big\langle P\psi^l_{\mu_i,\xi_i}, P\psi^h_{\mu_j,\xi_j}\Big\rangle = 0.$$

Moreover, by Lemma 5.3, we conclude that c_{il} are zero. We finish the proof of this theorem. □

3. The finite dimensional reduction

In this section, we prove Proposition 2.1. Let $L_{\mu,\xi} : E^{\perp}_{\mu,\xi} \to E^{\perp}_{\mu,\xi}$ be the linear operator defined by

$$L_{\mu,\xi}(\phi) = \phi - \Pi^{\perp}_{\mu,\xi}\Big(i^*[f_\varepsilon'(V)\phi]\Big), \tag{3.1}$$

where V is defined in (2.5). In the following, we establish the invertibility of $L_{\mu,\xi}$ on $E^{\perp}_{\mu,\xi}$.

Lemma 3.1. *There exist $\varepsilon_0 > 0$ and $C > 0$ such that for any $\xi \in \Omega$, $\bar{d} \in \mathbb{R}^k_+$, $\bar{\sigma} \in (\mathbb{R}^n_+)^k$ satisfying (2.4), for $\varepsilon \in (0,\varepsilon_0)$, it holds*

$$\|L_{\mu,\xi}(\phi)\| \geq C\|\phi\|, \quad \forall \phi \in E^{\perp}_{\mu,\xi}. \tag{3.2}$$

Proof. We argue by contradiction. Assume there exist sequences $\varepsilon_m \to 0$, $\xi \in \Omega$, $\bar{\sigma}_m \in (\mathbb{R}^n_+)^k$ and $\bar{d}_m = (d_{1m},\cdots,d_{km}) \in \mathbb{R}^k_+$ with $\xi_m \to \xi \in \Omega$, $\sigma_{im} \to \sigma_i$ and $d_{im} \to d_i > 0$, $i = 1,\cdots,k$, ϕ_m, $h_m \in \Lambda^{\perp}_{\mu_m,\xi_m}$ such that

$$L_{\mu_m,\xi_m}(\phi_m) = h_m, \quad \|\phi_m\| = 1 \quad \text{and} \quad \|h_m\| \to 0. \tag{3.3}$$

From (3.1), there exists $\omega_m \in E_{\mu_m,\xi_m}$ such that

$$\phi_m - i^*[f_\varepsilon'(V_m)\phi_m] = h_m + \omega_m, \tag{3.4}$$

where $V_m = V(\bar{d}_m, \bar{\sigma}_m, \xi_m) = \sum\limits_{i=1}^{k} PU_{\mu_{im},\xi_{im}}$.

Step 1. We prove that

$$\|\omega_m\| \to 0. \tag{3.5}$$

Let $\omega_m = \sum\limits_{i=1}^{k}\sum\limits_{l=0}^{n} c^{il}_m P\psi^l_{\mu^i_m,\xi^i_m}$, we multiply (3.4) by $P\psi^h_{\mu^l_m,\xi^l_m}$, and integrating in Ω, then

$$\sum_{i=1}^{k}\sum_{l=0}^{n} c^{il}_m \langle P\psi^l_{\mu^i_m,\xi^i_m}, P\psi^h_{\mu^j_m,\xi^j_m}\rangle = \int_\Omega f_\varepsilon'(V_m)\phi_m P\psi^h_{\mu^j_m,\xi^j_m}\,dx. \tag{3.6}$$

From Lemma 5.3, we obtain

$$\sum_{i=1}^{k}\sum_{l=0}^{n} c^{il}_m \langle P\psi^l_{\mu^i_m,\xi^i_m}, P\psi^h_{\mu^j_m,\xi^j_m}\rangle$$

$$=c_m^{jh}\Big(c_h(1+o(1))\Big)+O(1)\sum_{l=0,l\neq h}^{n}c_m^{jl}+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n}{n-2}}\Big)\sum_{i=1,i\neq j}^{k}\sum_{l=0}^{n}c_m^{il}. \tag{3.7}$$

On the other hand, by (5.4), (5.6), (5.7), (5.9), (5.11), (5.13) and the orthogonality condition $\langle\phi_m, P\psi^h_{\mu_m^j,\xi_m^j}\rangle=0$, we have

$$\begin{aligned}
&\int_\Omega f_\varepsilon'(V_m)\phi_m P\psi^h_{\mu_m^j,\xi_m^j}dx\\
=&\int_\Omega\Big(f_\varepsilon'(V_m)-f_0'(V_m)\Big)\phi_m\Big(P\psi^h_{\mu_m^j,\xi_m^j}-\psi^h_{\mu_m^j,\xi_m^j}\Big)dx+\int_\Omega\Big(f_\varepsilon'(V_m)-f_0'(V_m)\Big)\phi_m\psi^h_{\mu_m^j,\xi_m^j}dx\\
&+\int_\Omega\Big(f_0'(V_m)-\sum_{i=1}^{k}(-1)^i f_0'(PU_{\mu_m^i,\xi_m^i})\Big)\phi_m P\psi^h_{\mu_m^j,\xi_m^j}dx\\
&+\sum_{i=1}^{k}(-1)^i\int_\Omega\Big(f_0'(PU_{\mu_m^i,\xi_m^i})-f_0'(U_{\mu_m^i,\xi_m^i})\Big)\phi_m P\psi^h_{\mu_m^j,\xi_m^j}dx\\
\le&\Big|f_\varepsilon'(V_m)-f_0'(V_m)\Big|_{\frac{n}{2}}|\phi_m|_{\frac{2n}{n-2}}\Big|P\psi^h_{\mu_m^j,\xi_m^j}-\psi^h_{\mu_m^j,\xi_m^j}\Big|_{\frac{2n}{n-2}}\\
&+\Big|f_\varepsilon'(V_m)-f_0'(V_m)\Big|_{\frac{n}{2}}|\phi_m|_{\frac{2n}{n-2}}|\psi^h_{\mu_m^j,\xi_m^j}|_{\frac{2n}{n-2}}\\
&+\Big|f_0'(V_m)-\sum_{i=1}^{k}(-1)^i f_0'(PU_{\mu_m^i,\xi_m^i})\Big|_{\frac{n}{2}}|P\psi^h_{\mu_m^j,\xi_m^j}|_{\frac{2n}{n-2}}|\phi_m|_{\frac{2n}{n-2}}\\
&+\Big|f_0'(PU_{\mu_m^i,\xi_m^i})-f_0'(U_{\mu_m^i,\xi_m^i})\Big|_{\frac{n}{2}}|P\psi^h_{\mu_m^j,\xi_m^j}|_{\frac{2n}{n-2}}|\phi_m|_{\frac{2n}{n-2}}=O\Big(\varepsilon\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big).
\end{aligned}\tag{3.8}$$

Consequently, from (3.6)-(3.8), we obtain (3.5).

Step 2. We prove that

$$\liminf_{m\to\infty}\int_\Omega f_\varepsilon'(V_m)u_m^2dx=C>0, \tag{3.9}$$

where u_m satisfies

$$\begin{cases}-\Delta u_m=f_\varepsilon'(V_m)u_m+f_\varepsilon'(V_m)(h_m+\omega_m) & \text{in }\Omega,\\ u_m=0 & \text{on }\partial\Omega,\end{cases} \tag{3.10}$$

with

$$u_m=\phi_m-h_m-\omega_m,\qquad \|u_m\|\to 1. \tag{3.11}$$

We prove that

$$\liminf_{m\to\infty}\|u_m\|=C>0. \tag{3.12}$$

From (3.10), there holds

$$u_m=i^*\Big[f_\varepsilon'(V_m)u_m+f_\varepsilon'(V_m)(h_m+\omega_m)\Big]. \tag{3.13}$$

Moreover, by (2.1), (5.9), (5.11) and (5.13), we get

$$\begin{aligned}
|u_m|_{\frac{2n}{n+2}}\le&C\Big(|f_\varepsilon'(V_m)u_m|_{\frac{2n}{n+2}}+|f_\varepsilon'(V_m)(h_m+\omega_m)|_{\frac{2n}{n+2}}\Big)\\
\le&C|f_\varepsilon'(V_m)-f_0'(V_m)|_{\frac{n}{2}}|u_m|_{\frac{2n}{n-2}}+C\Big|f_0'(V_m)-\sum_{i=1}^{k}(-1)^i f_0'(PU_{\mu_m^i,\xi_m^i})\Big|_{\frac{n}{2}}|u_m|_{\frac{2n}{n-2}}
\end{aligned}$$

$$
\begin{aligned}
&+C|f_\varepsilon'(V_m)-f_0'(V_m)|_{\frac{n}{2}}|h_m+\omega_m|_{\frac{2n}{n-2}}\\
&+C\Big|f_0'(V_m)-\sum_{i=1}^{k}(-1)^i f_0'(PU_{\mu_m^i,\xi_m^i})\Big|_{\frac{n}{2}}|h_m+\omega_m|_{\frac{2n}{n-2}}\\
&+C\Big|\sum_{i=1}^{k}(-1)^i\Big(f_0'(PU_{\mu_m^i,\xi_m^i})-f_0'(U_{\mu_m^i,\xi_m^i})\Big)\Big|_{\frac{n}{2}}|h_m+\omega_m|_{\frac{2n}{n-2}}\le C\|u_m\|+o(1).
\end{aligned}\tag{3.14}
$$

It follows that $|u_m|_{\frac{2n}{n+2}}\to 0$ provided that $\|u_m\|\to 0$, this contradicts with (3.11). Therefore, (3.12) holds.

We multiply (3.13) by u_m, that is

$$
\|u_m\|^2=\int_\Omega f_\varepsilon'(V_m)u_m^2dx+\int_\Omega f_\varepsilon'(V_m)(h_m+\omega_m)u_mdx.\tag{3.15}
$$

By (3.3) and (3.5), one has

$$
\begin{aligned}
\int_\Omega f_\varepsilon'(V_m)(h_m+\omega_m)u_mdx\le&|f_\varepsilon'(V_m)|_{\frac{n}{2}}|h_m+\omega_m)|_{\frac{2n}{n-2}}|u_m|_{\frac{2n}{n-2}}\\
\le&\|h_m+\omega_m\|\|u_m\|=o(1).
\end{aligned}\tag{3.16}
$$

Therefore (3.9) follows by (3.11), (3.12), (3.15) and (3.16).

Step 3. Let us prove that a contradiction arises, by showing that

$$
\int_\Omega f_\varepsilon'(V_m)u_m^2dx=o(1).\tag{3.17}
$$

In order to deal with this conclusion, we decompose $B(\xi,\rho)$ into the union of non-overlapping annuli, that is $B(\xi,\rho)=\bigcup_{i=1}^{k}\mathcal{A}_i$, where

$$
\mathcal{A}_i=B(\xi,\sqrt{\mu_i\mu_{i-1}})\setminus B(\xi,\sqrt{\mu_i\mu_{i+1}}),\quad i=1,\cdots,k,\tag{3.18}
$$

with $\mu_0=\frac{\rho^2}{\mu_1}$ and $\mu_{k+1}=0$. We set a smooth cut-off function χ_m^i as

$$
\chi_m^i(x)=\begin{cases}1 & \text{if } \sqrt{\mu_m^i\mu_m^{i+1}}\le|x-\xi_m|\le\sqrt{\mu_m^i\mu_m^{i-1}},\\ 0 & \text{if } |x-\xi_m|\le\frac{\sqrt{\mu_m^i\mu_m^{i+1}}}{2} \text{ or } |x-\xi_m|\ge 2\sqrt{\mu_m^i\mu_m^{i-1}},\end{cases}\tag{3.19}
$$

and

$$
|\nabla\chi_m^i(x)|\le\frac{2}{\sqrt{\mu_m^i\mu_m^{i-1}}}\quad\text{and}\quad|\nabla^2\chi_m^i(x)|\le\frac{4}{\mu_m^i\mu_m^{i-1}},\quad\text{for any } i=1,\cdots,k.\tag{3.20}
$$

We define

$$
\tilde{u}_m^i(y)=(\mu_m^i)^{\frac{n-2}{2}}u_m(\mu_m^iy+\xi_m)\chi_m^i(\mu_m^iy+\xi_m).\tag{3.21}
$$

First, the following results will be showed in Step 4,

$$
\tilde{u}_m^i\to 0\quad\text{weakly in } D^{1,2}(\mathbb{R}^n),\quad \tilde{u}_m^i\to 0\quad\text{strongly in } L^q_{loc}(\mathbb{R}^n) \text{ for any } q\in[2,2^*).\tag{3.22}
$$

Let us prove (3.17). There holds

$$
\int_\Omega f_\varepsilon'(V_m)u_m^2dx=\int_{\Omega\setminus B(\xi,\rho)}f_\varepsilon'(V_m)u_m^2dx+\sum_{i=1}^{k}\int_{\mathcal{A}_i}f_\varepsilon'(V_m)u_m^2dx,
$$

where

$$\int_{\Omega\setminus B(\xi,\rho)} f_\varepsilon'(V_m)u_m^2dx \le C\sum_{i=1}^k(\mu_i^n)^2\int_{\Omega\setminus B(\xi,\rho)} u_m^2dx = o(1).$$

Since $(\frac{1}{1+|x|^2})^2 \in L^{\frac{n}{2}}(\mathbb{R}^n)$ and (3.22) hold, we conclude that $\int_{(\mathcal{A}_m^i-\xi_m)/\mu_m^i}(\frac{1}{1+|y-\sigma_{in}|^2})^{\frac{n-2}{2}(p-1)}(\tilde{u}_m^i)^2dy \to 0$. On the other hand, we set $x-\xi_m=\mu_m^i y$ and by a fact that, let $h\in L^1_{rad}(\mathbb{R}^n)$, performing the proper change of variable: for any $i\neq l$,

$$\int_{\frac{\mathcal{A}_m^i-\xi_m}{\mu_m^i}} h(|x|)dx = \begin{cases} O\Big((\frac{\mu_l}{\mu_i})^{\frac{n}{2}}\Big) & \text{if } i\le l-1<l,\\ O\Big((\frac{\mu_i}{\mu_l})^{\frac{n}{2}}\Big) & \text{if } i\ge l-1>l.\end{cases} \tag{3.23}$$

By (3.23) and the choice of μ_i in (2.2), we deduce that

$$\text{if } h\in L^1_{rad}(\mathbb{R}^n),\ i\neq l,\quad \int_{\frac{\mathcal{A}_m^l-\xi_m}{\mu_m^i}} h(|x|)dx = O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n}{n-2}}\Big). \tag{3.24}$$

Then, there holds

$$\int_{\frac{\mathcal{A}_m^i-\xi_m}{\mu_m^i}} U_{\mu_{im},\xi_{im}}^{(p-1)\frac{n}{2}}dx = O\Big(\int_{\frac{\sqrt{\mu_m^i\mu_m^{i+1}}}{\mu_m^i}\le|y|\le\frac{\sqrt{\mu_m^i\mu_m^{i-1}}}{\mu_m^i}}\frac{1}{(1+|y-\sigma_{im}|^2)^n}dy\Big).$$

Consquently, from (5.13), we have

$$\begin{aligned}
&\int_{\mathcal{A}_m^i} f_\varepsilon'(V_m)u_m^2dx\\
=&\int_{\mathcal{A}_m^i}\Big(f_\varepsilon'(V_m)-f_0'(V_m)\Big)u_m^2dx+\int_{\mathcal{A}_m^i}f_0'(V_m)u_m^2dx\\
\le& C|f_\varepsilon'(V_m)-f_0'(V_m)|_{\frac{n}{2}}|u_m|^2_{\frac{n}{n-2}}+C\sum_{i=1}^k\int_{\mathcal{A}_m^i}U_{\mu_{im},\xi_{im}}^{p-1}u_m^2dx\\
\le& C\varepsilon\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|+C(\mu_m^i)^{2-\frac{n-2}{2}(p-1)}\int_{\frac{\mathcal{A}_m^i-\xi_m}{\mu_m^i}}\Big(\frac{1}{1+|y-\sigma_{im}|^2}\Big)^{\frac{n-2}{2}(p-1)}(\tilde{u}_m^i)^2dy\\
&+C\sum_{j=1,j\neq i}^k\Big(\int_{\mathcal{A}_m^i}U_{\mu_{jm},\xi_{jm}}^{(p-1)\frac{n}{2}}dx\Big)^{\frac{2}{n}}|u_m|^2_{\frac{2n}{n-2}}=o(1).
\end{aligned}$$

Step 4. We prove (3.22).

From the definition of $\tilde{u}_m^i$, $i=1,\cdots,k$, in (3.21), when $x-\xi_m=\mu_m^i y$, we get

$$\nabla\tilde{u}_m^i(y)=(\mu_m^i)^{\frac{n}{2}}\Big[\Big(\nabla u_m(x)\Big)\chi_m^i(x)+u_m(x)\Big(\nabla\chi_m^i(x)\Big)\Big], \tag{3.25}$$

and

$$\Delta\tilde{u}_m^i(y)=(\mu_m^i)^{\frac{n+2}{2}}\Big[\Big(\Delta u_m(x)\Big)\chi_m^i(x)+2\nabla u_m(x)\nabla\chi_m^i(x)+u_m(x)\Big(\Delta\chi_m^i(x)\Big)\Big]. \tag{3.26}$$

Then, from (3.19), (3.20) and (3.25), it holds that $\|\tilde{u}_m^i\|_{D^{1,2}(\mathbb{R}^n)}\le C$. It follows that, up to a subsequence,

$$\tilde{u}_m^i\to\tilde{u}^i\quad\text{weakly in } D^{1,2}(\mathbb{R}^n),\quad \tilde{u}_m^i\to\tilde{u}^i\quad\text{strongly in } L^q_{loc}(\mathbb{R}^n) \text{ for any } q\in[2,2^*).$$

Next, we show that $\tilde{u}^i$ is the solution of the following problem

$$-\Delta\tilde{u}^i=f_0'(U_{1,\sigma_i})\tilde{u}^i\quad\text{in } \mathbb{R}^n, \tag{3.27}$$

and satisfies the orthogonality conditions

$$\int_{\mathbb{R}^n} \nabla\psi^h_{1,\sigma_i}\nabla\tilde{u}^i dx = 0, \quad h = 0, 1, \cdots, n. \tag{3.28}$$

It follows that $\tilde{u}^i = 0$. This is a contradiction by the result of [10], which concludes the proof.

Step 5. We prove (3.27) and (3.28).

(1) Let us prove (3.27). By (3.25) and (3.26), if $x - \xi_m = \mu^i_m y$, $y \in \Omega^i_m = \frac{\Omega - \xi_m}{\mu^i_m}$, we have

$$\begin{cases} -\Delta\tilde{u}^i_m(y) = (\mu^i_m)^2 f'_\varepsilon\Big(V_m(x)\Big)\tilde{u}^i_m(y) + (\mu^i_m)^{\frac{n+2}{2}} f'_\varepsilon\Big(V_m(x)\Big)\Big(h_m(x) + \omega_m(x)\Big)\chi^i_m(x) \\ \qquad\qquad +2(\mu^i_m)^{\frac{n+2}{2}}\nabla u_m(x)\nabla\chi^i_m(x) + 2(\mu^i_m)^{\frac{n+2}{2}} u_m(x)\Delta\chi^i_m(x), \\ \tilde{u}^i_m = 0 \qquad \text{on } \partial\Omega^i_m. \end{cases}$$

Therefore, if $\varpi \in C^\infty_0(\mathbb{R}^n)$, one has

$$\begin{aligned} &\int_{\mathbb{R}^n} \nabla\tilde{u}^i_m(y)\nabla\varpi(y)dy \\ =&\int_{\mathbb{R}^n} (\mu^i_m)^2 f'_\varepsilon\Big(V_m(\mu^i_m y + \xi_m)\Big)\tilde{u}^i_m(y)\varpi(y)dy + \int_{\mathbb{R}^n} (\mu^i_m)^{\frac{n+2}{2}} \\ &\times f'_\varepsilon\Big(V_m(\mu^i_m y + \xi_m)\Big)\Big(h_m(\mu^i_m y + \xi_m) + \omega_m(\mu^i_m y + \xi_m)\Big)\chi^i_m(\mu^i_m y + \xi_m)\varpi(y)dy \\ &+ 2(\mu^i_m)^{\frac{n+2}{2}} \int_{\mathbb{R}^n} \Big(\nabla u_m(\mu^i_m y + \xi_m)\nabla\chi^i_m(y) + u_m(\mu^i_m y + \xi_m)\Delta\chi^i_m(\mu^i_m y + \xi_m)\Big)\varpi(y)dy. \end{aligned} \tag{3.29}$$

From (3.19) and (3.20), we deduce that the second and the third term tends to 0. For the first term, if $\frac{\sqrt{\mu^i_m\mu^{i+1}_m}}{2} \le |\mu^i_m y| \le 2\sqrt{\mu^i_m\mu^{i-1}_m}$, there holds

$$\begin{aligned} f'_\varepsilon\Big(V_m(\mu^i_m y + \xi_m)\Big) =& f'_\varepsilon\Big(PU_{\mu_{in},\xi_{in}}(\mu^i_m y + \xi_m) + \sum_{i=1, j\neq i}^{k} PU_{\mu_{jm},\xi_{jm}}(\mu^i_m y + \xi_m)\Big) \\ =& f'_\varepsilon\Big((\mu^i_m)^{-\frac{n-2}{2}} U(y - \sigma_{in}) + U_{\mu_{jm},\xi_{jm}}(\mu^i_m y + \xi_m) + o(1)\Big), \end{aligned} \tag{3.30}$$

where

$$U_{\mu_{jm},\xi_{jm}}(\mu^i_m y + \xi_m) = \begin{cases} O\Big((\mu^j_m)^{-\frac{n-2}{2}}\Big) & \text{if } i > j, \\ O\Big(\frac{(\mu^j_m)^{\frac{n-2}{2}}}{(\mu^i_m)^{n-2}}\Big|y - \frac{\mu^j_m}{\mu^i_m}\sigma^j_m\Big|^{-(n-2)}\Big) & \text{if } i < j. \end{cases} \tag{3.31}$$

Moreover, by (3.30), (3.31) and Lebesgue's dominated convergence theorem, it holds that

$$\int_{\mathbb{R}^n} (\mu^i_m)^2 f'_\varepsilon\Big(V_m(\mu^i_m y + \xi_m)\Big)\tilde{u}^i_m(y)\varpi(y)dy \to \int_{\mathbb{R}^n} f'_0\Big(U(y - \sigma_i)\Big)\tilde{u}^i(y)\varpi(y)dy.$$

Then, (3.27) follows by passing to the limit in (3.29).

(2) Let us prove (3.28). We set $x - \xi_m = \mu^i_m y$, then

$$\begin{aligned} \int_{\mathbb{R}^n} \nabla\psi^h_{1,\sigma^i_m}(y)\nabla\tilde{u}^i_m(y)dy =& \int_{\mathbb{R}^n} f'_0\Big(U_{1,\sigma^i_m}(y)\Big)\psi^h_{1,\sigma^i_m}(y)\tilde{u}^i_m(y)dy \\ =&\mu^i_m \int_{\frac{\sqrt{\mu^i_m\mu^{i+1}_m}}{2} \le |x-\xi| \le 2\sqrt{\mu^i_m\mu^{i-1}_m}} f'_0\Big(U_{\mu^i_m,\xi^i_m}(x)\Big)\psi^h_{\mu^i_m,\xi^i_m}(x)u_m(x)\chi^i_m(x)dx \\ =&\mu^i_m\Big(\int_{\mathcal{A}^i_m} f'_0\Big(U_{\mu^i_m,\xi^i_m}(x)\Big)\psi^h_{\mu^i_m,\xi^i_m}(x)u_m(x)dx + o(1)\Big). \end{aligned} \tag{3.32}$$

Now, we show that

$$\mu_m^i \int_{\mathcal{A}_m^i} f_0'\Big(U_{\mu_m^i,\xi_m^i}(x)\Big)\psi^h_{\mu_m^i,\xi_m^i}(x)u_m(x)dx = o(1). \tag{3.33}$$

Therefore, (3.28) follows by (3.32) and (3.33), taking into account that $\sigma_m^i \to \sigma_i$. From (3.5) and (3.11), one has

$$\mu_m^i \int_\Omega \nabla P\psi^h_{\mu_m^i,\xi_m^i}(x)\nabla u_m(x)dx = o(1). \tag{3.34}$$

On the other hand,

$$\begin{aligned}\int_\Omega \nabla P\psi^h_{\mu_m^i,\xi_m^i}(x)\nabla u_m(x)dx &= \int_\Omega f_0'\Big(U_{\mu_m^i,\xi_m^i}(x)\Big)\psi^h_{\mu_m^i,\xi_m^i}(x)u_m(x)dx\\ &= \int_{\Omega\setminus B(\xi_m,\rho)}\cdots dx + \sum_{l=1,\ l\neq i}^{k}\int_{\mathcal{A}_m^l}\cdots dx + \int_{\mathcal{A}_m^i}\cdots dx\\ &= \int_{\mathcal{A}_m^i}\cdots dx + o\Big(\frac{1}{\mu_m^i}\Big),\end{aligned}$$

where

$$\begin{aligned}&\int_{\Omega\setminus B(\xi_m,\rho)}\Big|f_0'\Big(U_{\mu_m^i,\xi_m^i}(x)\Big)\psi^h_{\mu_m^i,\xi_m^i}(x)u_m(x)\Big|dx\\ \le& C|\psi^h_{\mu_m^i,\xi_m^i}|_{\frac{2n}{n-2}}|u_m|_{\frac{2n}{n-2}}\Big(\int_{\Omega\setminus B(\xi_m,\rho)} U^{\frac{2n}{n-2}}_{\mu_m^i,\xi_m^i}dx\Big)^{\frac{2}{n}} = O(\mu_m^i).\end{aligned} \tag{3.35}$$

If $l\neq i$, by (3.24), there is a fact that $\int_{\mathcal{A}_m^i} U^{\frac{2n}{n-2}}_{\mu_m^i,\xi_m^i}dx = O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n}{n-2}}\Big)$, then

$$\begin{aligned}&\int_{\mathcal{A}_m^l}\Big|f_0'\Big(U_{\mu_m^i,\xi_m^i}(x)\Big)\psi^h_{\mu_m^i,\xi_m^i}(x)u_m(x)\Big|dx\\ \le& C|\psi^h_{\mu_m^i,\xi_m^i}|_{\frac{2n}{n-2}}|u_m|_{\frac{2n}{n-2}}\Big(\int_{\mathcal{A}_m^l} U^{\frac{2n}{n-2}}_{\mu_m^i,\xi_m^i}dx\Big)^{\frac{2}{n}} = O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n}{n-2}}\Big).\end{aligned}$$

We finish the proof of this lemma. □

Now, by means of the previous result, we show the following proof.

Proof of Proposition 2.1: First of all, we point out that ϕ solves equation (2.11) if and only if ϕ is a fixed point of the map $T_{\mu,\xi}: E^\perp_{\mu,\xi}\to E^\perp_{\mu,\xi}$ defined by

$$\begin{aligned}T_{\mu,\xi}(\phi) =& L^{-1}_{\mu,\xi}\Pi^\perp_{\mu,\xi}i^*\Big[\Big(f_\varepsilon(V+\phi)-f_\varepsilon(V)-f_\varepsilon'(V)\phi\Big)\\ &+\Big(f_\varepsilon'(V)-\sum_{i=1}^k(-1)^i f_0'(PU_{\mu_i,\xi_i})\Big)\phi+\Big(\sum_{i=1}^k(-1)^i f_0'(PU_{\mu_i,\xi_i})-\sum_{i=1}^k(-1)^i f_0'(U_{\mu_i,\xi_i})\Big)\phi\\ &+\Big(f_\varepsilon(V)-\sum_{i=1}^k(-1)^i f_0(PU_{\mu_i,\xi_i})\Big)+\Big(\sum_{i=1}^k(-1)^i f_0(PU_{\mu_i,\xi_i})-\sum_{i=1}^k(-1)^i f_0(U_{\mu_i,\xi_i})\Big)\Big].\end{aligned}$$

From Lemma 3.1 and Sobolev inequality, we have

$$\|T_{\mu,\xi}(\phi)\| \le C\Big|f_\varepsilon(V+\phi)-f_\varepsilon(V)-f_\varepsilon'(V)\phi\Big|_{\frac{2n}{n+2}}$$

$$
\begin{aligned}
&+C\Big|\Big(f_\varepsilon'(V)-\sum_{i=1}^{k}(-1)^i f_0'(PU_{\mu_i,\xi_i})\Big)\phi\Big|_{\frac{2n}{n+2}}+C\Big|\Big(f_0'(V)-\sum_{i=1}^{k}(-1)^i f_0'(U_{\mu_i,\xi_i})\Big)\phi\Big|_{\frac{2n}{n+2}}\\
&+C\Big|f_\varepsilon(V)-\sum_{i=1}^{k}(-1)^i f_0(PU_{\mu_i,\xi_i})\Big|_{\frac{2n}{n+2}}+C\Big|\sum_{i=1}^{k}(-1)^i\Big(f_0(PU_{\mu_i,\xi_i})-f_0(U_{\mu_i,\xi_i})\Big)\Big|_{\frac{2n}{n+2}}\\
=&H_1+\cdots+H_5.
\end{aligned}
$$

Estimate of H_1: From the mean value theorem, we choose $t=t(x)\in(0,1)$, then

$$
H_1=\Big|f_\varepsilon(V+\phi)-f_\varepsilon(V)-f_\varepsilon'(V)\phi)\Big|_{\frac{2n}{n+2}}=\Big|f_\varepsilon(V+t\phi)-f_\varepsilon'(V)\phi)\Big|_{\frac{2n}{n+2}}. \tag{3.36}
$$

When $n<6$, Lemma 5.5 follows that

$$
H_1\le C\Big(|\phi|^p_{\frac{2n}{n+2}}+|U^{p-2}_{\mu_i,\xi_i}\phi^2|_{\frac{2n}{n+2}}\Big)\le C\Big(|\phi|^{p-2}_{\frac{2n}{n-2}}+|U_{\mu_i,\xi_i}|^{p-2}_{p-2}\Big)|\phi|^2_{\frac{2n}{n-2}}=C\Big(|\phi|^{p-2}+1\Big)\|\phi\|^2.
$$

When $n=6$, by Sobolev inequality, one has

$$
H_1\le C\Big(\Big||\phi|^p\Big|_{\frac{2n}{n+2}}+|\phi^2|_{\frac{2n}{n+2}}\Big)=C\Big(|\phi|^p_{\frac{2n}{n-2}}+\Big(\int_\Omega|\phi|^{\frac{2n}{n-2}}dx\Big)^{\frac{n+2}{2n}}\Big)=2C|\phi|^p_{p+1}\le 2C\|\phi\|^2.
$$

When $n>6$, there holds

$$
\begin{aligned}
H_1\le&C\Big(|\phi|^p_{\frac{2n}{n+2}}+\varepsilon|U^{p-1}_{\mu_i,\xi_i}\phi|_{\frac{2n}{n+2}}\Big)=C\Big(|\phi|^p_{\frac{2n}{n-2}}+\Big(\int_\Omega(U^{p-1}_{\mu_i,\xi_i}|\phi|)^{\frac{2n}{n+2}}dx\Big)^{\frac{n+2}{2n}}\Big)\\
\le&C\Big(|\phi|^p_{\frac{2n}{n-2}}+\varepsilon|U_{\mu_i,\xi_i}|^{p-1}_{\frac{2n}{n-2}}|\phi|_{\frac{2n}{n-2}}\Big)=C\Big(|\phi|^{p-1}_{\frac{2n}{n-2}}+\varepsilon|U_{\mu_i,\xi_i}|^{p-1}_{\frac{2n}{n-2}}\Big)|\phi|_{\frac{2n}{n-2}}\\
\le&C(\|\phi\|^{p-1}+\varepsilon)\|\phi\|.
\end{aligned}
$$

Sum up these estimates, we have

$$
H_1\le\begin{cases}C(|\phi|^{p-2}+1)\|\phi\|^2 & \text{if } 3\le n\le 5,\\ C\|\phi\|^2 & \text{if } n=6,\\ C(\|\phi\|^{p-1}+\varepsilon)\|\phi\| & \text{if } n\ge 7.\end{cases} \tag{3.37}
$$

Estimate of H_2: From Hölder's inequality and (5.12), we get

$$
\begin{aligned}
H_2=&\Big|\Big(f_\varepsilon'(V)-\sum_{i=1}^{k}(-1)^i f_0'(PU_{\mu_i,\xi_i})\Big)\phi\Big|_{\frac{2n}{n+2}}\\
\le&\Big|f_\varepsilon'(V)-\sum_{i=1}^{k}(-1)^i f_0'(PU_{\mu_i,\xi_i})\Big|_{\frac{n}{2}}|\phi|_{\frac{2n}{n-2}}\le C\varepsilon\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\|\phi\|.
\end{aligned}
$$

Estimate of H_3: By Hölder's inequality, (5.9) and (5.11), there holds

$$
\begin{aligned}
H_3=&\Big|\Big(f_0'(V)-\sum_{i=1}^{k}(-1)^i f_0'(U_{\mu_i,\xi_i})\Big)\phi\Big|_{\frac{2n}{n+2}}\\
=&\Big|\Big(f_0'(V)-\sum_{i=1}^{k}(-1)^i f_0'(PU_{\mu_i,\xi_i})\Big)\phi\Big|_{\frac{2n}{n+2}}+\Big|\sum_{i=1}^{k}(-1)^i\Big(f_0'(PU_{\mu_i,\xi_i})-f_0'(U_{\mu_i,\xi_i})\Big)\phi\Big|_{\frac{2n}{n+2}}\\
\le&\Big|f_0'(V)-\sum_{i=1}^{k}(-1)^i f_0'(U_{\mu_i,\xi_i})\Big|_{\frac{n}{2}}|\phi|_{\frac{2n}{n-2}}+k\Big|f_0'(PU_{\mu_i,\xi_i})-f_0'(U_{\mu_i,\xi_i})\Big|_{\frac{n}{2}}|\phi|_{\frac{2n}{n-2}}
\end{aligned}
$$

$$\leq \begin{cases} O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\|\phi\|\Big) & \text{if } 3\leq n\leq 5,\\ \Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\|\phi\|\Big) & \text{if } n=6,\\ O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{-n+8}{n-2}}\|\phi\|\Big) & \text{if } n\geq 7. \end{cases}$$

Estimate of H_4 and H_5: From (5.12) and (5.8), one has

$$|H_4|_{\frac{2n}{n+2}}+|H_5|_{\frac{2n}{n+2}}=O(R_\varepsilon),$$

and R_ε satisfies

$$R_\varepsilon=\begin{cases} O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big) & \text{if } 3\leq n\leq 6,\\ O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n+2}{2(n-2)}}\Big) & \text{if } n\geq 7. \end{cases}$$

From H_1-H_5, there is a constant $C^*>0$ and $\mu_0>0$ such that for each $\mu\in(0,\mu_0)$, we obtain

$$\|T_{\mu,\xi}(\phi)\|\leq C^*R_\varepsilon \quad \text{for every } \phi\in\tilde{B}=\{\phi\in E^{\perp}_{\mu,\xi}:\|\phi\|\leq C^*R_\varepsilon\}.$$

Next, we prove that $T_{\mu,\xi}$ is a contraction map. If ϕ_1, $\phi_2\in\tilde{B}$, then

$$\begin{aligned} &\|T_{\mu,\xi}(\phi_2)-T_{\mu,\xi}(\phi_1)\|\\ \leq& C\Big|f_\varepsilon(V+\phi_2)-f_\varepsilon(V+\phi_1)-f'_\varepsilon(V)(\phi_2-\phi_1)\Big|_{\frac{2n}{n+2}}\\ &+C\Big|\Big(f'_\varepsilon(V)-\sum_{i=1}^{k}(-1)^i f'_0(PU_{\mu_i,\xi_i})\Big)(\phi_2-\phi_1)\Big|_{\frac{2n}{n+2}}\\ &+C\Big|\Big(\sum_{i=1}^{k}(-1)^i f'_0(PU_{\mu_i,\xi_i})-\sum_{i=1}^{k}(-1)^i f'_0(U_{\mu_i,\xi_i})\Big)(\phi_2-\phi_1)\Big|_{\frac{2n}{n+2}}=K_1+K_2+K_3. \end{aligned}$$

Estimate of K_1: Similar to the computations of H_1-H_3. By mean value theorem, we choose $\varrho=\varrho(x)\in(0,1)$ and $\phi_\varrho=(1-\varrho)\phi_1+\varrho\phi_2$, then

$$\begin{aligned} K_1=&\Big|f_\varepsilon(V+\phi_2)-f_\varepsilon(V+\phi_1)-f'_\varepsilon(V)(\phi_2-\phi_1)\Big|_{\frac{2n}{n+2}}\\ =&\Big|\Big(f'_\varepsilon(V+\phi_\varrho)-f'_\varepsilon(V)\Big)(\phi_2-\phi_1)\Big|_{\frac{2n}{n+2}}. \end{aligned}$$

When $n<6$, by Lemma 5.5 and Hölder's inequality, we get

$$\begin{aligned} K_1\leq& C\Big(\Big||\phi_\varrho|^{p-1}(\phi_2-\phi_1)\Big|_{\frac{2n}{n+2}}+\Big|\Big(\sum_{i=1}^{k}U_{\mu_i,\xi_i}\Big)^{p-2}\phi_\varrho(\phi_2-\phi_1)\Big|_{\frac{2n}{n+2}}\Big)\\ \leq& C\Big(|\phi_\varrho|^{p-1}_{\frac{2n}{n-2}}|\phi_2-\phi_1|_{\frac{2n}{n-2}}+\Big(\int_\Omega\Big[\Big(\sum_{i=1}^{k}U_{\mu_i,\xi_i}\Big)^{p-2}\phi_\varrho(\phi_2-\phi_1)\Big]^{\frac{2n}{n+2}}dx\Big)^{\frac{n+2}{2n}}\Big)\\ \leq& C\Big(|\phi_\varrho|^{p-1}_{\frac{2n}{n-2}}+\sum_{i=1}^{k}|U_{\mu_i,\xi_i}|^{p-2}_{\frac{2n}{n-2}}|\phi_\varrho|_{\frac{2n}{n-2}}\Big)|\phi_2-\phi_1|_{\frac{2n}{n-2}}. \end{aligned}$$

When $n=6$, we have

$$K_1\leq C|\phi_\varrho|^{p-1}_{\frac{2n}{n-2}}|\phi_2-\phi_1|_{\frac{2n}{n-2}}.$$

When $n>6$, there holds

$$\begin{aligned}K_1 \leq & C\Big(\big||\phi_\varrho|^{p-1}(\phi_2-\phi_1)\big|_{\frac{2n}{n+2}}+\varepsilon\Big|\Big(\sum_{i=1}^k U_{\mu_i,\xi_i}\Big)^{p-1}\phi_\varrho(\phi_2-\phi_1)\Big|_{\frac{2n}{n+2}}\Big)\\ \leq & C\Big[|\phi_\varrho|^{p-1}_{\frac{2n}{n-2}}|\phi_2-\phi_1|_{\frac{2n}{n-2}}+\Big(\varepsilon\int_\Omega\Big[\Big(\sum_{i=1}^k U_{\mu_i,\xi_i}\Big)^{p-1}\phi_\varrho(\phi_2-\phi_1)\Big]^{\frac{2n}{n+2}}dx\Big)^{\frac{n+2}{2n}}\Big]\\ \leq & C\Big(|\phi_\varrho|^{p-1}_{\frac{2n}{n-2}}+\varepsilon\Big)|\phi_2-\phi_1|_{\frac{2n}{n-2}}.\end{aligned}$$

Hence, by Sobolev inequality, we obtain

$$K_1\leq C\Big(|\phi_\varrho|^{p-1}+\max\{\|\phi_\varrho\|,\varepsilon\}\Big)\|\phi_2-\phi_1\|.$$

Estimate of K_2: Similar to the proof of H_2 and H_3, from (5.13) and (5.11), there holds

$$\begin{aligned}K_2=&\Big|\Big(f_\varepsilon'(V)-\sum_{i=1}^k(-1)^i f_0'(PU_{\mu_i,\xi_i})\Big)(\phi_1-\phi_2)\Big|_{\frac{2n}{n+2}}\\ \leq&\Big|f_\varepsilon'(V)-f_0'(V)\Big|_{\frac{n}{2}}|\phi_2-\phi_1|_{\frac{2n}{n-2}}+\Big|f_0'(V)-\sum_{i=1}^k(-1)^i f_0'(PU_{\mu_i,\xi_i})\Big|_{\frac{n}{2}}|\phi_2-\phi_1|_{\frac{2n}{n-2}}\\ \leq& C\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{-n+8}{n-2}}\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\|\phi_2-\phi_1\|.\end{aligned}$$

Estimate of K_3: By (5.9), one has

$$\begin{aligned}K_3 &= \Big|\Big(\sum_{i=1}^k(-1)^i f_0'(PU_{\mu_i,\xi_i})-\sum_{i=1}^k(-1)^i f_0'(U_{\mu_i,\xi_i})\Big)(\phi_2-\phi_1)\Big|_{\frac{2n}{n+2}}\\ &\leq k\Big|f_0'(PU_{\mu_i,\xi_i})-f_0'(U_{\mu_i,\xi_i})\Big|_{\frac{n}{2}}|\phi_2-\phi_1|_{\frac{2n}{n-2}}\\ &\leq\begin{cases}O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\|\phi_2-\phi_1\|\Big) & \text{if } 3\leq n\leq 5,\\ O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|^{\frac12}\|\phi_2-\phi_1\|\Big) & \text{if } n=6,\\ O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{2}{n-2}}\|\phi_2-\phi_1\|\Big) & \text{if } n\geq 7.\end{cases}\end{aligned}$$

From K_1-K_3, if ε is sufficient small, there exists a constant $L^*\in(0,1)$ such that

$$\|T_{\mu,\xi}(\phi_2)-T_{\mu,\xi}(\phi_1)\|\leq L^*\|\phi_2-\phi_1\|.$$

It follows that $T_{\mu,\xi}$ is a contraction mapping from $\tilde B$ to $\tilde B$, then, it has a unique fixed point $\phi\in\tilde B$. This concludes the proof. □

4. Proof of Proposition 2.2

This section is devoted to prove Proposition 2.2.

Proof of Part a. We consider the following perturbation problem

$$\begin{cases}-\Delta(V+\phi)=f_\varepsilon(V+\phi)+\sum\limits_{i=0}^k\sum\limits_{l=0}^n c_{il}U^{p-1}_{\mu\varepsilon i,\xi\varepsilon i}P\psi^l_{\mu\varepsilon i,\xi\varepsilon i} & \text{in }\Omega,\\ \sum\limits_{i=1}^k\int_\Omega U^{p-1}_{\mu\varepsilon i,\xi\varepsilon i}P\psi^l_{\mu\varepsilon i,\xi\varepsilon i}\phi dx=0 & \text{for } l=0,1,\cdots,n.\end{cases}\tag{4.1}$$

From (2.15), we have

$$\int_\Omega \Delta(V+\phi)P\psi^h_{\mu_j,\xi_j}dx + \int_\Omega f_\varepsilon(V+\phi)P\psi^h_{\mu_j,\xi_j}dx. = 0, \tag{4.2}$$

Thus, by (4.1) and (4.2), we obtain

$$c_{il}\int_\Omega U^{p-1}_{\mu_{\varepsilon i},\xi_{\varepsilon i}} P\psi^l_{\mu_{\varepsilon i},\xi_{\varepsilon i}} P\psi^h_{\mu_j,\xi_j}dx = 0,$$

which means that $c_{il}=0$ for $i=1,\cdots,k$ and $l=0,1,\cdots,n$. Then $V+\phi$ is a solution of problem (1.1).

Proof of Part b. There holds

$$\begin{aligned}
&\Big\langle V+\phi-i^*[f_\varepsilon(V+\phi)], P\psi^h_{\mu_j,\xi_j}\Big\rangle\\
=&\sum_{i=1}^k\langle PU_{\mu_i,\xi_i}, P\psi^h_{\mu_j,\xi_j}\rangle - \int_\Omega f_\varepsilon(V+\phi)P\psi^h_{\mu_j,\xi_j}dx\\
=&\sum_{i=1}^k\int_\Omega (-1)^i f_0(U_{\mu_i,\xi_i})P\psi^h_{\mu_j,\xi_j}dx - \int_\Omega f_\varepsilon(V+\phi)P\psi^h_{\mu_j,\xi_j}dx\\
=&\sum_{i=1}^k\int_\Omega (-1)^i\Big[f_0(U_{\mu_i,\xi_i})-f_0(PU_{\mu_i,\xi_i})\Big]\psi^h_{\mu_j,\xi_j}dx\\
&+\sum_{i=1}^k\int_\Omega (-1)^i\Big[f_0(U_{\mu_i,\xi_i})-f_0(PU_{\mu_i,\xi_i})\Big](P\psi^h_{\mu_j,\xi_j}-\psi^h_{\mu_j,\xi_j})dx\\
&+\int_\Omega\Big[\sum_{i=1}^k(-1)^i f_0(PU_{\mu_i,\xi_i})-f_\varepsilon(V)\Big]\psi^h_{\mu_j,\xi_j}dx\\
&+\int_\Omega\Big[\sum_{i=1}^k(-1)^i f_0(PU_{\mu_i,\xi_i})-f_\varepsilon(V)\Big](P\psi^h_{\mu_j,\xi_j}-\psi^h_{\mu_j,\xi_j})dx\\
&-\int_\Omega\Big[f_\varepsilon(V+\phi)-f_\varepsilon(V)-f'_\varepsilon(V)\phi\Big]P\psi^h_{\mu_j,\xi_j}dx - \int_\Omega [f'_\varepsilon(V)-f'_0(V)]\phi P\psi^h_{\mu_j,\xi_j}dx\\
&-\int_\Omega\Big[f'_0(V)-\sum_{i=1}^k(-1)^i f'_0(U_{\mu_i,\xi_i})\Big]\phi P\psi^h_{\mu_j,\xi_j}dx\\
&-\sum_{i=1}^k\int_\Omega (-1)^i f'_0(U_{\mu_i,\xi_i})\phi(P\psi^h_{\mu_j,\xi_j}-\psi^h_{\mu_j,\xi_j})dx - \sum_{i=1}^k\int_\Omega (-1)^i f'_0(U_{\mu_i,\xi_i})\phi\psi^h_{\mu_j,\xi_j}dx\\
=&P_1+\cdots,P_9.
\end{aligned}$$

Estimate of P_1: It holds

$$\begin{aligned}
P_1=&\sum_{i=1}^k\int_\Omega (-1)^i\Big[f_0(U_{\mu_i,\xi_i})-f_0(PU_{\mu_i,\xi_i})\Big]\psi^h_{\mu_j,\xi_j}dx\\
=&-\sum_{i=1}^k\int_\Omega (-1)^i f'_0(U_{\mu_i,\xi_i})(PU_{\mu_i,\xi_i}-U_{\mu_i,\xi_i})\psi^h_{\mu_j,\xi_j}dx\\
&-\sum_{i=1}^k\int_\Omega (-1)^i\Big[f_0(PU_{\mu_i,\xi_i})-f_0(U_{\mu_i,\xi_i})-f'_0(U_{\mu_i,\xi_i})(PU_{\mu_i,\xi_i}-U_{\mu_i,\xi_i})\Big]\psi^h_{\mu_j,\xi_j}dx.
\end{aligned}$$

If $h=0$, by (5.1) and (2.9), we have

$$\begin{aligned}
&-\int_\Omega f_0'(U_{\mu_i,\xi_i})(PU_{\mu_i,\xi_i}-U_{\mu_i,\xi_i})\psi^0_{\mu_j,\xi_j}dx\\
=\;& p\alpha_n\mu_i^{\frac{n-2}{2}}\int_\Omega U^{p-1}_{\mu_i,\xi_i}H(x,\xi_i)\psi^0_{\mu_j,\xi_j}dx\\
=\;& p\alpha_n\mu_i^{n-2}\Big(H(\xi_i,\xi_i)+O(\mu_i)\Big)\int_{\frac{\Omega-\xi_i}{\mu_i}}\frac{1}{(1+|y|^2)^2}\psi^0\Big(\frac{\mu_i y+\xi_i-\xi_j}{\mu_j}\Big)dy\\
=\;& \begin{cases} p\alpha_n\mu_i^{n-2}\Big(H(\xi_i,\xi_i)+O(\mu_i)\Big)\int_{\mathbb{R}^n}U^{p-1}(y)\psi^0(y)dy & \text{if } j=i,\\ \frac{(n-2)\alpha_n^2 p}{2}\mu_i^{n-2}\Big(H(\xi_i,\xi_i)+O(\mu_i)\Big)\int_{\frac{\Omega-\xi_i}{\mu_i}}\frac{1}{(1+|y|^2)^2}\frac{|\mu_i y+\xi_i-\xi_j|^2-\mu_j^2}{(\mu_j^2+|\mu_i y+\xi_i-\xi_j|^2)^{\frac{n}{2}}}dy & \text{if } j>i,\end{cases}\\
=\;& \begin{cases} \alpha_n a_1 H(\xi,\xi)\mu_i^{n-2}+O(\mu_i^{n-1}) & \text{if } j=i \text{ and } h=0,\\ CH(\xi,\xi)+O(\mu_i^{n-1}) & \text{if } j>i \text{ and } h=0.\end{cases}
\end{aligned}$$

If $h=1,\cdots,n$ and $j=i$, we set $\partial_{\xi_i^h}\varphi(\xi)=\frac{\partial\varphi(\xi)}{\partial\xi_i^h}$, by (2.10), one has

$$\begin{aligned}
&-\int_\Omega f_0'(U_{\mu_i,\xi_i})(PU_{\mu_i,\xi_i}-U_{\mu_i,\xi_i})\psi^h_{\mu_j,\xi_j}dx\\
=&p\alpha_n\mu_i^{\frac{n-2}{2}}\int_\Omega U^{p-1}_{\mu_i,\xi_i}H(x,\xi_i)\psi^h_{\mu_i,\xi_i}dx\\
=&\alpha_n\mu_i^{\frac{n}{2}}\int_\Omega H(x,\xi_i)\frac{\partial U^p_{\mu_i,\xi_i}}{\partial\xi_i^h}dx\\
=&\alpha_n\mu_i^{\frac{n}{2}}\Big[\frac{\partial}{\partial\xi_i^h}\int_\Omega U^p_{\mu_i,\xi_i}H(x,\xi_i)dx-\int_\Omega U^p_{\mu_i,\xi_i}\frac{\partial H(x,\xi_i)}{\partial\xi_i^h}dx\Big]\\
=&\alpha_n\mu_i^{\frac{n}{2}}\Big[\mu_i^{\frac{n-2}{2}}\frac{\partial}{\partial\xi_i^h}\int_{\frac{\Omega-\xi_i}{\mu_i}}U^p_{\mu_i,\xi_i}H(\mu_i y+\xi_i,\xi_i)dy-\mu_i^{\frac{n-2}{2}}\int_{\frac{\Omega-\xi_i}{\mu_i}}U^p_{\mu_i,\xi_i}\frac{\partial H(\mu_i y+\xi_i,\xi_i)}{\partial\xi_i^h}dy\Big]\\
=&\alpha_n a_2\mu_i^{n-1}\Big[\frac{\partial(H(\xi_i,\xi_i))}{\partial\xi_i^h}-\frac{\partial H(\xi_i,\xi_i)}{\partial\xi_i^h}+O(\mu_i)\Big]\\
=&\alpha_n a_2\mu_i^{n-1}\Big(\frac{1}{2}\partial_{\xi_h}\varphi(\xi)+O(\mu_i)\Big).
\end{aligned}$$

If $h=1,\cdots,n$ and $j>i$, by Lemma 5.1 and (2.9), one has

$$\begin{aligned}
&\int_\Omega f_0'(U_{\mu_i,\xi_i})(PU_{\mu_i,\xi_i}-U_{\mu_i,\xi_i})\psi^h_{\mu_j,\xi_j}dx\\
=&p\alpha_n\mu_i^{\frac{n-2}{2}}\int_\Omega U^{p-1}_{\mu_i,\xi_i}H(x,\xi_i)\psi^h_{\mu_j,\xi_j}dx\\
=&(n-2)p\alpha_n^2\mu_i^{\frac{n-2}{2}}\mu_j^{\frac{n}{2}}\int_\Omega\frac{\mu_i^2}{(\mu_i^2+|x-\xi_i|^2)^2}H(x,\xi_i)\frac{(x-\xi_j)_h}{(\mu_j^2+|x-\xi_j|^2)^{\frac{n}{2}}}dx\\
=&O(\mu_i^{\frac{n-2}{2}}\mu_j^{\frac{n}{2}})=O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n+1}{n-2}}\Big).
\end{aligned}$$

On the other hand, by (5.6), (5.7) and (5.10), we get

$$\int_\Omega\Big[f_0(PU_{\mu_i,\xi_i})-f_0(U_{\mu_i,\xi_i})-f_0'(U_{\mu_i,\xi_i})(PU_{\mu_i,\xi_i}-U_{\mu_i,\xi_i})\Big]\psi^h_{\mu_j,\xi_j}dx$$

$$\leq \Big| f_0(PU_{\mu_i,\xi_i}) - f_0(U_{\mu_i,\xi_i}) - f_0'(U_{\mu_i,\xi_i})(PU_{\mu_i,\xi_i} - U_{\mu_i,\xi_i}) \Big|_{\frac{n}{2}} |\psi^h_{\mu_j,\xi_j}|_{\frac{n}{n-2}}$$

$$= o\Big(\Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big)^{\frac{n-1}{n-2}} \Big) \quad \text{for } h = 0, \cdots, n.$$

Therefore, we get

$$P_1 = \begin{cases} \alpha_n a_1 \sum\limits_{i=1}^k \mu_i^{n-2} \Big(H(\xi_i,\xi_i) + O(\mu_i) \Big) + o(\mu_i^{n-2}) & \text{if } j = i \text{ and } h = 0, \\ \sum\limits_{i=1}^k O\Big(H(\xi_i,\xi_i) + O(\mu_i) \Big) + o(\mu_i^{n-2}) & \text{if } j > i \text{ and } h = 0, \\ \alpha_n a_2 \sum\limits_{i=1}^k \mu_i^{n-1} \Big(\frac{1}{2} \partial_{\xi_h} \varphi(x) + O(\mu_i) \Big) + o(\mu_i^{n-1}) & \text{if } j = i \text{ and } h = 1, \cdots, n, \\ O(\sum\limits_{i=1}^k \mu_i^{\frac{n-2}{2}} \mu_j^{\frac{n}{2}}) + o(\mu_i^{n-1}) & \text{if } j > i \text{ and } h = 1, \cdots, n, \end{cases}$$

$$= \begin{cases} \alpha_n a_1 H(\xi,\xi) \frac{\varepsilon}{|\ln \varepsilon|^2} d_1^{n-2} + o\Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big) & \text{if } j = i \text{ and } h = 0, \\ C H(\xi,\xi) + o\Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big) & \text{if } j > i \text{ and } h = 0, \\ \frac{1}{2} \alpha_n a_2 \partial_{\xi_h} \varphi(\xi) \Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big)^{\frac{n-1}{n-2}} d_1^{n-2} + o\Big(\Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big)^{\frac{n-1}{n-2}} \Big) & \text{if } j = i \text{ and } h = 1, \cdots, n, \\ O\Big(\Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big)^{\frac{n-1}{n-2}} \Big) & \text{if } j > i \text{ and } h = 1, \cdots, n. \end{cases}$$

Estimate of P_2: By (5.8) and (5.4), we deduce

$$P_2 = \sum_{i=1}^k \int_\Omega \Big| (-1)^i \Big[f_0(U_{\mu_i,\xi_i}) - f_0(PU_{\mu_i,\xi_i}) \Big] (P\psi^h_{\mu_j,\xi_j} - \psi^h_{\mu_j,\xi_j}) \Big| dx$$

$$\leq C \sum_{i=1}^k \Big| f_0(U_{\mu_i,\xi_i}) - f_0(PU_{\mu_i,\xi_i}) \Big|_{\frac{2n}{n+2}} \Big| P\psi^h_{\mu_j,\xi_j} - \psi^h_{\mu_j,\xi_j} \Big|_{\frac{2n}{n-2}}$$

$$= \begin{cases} o\Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big) & \text{if } h = 0, \\ o\Big(\Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big)^{\frac{n-1}{n-2}} \Big) & \text{if } h = 1, \cdots, n. \end{cases}$$

Estimate of P_3: The main proof of P_3 shows in Lemma 5.7, the final result is

$$P_3 = \int_\Omega \Big[\sum_{i=1}^k (-1)^i f_0(PU_{\mu_i,\xi_i}) - f_\varepsilon(V) \Big] \psi^h_{\mu_j,\xi_j} dx$$

$$= \begin{cases} \alpha_n a_1 \frac{\varepsilon}{|\ln \varepsilon|^2} H(\xi,\xi) d_1^{n-2} + a_3 \frac{\varepsilon}{|\ln \varepsilon|^2} \sum\limits_{i=1}^{k-1} \Big(\frac{d_{i+1}}{d_i} \Big)^{\frac{n-2}{2}} g(\sigma_i) - \frac{2k^2}{(n-2)^2} a_4 \varepsilon \Big| \ln \frac{\varepsilon}{|\ln \varepsilon|^2} \Big| \\ \quad - a_4 \frac{\varepsilon}{|\ln \varepsilon|^2} \sum\limits_{i=1}^k \frac{2}{2i-1} |\ln d_i| + o\Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big) \quad \text{if } h = 0, \\ \frac{1}{2} \alpha_n a_2 \Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big)^{\frac{n-1}{n-2}} d_1^{n-1} \partial_{\xi_h} \varphi(\xi) + o\Big(\Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big)^{\frac{n-1}{n-2}} \Big) \quad \text{if } h = 1, \cdots, n, \end{cases}$$

Estimate of P_4: From (5.12) and (5.4), we get

$$P_4 = \int_\Omega \Big| \Big[\sum_{i=1}^k (-1)^i f_0(PU_{\mu_i,\xi_i}) - f_\varepsilon(V) \Big] (P\psi^h_{\mu_j,\xi_j} - \psi^h_{\mu_j,\xi_j}) \Big| dx$$

$$\begin{aligned}
&\leq C\Big|f_\varepsilon(V)-\sum_{i=1}^{k}(-1)^i f_0(PU_{\mu_i,\xi_i})\Big|_{\frac{2n}{n+2}}\Big|P\psi^h_{\mu_j,\xi_j}-\psi^h_{\mu_j,\xi_j}\Big|_{\frac{2n}{n-2}}\\
&=\begin{cases} o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } h=0 \text{ and } 3\leq n\leq 4,\\ o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } h=0 \text{ and } n\geq 5,\\ o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } h=1,\cdots,n \text{ and } n\geq 3.\end{cases}
\end{aligned}$$

Estimate of P_5: From (5.4), (5.6), (5.7), (3.36) and (3.37), we have

$$\begin{aligned}
P_5 &= \int_\Omega \Big|\Big[f_\varepsilon(V+\phi)-f_\varepsilon(V)-f'_\varepsilon(V)\phi\Big]P\psi^h_{\mu_j,\xi_j}\Big|dx\\
&= O\Big(\Big|f_\varepsilon(V+\phi)-f_\varepsilon(V)-f'_\varepsilon(V)\phi)\Big|_{\frac{2n}{n+2}}|P\psi^h_{\mu_j,\xi_j}|_{\frac{2n}{n-2}}\Big)\\
&\leq \begin{cases} O(1+\|\phi\|^{p-2}|)\|\phi\|^2 & \text{if } 3\leq n\leq 5,\\ O(\|\phi\|^2) & \text{if } n=6,\\ O(\varepsilon+\|\phi\|^{p-1}|)\|\phi\|^2 & \text{if } n\geq 7,\end{cases}\\
&= \begin{cases} o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } n=3,\\ o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } n\geq 4.\end{cases}
\end{aligned}$$

Estimate of P_6: By (5.13) and (2.13), it holds

$$\begin{aligned}
P_6 &= \int_\Omega \Big|[f'_\varepsilon(V)-f'_0(V)]\phi P\psi^h_{\mu_j,\xi_j}\Big|dx\\
&= O\Big(|f'_\varepsilon(V)-f'_0(V)|_{\frac{n}{2}}|\phi|_{\frac{2n}{n-2}}|P\psi^h_{\mu_j,\xi_j}|_{\frac{2n}{n-2}}\Big)\\
&= \begin{cases} o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } h=0,\\ o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } h=1,\cdots,n..\end{cases}
\end{aligned}$$

Estimate of P_7: By (5.11), (2.13), (5.7), (5.6), (5.4), (5.9), there holds

$$\begin{aligned}
P_7 &= \int_\Omega \Big|\Big[f'_0(V)-\sum_{i=1}^{k}(-1)^i f'_0(U_{\mu_i,\xi_i})\Big]\phi P\psi^h_{\mu_j,\xi_j}\Big|dx\\
&= O\Big(\Big|\Big[f'_0(V)-\sum_{i=1}^{k}(-1)^i f'_0(U_{\mu_i,\xi_i})\Big]P\psi^h_{\mu_j,\xi_j}\Big|_{\frac{2n}{n+2}}\|\phi\|\Big)\\
&= O\Big(\Big|f'_0(V)-\sum_{i=1}^{k}(-1)^i f'_0(PU_{\mu_i,\xi_i})\Big|_{\frac{n}{2}}|P\psi^h_{\mu_j,\xi_j}|_{\frac{2n}{n+2}}\|\phi\|\Big)\\
&+ O\Big(\sum_{i=1}^{k}(-1)^i\Big|f'_0(PU_{\mu_i,\xi_i})-f'_0(U_{\mu_i,\xi_i})\Big|_{\frac{n}{2}}|P\psi^h_{\mu_j,\xi_j}|_{\frac{2n}{n+2}}\|\phi\|\Big)
\end{aligned}$$

$$= \begin{cases} o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } h=0, \\ o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } h=1,\cdots,n. \end{cases}$$

Estimate of P_8: For $h=0$, by (5.5), (2.13) and (5.4), it follows that

$$\begin{aligned} P_8 &= \sum_{i=1}^{k}\int_\Omega \Big|(-1)^i f_0'(U_{\mu_i,\xi_i})\phi(P\psi^0_{\mu_j,\xi_j}-\psi^0_{\mu_j,\xi_j})\Big|dx \\ &= O\Big(\sum_{i=1}^{k}\Big|f_0'(U_{\mu_i,\xi_i})\Big|_{\frac{n}{2}}|\phi|_{\frac{2n}{n-2}}\Big|P\psi^0_{\mu_j,\xi_j}-\psi^0_{\mu_j,\xi_j}\Big|_{\frac{2n}{n-2}}\Big) \\ &= O\Big(\sum_{i=1}^{k}\mu_i^{\frac{n-2}{2}}\|\phi\|\Big) = \begin{cases} o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } n=3, \\ o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } n\geq 4, \end{cases} \end{aligned}$$

and for $h=1,\cdots,n$, we obtain

$$\begin{aligned} P_8 &= \sum_{i=1}^{k}\int_\Omega \Big|(-1)^i f_0'(U_{\mu_i,\xi_i})\phi(P\psi^h_{\mu_j,\xi_j}-\psi^h_{\mu_j,\xi_j})\Big|dx \\ &= O\Big(\Big|\sum_{i=1}^{k}f_0'(U_{\mu_i,\xi_i})\Big|_{\frac{n}{2}}|\phi|_{\frac{2n}{n-2}}|P\psi^h_{\mu_j,\xi_j}-\psi^h_{\mu_j,\xi_j}|_{\frac{2n}{n-2}}\Big) \\ &= O\Big(\sum_{i=1}^{k}\mu_i^{\frac{n}{2}}\|\phi\|\Big) = \begin{cases} o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } 3\leq n\leq 5, \\ o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } n\geq 6. \end{cases} \end{aligned}$$

Estimate of P_9: We use ϕ to multiply (2.7) and integral in the Ω, then

$$\int_\Omega f_0'(U_{\mu_i,\xi_i})\phi\psi^h_{\mu_j,\xi_j}dx = 0.$$

From P_1-P_9, we complete the proof. □

5. Appendix

We collect some well known estimates.

Lemma 5.1. *[42] Let $\xi\in\Omega$, $\mu>0$ is small, there holds*

$$PU_{\mu,\xi}(x) = U_{\mu,\xi}(x) - \alpha_n\mu^{\frac{n-2}{2}}H(x,\xi) + O(\mu^{\frac{n+2}{2}}), \tag{5.1}$$

$$P\psi^0_{\mu,\xi}(x) = \psi^0_{\mu,\xi}(x) - \frac{n-2}{2}\alpha_n\mu^{\frac{n-2}{2}}H(x,\xi) + O(\mu^{\frac{n+4}{2}}), \tag{5.2}$$

$$P\psi^h_{\mu,\xi}(x) = \psi^h_{\mu,\xi}(x) - \alpha_n\mu^{\frac{n}{2}}\partial_{\xi_h}H(x,\xi) + O(\mu^{\frac{n+2}{2}}), \tag{5.3}$$

as $\mu\to 0$ uniformly with respect to ξ in compact subsets of Ω, where $h=1,\cdots,n$ and α_n is given in (1.3). Moreover,

$$|P\psi^h_{\mu_j,\xi_j}-\psi^h_{\mu_j,\xi_j}|_{\frac{2n}{n-2}} = \begin{cases} O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{1}{2}}\Big) & \text{if } h=0, \\ O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n}{2(n-2)}}\Big) & \text{if } h=1,\cdots,n, \end{cases} \tag{5.4}$$

for $j = 1, \cdots, k$.

Lemma 5.2. *[18] There holds*

$$\int_\Omega U^q_{\mu,\xi}(x)dx = \begin{cases} O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{q}{2}}\Big) & \text{if } 0 < q < \frac{n}{n-2}, \\ O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{n}{2(n-2)}}\big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\big|\Big) & \text{if } q = \frac{n}{n-2}, \\ O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{n}{n-2}-\frac{q}{2}}\Big) & \text{if } \frac{n}{n-2} < q \le \frac{2n}{n-2}, \end{cases} \tag{5.5}$$

$$\int_\Omega |\psi^0_{\mu,\xi}(x)|^q dx = \begin{cases} O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{q}{2}}\Big) & \text{if } 0 < q < \frac{n}{n-2}, \\ O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{n}{2(n-2)}}\big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\big|\Big) & \text{if } q = \frac{n}{n-2}, \\ O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{n}{n-2}-\frac{q}{2}}\Big) & \text{if } \frac{n}{n-2} < q \le \frac{2n}{n-2}, \end{cases} \tag{5.6}$$

and

$$\int_\Omega |\psi^h_{\mu,\xi}(x)|^q dx = \begin{cases} O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{nq}{2(n-2)}}\Big) & \text{if } 0 < q < \frac{n}{n-1}, \\ O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{n^2}{2(n-1)(n-2)}}\big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\big|\Big) & \text{if } q = \frac{n}{n-1}, \\ O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{n}{n-2}-\frac{q}{2}}\Big) & \text{if } \frac{n}{n-1} < q \le \frac{2n}{n-2}, \end{cases} \tag{5.7}$$

for $h = 1, \cdots, n$. *Moreover,*

$$\Big|f_0(PU_{\mu,\xi}) - f_0(U_{\mu,\xi})\Big|_{\frac{2n}{n+2}} = \begin{cases} O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } 3 \le n \le 5, \\ O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\big|^{\frac{2}{3}}\Big) & \text{if } n = 6, \\ O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{n+2}{2(n-2)}}\Big) & \text{if } n \ge 7, \end{cases} \tag{5.8}$$

$$\Big|f_0'(PU_{\mu,\xi}) - f_0'(U_{\mu,\xi})\Big|_{\frac{n}{2}} = \begin{cases} O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } n = 3, \\ O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\big|^{\frac{1}{2}}\Big) & \text{if } n = 4, \\ O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{2}{n-2}}\Big) & \text{if } n \ge 5, \end{cases} \tag{5.9}$$

$$\begin{aligned} &\Big|f_0(PU_{\mu,\xi}) - f_0(U_{\mu,\xi}) - f_0'(U_{\mu,\xi})(PU_{\mu,\xi} - U_{\mu,\xi})\Big|_{\frac{n}{2}} \\ &= \begin{cases} O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{5}{2}}\Big) & \text{if } n = 3, \\ O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{3}{2}}\big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\big|^{\frac{1}{2}}\Big) & \text{if } n = 4, \\ O\Big(\big(\frac{\varepsilon}{|\ln\varepsilon|^2}\big)^{\frac{n+2}{2(n-2)}}\Big) & \text{if } n \ge 5. \end{cases} \end{aligned} \tag{5.10}$$

Lemma 5.3. *It holds*

$$\Big\langle P\psi^l_{\mu_i,\xi_i}, P\psi^h_{\mu_j,\xi_j}\Big\rangle = \begin{cases} o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n}{n-2}}\Big) & \text{if } j>i,\\ O(1) & \text{if } l\neq h,\\ c_h(1+o(1)) & \text{if } i=j,\ l=h,\end{cases}$$

for some positive constants c_0 and $c_1=\cdots,c_n$, where $i,\ j=1,\cdots,k$ and $h,\ l=0,\cdots,n$,

Proof. We have

$$\Big\langle P\psi^l_{\mu_i,\xi_i}, P\psi^h_{\mu_j,\xi_j}\Big\rangle = \int_\Omega f_0'(U_{\mu_i,\xi_i})\psi^l_{\mu_i,\xi_i}\psi^h_{\mu_j,\xi_j}dx + \int_\Omega f_0'(U_{\mu_i,\xi_i})\psi^l_{\mu_i,\xi_i}(P\psi^h_{\mu_j,\xi_j}-\psi^h_{\mu_j,\xi_j})dx.$$

From (5.5), (5.7) and (5.4), there holds

$$\begin{aligned}&\int_\Omega f_0'(U_{\mu_i,\xi_i})\psi^l_{\mu_i,\xi_i}(P\psi^h_{\mu_j,\xi_j}-\psi^h_{\mu_j,\xi_j})dx\\ &\le |f_0'(U_{\mu_i,\xi_i})|_{\frac{n}{2}}|\psi^l_{\mu_i,\xi_i}|_{\frac{2n}{n-2}}|P\psi^h_{\mu_j,\xi_j}-\psi^h_{\mu_j,\xi_j}|_{\frac{2n}{n-2}} = o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\Big).\end{aligned}$$

On the other hand, if $l,\ h=1,\cdots,n$, the change of variables $x-\xi_i=\mu_i y$ shows that

$$\begin{aligned}&\int_\Omega f_0'(U_{\mu_i,\xi_i})\psi^l_{\mu_i,\xi_i}\psi^h_{\mu_j,\xi_j}dx\\ =\ & (n-2)^2\alpha_n^{\frac{2n}{n-2}}\mu_i^{\frac{n+4}{2}}\mu_j^{\frac{n}{2}}\int_\Omega \frac{(x-\xi_i)_l}{(\mu_i^2+|x-\xi_i|^2)^{\frac{n+4}{2}}}\frac{(x-\xi_j)_h}{(\mu_j^2+|x-\xi_j|^2)^{\frac{n}{2}}}dx\\ =\ & (n-2)^2\alpha_n^{\frac{2n}{n-2}}\mu_i^{\frac{n-2}{2}}\mu_j^{\frac{n}{2}}\int_{\frac{\Omega-\xi_i}{\mu_i}} \frac{y_l}{(1+|y|^2)^{\frac{n+4}{2}}}\frac{(\mu_i y+\xi_i-\xi_j)_h}{(\mu_j^2+|\mu_i y+\xi_i-\xi_j|^2)^{\frac{n}{2}}}dy\\ =\ & \begin{cases} O\Big((\frac{\mu_j}{\mu_i})^{\frac{n}{2}}\Big) & \text{if } j>i,\\ O(1) & \text{if } i=j,\ l\neq h,\\ c_h(1+o(1)) & \text{if } i=j,\ l=h.\end{cases}\end{aligned}$$

If $l=1,\cdots,n$ and $h=0$, we have

$$\begin{aligned}&\int_\Omega f_0'(U_{\mu_i,\xi_i})\psi^l_{\mu_i,\xi_i}\psi^0_{\mu_j,\xi_j}dx\\ =\ & \frac{(n-2)^2}{2}\alpha_n^{\frac{2n}{n-2}}\mu_i^{\frac{n+4}{2}}\mu_j^{\frac{n-2}{2}}\int_\Omega \frac{(x-\xi_i)^l}{(\mu_i^2+|x-\xi_i|^2)^{\frac{n+4}{2}}}\frac{|x-\xi_j|^2-\mu_j^2}{(\mu_j^2+|x-\xi_j|^2)^{\frac{n}{2}}}dx\\ =\ & (n-2)^2\alpha_n^{\frac{2n}{n-2}}\mu_i^{\frac{n-2}{2}}\mu_j^{\frac{n-2}{2}}\int_{\frac{\Omega-\xi_i}{\mu_i}} \frac{y^l}{(1+|y|^2)^{\frac{n+4}{2}}}\frac{|\mu_i y+\xi_i-\xi_j|^2-\mu_j^2}{(\mu_j^2+|\mu_i y+\xi_i-\xi_j|^2)^{\frac{n}{2}}}dy\\ =\ & o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\Big).\end{aligned}$$

Finally, if $l=0$ and $h=0$, one has

$$\begin{aligned}&\int_\Omega f_0'(U_{\mu_i,\xi_i})\psi^0_{\mu_i,\xi_i}\psi^0_{\mu_j,\xi_j}dx\\ =\ & \frac{(n-2)^2}{4}\alpha_n^{\frac{2n}{n-2}}\mu_i^{\frac{n-2}{2}}\mu_j^{\frac{n-2}{2}}\int_\Omega \frac{|x-\xi_i|^2-\mu_i^2}{(\mu_i^2+|x-\xi_i|^2)^{\frac{n}{2}}}\frac{|x-\xi_j|^2-\mu_j^2}{(\mu_j^2+|x-\xi_j|^2)^{\frac{n}{2}}}dx\end{aligned}$$

$$= (n-2)^2\alpha_n^{\frac{2n}{n-2}}\mu_i^{-\frac{n-2}{2}}\mu_j^{\frac{n-2}{2}}\int_{\frac{\Omega-\xi_i}{\mu_i}}\frac{(|y-\sigma_i|^2-1)}{(1+|y-\sigma_i|^2)^{\frac{n}{2}}}\frac{|\mu_i y+\xi_i-\xi_j|^2-\mu_j^2}{(\mu_j^2+|\mu_i y+\xi_i-\xi_j|^2)^{\frac{n}{2}}}dy$$

$$= \begin{cases} o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\Big) & \text{if } j>i,\\ c_0(1+o(1)) & \text{if } i=j. \end{cases}$$

Therefore, this lemma follows from above estimates.

□

Lemma 5.4. *[37] Let $\xi\in\Omega$, there holds*

$$P\psi^0_{\mu,\xi}(x)=\frac{n-2}{2}a_2\mu^{\frac{n-2}{2}}G(x,\xi)+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big),\quad x\in\Omega,$$

and

$$P\psi^h_{\mu,\xi}(x)=a_2\mu^{\frac{n}{2}}\frac{\partial G}{\partial\xi_h}(x,\xi)+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big)\ \text{ if } h=1,\cdots,n,\quad x\in\Omega,$$

as $\varepsilon\to0$ uniformly on compact sets of $\Omega\setminus\{\xi\}$, where a_2 is given in Proposition 2.2.

Lemma 5.5. *[18] Let $\theta>0$ and $u=\sum_{i=1}^k u_i$, $i=1,\cdots,n$, if $\varepsilon>0$ small enough, for $u,u_i,v\in\mathbb{R}$, $p=2^*-1$, it holds that*

(1) $|f_\varepsilon(u)-f_0(u)|\le\varepsilon|u|^p\ln\ln(e+|u|)$,

(2) $f_\varepsilon'(u)\le C|u|^{p-1}$,

(3) $|f_\varepsilon'(u)-f_0'(u)|\le\varepsilon|u|^{p-1}\Big(p\ln\ln(e+|u|)+\frac{1}{\ln(e+|u|)}\Big)$,

(4)

$$|f_\varepsilon'(u+v)-f_\varepsilon'(u)|\le\begin{cases}C(|u|^{p-2}+|v|^{p-2})|v| & \text{if } n\le6,\\ C(|v|^{p-1}+\varepsilon|u|^{p-1}) & \text{if } n>6,\end{cases}$$

(5) $\ln\ln(e+\mu^{-\theta}u)=\ln\ln(\mu^{-\theta})+\ln\Big(1+\frac{\ln(e^{1-\theta|\ln\mu|}+u)}{\theta|\ln\mu|}\Big)$,

(6) $\lim_{\mu\to0}\Big(|\ln\mu|\ln\Big(1+\frac{\ln(e^{1-\theta|\ln\mu|}+u)}{\theta|\ln\mu|}\Big)\Big)=\frac{1}{\theta}\ln u$,

where C is a positive constant.

Lemma 5.6. *There holds*

$$\Big|f_0'(V)-\sum_{i=1}^k(-1)^i f_0'(PU_{\mu_i,\xi_i})\Big|_{\frac{n}{2}}=\begin{cases}O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } 3\le n\le5,\\ O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big) & \text{if } n=6,\\ O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{-n+8}{n-2}}\Big) & \text{if } n\ge7,\end{cases}\tag{5.11}$$

$$\Big|f_\varepsilon(V)-\sum_{i=1}^k(-1)^i f_0(PU_{\mu_i,\xi_i})\Big|_{\frac{2n}{n+2}}=\begin{cases}O\Big(\varepsilon\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big) & \text{if } 3\le n\le6,\\ O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n+2}{2(n-2)}}\Big) & \text{if } n\ge7,\end{cases}\tag{5.12}$$

$$|f_\varepsilon'(V)-f_0'(V)|_{\frac{n}{2}}=O\Big(\varepsilon\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big).\tag{5.13}$$

Proof. Let us estimate (5.11). One has

$$\begin{aligned}
&\int_\Omega \Big|f_0'(V)-\sum_{i=1}^k(-1)^i f_0'(PU_{\mu_i,\xi_i})\Big|^{\frac{n}{2}}dx\\
=&\int_{\Omega\setminus B(\xi,\rho)}\Big|V^{p-1}-\sum_{i=1}^k(-1)^i(PU_{\mu_i,\xi_i})^{p-1}\Big|^{\frac{n}{2}}dx+\sum_{l=1}^k\int_{\mathcal{A}_l}\Big|V^{p-1}-\sum_{i=1}^k(-1)^i(PU_{\mu_i,\xi_i})^{p-1}\Big|^{\frac{n}{2}}dx.
\end{aligned}$$

We estimate the first term

$$\begin{aligned}
&\int_{\Omega\setminus B(\xi,\rho)}\Big|V^{p-1}-\sum_{i=1}^k(-1)^i(PU_{\mu_i,\xi_i})^{p-1}\Big|^{\frac{n}{2}}dx\\
\le&\sum_{i=1}^k\int_{\Omega\setminus B(\xi,\rho)}U_{\mu_i,\xi_i}^{(p-1)\frac{n}{2}}dx\le C\sum_{i=1}^k\mu_i^n=O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n}{n-2}}\Big).
\end{aligned}$$

For any l, by the mean value theorem, there exists $t=t(x)\in[0,1]$ such that

$$\begin{aligned}
&\int_{\mathcal{A}_l}\Big|V^{p-1}-\sum_{i=1}^k(-1)^i(PU_{\mu_i,\xi_i})^{p-1}\Big|^{\frac{n}{2}}dx\\
=&\int_{\mathcal{A}_l}\Big|\Big((-1)^lPU_{\mu_l,\xi_l}+\sum_{i\neq l}^k(-1)^iPU_{\mu_i,\xi_i}\Big)^{p-1}-(-1)^l(PU_{\mu_l,\xi_l})^{p-1}-\sum_{i\neq l}^k(-1)^i(PU_{\mu_i,\xi_i})^{p-1}\Big|^{\frac{n}{2}}dx\\
\le&C\int_{\mathcal{A}_l}\Big|\Big((-1)^lPU_{\mu_l,\xi_l}+t\sum_{i\neq l}^k(-1)^iPU_{\mu_i,\xi_i}\Big)^{p-2}\sum_{i\neq l}^k(-1)^iPU_{\mu_i,\xi_i}\Big|^{\frac{n}{2}}dx+C\sum_{i\neq l}^k\int_{\mathcal{A}_l}|PU_{\mu_i,\xi_i}|^{(p-1)\frac{n}{2}}dx\\
\le&C\int_{\mathcal{A}_l}\Big|(-1)^{l+i}(PU_{\mu_l,\xi_l})^{p-2}\sum_{i\neq l}^kPU_{\mu_i,\xi_i}\Big|^{\frac{n}{2}}dx+C\sum_{i\neq l}^k\int_{\mathcal{A}_l}|PU_{\mu_i,\xi_i}|^{(p-1)\frac{n}{2}}dx\\
\le&C\sum_{i\neq l}^k\int_{\mathcal{A}_l}\Big|U_{\mu_l,\xi_l}^{p-2}U_{\mu_i,\xi_i}\Big|^{\frac{n}{2}}dx+C\sum_{i\neq l}^k\int_{\mathcal{A}_l}|U_{\mu_i,\xi_i}|^{(p-1)\frac{n}{2}}dx.
\end{aligned}$$

If $i\neq l$, by (3.24), let $x-\xi=\mu_i y$, then

$$\begin{aligned}
\int_{\mathcal{A}_l}|U_{\mu_i,\xi_i}|^{(p-1)\frac{n}{2}}dx\le&C\int_{\mathcal{A}_l}\Big(\frac{\mu_i^{\frac{n-2}{2}}}{(\mu_i^2+|x-\xi_i|^2)^{\frac{n-2}{2}}}\Big)^{(p-1)\frac{n}{2}}dx\\
=&C\mu_i^{n-\frac{n-2}{2}(p-1)\frac{n}{2}}\int_{\frac{\mathcal{A}_l}{\mu_i}}\frac{1}{(1+|y-\sigma_i|^2)^n}dy=O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n}{n-2}}\Big).
\end{aligned}$$

If $n>6$, by (3.24), one has

$$\begin{aligned}
&\int_{\mathcal{A}_l}\Big|U_{\mu_l,\xi_l}^{p-2}U_{\mu_i,\xi_i}\Big|^{\frac{n}{2}}dx\\
\le&\ C\int_{\mathcal{A}_l}\Big(\frac{\mu_l^{\frac{-n+6}{2}}}{(\mu_l^2+|x-\xi_l|^2)^{\frac{-n+6}{2}}}\Big)^{\frac{n}{2}}\Big(\frac{\mu_i^{\frac{n-2}{2}}}{(\mu_i^2+|x-\xi_i|^2)^{\frac{n-2}{2}}}\Big)^{\frac{n}{2}}dx\\
=&\ C\mu_i^{n-\frac{n-2}{2}\frac{n}{2}}\mu_l^{\frac{-n+6}{2}\frac{n}{2}}\int_{\frac{\mathcal{A}_l}{\mu_i}}\frac{1}{(\mu_l^2+|\mu_i y-\mu_l\sigma_l|^2)^{\frac{-n+6}{2}\frac{n}{2}}}\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n-2}{2}\frac{n}{2}}}dy
\end{aligned}$$

$$
\begin{aligned}
&= \begin{cases} O(\mu_i^{n-\frac{n-2}{2}\frac{n}{2}+(n-6)\frac{n}{2}}\mu_l^{\frac{-n+6}{2}\frac{n}{2}})\int_{\frac{\mathcal{A}_l}{\mu_i}} \frac{1}{|y-\frac{\mu_l}{\mu_i}\sigma_l|^{\frac{n(-n+6)}{2}}}\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n-2}{2}\frac{n}{2}}}dy & \text{if } l>i,\\ O(\mu_i^{n-\frac{n-2}{2}\frac{n}{2}}\mu_l^{-\frac{-n+6}{2}\frac{n}{2}})\int_{\frac{\mathcal{A}_l}{\mu_i}} \frac{1}{(1+|y-\sigma_i|^2)^{\frac{n-2}{2}\frac{n}{2}}}dy & \text{if } l<i, \end{cases}\\
&= \begin{cases} O\Big(\mu_i^{n-\frac{n-2}{2}\frac{n}{2}}(\frac{\mu_l}{\mu_i})^{\frac{-n+6}{2}\frac{n}{2}}(\frac{\varepsilon}{|\ln\varepsilon|^2})^{\frac{n}{n-2}}\Big) & \text{if } l>i,\\ O\Big((\frac{\mu_i}{\mu_l})^{\frac{-n+6}{2}\frac{n}{2}}(\frac{\varepsilon}{|\ln\varepsilon|^2})^{\frac{n}{n-2}}\Big) & \text{if } l<i, \end{cases}\\
&= O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{-n+6}{n-2}\frac{n}{2}+\frac{n}{n-2}}\Big).
\end{aligned}
$$

If $n<6$, it holds

$$
\begin{aligned}
&\int_{\mathcal{A}_l}\Big|U_{\mu_l,\xi_l}^{p-2}U_{\mu_i,\xi_i}\Big|^{\frac{n}{2}}dx\\
\le\ & C\int_{\mathcal{A}_l}\Big(\frac{\mu_l^{\frac{-n+6}{2}}}{(\mu_l^2+|x-\xi_l|^2)^{\frac{-n+6}{2}}}\Big)^{\frac{n}{2}}\Big(\frac{\mu_i^{\frac{n-2}{2}}}{(\mu_i^2+|x-\xi_i|^2)^{\frac{n-2}{2}}}\Big)^{\frac{n}{2}}dx\\
=\ & C\mu_l^{n-\frac{-n+6}{2}\frac{n}{2}}\mu_i^{\frac{n-2}{2}\frac{n}{2}}\int_{\frac{\mathcal{A}_l}{\mu_l}}\frac{1}{(1+|y-\sigma_l|^2)^{\frac{-n+6}{2}\frac{n}{2}}}\frac{1}{(\mu_i^2+|\mu_l y-\mu_i\sigma_i|^2)^{\frac{n-2}{2}\frac{n}{2}}}dy\\
=\ & \begin{cases} O(\mu_l^{n-\frac{-n+6}{2}\frac{n}{2}}\mu_i^{-\frac{n-2}{2}\frac{n}{2}})\int_{\frac{\mathcal{A}_l}{\mu_l}}\frac{1}{(1+|y-\sigma_i|^2)^{\frac{-n+6}{2}\frac{n}{2}}}dy & \text{if } l>i,\\ O(\mu_l^{n-\frac{-n+6}{2}\frac{n}{2}-(n-2)\frac{n}{2}}\mu_i^{\frac{n-2}{2}\frac{n}{2}})\int_{\frac{\mathcal{A}_l}{\mu_l}}\frac{1}{|y-\frac{\mu_i}{\mu_l}\sigma_l|^{\frac{n(n-2)}{2}}}\frac{1}{(1+|y-\sigma_l|^2)^{\frac{-n+6}{2}\frac{n}{2}}}dy & \text{if } l<i, \end{cases}\\
=\ & \begin{cases} O((\frac{\mu_l}{\mu_i})^{\frac{n-2}{2}\frac{n}{2}}) & \text{if } l>i,\\ O((\frac{\mu_i}{\mu_l})^{\frac{n-2}{2}\frac{n}{2}}) & \text{if } l<i, \end{cases}\\
=\ & O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n}{2}}\Big).
\end{aligned}
$$

A similar estimate can be obtained for $n=6$, we prove that

$$
\int_{\mathcal{A}_l}\Big|U_{\mu_l,\xi_l}^{p-2}U_{\mu_i,\xi_i}\Big|^3dx = O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^3\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|^3\Big).
$$

Thus, (5.11) holds.

Let us now estimate (5.12), one has

$$
\begin{aligned}
&\Big|f_\varepsilon(V)-\sum_{i=1}^k(-1)^i f_0(PU_{\mu_i,\xi_i})\Big|_{\frac{2n}{n+2}}\\
=&\Big|f_\varepsilon(V)-f_0(V)\Big|_{\frac{2n}{n+2}}+\Big|f_0(V)-\sum_{i=1}^k(-1)^i f_0(PU_{\mu_i,\xi_i})\Big|_{\frac{2n}{n+2}}. \qquad (5.14)
\end{aligned}
$$

Similar to the proof of (5.11), we obtain

$$\Big|f_0(V)-\sum_{i=1}^k(-1)^i f_0(PU_{\mu_i,\xi_i})\Big|_{\frac{2n}{n+2}}=\begin{cases}O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } 3\le n\le 5,\\ O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big) & \text{if } n=6,\\ O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n+2}{2(n-2)}}\Big) & \text{if } n\ge 7.\end{cases} \tag{5.15}$$

On the other hand, by Lemma 5.5, there holds

$$\begin{aligned}\int_\Omega\Big|f_\varepsilon(V)-f_0(V)\Big|^{\frac{2n}{n+2}}dx\le&\varepsilon\int_\Omega|V^p\ln\ln(e+V)|^{\frac{2n}{n+2}}dx\\ \le&\varepsilon\int_\Omega\Big|\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big|^{\frac{2n}{n+2}}dx\\ \le&\varepsilon\int_{\Omega\setminus B(\xi,\rho)}\Big|\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big|^{\frac{2n}{n+2}}dx\\ &+\varepsilon\sum_{l=1}^k\int_{\mathcal{A}_l}\Big|\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big|^{\frac{2n}{n+2}}dx.\end{aligned} \tag{5.16}$$

We now observe that

$$\begin{aligned}&\int_{\Omega\setminus B(\xi,\rho)}\Big|\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big|^{\frac{2n}{n+2}}dx\\ \le&C\sum_{i=1}^k\int_{\Omega\setminus B(\xi,\rho)}\Big|U^p_{\mu_i,\xi_i}\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big|^{\frac{2n}{n+2}}dx\\ \le&C\sum_{i=1}^k\mu_i^n\Big|\ln\ln\Big(e+\sum_{i=1}^k(-1)^i\mu_i^{\frac{n-2}{2}}\Big)\Big|^{\frac{2n}{n+2}}dy\\ \le&C\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n}{n-2}}\Big(\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big)^{\frac{2n}{n+2}}.\end{aligned}$$

For the second integral in (5.16), from (3.18), and let $x-\xi=\mu_l y$, then

$$\begin{aligned}&\int_{\mathcal{A}_l}\Big|\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big|^{\frac{2n}{n+2}}dx\\ =&\int_{\mathcal{A}_l}\Big|\Big((-1)^lU_{\mu_l,\xi_l}+\sum_{i\ne l}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\ln\ln\Big(e+(-1)^lU_{\mu_l,\xi_l}+\sum_{i\ne l}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big|^{\frac{2n}{n+2}}dx\\ =&\int_{\mathcal{A}_l}\Big|\Big[(-1)^lU^p_{\mu_l,\xi_l}+O\Big(\sum_{i\ne l}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big]\ln\ln\Big(e+(-1)^lU_{\mu_l,\xi_l}+\sum_{i\ne l}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big|^{\frac{2n}{n+2}}dx\\ =&\int_{\mathcal{A}_l}\Big|(-1)^lU^p_{\mu_l,\xi_l}\ln\ln\Big(e+(-1)^lU_{\mu_l,\xi_l}+\sum_{i\ne l}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big|^{\frac{2n}{n+2}}dx\\ &+C\sum_{i\ne l}^k\int_{\mathcal{A}_l}U_{\mu_i,\xi_i}\Big|\ln\ln\Big(e+(-1)^lU_{\mu_l,\xi_l}+\sum_{i\ne l}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big|^{\frac{2n}{n+2}}dx.\end{aligned} \tag{5.17}$$

For $i>l$, by Lemma 5.5, we have

$$\begin{aligned}
&\int_{\mathcal{A}_l}\Big|U^p_{\mu_l,\xi_l}\ln\ln\Big(e+(-1)^lU_{\mu_l,\xi_l}+\sum_{i\neq l}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big|^{\frac{2n}{n+2}}dx\\
=&\int_{\mathcal{A}_l}\frac{\alpha_n^{\frac{2n}{n-2}}\mu_l^n}{(\mu_l^2+|x-\xi_l|^2)^n}\Big|\ln\ln\Big(e+(-1)^l\frac{\alpha_n\mu_l^{\frac{n-2}{2}}}{(\mu_l^2+|x-\xi_l|^2)^{\frac{n-2}{2}}}+\sum_{i\neq l}^k(-1)^i\frac{\alpha_n\mu_i^{\frac{n-2}{2}}}{(\mu_i^2+|x-\xi_i|^2)^{\frac{n-2}{2}}}\Big)\Big|^{\frac{2n}{n+2}}dx\\
=&\int_{\frac{\mathcal{A}_l}{\mu_l}}\frac{\alpha_n^{\frac{2n}{n-2}}}{(1+|y-\sigma_l|^2)^n}\Big|\ln\ln\Big(e+(-1)^l\mu_l^{-\frac{n-2}{2}}\frac{\alpha_n}{(1+|y-\sigma_l|^2)^{\frac{n-2}{2}}}+o(\varepsilon)\Big)\Big|^{\frac{2n}{n+2}}dy\\
=&\int_{\frac{\mathcal{A}_l}{\mu_l}}\frac{\alpha_n^{\frac{2n}{n-2}}}{(1+|y-\sigma_l|^2)^n}\Big|\ln\ln\mu_l^{-\frac{n-2}{2}}+\ln\Big[1+\frac{\ln\Big(e^{1-\frac{n-2}{2}|\ln\mu_l|}+\frac{\alpha_n}{1+|y-\sigma_l|^2)^{\frac{n-2}{2}}}\Big)}{\frac{n-2}{2}|\ln\mu_l|}\Big]\Big|^{\frac{2n}{n+2}}dy+o(\varepsilon)\\
=&\int_{\frac{\mathcal{A}_l}{\mu_l}}\frac{\alpha_n^{\frac{2n}{n-2}}}{(1+|y-\sigma_l|^2)^n}\Big|\ln\ln\mu_l^{-\frac{n-2}{2}}+\frac{1}{|\ln\mu_l|}\frac{2}{n-2}\ln\frac{\alpha_n}{(1+|y-\sigma_l|^2)^{\frac{n-2}{2}}}\Big|^{\frac{2n}{n+2}}dy+o(\varepsilon)\\
\le&C(\ln|\ln\mu_1|)^{\frac{2n}{n+2}}=O\Big(\Big(\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big)^{\frac{2n}{n+2}}\Big).
\end{aligned}$$

In a same way, the estimate of second term in (5.17) is

$$\sum_{i\neq l}^k\int_{\mathcal{A}_l}U_{\mu_i,\xi_i}\Big|\ln\ln\Big(e+(-1)^lU_{\mu_l,\xi_l}+\sum_{i\neq l}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big|^{\frac{2n}{n+2}}dx=O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big).$$

Thus,

$$\Big|f_\varepsilon(V)-f_0(V)\Big|_{\frac{2n}{n+2}}=O\Big(\varepsilon\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big).\tag{5.18}$$

Combining (5.14), (5.15) and (5.18), we obtain (5.12). Similar to the proof of (5.16), the estimate (5.13) holds. □

Lemma 5.7. *There holds*

$$\begin{aligned}
P_3&=\int_\Omega\Big[\sum_{i=1}^k(-1)^if_0(PU_{\mu_i,\xi_i})-f_\varepsilon(V)\Big]\psi^h_{\mu_j,\xi_j}dx\\
&=\begin{cases}
\alpha_na_1\frac{\varepsilon}{|\ln\varepsilon|^2}H(\xi,\xi)d_1^{n-2}+a_3\frac{\varepsilon}{|\ln\varepsilon|^2}\sum\limits_{i=1}^{k-1}\Big(\frac{d_{i+1}}{d_i}\Big)^{\frac{n-2}{2}}g(\sigma_i)-\frac{2k^2}{(n-2)^2}a_4\varepsilon\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\\
\quad -a_4\frac{\varepsilon}{|\ln\varepsilon|^2}\sum\limits_{i=1}^k\frac{2}{2i-1}|\ln d_i|+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\quad\text{if } h=0,\\
\frac12\alpha_na_2\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}d_1^{n-1}\partial_{\xi_h}\varphi(\xi)+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big)\quad\text{if } h=1,\cdots,n,
\end{cases}
\end{aligned}$$

where a_1-a_4 and $g(\sigma_i)$ are given in Proposition 2.2 for $i=1,\cdots,k$.

Proof. We have

$$P_3=\int_\Omega\Big[\sum_{i=1}^k(-1)^if_0(PU_{\mu_i,\xi_i})-f_\varepsilon(V)\Big]\psi^h_{\mu_j,\xi_j}dx$$

$$
\begin{aligned}
&= \int_\Omega \Big[\sum_{i=1}^k (-1)^i f_0(U_{\mu_i,\xi_i}) - f_0(V)\Big]\psi^h_{\mu_j,\xi_j} dx \\
&\quad + \int_\Omega \Big[\sum_{i=1}^k (-1)^i f_0(PU_{\mu_i,\xi_i}) - \sum_{i=1}^k (-1)^i f_0(U_{\mu_i,\xi_i})\Big]\psi^h_{\mu_j,\xi_j} dx \\
&\quad + \int_\Omega \Big[f_0(V) - f_\varepsilon(V)\Big]\psi^h_{\mu_j,\xi_j} dx = J_1 + J_2 + J_3.
\end{aligned}
$$

Estimate of J_1: One has

$$
\begin{aligned}
J_1 =& \int_\Omega \Big[\sum_{i=1}^k (-1)^i f_0(U_{\mu_i,\xi_i}) - f_0(V)\Big]\psi^h_{\mu_j,\xi_j} dx \\
=& \sum_{i=1}^k \int_\Omega (-1)^i f_0(U_{\mu_i,\xi_i})\psi^h_{\mu_j,\xi_j} dx - \int_\Omega f_0(V)\psi^h_{\mu_j,\xi_j} dx \\
=& \sum_{i=1}^k (-1)^i \int_\Omega f_0(U_{\mu_i,\xi_i}) P\psi^h_{\mu_j,\xi_j} dx + \sum_{i=1}^k (-1)^i \int_\Omega f_0(U_{\mu_i,\xi_i})(\psi^h_{\mu_j,\xi_j} - P\psi^h_{\mu_j,\xi_j}) dx \\
&- \int_\Omega f_0(V)\psi^h_{\mu_j,\xi_j} dx. \qquad (5.19)
\end{aligned}
$$

For the first term in (5.19), if $h = 0$ and $i \neq j$, from Lemma 5.4, we deduce

$$
\begin{aligned}
&\int_\Omega f_0(U_{\mu_i,\xi_i}) P\psi^0_{\mu_j,\xi_j} dx \\
=& \frac{n-2}{2} a_2 \alpha_n \mu_j^{\frac{n-2}{2}} \int_\Omega \frac{\mu_i^{\frac{n+2}{2}}}{(\mu_i^2 + |x-\xi_i|^2)^{\frac{n+2}{2}}} G(x,\xi_j) dx + o\Big(\frac{\varepsilon}{|\ln \varepsilon|^2}\Big) \\
=& \frac{n-2}{2} a_2 \alpha_n \mu_i^{\frac{n-2}{2}} \mu_j^{\frac{n-2}{2}} \int_{\frac{\Omega-\xi_i}{\mu_i}} \frac{1}{(1+|y|^2)^{\frac{n+2}{2}}} G(\mu_i y + \xi_i, \xi_j) dy + o\Big(\frac{\varepsilon}{|\ln \varepsilon|^2}\Big) \\
=& \frac{n-2}{2} a_2^2 \mu_j^{\frac{n-2}{2}} \mu_i^{\frac{n-2}{2}} G(\xi_i,\xi_j) + o\Big(\frac{\varepsilon}{|\ln \varepsilon|^2}\Big). \qquad (5.20)
\end{aligned}
$$

Let $h = 0$ and $i = j$, from (5.2) and (2.10), there holds

$$
\begin{aligned}
\int_\Omega f_0(U_{\mu_i,\xi_i}) P\psi^0_{\mu_i,\xi_i} dx =& \int_\Omega U^p_{\mu_i,\xi_i}\psi^0_{\mu_i,\xi_i} dx - \frac{n-2}{2}\alpha_n \mu_i^{\frac{n-2}{2}} \int_\Omega U^p_{\mu_i,\xi_i}\Big(H(x,\xi_i) + O(\mu_i^{\frac{n+4}{2}})\Big) dx \\
=& \mu_i^{n-\frac{n+2}{2}+1+\frac{n-2}{2}-(n-2)} \int_{\frac{\Omega-\xi_i}{\mu_i}} U^p(y)\Big(\frac{\partial U}{\partial \mu}\Big)_{|\mu=1} dy \\
&- \frac{n-2}{2}\mu_i^{n-2} H(\xi_i,\xi_i) \int_{\frac{\Omega-\xi_i}{\mu_i}} U^p(y) dy + O(\mu_i^{\frac{3n}{2}}) \\
=& -\frac{n-2}{2} a_2 \mu_i^{n-2} H(\xi_i,\xi_i) + O\Big(\frac{\varepsilon}{|\ln \varepsilon|^2}\Big). \qquad (5.21)
\end{aligned}
$$

If $h = 1, \cdots, n$ and $i \neq j$, from Lemma 5.4, we get

$$
\begin{aligned}
\int_\Omega f_0(U_{\mu_i,\xi_i}) P\psi^h_{\mu_j,\xi_j} dx =& \int_\Omega U^{2^*-1}_{\mu_i,\xi_i} P\psi^h_{\mu_j,\xi_j} dx \\
=& a_2 \mu_j^{\frac{n}{2}} \mu_i^{\frac{n-2}{2}} \int_{\frac{\Omega-\xi_i}{\mu_i}} \frac{1}{(1+|y|^2)^{\frac{n+2}{2}}} \frac{\partial G}{\partial \xi_j^h}(\mu_i y + \xi_i, \xi_j) dy + o\Big(\Big(\frac{\varepsilon}{|\ln \varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big)
\end{aligned}
$$

$$=a_2^2\mu_j^{\frac{n}{2}}\mu_i^{\frac{n-2}{2}}\frac{\partial G}{\partial \xi_j^h}(\xi_i,\xi_j)+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big). \tag{5.22}$$

If $h=1,\cdots,n$ and $i=j$, from (5.3) and (2.10), we have

$$\begin{aligned}\int_\Omega f_0(U_{\mu_i,\xi_i})P\psi^h_{\mu_i,\xi_i}dx=&\int_\Omega U^{2^*-1}_{\mu_i,\xi_i}P\psi^h_{\mu_i,\xi_i}dx\\ =&\int_\Omega U^{2^*-1}_{\mu_i,\xi_i}\psi^h_{\mu,\xi_i}(x)dx-\alpha_n\mu_i^{\frac{n}{2}}\int_\Omega U^{2^*-1}_{\mu_i,\xi_i}\partial_{\xi_i^h}H(x,\xi_i)dx+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big)\\ =&\mu_i^{n-\frac{n+2}{2}+1-\frac{n-2}{2}}\int_{\frac{\Omega-\xi_i}{\mu_i}}U^{2^*-1}(y)\psi^h(y)dy\\ &-\alpha_n\mu_i^{\frac{n}{2}-\frac{n+2}{2}+n}\partial_{\xi_i^h}H(\xi_i,\xi_i)\int_{\frac{\Omega-\xi_i}{\mu_i}}U^{2^*-1}(y)dy+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big)\\ =&\mu_i\int_{\frac{\Omega-\xi_i}{\mu_i}}U^{2^*-1}(y)\psi^h(y)dy\\ &-\alpha_n\mu_i^{n-1}\partial_{\xi_i^h}H(\xi_i,\xi_i)\int_{\frac{\Omega-\xi_i}{\mu_i}}U^{2^*-1}(y)dy+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big)\\ =&-\alpha_n a_2\mu_i^{n-1}\frac{\partial H}{\partial\xi_i^h}(\xi_i,\xi_i)+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big).\end{aligned} \tag{5.23}$$

It remains to estimate the second term in (5.19), if $h=0$, by (5.2), it holds

$$\begin{aligned}&\int_\Omega f_0(U_{\mu_i,\xi_i})(\psi^0_{\mu_j,\xi_j}-P\psi^0_{\mu_j,\xi_j})dx\\ =&\frac{n-2}{2}\alpha_n\mu_i^{\frac{n-2}{2}}\mu_j^{\frac{n-2}{2}}\int_{\frac{\Omega-\xi_i}{\mu_i}}\frac{1}{(1+|y|^2)^{\frac{n+2}{2}}}H(\mu_i y+\xi_i,\xi_j)dy+o(\mu_i^{\frac{n-2}{2}}\mu_j^{\frac{n+4}{2}})\\ =&\begin{cases}\frac{n-2}{2}a_2\mu_i^{\frac{n-2}{2}}\mu_j^{\frac{n-2}{2}}H(\xi_i,\xi_j)+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } i>j,\\ \frac{n-2}{2}a_2\mu_i^{n-2}H(\xi_i,\xi_i)+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } i=j.\end{cases}\end{aligned} \tag{5.24}$$

If $h=1,\cdots,n$, by (5.3), we obtain

$$\begin{aligned}&\int_\Omega f_0(U_{\mu_i,\xi_i})(\psi^h_{\mu_j,\xi_j}-P\psi^h_{\mu_j,\xi_j})dx\\ =&\alpha_n\mu_i^{\frac{n-2}{2}}\mu_j^{\frac{n}{2}}\int_{\frac{\Omega-\xi_i}{\mu_i}}\frac{1}{(1+|y|^2)^{\frac{n+2}{2}}}\partial_{\xi_j^h}H(\mu_i y+\xi_i,\xi_j)dy+o(\mu_i^{\frac{n-2}{2}}\mu_j^{\frac{n+2}{2}})\\ =&\begin{cases}a_2\mu_i^{\frac{n-2}{2}}\mu_j^{\frac{n}{2}}\partial_{\xi_j^h}H(\xi_i,\xi_j)+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } i>j,\\ a_2\mu_i^{n-1}\partial_{\xi_i^h}H(\xi_i,\xi_i)+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } i=j.\end{cases}\end{aligned} \tag{5.25}$$

For the last integral in (5.19),

$$\int_\Omega f_0(V)\psi^h_{\mu_j,\xi_j}dx=\int_{\Omega\setminus B(\xi,\rho)}f_0(V)\psi^h_{\mu_j,\xi_j}dx+\sum_{l=1}^k\int_{\mathcal{A}_l}f_0(V)\psi^h_{\mu_j,\xi_j}dx. \tag{5.26}$$

If $h=0$, by (2.9), a direct computation shows that

$$\begin{aligned}\int_{\Omega\setminus B(\xi,\rho)}\Big|f_0(V)\psi^0_{\mu_j,\xi_j}\Big|dx \leq& C\frac{n-2}{2}\alpha_n\mu_j^{\frac{n-2}{2}}\sum_{i=1}^k\int_{\Omega\setminus B(\xi,\rho)}\Big|U^p_{\mu_i,\xi_i}\frac{|x-\xi_j|^2-\mu_j^2}{(\mu_j^2+|x-\xi_j|^2)^{\frac{n}{2}}}\Big|dx\\ \leq& C\frac{n-2}{2}\alpha_n\mu_j^{\frac{n-2}{2}}\sum_{i=1}^k\int_{\Omega\setminus B(\xi,\rho)}\Big|\frac{\mu_i^{\frac{n+2}{2}}}{(\mu_i^2+|x-\xi_i|^2)^{\frac{n+2}{2}}}\frac{|x-\xi_j|^2-\mu_j^2}{(\mu_j^2+|x-\xi_j|^2)^{\frac{n}{2}}}\Big|dx\\ =& o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big).\end{aligned}$$

If $h=1,\cdots,n$, we obtain

$$\begin{aligned}\int_{\Omega\setminus B(\xi,\rho)}\Big|f_0(V)\psi^0_{\mu_j,\xi_j}\Big|dx \leq& C(n-2)\alpha_n\mu_j^{\frac{n}{2}}\sum_{i=1}^k\int_{\Omega\setminus B(\xi,\rho)}\Big|(-1)^iU^p_{\mu_i,\xi_i}\frac{x^h-\xi_j^h}{(\mu_j^2+|x-\xi_j|^2)^{\frac{n}{2}}}\Big|dx\\ \leq& C\frac{n-2}{2}\alpha_n\mu_j^{\frac{n}{2}}\sum_{i=1}^k\int_{\Omega\setminus B(\xi,\rho)}\Big|\frac{\mu_i^{\frac{n+2}{2}}}{(\mu_i^2+|x-\xi_i|^2)^{\frac{n+2}{2}}}\frac{x^h-\xi_j^h}{(\mu_j^2+|x-\xi_j|^2)^{\frac{n}{2}}}\Big|dx\\ =& o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big).\end{aligned}$$

On the other hand, we estimate the second term in (5.26),

$$\begin{aligned}&\int_{\mathcal{A}_l}\Big|f_0(V)\psi^h_{\mu_j,\xi_j}\Big|dx\\ =&\int_{\mathcal{A}_l}\Big|\Big((-1)^lPU_{\mu_l,\xi_l}+\sum_{i\neq l}^k(-1)^iPU_{\mu_i,\xi_i}\Big)^p\psi^h_{\mu_j,\xi_j}\Big|dx\\ =&\int_{\mathcal{A}_l}\Big|(-1)^l(PU_{\mu_l,\xi_l})^p\psi^h_{\mu_j,\xi_j}\Big|dx+O\Big(\int_{\mathcal{A}_l}\Big|\Big(\sum_{i\neq l}^k(-1)^iPU_{\mu_i,\xi_i}\Big)\psi^h_{\mu_j,\xi_j}\Big|dx\Big)\\ =&\int_{\mathcal{A}_l}\Big|(-1)^l\Big((PU_{\mu_l,\xi_l})^p-U^p_{\mu_l,\xi_l}\Big)\psi^h_{\mu_j,\xi_j}\Big|dx+\int_{\mathcal{A}_l}|(-1)^lU^p_{\mu_l,\xi_l}\psi^h_{\mu_j,\xi_j}|dx\\ &+O\Big(\int_{\mathcal{A}_l}\Big|\Big(\sum_{i\neq l}^k(-1)^iPU_{\mu_i,\xi_i}\Big)\psi^h_{\mu_j,\xi_j}\Big|dx\Big).\end{aligned}\tag{5.27}$$

On the fixed annulus $\mathcal{A}_l$, by (5.1), for $h=0$, let $x-\xi=\mu_l y$, then

$$\begin{aligned}&\int_{\mathcal{A}_l}\Big|\Big((PU_{\mu_l,\xi_l})^p-U^p_{\mu_l,\xi_l}\Big)\psi^0_{\mu_j,\xi_j}\Big|dx\\ =&\int_{\mathcal{A}_l}\Big|\Big[\Big(U_{\mu_l,\xi_l}-\alpha_n\mu_l^{\frac{n-2}{2}}H(x,\xi_l)+O(\mu_l^{\frac{n+2}{2}})\Big)^p-U^p_{\mu_l,\xi_l}\Big]\psi^0_{\mu_j,\xi_j}\Big|dx\\ =&\frac{(n-2)\alpha_n}{2}\mu_l^{\frac{n-2}{2}+n}\mu_j^{\frac{n-2}{2}}O\left(-\alpha_nH(\xi_l,\xi_l)+O(\mu_l^2)\right)\int_{\frac{\mathcal{A}_l}{\mu_l}}\left|\frac{\Big[|y-\sigma_j\frac{\mu_j}{\mu_l}|^2-(\frac{\mu_j}{\mu_l})^2\Big]\mu_l^2}{\Big((\frac{\mu_j}{\mu_l})^2+|y-\sigma_j\frac{\mu_j}{\mu_l}|^2\Big)^{\frac{n}{2}}\mu_l^n}\right|dy\\ =&\begin{cases}O(\mu_j^{\frac{n+2}{2}}\mu_l^{\frac{n-2}{2}}H(\xi_l,\xi_l)) & \text{if } j>l,\\ O(\mu_l^nH(\xi_l,\xi_l)) & \text{if } j=l,\end{cases}\end{aligned}$$

$$= o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big).$$

Similarly, for $h=1,\cdots,n$, we get

$$\begin{aligned}
&\int_{\mathcal{A}_l}\Big|\Big((PU_{\mu_l,\xi_l})^p-U^p_{\mu_l,\xi_l}\Big)\psi^h_{\mu_j,\xi_j}\Big|dx\\
=&\int_{\mathcal{A}_l}\Big|\Big[\Big(U_{\mu_l,\xi_l}-\alpha_n\mu_l^{\frac{n-2}{2}}H(x,\xi_l)+O(\mu_l^{\frac{n+2}{2}})\Big)^p-U^p_{\mu_l,\xi_l}\Big]\psi^h_{\mu_j,\xi_j}\Big|dx\\
=&-(n-2)\alpha_n\mu_j^{\frac{n}{2}}\int_{\mathcal{A}_l}\Big|O\Big(-\alpha_n\mu_l^{\frac{n-2}{2}}H(x,\xi_l)+O(\mu_l^{\frac{n+2}{2}})\Big)\frac{(x-\xi_j)_h}{(\mu_j^2+|x-\xi_j|^2)^{\frac{n}{2}}}\Big|dx\\
=&\begin{cases}O\Big(\mu_j^{\frac{n}{2}}\mu_l^{\frac{n}{2}}H(\xi_l,\xi_l)\Big) & \text{if } j>l,\\ O\Big(\mu_l^nH(\xi_l,\xi_l)\Big) & \text{if } j=l,\end{cases}\\
=&o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big).
\end{aligned}$$

The second term in (5.27). By (5.1), for $h=0$, if $j>l$, let $x-\xi=\mu_l y$, then

$$\begin{aligned}
&\int_{\mathcal{A}_l}|U^p_{\mu_l,\xi_l}\psi^0_{\mu_j,\xi_j}|dx\\
=&\frac{n-2}{2}\alpha_n^{p+1}\mu_j^{\frac{n-2}{2}}\int_{\frac{\mathcal{A}_l}{\mu_l}}\Big|\frac{\mu_l^{-\frac{n+2}{2}}}{(1+|y-\sigma_l|^2)^{\frac{n+2}{2}}}\frac{\Big(|y-\sigma_j\frac{\mu_j}{\mu_l}|^2-(\frac{\mu_j}{\mu_l})^2\Big)\mu_l^2}{\Big((\frac{\mu_j}{\mu_l})^2+|y-\sigma_j\frac{\mu_j}{\mu_l}|^2\Big)^{\frac{n}{2}}\mu_l^n}\mu_l^n\Big|dy\\
=&\frac{n-2}{2}\alpha_n^{p+1}\Big(\frac{\mu_j}{\mu_l}\Big)^{\frac{n-2}{2}}\Big(\int_{\mathbb{R}^n}\Big|\frac{y^{2-n}}{(1+|y-\sigma_l|^2)^{\frac{n+2}{2}}}\Big|dy+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\Big)\\
=&\frac{n-2}{2}\alpha_n^{p+1}\Big(\frac{\mu_{l+1}}{\mu_l}\Big)^{\frac{n-2}{2}}\Big(\int_{\mathbb{R}^n}\Big|\frac{y^{2-n}}{(1+|y-\sigma_l|^2)^{\frac{n+2}{2}}}\Big|dy+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\Big).
\end{aligned}$$

For $h=1,\cdots,n$, if $j>l$, let $x-\xi=\mu_l y$, there holds

$$\begin{aligned}
\int_{\mathcal{A}_l}|U^p_{\mu_l,\xi_l}\psi^h_{\mu_j,\xi_j}|dx=&(n-2)\alpha_n^{p+1}\mu_j^{\frac{n}{2}}\int_{\frac{\mathcal{A}_l}{\mu_l}}\Big|\frac{\mu_l^{-\frac{n+2}{2}}}{(1+|y-\sigma_l|^2)^{\frac{n+2}{2}}}\frac{(y_h-\sigma_j\frac{\mu_j}{\mu_l})\mu_l}{\Big((\frac{\mu_j}{\mu_l})^2+|y-\sigma_j\frac{\mu_j}{\mu_l}|^2\Big)^{\frac{n}{2}}\mu_l^n}\mu_l^n\Big|dy\\
=&o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big).
\end{aligned}$$

The last term in (5.27). For $h=0$ and $j>i$,

$$\begin{aligned}
&\int_{\mathcal{A}_l}\Big|\Big(\sum_{i\neq l}^k(-1)^iPU_{\mu_i,\xi_i}\Big)\psi^0_{\mu_j,\xi_j}\Big|dx\\
\leq&\sum_{i\neq l}^k\int_{\mathcal{A}_l}|U_{\mu_i,\xi_i}\psi^0_{\mu_j,\xi_j}|dx\\
=&\frac{n-2}{2}\alpha_n^2\sum_{i\neq l}^k\mu_i^{-\frac{n-2}{2}}\mu_j^{\frac{n-2}{2}}\int_{\frac{\mathcal{A}_l}{\mu_i}}\Big|\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n-2}{2}}}\frac{\Big(|y-\sigma_j\frac{\mu_j}{\mu_i}|^2-(\frac{\mu_j}{\mu_i})^2\Big)\mu_i^2}{\Big((\frac{\mu_j}{\mu_i})^2+|y-\sigma_j\frac{\mu_j}{\mu_i}|^2\Big)^{\frac{n}{2}}}\Big|dy
\end{aligned}$$

$$=o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big).$$

For $h=1,\cdots,n$ and $j>i$,

$$\begin{aligned}
&\int_{\mathcal{A}_l}\Big|\Big(\sum_{i\neq l}^{k}(-1)^i PU_{\mu_i,\xi_i}\Big)\psi^h_{\mu_j,\xi_j}\Big|dx\\
\leq&\sum_{i\neq l}^{k}\int_{\mathcal{A}_l}|U_{\mu_i,\xi_i}\psi^h_{\mu_j,\xi_j}|dx\\
=&(n-2)\alpha_n^2\sum_{i\neq l}^{k}\mu_j^{\frac{n}{2}}\int_{\frac{\mathcal{A}_l}{\mu_l}}\Big|\frac{\mu_i^{-\frac{n-2}{2}}}{(1+|y-\sigma_i|^2)^{\frac{n-2}{2}}}\frac{(y^h-\sigma_j\frac{\mu_j}{\mu_l})\mu_i}{\Big((\frac{\mu_j}{\mu_i})^2+|y-\sigma_j\frac{\mu_j}{\mu_i}|^2\Big)^{\frac{n}{2}}\mu_i^n}\Big|\mu_i^n dy\\
=&o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big).
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
&\int_\Omega f_0(V)\psi^h_{\mu_j,\xi_j}dx\\
=&\begin{cases}
a_3\Big(\frac{\mu_{l+1}}{\mu_l}\Big)^{\frac{n-2}{2}}g(\sigma_l)+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } l=1,\cdots,k-1,\ h=0,\\
o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } l=k,
\end{cases}
\end{aligned}\tag{5.28}$$

where a_3 and $g(\sigma_l)$ are defined in Proposition 2.2. From (5.19)-(5.25) and (5.28), we obtain

$$J_1=\begin{cases}
-a_3\dfrac{\varepsilon}{|\ln\varepsilon|^2\sum_{l=1}^{k-1}\Big(\frac{d_{l+1}}{d_l}\Big)^{\frac{n-2}{2}}}g(\sigma_l)+o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } l=1,\cdots,k-1,\ h=0,\\
o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } l=k,\ h=1,\cdots,n.
\end{cases}\tag{5.29}$$

Estimate of J_2: By Lemma 5.1, there holds

$$\begin{aligned}
J_2=&\sum_{i=1}^{k}\int_\Omega(-1)^i\Big[f_0(PU_{\mu_i,\xi_i})-f_0(U_{\mu_i,\xi_i})\Big]\psi^h_{\mu_j,\xi_j}dx\\
=&\sum_{i=1}^{k}\int_\Omega(-1)^i f_0'(U_{\mu_i,\xi_i})(PU_{\mu_i,\xi_i}-U_{\mu_i,\xi_i})\psi^h_{\mu_j,\xi_j}dx\\
&+\sum_{i=1}^{k}\int_\Omega(-1)^i\Big[f_0(PU_{\mu_i,\xi_i})-f_0(U_{\mu_i,\xi_i})-f_0'(U_{\mu_i,\xi_i})(PU_{\mu_i,\xi_i}-U_{\mu_i,\xi_i})\Big]\psi^h_{\mu_j,\xi_j}dx\\
=&(2^*-1)\sum_{i=1}^{k}\mu_i^{\frac{n-2}{2}}\int_\Omega U_{\mu_i,\xi_i}^{2^*-2}\Big(-\alpha_n H(x,\xi_i)+O(\mu_i^2)\Big)\psi^h_{\mu_j,\xi_j}dx\\
&-\int_\Omega(-1)^i\Big[f_0(PU_{\mu_i,\xi_i})-f_0(U_{\mu_i,\xi_i})-f_0'(U_{\mu_i,\xi_i})(PU_{\mu_i,\xi_i}-U_{\mu_i,\xi_i})\Big]\psi^h_{\mu_j,\xi_j}dx.
\end{aligned}$$

Moreover, for $l=0,\cdots,n$, by Hölder inequality, (5.6), (5.7) and (5.10), one has

$$\int_\Omega\Big|(-1)^i\Big[f_0(PU_{\mu_i,\xi_i})-f_0(U_{\mu_i,\xi_i})-f_0'(U_{\mu_i,\xi_i})(PU_{\mu_i,\xi_i}-U_{\mu_i,\xi_i})\Big]\psi^h_{\mu_j,\xi_j}\Big|dx$$

$$\leq \Big|f_0(PU_{\mu_i,\xi_i}) - f_0(U_{\mu_i,\xi_i}) - f_0'(U_{\mu_i,\xi_i})(PU_{\mu_i,\xi_i} - U_{\mu_i,\xi_i})\Big|_{\frac{n}{2}} |\psi^h_{\mu_j,\xi_j}|_{\frac{n}{n-2}} = o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big).$$

If $h=0$ and $j=i$, by (2.9), we get

$$\begin{aligned}
&(2^*-1)\alpha_n \sum_{i=1}^k \mu_i^{\frac{n-2}{2}} \int_\Omega U^{2^*-2}_{\mu_i,\xi_i} H(x,\xi_i)\psi^0_{\mu_i,\xi_i} dx\\
=&(2^*-1)\alpha_n \sum_{i=1}^k \mu_i^{\frac{3(n-2)}{2}} \mu_i^{-\frac{n-2}{2}} \int_{\frac{\Omega-\xi_i}{\mu_i}} U^{2^*-2}(y) H(\mu_i y+\xi_i,\xi_i)\psi^0\Big(\frac{x-\xi_i}{\mu_i}\Big) dy\\
=&\alpha_n a_1 \sum_{i=1}^k \mu_i^{n-2} H(\xi_i,\xi_i) + o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big),
\end{aligned}$$

where a_1 is given in Proposition 2.2. If $h=1,\cdots,n$ and $j=i$, by (2.10), one has

$$\begin{aligned}
&(2^*-1)\alpha_n \sum_{i=1}^k \mu_i^{\frac{n-2}{2}} \int_\Omega U^{2^*-2}_{\mu_i,\xi_i}(x) H(x,\xi_i)\psi^h_{\mu_i,\xi_i} dx\\
=&\alpha_n \sum_{i=1}^k \mu_i^{\frac{n}{2}} \int_\Omega H(x,\xi_i) \frac{\partial}{\partial \xi_i^h} U^{2^*-1}_{\mu_i,\xi_i}(x) dx\\
=&\alpha_n \sum_{i=1}^k \mu_i^{\frac{n}{2}} \Big(\mu_i^{\frac{n-2}{2}} \frac{\partial}{\partial \xi_i^h} \int_{\frac{\Omega-\xi_i}{\mu_i}} U^{2^*-1}_{\mu_i,\xi_i}(y) H(\mu_i y+\xi_i,\xi_i) dy - \mu_i^{\frac{n-2}{2}} \int_{\frac{\Omega-\xi_i}{\mu_i}} U^{2^*-1}_{\mu_i,\xi_i} \frac{\partial H(\mu_i y+\xi_i,\xi_i)}{\partial \xi_i^h} dy\Big)\\
=&\alpha_n a_2 \sum_{i=1}^k \mu_i^{n-1}\Big(\frac{\partial(H(\xi_i,\xi_i))}{\partial \xi_i^h} - \frac{\partial H(\xi_i,\xi_i)}{\partial \xi_i^h} + O(\mu_i)\Big)\\
=&\frac{1}{2}\alpha_n a_2 \sum_{i=1}^k \mu_i^{n-1} \partial_{\xi_i^h}\rho(\xi_i) + o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big).
\end{aligned}$$

In a same way, if $h=1,\cdots,n$ and $j\neq i$, we have

$$(2^*-1)\alpha_n \sum_{i=1}^k \mu_i^{\frac{n-2}{2}} \int_\Omega U^{2^*-2}_{\mu_i,\xi_i}(x) H(x,\xi_i)\psi^h_{\mu_j,\xi_j} dx = o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big).$$

If $h=0$ and $j\neq i$,

$$(2^*-1)\alpha_n \sum_{i=1}^k \mu_i^{\frac{n-2}{2}} \int_\Omega U^{2^*-2}_{\mu_i,\xi_i} H(x,\xi_i)\psi^0_{\mu_j,\xi_j} dx = o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big).$$

As a consequence, there holds

$$J_2 = \begin{cases} \alpha_n a_1 d_1 \frac{\varepsilon}{|\ln\varepsilon|^2} H(\xi,\xi) + o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } h=0,\\ \frac{1}{2}\alpha_n a_2 \Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}} \partial_{\xi_h}\rho(\xi) + o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } h=1,\cdots,n. \end{cases}$$

Estimate of J_3: By Taylor expansion with respect to ε, we have

$$\begin{aligned}
J_3 =& \int_\Omega \Big[f_0(V) - f_\varepsilon(V)\Big]\psi^h_{\mu_j,\xi_j} dx\\
=&\varepsilon \int_\Omega V^p \ln\ln(e+V)\psi^h_{\mu_j,\xi_j} dx - \varepsilon^2 \int_\Omega V^p \Big(\ln\ln(e+V)\Big)^2 \psi^h_{\mu_j,\xi_j} dx. \qquad (5.30)
\end{aligned}$$

For the second term in (5.30), from Lemma 5.5 and the annulus given in (3.18), it holds

$$
\begin{aligned}
&\int_\Omega \Big|V^p\Big(\ln\ln(e+V)\Big)^2\psi^h_{\mu_j,\xi_j}\Big|dx\\
\le&\int_\Omega \Big|\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\Big[\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big]^2\psi^h_{\mu_j,\xi_j}\Big|dx\\
=&\int_{B(\xi,\rho)} \Big|\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\Big[\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big]^2\psi^h_{\mu_j,\xi_j}\Big|dx+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\\
=&\sum_{i=1}^k\int_{\mathcal{A}_i}\Big|\Big((-1)^iU_{\mu_i,\xi_i}+(-1)^j\sum_{j\neq i}^kU_{\mu_j,\xi_j}\Big)^p\\
&\times\Big[\ln\ln\Big(e+(-1)^iU_{\mu_i,\xi_i}+(-1)^j\sum_{j\neq i}^kU_{\mu_j,\xi_j}\Big)\Big]^2\psi^h_{\mu_j,\xi_j}\Big|dx+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big),
\end{aligned}
\tag{5.31}
$$

and on each annulus $\mathcal{A}_i$, we change variable setting $\mu_i y=x-\xi$, for $h=0$, by (2.9), then

$$
\begin{aligned}
&\int_{\mathcal{A}_i}\Big|\Big((-1)^iU_{\mu_i,\xi_i}+(-1)^j\sum_{j\neq i}^kU_{\mu_j,\xi_j}\Big)^p\Big[\ln\ln\Big(e+(-1)^iU_{\mu_i,\xi_i}+(-1)^j\sum_{j\neq i}^kU_{\mu_j,\xi_j}\Big)\Big]^2|\psi^0_{\mu_j,\xi_j}|\Big|dx\\
=&\frac{(n-2)\alpha_n^{p+1}}{2}\mu_j^{\frac{n-2}{2}}\int_{\frac{\mathcal{A}_i}{\mu_i}}\Big|\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n-2}{2}}}+\mu_i^{\frac{n-2}{2}}\sum_{j\neq i}^k\frac{\mu_j^{\frac{n-2}{2}}}{(\mu_j^2+|\mu_iy-\mu_j\sigma_j|^2)^{\frac{n-2}{2}}}\Big|^p\\
&\times\Big|\ln\ln\Big[\alpha_n\mu_i^{-\frac{n-2}{2}}\Big(e+\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n-2}{2}}}+\mu_i^{\frac{n-2}{2}}\sum_{j\neq i}^k\frac{\mu_j^{\frac{n-2}{2}}}{(\mu_j^2+|\mu_iy-\mu_j\sigma_j|^2)^{\frac{n-2}{2}}}\Big)\Big]\Big|^2\\
&\times\Big|\frac{|\mu_iy-\mu_j\sigma_j|^2-\mu_j^2}{(\mu_j^2+|\mu_iy-\mu_j\sigma_j|^2)^{\frac{n}{2}}}\Big|dy\\
=&O\Big(\Big|\ln|\ln\mu_i|\Big|^2+\Big(\frac{\mu_i}{\mu_{i-1}}\Big)^{\frac{n}{2}}+\Big(\frac{\mu_{i+1}}{\mu_i}\Big)^{\frac{n}{2}}\Big)=O(\ln|\ln\mu_i|)=O\Big(\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big).
\end{aligned}
$$

Similarly, for $h=1,\cdots,n$, we have

$$
\begin{aligned}
&\int_{\mathcal{A}_i}\Big((-1)^iU_{\mu_i,\xi_i}+(-1)^j\sum_{j\neq i}^kU_{\mu_j,\xi_j}\Big)^p\Big[\ln\ln\Big(e+(-1)^iU_{\mu_i,\xi_i}+(-1)^j\sum_{j\neq i}^kU_{\mu_j,\xi_j}\Big)\Big]^2|\psi^h_{\mu_j,\xi_j}|dx\\
=&O\Big(\ln|\ln\mu_i|+\Big(\frac{\mu_i}{\mu_{i-1}}\Big)^{\frac{n+2}{2}}+\Big(\frac{\mu_{i+1}}{\mu_i}\Big)^{\frac{n+2}{2}}\Big)=O(\ln|\ln\mu_i|)=O\Big(\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big).
\end{aligned}
$$

Thus the second term in (5.30) becomes

$$
\int_\Omega\Big|V^p\Big(\ln\ln(e+V)\Big)^2\psi^h_{\mu_j,\xi_j}\Big|dx=O((\ln|\ln\mu_i|)^2)=O\Big(\Big(\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big)^2\Big).
$$

Consequently,

$$
\begin{aligned}
J_3=&\int_\Omega\Big(f_0(V)-f_\varepsilon(V)\Big)\psi^h_{\mu_j,\xi_j}dx\\
=&\varepsilon\int_\Omega V^p\Big(\ln\ln(e+V)\Big)\psi^h_{\mu_j,\xi_j}dx+O\Big(\varepsilon^2\Big(\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big)^2\Big).
\end{aligned}
\tag{5.32}
$$

Moreover,

$$\begin{aligned}
&\int_\Omega V^p\Big(\ln\ln(e+V)\Big)\psi^h_{\mu_j,\xi_j}dx\\
&=\int_\Omega\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\Big[\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big]\psi^h_{\mu_j,\xi_j}dx\\
&\quad-\Big[\int_\Omega\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\Big[\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big]-V^p\ln\ln(e+V)\Big]\psi^h_{\mu_j,\xi_j}dx. \qquad (5.33)
\end{aligned}$$

Let us set $h(u)=u^p\ln\ln(e+u)$, by the mean value theorem, one has

$$0\le h(u)-h(v)\le Cu^{p-1}\Big(\ln\ln(e+u)+1\Big)(u-v)\quad\text{for } 0\le v\le u.$$

Then

$$\begin{aligned}
&\int_\Omega\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^{p-1}\Big(\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)+1\Big)\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}-V\Big)\psi^h_{\mu_j,\xi_j}dx\\
=&\sum_{i=1}^k\int_{\mathcal{A}_i}\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^{p-1}\Big(\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)+1\Big)\sum_{i=1}^k(U_{\mu_i,\xi_i}-PU_{\mu_i,\xi_i})\psi^h_{\mu_j,\xi_j}dx\\
&+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big).
\end{aligned}$$

Moreover, on each annulus $\mathcal{A}_i$, $i=1,\cdots,k$, if $h=0$, by Lemma 5.1 , (2.9) and the Lemma 5.5, we change variable setting $\mu_iy=x-\xi$, then

$$\begin{aligned}
&\int_{\mathcal{A}_i}\Big|\Big((-1)^iU_{\mu_i,\xi_i}+(-1)^j\sum_{j\neq i}^kU_{\mu_j,\xi_j}\Big)^{p-1}\Big(\ln\ln\Big(e+(-1)^iU_{\mu_i,\xi_i}+(-1)^j\sum_{j\neq i}^kU_{\mu_j,\xi_j}\Big)+1\Big)\\
&\times\sum_{i=1}^k(-1)^i(U_{\mu_i,\xi_i}-PU_{\mu_i,\xi_i})\psi^0_{\mu_j,\xi_j}\Big|dx\\
=\;&\alpha_n\sum_{i=1}^k\mu_i^{\frac{n-2}{2}}\int_{\mathcal{A}_i}\Big|(-1)^iU^{p-1}_{\mu_i,\xi_i}\Big(\ln\ln(e+(-1)^iU_{\mu_i,\xi_i})+1\Big)(H(x,\xi_i)+O(\mu_i^2))\psi^0_{\mu_j,\xi_j}\Big|dx\\
&+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\\
=\;&\alpha_n\sum_{i=1}^k\mu_i^{\frac{n-2}{2}}\int_{\frac{\mathcal{A}_i}{\mu_i}}\Big|\frac{\mu_i^{-2}}{(1+|y-\sigma_i|^2)^2}\Big(\ln\ln\Big(e+\frac{\mu_i^{-\frac{n-2}{2}}}{(1+|y-\sigma_i|^2)^{\frac{n-2}{2}}}\Big)+1\Big)\\
\times\;&\Big(H(\xi_i+\mu_iy-\mu_i\sigma_i,\xi_i)+O(\mu_i^2)\Big)\frac{|\mu_iy-\mu_j\sigma_j|^2-\mu_j^2}{\Big(\mu_j^2+|\mu_iy-\mu_j\sigma_j|^2\Big)^{\frac{n}{2}}}\Big|dy+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\\
\le\;&\begin{cases}
\alpha_n\sum\limits_{i=1}^k\mu_i^{\frac{n-2}{2}}\Big(H(\xi_i,\xi_i)+O(\mu_i^2)\Big)\mu_j^{-(n-2)}(\ln|\ln\mu_i|+1)\\
\quad\times\int_{\frac{\mathcal{A}_i}{\mu_i}}\frac{1}{(1+|y-\sigma_i|^2)^2}\frac{1}{|y|^{n-2}}dy+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\quad\text{if } j<i,\\
\alpha_n\sum\limits_{i=1}^k\mu_i^{-\frac{n-2}{2}}(\ln|\ln\mu_i|+1)\Big(H(\xi_i,\xi_i)+O(\mu_i^2)\Big)\\
\quad\times\int_{\frac{\mathcal{A}_i}{\mu_i}}\frac{1}{(1+|y-\sigma_i|^2)^2}\frac{1}{|y-\frac{\mu_j}{\mu_i}\sigma_j|^{n-2}}dy+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\quad\text{if } j>i,
\end{cases}
\end{aligned}$$

$$
\begin{aligned}
&= \begin{cases} O\Big(\big(\frac{\mu_i}{\mu_j}\big)^{\frac{n-2}{2}} \mu_j^{-\frac{n-2}{2}} \ln|\ln\mu_i|\Big) & \text{if } h=0,\\ O\Big(\mu_i^{-\frac{n-2}{2}} \ln|\ln\mu_i|\Big) & \text{if } h=1,\cdots,n,\end{cases}\\
&= O\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big).
\end{aligned}
$$

By the same argument, for $h=1,\cdots,n$, we have

$$
\begin{aligned}
&\int_{\mathcal{A}_i}\Big|\Big((-1)^i U_{\mu_i,\xi_i}+(-1)^j\sum_{j\neq i}^k U_{\mu_j,\xi_j}\Big)^{p-1}\Big(\ln\ln\Big(e+(-1)^i U_{\mu_i,\xi_i}+(-1)^j\sum_{j\neq i}^k U_{\mu_j,\xi_j}\Big)+1\Big)\\
&\quad\times\sum_{i=1}^k(-1)^i(U_{\mu_i,\xi_i}-PU_{\mu_i,\xi_i})\psi^h_{\mu_j,\xi_j}\Big|dx\\
=&O\Big(\sum_{i=1}^k\mu_i^{\frac{n-2}{2}}\ln|\ln\mu_i|\Big)=O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac12}\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big).
\end{aligned}
$$

Therefore, the second term in (5.33) becomes

$$
\begin{aligned}
&\int_\Omega\Big|\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\Big(\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big)\psi^h_{\mu_j,\xi_j}\Big|dx-\int_\Omega\Big|V^p\ln\ln(e+V)\psi^h_{\mu_j,\xi_j}\Big|dx\\
=&O\Big(\sum_{i=1}^k\mu_i^{\frac{n-2}{2}}\ln|\ln\mu_i|\Big)=O\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac12}\ln\Big|\ln\frac{\varepsilon}{|\ln\varepsilon|^2}\Big|\Big).
\end{aligned}
$$

For the first term in (5.33), if $j=i$, from Lemma 5.5, (2.9), let $x-\xi=\mu_i y$, then

$$
\begin{aligned}
&\int_\Omega\Big|\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\Big[\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\Big]\psi^h_{\mu_i,\xi_i}\Big|dx\\
=&\sum_{i=1}^k\int_{\mathcal{A}_i}\Big|\Big((-1)^iU_{\mu_i,\xi_i}+(-1)^j\sum_{j\neq i}^kU_{\mu_j,\xi_j}\Big)^p\\
&\quad\times\Big[\ln\ln\Big(e+(-1)^iU_{\mu_i,\xi_i}+(-1)^j\sum_{j\neq i}^kU_{\mu_j,\xi_j}\Big)\Big]\psi^h_{\mu_i,\xi_i}\Big|dx+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\\
=&\mu_1^{n-\frac{n+2}{2}-\frac{n-2}{2}}\int_{\frac{\mathcal{A}_i}{\mu_i}}\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}\Big|\ln\ln\Big(e+(-1)^i\mu_i^{-\frac{n-2}{2}}\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n-2}{2}}}\\
&\quad+(-1)^j\sum_{j\neq i}^k\frac{\mu_j^{\frac{n-2}{2}}}{\Big((\frac{\mu_j}{\mu_i})^2+|y-\frac{\sigma_i}{\mu_i}|^2\Big)^{\frac{n+2}{2}}\mu_i^{n-2}}\Big)\psi^h(y)\Big|dy+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)\\
=&\sum_{i=1}^k\ln\Big|\ln\mu_i^{-\frac{n-2}{2}}\Big|\int_{\frac{\mathcal{A}_i}{\mu_i}}\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}\psi^h(y)dy+\sum_{i=1}^k\frac{1}{|\ln\mu_i|}\int_{\frac{\mathcal{A}_i}{\mu_i}}\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}\\
&\quad\times\Big[|\ln\mu_i|\ln\Big(1+\frac{\ln\Big[e^{1-\frac{n-2}{2}|\ln\mu_i|}+\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}\Big]}{\frac{n-2}{2}|\ln\mu_i|}\Big)\Big]\psi^h(y)dy+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big).
\end{aligned}
\tag{5.34}
$$

Moreover, let $\Lambda(y)=\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}|\ln\mu_i|\ln\Big(1+\frac{\ln\Big(e^{1-\frac{n-2}{2}|\ln\mu_i|}+\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}\Big)}{\frac{n-2}{2}|\ln\mu_i|}\Big)\psi^h(y)$ for $h=1,\cdots,n$. Since $\psi(y)$ is a odd function, we deduce $\int_{\mathbb{R}^n}\Lambda(y)dy=0$. Further, by Lemma 5.5, there holds

$$\begin{aligned}
&\int_{\mathbb{R}^n}\Lambda(y)dy-\int_{\frac{\mathcal{A}_i}{\mu_i}}\Lambda(y)dy=\int_{\mathbb{R}^n\setminus\frac{\mathcal{A}_i}{\mu_i}}\Lambda(y)dy\\
\le& C|\ln\mu_i|\ln\Big(1+\frac{\ln\Big(e^{1-\frac{n-2}{2}|\ln\mu_i|}+\alpha_n\Big)}{\frac{n-2}{2}|\mu_i|}\Big)\int_{\mathbb{R}^n\setminus\frac{\mathcal{A}_i}{\mu_i}}\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}|\psi^h(y)|dy\\
\le& C\Big(\frac{2}{n-2}\ln\alpha_n+o(1)\Big)\mu_i^{n+1}=O(\mu_i^{n+1})=o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big).
\end{aligned}$$

Hence, for μ_i small enough, we conclude that

$$\begin{aligned}
&\int_\Omega\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\psi^h_{\mu_j,\xi_j}dx\\
=&O\Big(\sum_{i=1}^k\mu_i^{n+1}\ln\Big|\ln\mu_i^{-\frac{n-2}{2}}\Big|\Big)=o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big),\quad\text{for } h=1,\cdots,n.
\end{aligned}\tag{5.35}$$

When $h=0$, since U_{μ_i,ξ_i} is the unique positive solution of problem (1.4) and also, $\psi^0_{\mu_i,\xi_i}$ solves (2.7), then

$$\int_{\mathbb{R}^n}U^p_{\mu_i,\xi_i}\psi^0_{\mu_i,\xi_i}dx=\int_{\mathbb{R}^n}\nabla U_{\mu_i,\xi_i}\nabla\psi^0_{\mu_i,\xi_i}dx=p\int_{\mathbb{R}^n}U^p_{\mu_i,\xi_i}\psi^0_{\mu_i,\xi_i}dx,$$

which follows that

$$\langle U_{\mu_i,\xi_i},\psi^0_{\mu_i,\xi_i}\rangle=\int_{\mathbb{R}^n}U^p_{\mu_i,\xi_i}\psi^0_{\mu_i,\xi_i}dx=0.$$

Thus, $\int_{\mathbb{R}^n}\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}\psi^0(y)dx=0$ holds, also, $\int_{\mathbb{R}^n\setminus\frac{\mathcal{A}_i}{\mu_i}}\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}\psi^0dy=0$. From (5.34) and Lemma 5.5, there holds

$$\begin{aligned}
&\int_\Omega\Big|\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\psi^0_{\mu_i,\xi_i}\Big|dx\\
=&\sum_{i=1}^k\frac{1}{|\ln\mu_i|}\frac{2}{n-2}\int_{\mathbb{R}^n}\Big|\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}\ln\Big(\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}}\Big)\psi^0(y)\Big|dy+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
&\int_\Omega\Big|\Big(\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)^p\ln\ln\Big(e+\sum_{i=1}^k(-1)^iU_{\mu_i,\xi_i}\Big)\psi^h_{\mu_i,\xi_i}\Big|dx\\
=&\begin{cases}\frac{2}{n-2}a_4\sum\limits_{i=1}^k\frac{\varepsilon}{|\ln\mu_i|}+O\Big(\frac{1}{|\ln\mu_1|}\Big) & \text{if } h=0,\\ o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } h=1,\cdots,n.\end{cases}\\
=&\begin{cases}\frac{2}{n-2}a_4\frac{\varepsilon}{|\ln\varepsilon|^2}+\frac{2}{n-2}a_4\sum\limits_{i=1}^k|\ln d_i|+o\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big) & \text{if } h=0,\\ o\Big(\Big(\frac{\varepsilon}{|\ln\varepsilon|^2}\Big)^{\frac{n-1}{n-2}}\Big) & \text{if } h=1,\cdots,n.\end{cases}
\end{aligned}$$

By the same argument, if $j \neq i$, one has

$$\int_\Omega \Big| \Big(\sum_{i=1}^k (-1)^i U_{\mu_i,\xi_i} \Big)^p \ln\ln \Big(e + \sum_{i=1}^k (-1)^i U_{\mu_i,\xi_i} \Big) \psi^h_{\mu_i,\xi_i} \Big| dx = o\Big(\Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big)^{\frac{n-1}{n-2}} \Big).$$

Consequently,

$$\begin{aligned} J_3 &= \int_\Omega \Big(f_0(V) - f_\varepsilon(V) \Big) \psi^h_{\mu_j,\xi_j} dx \\ &= \begin{cases} -\frac{2}{n-2} a_4 \frac{\varepsilon}{|\ln \varepsilon|^2} - \frac{2}{n-2} a_4 \sum\limits_{i=1}^k |\ln d_i| + o\Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big) & \text{if } h = 0, \\ O\Big(\Big(\frac{\varepsilon}{|\ln \varepsilon|^2} \Big)^{\frac{1}{2}} \ln \Big| \ln \frac{\varepsilon}{|\ln \varepsilon|^2} \Big| \Big) & \text{if } h = 1, \cdots, n. \end{cases} \end{aligned}$$

Combining J_1-J_3, the proof of this lemma is completed.

Finally, let us state that a_4 is a positive constant. From (2.8), polar coordinates, integrating by parts, changing variables ($s = r^2$, $dr = \frac{1}{2} s^{-\frac{1}{2}} ds$), and a fact that $|\partial B_1(0)| = \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})}$, we obtain

$$\begin{aligned} a_4 =& - \int_{\mathbb{R}^n} \frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}} \ln \Big(\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}} \Big) \psi^0(y) dy \\ =& - \int_{B_1(0)} \frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}} \ln \Big(\frac{1}{(1+|y-\sigma_i|^2)^{\frac{n+2}{2}}} \Big) \psi^0(y) dy \\ =& - \frac{(n-2)\alpha_n^{2^*}}{2} |\partial B_1(0)| \int_0^\infty \frac{r^{n-1}}{(1+r^2)^{\frac{n+2}{2}}} \ln \Big(\frac{1}{(1+r^2)^{\frac{n+2}{2}}} \Big) \frac{r^2-1}{(1+r^2)^{\frac{n}{2}}} dr \\ =& \frac{(n-2)^2 \alpha_n^{2^*}}{4} \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \int_0^\infty \frac{r^{n-1}(r^2-1)}{(1+r^2)^{n+1}} \ln(1+r^2) dr \\ =& \frac{(n-2)^2 \alpha_n^{2^*}}{2} \frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \int_0^\infty \frac{1}{n} \Big(\frac{r}{1+r^2} \Big)^n \frac{2r}{1+r^2} dr \\ =& \frac{(n-2)^2 \alpha_n^{2^*}}{2n} \frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \int_0^\infty \frac{s^{\frac{n}{2}}}{(1+s)^{n+1}} ds = \frac{(n-2)^2 \alpha_n^{2^*}}{2n} \frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} B(\frac{n}{2}+1, \frac{n}{2}) ds \\ =& \frac{(n-2)^2 \alpha_n^{2^*}}{2n} \frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} \frac{\Gamma(\frac{n}{2}+1)\Gamma(\frac{n}{2})}{\Gamma(n+1)} = \frac{\Gamma(\frac{n}{2})\pi^{\frac{n}{2}}}{4\Gamma(n+1)} n^{\frac{n}{2}} (n-2)^{\frac{n+4}{2}}. \end{aligned}$$

□

Acknowledgments. The authors were supported by National Natural Science Foundation of China 11971392.

References

[1] A. Bahri, J. Coron, On a nonlinear elliptic equation involving the critical Sobolev exponent: the effect of the topology of the domain, *Comm. Pure Appl. Math.*, **41**(3), (1988), 253-294.

[2] A. Bahri, Y. Li, O. Rey, On a variational problem with lack of compactness: the topological effect of the critical points as infinity, *Calc. Var. Partial Differ. Equ.*, **3**(1), (1995), 67-93.

[3] T. Bartsch, T. D'Aprile, A. Pistoia, Multi-bubble nodal solutions for slightly subcritical elliptic problems in domains with symmetries, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **30**(6), (2013), 1027-1047.

[4] T. Bartsch, A. Micheletti, A. Pistoia, On the existence and the profile of nodal solutions of elliptic equations involving critical growth, *Calc. Var. Partial Differ. Equ.*, **26**, (2006), 265-282.

[5] T. Bartsch, Z. Wang, On the existence of sign changing solutions for semilinear Dirichlet problems, *Topol. Methods Nonlinear Anal.*, **7**, (1996), 115-131.
[6] T. Bartsch, Critical point theory on partially ordered Hilbert spaces, *J. Funct. Anal.*, **186**, (2001), 117-152.
[7] T. Bartsch, T. Weth, A note on additional properties of sign-changing solutions to superlinear elliptic equations, *Topol. Methods Nonlinear Anal.*, **22**, (2003), 1-14.
[8] M. Ben Ayed, K. EI Mehdi, F. Pacella, Classification of low energy sign-changing solutions of an almost critical problem, *J. Funct. Anal.*, **50**, (2007), 347-373.
[9] M. Ben Ayed, H. Fourti, Multispike solutions for a slightly subcritical elliptic problem with non-power nonlinearity, *Discrete Contin. Dyn. Syst.*, **43**(2), (2023), 661-687.
[10] E. Bianchi, H. Egnell, A note on the Sobolev inequality, *J. Funct. Anal.*, **100**, (1991), 18-24.
[11] A. Castro, J. Cossio, J. Neuberger, A sign-changing solution for a superlinear Dirichlet problem, *Rocky Mountain J. Math.*, **27**, (1997), 1041-1053.
[12] A. Castro, R. Pardo, A priori bounds for positive solutions of subcritical elliptic equations, *Rev. Mat. Complut.*, **28**, (2015), 715-731.
[13] A. Castro, R. Pardo, A priori estimates for positive solutions to subcritical elliptic problems in a class of non-convex regions, *Discrete Contin. Dyn. Syst. Ser. B*, **22**(3), (2017), 783-790.
[14] W. Chen, J. Dávila, I. Guerra, Bubble tower solutions for a supercritical elliptic problem in $\mathbb{R}^N$, *Ann. Sc. Norm. Super. Pisa Cl. Sci.*, **15**(5), (2016), 85-116.
[15] W. Chen, S. Deng, Sign-changing bubble tower solutions for a supercritical elliptic problem with the Hénon term, *Nonlinearity*, **30**(12), (2017), 4344-4368.
[16] W. Chen, X. Huang, Sign-changing bubble tower solutions for a Paneitz-type problem, arXiv preprint arXiv:2202.06006, 2022.
[17] W. Chen, W. Long, J. Yang, Sign-changing bubble tower solutions for a critical fractional problem, *Nonlinear Anal.*, **223**, (2022), 113054.
[18] M. Clapp, R. Pardo, A. Pistoia, A. Saldaña, A solution to a slightly subcritical elliptic problem with non-power nonlinearity, *J. Differential Equations*, **275**, (2021), 418-446.
[19] M. Clapp, T. Weth, Minimal nodal solutions of the pure critical exponent problem on a symmetric domain, *Calc. Var. Partial Differ. Equ.*, **21**(1), (2004), 1-14.
[20] A. Contreras, M. del Pino, Nodal bubble-tower solutions to radial elliptic problems near criticality, *Discrete Contin. Dyn. Syst.*, **16**(3), (2006), 525-539.
[21] G. Cora, A. Iacopetti, Sign-changing bubble-tower solutions to fractional semilinear elliptic problems, *Discrete Contin. Dyn. Syst.*, **39**, (2019)(10), 6149-6173.
[22] M. Cuesta, R. Pardo, Positive solutions for slightly subcritical elliptic problems via Orlicz spaces, *Milan J. Math.*, **90**(1), (2022), 229-255.
[23] Y. Dammak, R. Ghoudi, Sign-changing tower of bubbles to an elliptic subcritical equation, *Commun. Contemp. Math.*, **21**(7), (2019), 1850052, 38 pp.
[24] M. del Pino, J. Dolbeault, M. Musso, "Bubble-tower" radial solutions in the slightly supercritical Brezis-Nirenberg problem, *J. Differential Equations*, **193**(2), (2003), 280-306.
[25] M. del Pino, P. Felmer, M. Musso, Two-bubble solutions in the super-critical Bahri-Coron's problem, *Calc. Var. Partial Differ. Equ.*, **16**(2), (2003), 113-145.
[26] Y. Deng, S. Peng, X. Zhang, Y. Zhou, A class of supercritical Sobolev type inequalities with logarithm and related elliptic equations, *J. Differential Equations*, **341**, (2022), 150-188.
[27] W. Ding, On a conformally invariant elliptic equation on $\mathbb{R}^N$, *Comm. Math. Phys.*, **107**, (1986), 331-335.
[28] M. Flucher, J. Wei, Semilinear Dirichlet problem with nearly critical exponent, asymptotic location of hot spots, *Manuscripta Math.*, **94**, (1997), 337-346.
[29] Y. Ge, M. Musso, A. Pistoia, Sign changing tower of bubbles for an elliptic problem at the critical exponent in pierced non-symmetric domains, *Comm. Partial Differential Equations*, **35**(8), (2010), 1419-1457.
[30] M. Grossi, F. Takahashi, Nonexistence of multi-bubble solutions to some elliptic equations on convex domains, *J. Funct. Anal.*, **259**, (2010), 904-917.
[31] Z. Han, Asymptotic approach to singular solutions for nonlinear elliptic equations involving critical Sobolev exponent, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **8**, (1991), 159-174.
[32] E. Hebey, M. Vaugon, Existence and multiplicity of nodal solutions for nonlinear elliptic equations with critical Sobolev growth, *J. Funct. Anal.*, **119**, (1994), 298-318.
[33] J. Kazdan, F. Warner, Remarks on some quasilinear elliptic equations, *Comm. Pure Appl. Math.*, **28**, (1975), 567-597.
[34] Y. Li, Prescribing scalar curvature on $\mathcal{S}^n$ and related problems. I, *J. Differential Equations*, **120**(2), (1995), 319-410.

[35] Z. Liu, Z. Liu, W. Xu, Non-power type perturbation for the critical Hénon problem, *J. Math. Phys.*, **64**(2), (2023), Paper No. 021509, 19 pp.
[36] N. Mavinga, R. Pardo, A priori bounds and existence of positive solutions for semilinear elliptic systems, *J. Math. Anal. Appl.*, **449**(2), (2017), 1172-1188.
[37] M. Musso, A. Pistoia, Multispike solutions for a nonlinear elliptic problem involving critical Sobolev exponent, *Indiana Univ. Math. J.*, **5**, (2002), 541-579.
[38] M. Musso, A. Pistoia, Tower of bubbles for almost critical problems in general domains, *J. Math. Pures Appl.*, **93**, (2010), 1-40.
[39] R. Pardo, On the existence of a priori bounds for positive solutions of elliptic problems, II, *Rev. Integr. Temas Mat.*, **37**(1), (2019), 113-148.
[40] A. Pistoia, T. Weth, Sign-changing bubble tower solutions in a slightly subcritical semilinear Dirichlet problem, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **24**(2), (2007), 325-340.
[41] S. Pohozaev, On the eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$, *Dokl. Akad. Nauk*, **165**, (1965), 36-39, (Russian).
[42] O. Rey, The role of the Green's function in a non-linear elliptic equation involving the critical Sobolev exponent, *J. Funct. Anal.*, **89**, (1990), 1-52.
[43] O. Rey, Blow-up points of solutions to elliptic equations with limiting nonlinearity, *Differential Integral Equations*, **4**, (1991), 1155-1167.

(S. Deng) School of Mathematics and Statistics
Southwest University
Chongqing 400715, People's Republic of China.
Email address: shbdeng@swu.edu.cn

(F. Yu) School of Mathematics and Statistics
Southwest University
Chongqing 400715, People's Republic of China.
Email address: fangyumath@163.com