arXiv:2501.07281v2 [math.CA] 16 Jan 2025

Asymptotics of the Humbert functions Ψ_1 and Ψ_2

Peng-Cheng Hang [1], Malte Henkel [2,3], and Min-Jie Luo [*1]

[1] Department of Mathematics, School of Mathematics and Statistics, Donghua University, Shanghai 201620, People's Republic of China

[2] Laboratoire de Physique et Chimie Théoriques (CNRS UMR 7019), Université de Lorraine Nancy, B.P. 70239, F – 54506 Vandœuvre lès Nancy Cedex, France

[3] Centro de Física Teórica e Computacional, Universidade de Lisboa, Campo Grande, P–1749-016 Lisboa, Portugal

Abstract

A compilation of new results on the asymptotic behaviour of the Humbert functions Ψ_1 and Ψ_2, and also on the Appell function F_2, is presented. As a by-product, we confirm a conjectured limit which appeared recently in the study of the $1D$ Glauber-Ising model. We also propose two elementary asymptotic methods and confirm through some illustrative examples that both methods have great potential and can be applied to a large class of problems of asymptotic analysis. Finally, some directions of future research are pointed out in order to suggest ideas for further study.

2020 *Mathematics Subject Classification*: 33C65, 33C70, 41A60, 82C23.

Keywords and phases: asymptotic expansions, Humbert functions, Glauber-Ising model.

1. Introduction

When studying the limiting cases of the famous Appell hypergeometric functions F_1, F_2, F_3 and F_4, P. Humbert [25] introduced seven confluent hypergeometric functions of two variables which are denoted by Φ_1, Φ_2, Φ_3, Ψ_1, Ψ_2, Ξ_1 and Ξ_2. Humbert's pioneering contributions have inspired many important works. Here we want to mention Erdélyi's work on systems of linear partial differential equations satisfied by Humbert's functions (see [16] and [17]). Actually, the study of systems satisfied by Humbert's functions remains an active area of research. For more recent work, the interested readers may refer to [32] and [38].

The Humbert functions have a wide range of applications in various branches of mathematics and physics. Tuan *et al.* [41] introduced a useful integral transform with the function Φ_1 in the kernel and showed that it is an isomorphism in the space of entire functions of exponential type. The function Φ_2 occurs as an approximate solution to the Schrödinger equation for the three-body Coulomb problem [19]. The function Φ_3 is usually related to the generalized Hille-Hardy formula [31]. It also appears in Kumar's work on generalizing one of Ramanujan's transformation formula [28]. The functions Φ_3 and Ψ_2 have been found to be useful in the evaluation of the Voigt functions [40]. A multivariate generalization of Ψ_2 has been required in the study of non-central matrix-variate Dirichlet distributions [36]. Belafhal and Nebdi [5] succeeded in generating a novel donut beam which they called the Humbert beam because the field distribution of such a beam is expressed in terms of the Humbert function Ψ_1. Later, Chib *et al.* [14] introduced the donut Humbert beam of type-II since its field distribution is expressed in terms of the Humbert function Ψ_2.

Very recently, the second author found that the Humbert function Ψ_1 appears naturally in the two-time correlator of the $1D$ Glauber-Ising model at temperature $T = 0$ and as a by-product, he conjectured the following limit

*Corresponding author, Email: mathwinnie@live.com; mathwinnie@dhu.edu.cn

[24]

$$\lim_{z\to 0^+} \mathrm{e}^{-\xi^2/(2z)} z^{1/2} \Psi_1\left[1, \frac{1}{2}; \frac{3}{2}, \frac{1}{2}; -z, \frac{\xi^2}{2z}\right] = \frac{\pi}{2} \operatorname{erf}\left(\frac{\xi}{\sqrt{2}}\right), \tag{1.1}$$

where $\xi > 0$ is kept fixed, erf denotes the usual error function [34] and the Humbert function Ψ_1 is defined by [39, p. 26, Eq. (21)]

$$\Psi_1[a, b; c, c'; x, y] := \sum_{m,n=0}^{\infty} \frac{(a)_{m+n}(b)_m}{(c)_m (c')_n} \frac{x^m}{m!} \frac{y^n}{n!} \quad (|x| < 1,\ |y| < \infty) \tag{1.2}$$

with $a, b \in \mathbb{C}$ and $c, c' \notin \mathbb{Z}_{\leqslant 0}$. The conjectured limit (1.1) is of particular interest because in general, the behaviour of $\Psi_1\left[x, \frac{y}{x}\right]$ as $x \to 0$ is singular and is quite different from the usual behaviours of

$$\Psi_1[tx, y],\ \Psi_1[x, ty],\ \Psi_1[tx, ty], \quad t \to \infty,$$

which have been studied in detail in [21, 22, 23]. Therefore, as a motivation for this paper, we first verify the correctness of the limit (1.1) by establishing some asymptotic formulas for $\Psi_1[x, y]$ under the condition that

$$x \text{ or } y \text{ is small, and } 0 < \gamma_1 \leqslant |xy| \leqslant \gamma_2 < \infty. \tag{1.3}$$

Next, we extend our study to the Humbert function Ψ_2, since Ψ_2 is a confluent form of Ψ_1. More precisely, we have [39, p. 26, Eq. (26)]

$$\lim_{\varepsilon\to 0} \Psi_1[a, 1/\varepsilon; c, c'; \varepsilon x, y] = \Psi_2[a; c, c'; x, y].$$

The Humbert function Ψ_2 is defined by [39, p. 26, Eq. (22)]

$$\Psi_2[a; c, c'; x, y] := \sum_{m,n=0}^{\infty} \frac{(a)_{m+n}}{(c)_m (c')_n} \frac{x^m}{m!} \frac{y^n}{n!} \quad (|x| < \infty,\ |y| < \infty), \tag{1.4}$$

where $a \in \mathbb{C}$ and $c, c' \notin \mathbb{Z}_{\leqslant 0}$. To the best of our knowledge, there has been no investigation on the asymptotics of Ψ_2. Joshi and Arya [27, p. 499] proposed a bilateral inequality for Ψ_2, but as they themselves pointed out, the inequality is not simple and sharp. So we shall first study the asymptotics of $\Psi_2[x, y]$ under the condition (1.3) and then consider the asymptotics of Ψ_2 for one or two large arguments for completeness.

Finally, we turn to the Appell function F_2, which is defined by [39, p. 23, Eq. (3)]

$$F_2[a, b, b'; c, c'; x, y] := \sum_{m,n=0}^{\infty} \frac{(a)_{m+n} (b)_m (b')_n}{(c)_m (c')_n} \frac{x^m}{m!} \frac{y^n}{n!} \quad (|x| + |y| < 1), \tag{1.5}$$

where $a, b, b' \in \mathbb{C}$ and $c, c' \notin \mathbb{Z}_{\leqslant 0}$. We note that both Ψ_1 and Ψ_2 are confluent forms of F_2:

$$\lim_{\varepsilon\to 0} F_2[a, b, 1/\varepsilon; c, c'; x, \varepsilon y] = \Psi_1[a, b; c, c'; x, y],$$
$$\lim_{\varepsilon\to 0} F_2[a, 1/\varepsilon, 1/\varepsilon; c, c'; \varepsilon x, \varepsilon y] = \Psi_2[a; c, c'; x, y].$$

The method that we apply to Ψ_1 can be used to derive the asymptotic formula of F_2 under the condition (1.3).

Table 1 lists the results we prove in this paper and the corresponding methods we use.

Here, we would also like to briefly mention two specific examples that also demonstrate the importance of the particular type of asymptotic behaviours (when one variable tends to infinity while the other remains small) studied in this paper. When studying approximate solution of a system of three charged particles, Gasaneo *et al.* [18] considered similar asymptotic behaviours of other hypergeometric functions. Under certain restrictions, Carvalho e Silva and Srivastava [15] obtained asymptotic formulas of functions defined by general double series. However, the theorems given by Carvalho e Silva and Srivastava *cannot* be used to derive the results obtained in our paper.

Notation. By $f(n, z) = \mathcal{O}(a_n g(z))\ (z \in \Omega)$, or briefly $f(n, z) \ll a_n g(z)$, we mean that there exists a constant $K > 0$ independent of the summation index n and the variable z such that

$$|f(n, z)| \leqslant K |a_n| |g(z)|, \quad n \in \mathbb{Z}_{\geqslant 0},\ z \in \Omega.$$

In general, we use $f(z) \ll g(z)$ to mean that $f(z) = \mathcal{O}(g(z))$. Moreover, the generalized hypergeometric function ${}_pF_q$ is defined by (see, for example, [4] and [34])

$${}_pF_q\left[\begin{matrix} a_1, \cdots, a_p \\ b_1, \cdots, b_q \end{matrix}; z\right] \equiv {}_pF_q[a_1, \cdots, a_p; b_1, \cdots, b_q; z] := \sum_{n=0}^{\infty} \frac{(a_1)_n \cdots (a_p)_n}{(b_1)_n \cdots (b_q)_n} \frac{z^n}{n!}, \tag{1.6}$$

where $a_1, \cdots, a_p \in \mathbb{C}$ and $b_1, \cdots, b_q \in \mathbb{C} \setminus \mathbb{Z}_{\leqslant 0}$. Empty products and sums are taken as 1 and 0, respectively.

Function	Limiting Case	Method	Results
$\Psi_1\left[x,\frac{y}{x}\right]$	$x\to 0$	Olver's Laplace method	Theorems 3.2 and 3.4
$\Psi_1\left[\frac{x}{y},y\right]$	$y\to 0$	Series manipulation technique	Theorem 3.6
$\Psi_2[x,y]$	$\Re(y)\to-\infty$	Mellin-Barnes integral	Theorem 4.1
$\Psi_2[x,y]$	$\Re(y)\to+\infty$, or $x,y\to\infty$	Olver's Laplace method	Theorem 4.2
$\Psi_2\left[x,\frac{\beta}{x}\right]$	$x\to\infty$	Mellin-Barnes integral and Series manipulation technique	Theorem 4.6
$F_2\left[\frac{x}{y},y\right]$	$y\to 0$	Series manipulation technique	Corollary 5.2

Table 1. Results proved and methods used.

2. Preliminaries

We list some asymptotic formulas for the Kummer function ${}_1F_1$ and the modified Bessel function I_ν, which are useful in what follows. In this section, $\delta>0$ is taken to be small.

For $z\to\infty$ in $-\pi\leqslant\arg(-z)\leqslant\pi$, we have [29, Eq. (5.8)]

$$ {}_1F_1\left[\begin{matrix}a\\c\end{matrix};z\right]\sim\frac{\Gamma(c)}{\Gamma(c-a)}\sum_{n=0}^{\infty}\frac{(a)_n(1+a-c)_n}{n!}(-z)^{-a-n}+\frac{\Gamma(c)}{\Gamma(a)}\mathrm{e}^z\sum_{n=0}^{\infty}\frac{(1-a)_n(c-a)_n}{n!}z^{a-c-n}. \tag{2.1}$$

As $z\to\infty$ in $-\frac{\pi}{2}+\delta\leqslant\arg(z)\leqslant\frac{3\pi}{2}-\delta$, we have [34, Eq. (10.40.5)]

$$ I_\nu(z)=\frac{\mathrm{e}^z}{\sqrt{2\pi z}}\left(1+\mathcal{O}\left(z^{-1}\right)\right)+\mathrm{i}\mathrm{e}^{\pi\mathrm{i}\nu}\frac{\mathrm{e}^{-z}}{\sqrt{2\pi z}}\left(1+\mathcal{O}\left(z^{-1}\right)\right), \tag{2.2}$$

where

$$ I_\nu(z)=\left(\frac{1}{2}z\right)^\nu\sum_{k=0}^{\infty}\frac{(\frac{1}{4}z^2)^k}{k!\Gamma(\nu+k+1)}. \tag{2.3}$$

As $a\to\infty$ in $\left|\arg(a)\right|\leqslant\pi-\delta$, we have [34, Eq. (13.8.12)]

$$ {}_1F_1\left[\begin{matrix}a\\b\end{matrix};z\right]=\Gamma(b)\,\mathrm{e}^{\frac{1}{2}z}\left(\frac{z}{a}\right)^{\frac{1}{2}(1-b)}\frac{\Gamma(a+1-b)}{\Gamma(a)}\left\{I_{b-1}\left(2\sqrt{az}\right)\left(1+\mathcal{O}\left(a^{-1}\right)\right)-\sqrt{\frac{z}{a}}\,I_b\left(2\sqrt{az}\right)\left(1+\mathcal{O}\left(a^{-1}\right)\right)\right\}. \tag{2.4}$$

Choose a and z such that $\left|\arg(a)\right|\leqslant\pi-\delta$ and $0\leqslant\arg(z)\leqslant 2\pi$. Then

$$ -\frac{\pi}{2}+\frac{\delta}{2}\leqslant\arg(\sqrt{az})\leqslant\frac{3\pi}{2}-\frac{\delta}{2}. $$

It follows from (2.2) that as $a\to\infty$ with $z\neq 0$ fixed,

$$ I_\nu(2\sqrt{az})=\frac{1}{2\sqrt{\pi}}(az)^{-\frac{1}{4}}\left(\mathrm{e}^{2\sqrt{az}}\left(1+o(1)\right)+\mathrm{i}\mathrm{e}^{\pi\mathrm{i}\nu}\mathrm{e}^{-2\sqrt{az}}\left(1+o(1)\right)\right). $$

Using (2.4), we obtain that for large a in $\left|\arg(a)\right|\leqslant\pi-\delta$ and fixed $z\neq 0$ in $0\leqslant\arg(z)\leqslant 2\pi$,

$$ {}_1F_1\left[\begin{matrix}a\\b\end{matrix};z\right]=\Gamma(b)\frac{\Gamma(a+1-b)}{2\sqrt{\pi}\,\Gamma(a)}z^{\frac{1}{4}-\frac{1}{2}b}\mathrm{e}^{\frac{1}{2}z}a^{\frac{1}{2}b-\frac{3}{4}}\left(\mathrm{e}^{2\sqrt{az}}\left(1+o(1)\right)-\mathrm{i}\mathrm{e}^{\pi\mathrm{i}b}\mathrm{e}^{-2\sqrt{az}}\left(1+o(1)\right)\right). \tag{2.5}$$

Finally, we shall also need the Euler-type integral [34, Eq. (13.4.1)]

$$ {}_1F_1\left[\begin{matrix}a\\b\end{matrix};z\right]=\frac{\Gamma(b)}{\Gamma(a)\Gamma(b-a)}\int_0^1 t^{a-1}(1-t)^{b-a-1}\mathrm{e}^{zt}\mathrm{d}t,\quad \Re(b)>\Re(a)>0. \tag{2.6}$$

3. Humbert function Ψ_1

First of all, we list a couple of identities which are useful in what follows:

$$\Psi_1[a,b;c,c';x,y] = \sum_{n=0}^{\infty} \frac{(a)_k\,(b)_k}{(c)_k}\,{}_1F_1\left[\begin{matrix}a+k\\ c'\end{matrix};y\right]\frac{x^k}{k!} \tag{3.1a}$$

$$= \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)}\int_0^1 t^{b-1}(1-t)^{c-b-1}(1-xt)^{-a}\,{}_1F_1\left[\begin{matrix}a\\ c'\end{matrix};\frac{y}{1-xt}\right]\mathrm{d}t \tag{3.1b}$$

$$= \frac{1}{\Gamma(a)}\int_0^\infty u^{a-1}\mathrm{e}^{-u}\,{}_1F_1\left[\begin{matrix}b\\ c\end{matrix};xu\right]{}_0F_1\left[\begin{matrix}-\\ c'\end{matrix};yu\right]\mathrm{d}u. \tag{3.1c}$$

With the exception of (3.1a), they arise in several places in the literature (see [8]). For instance, (3.1b) is useful for numerical computations (see [21, Appendix A]), and (3.1c) is from [42]. The convergence conditions for these identities are given below, respectively,

$$\begin{aligned}
&(3.1a): \quad c,c'\notin\mathbb{Z}_{\leqslant 0},\ |x|<1, |y|<\infty;\\
&(3.1b): \quad \Re(c)>\Re(b)>0,\ c'\notin\mathbb{Z}_{\leqslant 0},\ x\notin[1,+\infty),\ y\in\mathbb{C};\\
&(3.1c): \quad \Re(a)>0,\ c,c'\notin\mathbb{Z}_{\leqslant 0},\ \Re(x)<1,\ y\in\mathbb{C}.
\end{aligned}$$

Moreover, Ψ_1 has an extension to the region (see [23, Section 2.2])

$$\mathbb{D}_{\Psi_1} := \left\{(x,y)\in\mathbb{C}^2\colon\ x\neq 1,\ |\arg(1-x)|<\pi,\ |y|<\infty\right\}.$$

Before giving the first asymptotic result of Ψ_1 under the condition (1.3) (see Theorem 3.2), we derive a useful integral representation.

Proposition 3.1. *If $\Re(a)>0$, $b\in\mathbb{C}$, $c,c'\in\mathbb{C}\setminus\mathbb{Z}_{\leqslant 0}$, $\Re(x)>0$ and $\Re\left(\frac{y}{x}\right)>0$, then*

$$\Psi_1\left[a,b;c,c';-x,\frac{y^2}{x}\right] = \frac{2\Gamma(c')}{\Gamma(a)}x^{-a}\left(\frac{y}{x}\right)^{1-c'}\int_0^\infty w^{2a-c'}\mathrm{e}^{-\frac{w^2}{x}}\,{}_1F_1\left[\begin{matrix}b\\ c\end{matrix};-w^2\right]I_{c'-1}\left(2\frac{y}{x}w\right)\mathrm{d}w. \tag{3.2}$$

Proof. Assume for the moment that $a>0$, $x>0$ and $y>0$. In view of the relation [35, Eq. (7.13.1.1)]

$${}_0F_1\left[\begin{matrix}-\\ b\end{matrix};z\right] = \Gamma(b)\,z^{\frac{1-b}{2}}I_{b-1}\left(2\sqrt{z}\right) \tag{3.3}$$

and the integral representation (3.1c), we obtain

$$\begin{aligned}
\Psi_1\left[a,b;c,c';-x,\frac{y^2}{x}\right] &= \frac{1}{\Gamma(a)}\int_0^\infty u^{a-1}\mathrm{e}^{-u}\,{}_1F_1\left[\begin{matrix}b\\ c\end{matrix};-xu\right]{}_0F_1\left[\begin{matrix}-\\ c'\end{matrix};\frac{y^2}{x}u\right]\mathrm{d}u\\
&= \frac{\Gamma(c')}{\Gamma(a)}\int_0^\infty u^{a-1}\mathrm{e}^{-u}\,{}_1F_1\left[\begin{matrix}b\\ c\end{matrix};-xu\right]\left(\frac{y^2}{x}u\right)^{\frac{1-c'}{2}}I_{c'-1}\left(2\sqrt{\frac{y^2}{x}u}\right)\mathrm{d}u\\
&= \frac{\Gamma(c')}{x\Gamma(a)}\int_0^\infty\left(\frac{v}{x}\right)^{a-1}\mathrm{e}^{-\frac{v}{x}}\,{}_1F_1\left[\begin{matrix}b\\ c\end{matrix};-v\right]\left(\frac{y^2}{x^2}v\right)^{\frac{1-c'}{2}}I_{c'-1}\left(2\frac{y}{x}\sqrt{v}\right)\mathrm{d}v\\
&= \frac{2\Gamma(c')}{\Gamma(a)}x^{-a}\left(\frac{y}{x}\right)^{1-c'}\int_0^\infty w^{2a-c'}\mathrm{e}^{-\frac{w^2}{x}}\,{}_1F_1\left[\begin{matrix}b\\ c\end{matrix};-w^2\right]I_{c'-1}\left(2\frac{y}{x}w\right)\mathrm{d}w.
\end{aligned}$$

The proof is then completed by using (2.2) and analytic continuation. □

Theorem 3.2. *Let $\Re(a)>0$, $b\in\mathbb{C}$ and $c,c'\in\mathbb{C}\setminus\mathbb{Z}_{\leqslant 0}$. Then for fixed $y>0$,*

$$\Psi_1\left[a,b;c,c';\pm x,\frac{y}{x}\right]\sim\frac{\Gamma(c')}{\Gamma(a)}\,{}_1F_1\left[\begin{matrix}b\\ c\end{matrix};\pm y\right]\left(\frac{y}{x}\right)^{a-c'}\mathrm{e}^{\frac{y}{x}} \tag{3.4}$$

as $x\to 0$ in $\left|\arg(x)\right|\leqslant\frac{\pi}{2}-\delta$, where $\delta\in\left(0,\frac{\pi}{2}\right]$ is fixed.

Proof. We only give the proof of the result for $\Psi_1\left[-x,\frac{y^2}{x}\right]$. To establish the result for $\Psi_1\left[+x,\frac{y^2}{x}\right]$, one may repeat the proof here by starting with

$$\Psi_1\left[a,b;c,c';+x,\frac{y^2}{x}\right]=\frac{2\Gamma(c')}{\Gamma(a)}x^{-a}\left(\frac{y}{x}\right)^{1-c'}\int_0^\infty w^{2a-c'}\mathrm{e}^{-\frac{w^2}{x}}\,{}_1F_1\left[\begin{matrix}b\\c\end{matrix};+w^2\right]I_{c'-1}\left(2\frac{y}{x}w\right)\mathrm{d}w.$$

Assume that $\Re(a)>0$, $\Re(x)>0$ and $y>0$. Then separate the integral in (3.2) into two parts, namely,

$$\Psi_1:=\Psi_1\left[a,b;c,c';-x,\frac{y^2}{x}\right]=\frac{2\Gamma(c')}{\Gamma(a)}x^{-a}\left(\frac{y}{x}\right)^{1-c'}(\mathfrak{I}_1+\mathfrak{I}_2),$$

where $\eta=\left|\frac{x}{y}\right|$ and

$$\mathfrak{I}_1=\int_0^\eta w^{2a-c'}\mathrm{e}^{-\frac{w^2}{x}}\,{}_1F_1\left[\begin{matrix}b\\c\end{matrix};-w^2\right]I_{c'-1}\left(2\frac{y}{x}w\right)\mathrm{d}w,$$
$$\mathfrak{I}_2=\int_\eta^\infty w^{2a-c'}\mathrm{e}^{-\frac{w^2}{x}}\,{}_1F_1\left[\begin{matrix}b\\c\end{matrix};-w^2\right]I_{c'-1}\left(2\frac{y}{x}w\right)\mathrm{d}w.$$

Note that $\eta=\mathscr{O}(x)$ as $x\to 0$. Thus, by (2.3), we have

$$\begin{aligned}|\mathfrak{I}_1|&\leqslant\int_0^\eta w^{\Re(2a-c')}\mathrm{e}^{-\frac{w^2}{\Re(x)}}\left|{}_1F_1\left[\begin{matrix}b\\c\end{matrix};-w^2\right]\right|\cdot\left|I_{c'-1}\left(2\frac{y}{x}w\right)\right|\mathrm{d}w\\&\ll\int_0^\eta w^{\Re(2a-c')}\left|\frac{y}{x}w\right|^{\Re(c')-1}\cdot\left|\sum_{k=0}^\infty\frac{(wy/x)^{2k}}{k!\Gamma(c'+k)}\right|\mathrm{d}w\\&\ll\eta^{1-\Re(c')}\int_0^\eta w^{\Re(2a)-1}\mathrm{d}w=\frac{1}{2\Re(a)}\eta^{\Re(2a-c')+1}\ll|x|^{\Re(2a-c')+1}.\end{aligned}$$

To estimate $\mathfrak{I}_2$, recall $\eta=\left|\frac{x}{y}\right|$ and use (2.2) to yield

$$I_{c'-1}\left(2\frac{y}{x}w\right)=\frac{\mathrm{e}^{2yw/x}}{\sqrt{2\pi\cdot 2yw/x}}\left(1+\mathscr{O}(\eta w^{-1})\right),\quad w\geqslant\eta.$$

Therefore,

$$\mathfrak{I}_2=\frac{1}{2\sqrt{\pi}}\left(\frac{x}{y}\right)^{\frac{1}{2}}\mathrm{e}^{\frac{y^2}{x}}\int_\eta^\infty w^{2a-c'-\frac{1}{2}}\mathrm{e}^{-\frac{1}{x}(w-y)^2}\,{}_1F_1\left[\begin{matrix}b\\c\end{matrix};-w^2\right]\left(1+\mathscr{O}(\eta w^{-1})\right)\mathrm{d}w.$$

Using Olver's Laplace method [33, Theorem I], we obtain

$$\mathfrak{I}_2\sim\frac{1}{2}xy^{2a-c'-1}\mathrm{e}^{\frac{y^2}{x}}\,{}_1F_1\left[\begin{matrix}b\\c\end{matrix};-y^2\right].$$

Hence (3.4) follows from the estimates on $\mathfrak{I}_1$ and $\mathfrak{I}_2$. □

Remark 3.3. *The limit* (1.1) *is a direct result of* (3.4). *Indeed,* (3.4) *implies that*

$$\lim_{z\to 0^+}\mathrm{e}^{-\xi^2/(2z)}z^{1/2}\Psi_1\left[1,\frac{1}{2};\frac{3}{2},\frac{1}{2};-z,\frac{\xi^2}{2z}\right]=\sqrt{\frac{\pi}{2}}\xi\cdot{}_1F_1\left[\begin{matrix}1/2\\3/2\end{matrix};-\frac{\xi^2}{2}\right],$$

which, in view of [34, Eq. (7.11.1) and (13.6.5)], *is equivalent to* (1.1).

We can further establish the complete leading-term behaviour of $\Psi_1\left[x,\frac{y}{x}\right]$ for small x, but we have to impose different conditions on the parameters.

Theorem 3.4. *Let* $a\in\mathbb{C}$, $c,c'\in\mathbb{C}\setminus\mathbb{Z}_{\leqslant 0}$ *and* $\Re(c)>\Re(b)>0$. *For any fixed* $y\in\mathbb{C}\setminus\{0\}$,

$$\Psi_1\left[a,b;c,c';x,\frac{y}{x}\right]\sim\frac{\Gamma(c')}{\Gamma(a)}\,{}_1F_1\left[\begin{matrix}b\\c\end{matrix};y\right]\left(\frac{y}{x}\right)^{a-c'}\mathrm{e}^{\frac{y}{x}}+\frac{\Gamma(c')}{\Gamma(c'-a)}\left(-\frac{y}{x}\right)^{-a}\tag{3.5}$$

as $x\to 0$ *such that* $-\pi\leqslant\arg\left(-\frac{y}{x}\right)\leqslant\pi$.

Proof. Fix $y \in \mathbb{C} \setminus \{0\}$ and write

$$\lambda = \lambda(u) \equiv \frac{y/x}{1+ux}.$$

Then as $x \to 0$, $\lambda = \frac{y}{x} - yu + \mathscr{O}(x)$ and $\lambda^{-1} = \mathscr{O}(x)$. Use of (3.1b) gives

$$\Psi_1\left[a,b;c,c';-x,\frac{y}{x}\right] = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)}\int_0^1 u^{b-1}(1-u)^{c-b-1}(1+ux)^{-a}\,g(\lambda)\mathrm{d}u, \tag{3.6}$$

where

$$g(\lambda) := {}_1F_1\left[\begin{matrix}a\\c'\end{matrix};\lambda\right] = {}_1F_1\left[\begin{matrix}a\\c'\end{matrix};\frac{y}{x} - yu + \mathscr{O}(x)\right].$$

Applying the expansion (2.1), we deduce that as $x \to 0$ in $-\pi \leqslant \arg\left(-\frac{y}{x}\right) \leqslant \pi$,

$$g(\lambda) = \frac{\Gamma(c')}{\Gamma(a)}\left(\frac{y}{x}\right)^{a-c'}\mathrm{e}^{\frac{y}{x}}\left(\mathrm{e}^{-yu}+\mathscr{O}(x)\right) + \frac{\Gamma(c')}{\Gamma(c'-a)}\left(-\frac{y}{x}\right)^{-a}(1+\mathscr{O}(x)).$$

Combing this with (3.6) and (2.6) completes the proof. □

Remark 3.5. *With the help of the integrals given by Brychkov and Saad* [9, Eq. (3.2) and (3.15)-(3.18)]*, one could try to find asymptotic formulas for the Appell function F_3 and the Humbert functions $\Phi_2, \Phi_3, \Xi_1, \Xi_2$ under the condition* (1.3)*. We plan to return to this elsewhere.*

Since $\Psi_1[x,y]$ is not symmetric in x and y, we are led to study the behaviour of $\Psi_1\left[\frac{x}{y},y\right]$ for small y.

Theorem 3.6. *Let $a,b \in \mathbb{C}$, $c,c' \in \mathbb{C}\setminus\mathbb{Z}_{\leqslant 0}$ and $a-b \notin \mathbb{Z}$. For fixed $x \in \mathbb{C}\setminus\{0\}$,*

$$\begin{aligned}\Psi_1\left[a,b;c,c';\frac{x}{y},y\right] &= \mathfrak{f}_c(b,a)\left(-\frac{x}{y}\right)^{-a}\sum_{m=0}^{N-1}a_1(m)y^m + \mathfrak{f}_c(a,b)\left(-\frac{x}{y}\right)^{-b}\sum_{m=0}^{N-1}a_2(m)y^m \\ &\quad + \mathscr{O}\left(|y|^{\Re(a)+N} + |y|^{\Re(b)+N}\right)\end{aligned} \tag{3.7}$$

as $y \to 0$ in $\left|\arg\left(-\frac{x}{y}\right)\right| < \pi$, where N is any positive integer,

$$\mathfrak{f}_\gamma(a,b) := \frac{\Gamma(\gamma)\Gamma(a-b)}{\Gamma(a)\Gamma(\gamma-b)} \tag{3.8a}$$

and

$$a_1(m) = \sum_{\substack{k,\ell\geqslant 0\\ k+\ell=m}} \frac{(-k)_\ell\,(a)_k\,(1-c+a)_k}{(c')_\ell\,(1-b+a)_k\,\ell!\,k!}x^{-k}, \tag{3.8b}$$

$$a_2(m) = \sum_{\substack{k,\ell\geqslant 0\\ k+\ell=m}} \frac{(a-b-k)_\ell\,(b)_k\,(1-c+b)_k}{(c')_\ell\,(1-a+b)_k\,\ell!\,k!}x^{-k}. \tag{3.8c}$$

Proof. Recall the series expansion [23, Theorem 3.8]: for $\left|\arg(-x)\right| < \pi$, $|x|>1$ and $|y|<\infty$,

$$\begin{aligned}\Psi_1[a,b;c,c';x,y] &= \mathfrak{f}_c(b,a)\,(-x)^{-a}\sum_{k=0}^{\infty}{}_1F_1\left[\begin{matrix}-k\\c'\end{matrix};y\right]\frac{(a)_k\,(1-c+a)_k}{(1-b+a)_k\,k!}\frac{1}{x^k} \\ &\quad + \mathfrak{f}_c(a,b)\,(-x)^{-b}\sum_{k=0}^{\infty}{}_1F_1\left[\begin{matrix}a-b-k\\c'\end{matrix};y\right]\frac{(b)_k\,(1-c+b)_k}{(1-a+b)_k\,k!}\frac{1}{x^k}.\end{aligned} \tag{3.9}$$

Denote the series on the right by $U_1(x,y)$ and $U_2(x,y)$, respectively. Then we have

$$U_1\left(\frac{x}{y},y\right) = \sum_{k=0}^{N-1}\sum_{\ell=0}^{\infty}\frac{(-k)_\ell}{(c')_\ell}\frac{y^\ell}{\ell!}\cdot\frac{(a)_k\,(1-c+a)_k}{(1-b+a)_k\,k!}\frac{y^k}{x^k} + \mathscr{O}\left(|y|^N\right)$$

$$= \sum_{m=0}^{N-1} a_1(m) y^m + \mathscr{O}\left(|y|^N\right), \tag{3.10}$$

and similarly

$$U_2\left(\frac{x}{y}, y\right) = \sum_{m=0}^{N-1} a_2(m) y^m + \mathscr{O}\left(|y|^N\right). \tag{3.11}$$

The expansion for $\Psi_1\left[\frac{x}{y}, y\right]$ therefore follows from (3.9)-(3.11). □

Remark 3.7. *As an application, notice that for $x \to \infty$ and $y > 0$ fixed, one has*

$$\Psi_1\left[1, \frac{1}{2}; \frac{3}{2}, \frac{1}{2}; -x, \frac{y}{x}\right] \sim \frac{\pi}{2} x^{-\frac{1}{2}} - x^{-1} + \mathscr{O}\left(x^{-\frac{3}{2}}\right).$$

This new asymptotic identity, especially the y-independence of the leading two terms, proves the large-time universality in the time-space correlator in the $1D$ Glauber-Ising model at temperature $T = 0$ [24].

4. Humbert function Ψ_2

Clearly, Ψ_2 satisfies the symmetry relation $\Psi_2[a; c, c'; x, y] = \Psi_2[a; c', c; y, x]$ and admits the series expansion

$$\Psi_2[a; c, c'; x, y] = \sum_{n=0}^{\infty} \frac{(a)_n}{(c')_n} {}_1F_1\left[\begin{matrix} a+n \\ c \end{matrix}; x\right] \frac{y^n}{n!}, \quad |x| < \infty, |y| < \infty. \tag{4.1}$$

Using the Kummer transformation [34, Eq. (13.2.39)], we get

$$\Psi_2[a; c, c'; x, y] = \mathrm{e}^x \sum_{n=0}^{\infty} \frac{(a)_n}{(c')_n} {}_1F_1\left[\begin{matrix} c-a-n \\ c \end{matrix}; -x\right] \frac{y^n}{n!}, \quad |x| < \infty, |y| < \infty. \tag{4.2}$$

Let us first study the asymptotic behaviour of $\Psi_2[x, y]$ as $y \to \infty$ in the left half-plane using the Mellin-Barnes integral representation of Ψ_2.

Theorem 4.1. (i) *Let $a \in \mathbb{C}$, $c, c' \in \mathbb{C} \setminus \mathbb{Z}_{\leqslant 0}$, $\delta \in \left(0, \frac{\pi}{2}\right]$ and*

$$\mathbb{V}_{\Psi_2} = \left\{(x, y) \in \mathbb{C}^2 : x \neq 0,\ y \neq 0,\ 0 \leqslant \arg(x) \leqslant 2\pi,\ \left|\arg(-y)\right| \leqslant \frac{\pi}{2} - \delta\right\}.$$

Then

$$\Psi_2[a; c, c'; x, y] = \frac{1}{2\pi \mathrm{i}} \frac{\Gamma(c')}{\Gamma(a)} \int_{\mathfrak{L}_{\mathrm{i}\sigma\infty}} {}_1F_1\left[\begin{matrix} a+s \\ c \end{matrix}; x\right] \frac{\Gamma(a+s)}{\Gamma(c'+s)} \Gamma(-s) \left(-y\right)^s \mathrm{d}s, \tag{4.3a}$$

where the path $\mathfrak{L}_{\mathrm{i}\sigma\infty}$, starting at $\sigma - \mathrm{i}\infty$ and ending at $\sigma + \mathrm{i}\infty$, is a vertical line intended if necessary to separate the poles of $\Gamma(a+s)$ from the poles of $\Gamma(-s)$.

(ii) *Let $a \in \mathbb{C}$ and $c, c', c' - a \in \mathbb{C} \setminus \mathbb{Z}_{\leqslant 0}$. When $(x, y) \in \mathbb{V}_{\Psi_2}$ and $|y| \to \infty$, we have*

$$\Psi_2[a; c, c'; x, y] \sim \frac{\Gamma(c')}{\Gamma(c'-a)} \sum_{n=0}^{\infty} {}_1F_1\left[\begin{matrix} -n \\ c \end{matrix}; x\right] \frac{(a)_n (1+a-c')_n}{n!} (-y)^{-a-n}. \tag{4.3b}$$

Proof. (i) The proof will be divided into two steps. Firstly, we prove the integral (4.3a) exists for $(x, y) \in \mathbb{V}_{\Psi_2}$. Secondly, we show by using Cauchy's residue theorem that the integral is equal to Ψ_2 when x and y are appropriately restricted. Then by analytic continuation, (4.3a) is valid throughout $\mathbb{V}_{\Psi_2}$.

Step 1. We can formally obtain (4.3a) from (4.1) with the help of Ramanujan's master theorem [1, p. 107]. To check the validity, we first denote the integrand by

$$\Psi(s) := {}_1F_1\left[\begin{matrix} a+s \\ c \end{matrix}; x\right] \frac{\Gamma(a+s)}{\Gamma(c'+s)} \Gamma(-s)(-y)^s. \tag{4.4}$$

Recall the estimate [23, Eq. (3.4)]: as $t \to \pm\infty$,

$$\left|\frac{\Gamma(a+\sigma+\mathrm{i}t)}{\Gamma(c'+\sigma+\mathrm{i}t)}\Gamma(-\sigma-\mathrm{i}t)(-y)^{\sigma+\mathrm{i}t}\right| = \mathscr{O}\left(|t|^{\Re(a-c')-\sigma-\frac{1}{2}}\,\mathrm{e}^{-\left(\frac{\pi}{2}\pm\arg(-y)\right)|t|}\right). \tag{4.5}$$

Using (2.5), we obtain that as $t \to \pm\infty$,

$$\left|{}_1F_1\left[\begin{matrix}a+\sigma+\mathrm{i}t\\ c\end{matrix};x\right]\right| = \mathscr{O}\left(|t|^{\frac{1}{4}-\frac{1}{2}\Re(c)}\,\mathrm{e}^{3\sqrt{|xt|}}\right). \tag{4.6}$$

Hence when $\left|\arg(-y)\right| \leqslant \frac{\pi}{2}-\delta$, the integrand $\Psi(s)$ decays exponentially as $|s|\to\infty$ along the path $\mathfrak{L}_{\mathrm{i}\sigma\infty}$. Therefore, the integral in (4.3a) is certainly convergent in $\mathbb{V}_{\Psi_2}$.

Step 2. Let us show that in a region $\widetilde{\mathbb{V}}_{\Psi_2} \subset \mathbb{V}_{\Psi_2}$, the integral in (4.3a) yields the series representation (4.1) of Ψ_2. Define

$$\widetilde{\mathbb{V}}_{\Psi_2} = \left\{(x,y)\in\mathbb{C}^2 : x\neq 0,\ 0<|y|<\frac{1}{\mathrm{e}},\ 0\leqslant\arg(x)\leqslant 2\pi,\ \left|\arg(-y)\right|\leqslant\frac{\pi}{2}-\delta\right\}.$$

We close the path of integration by the semi-circle $\mathfrak{C}$, that is parameterized by $s=(N+1/2)\mathrm{e}^{\mathrm{i}\theta}$ ($|\theta|\leqslant\pi/2$), and then we let $N\to\infty$ through integral values. Recall the estimate [23, p. 7]

$$\left|\frac{\Gamma(a+s)}{\Gamma(c'+s)}\Gamma(-s)(-y)^s\right| = \left(N^{\Re(a-c')-\frac{1}{2}-\left(N+\frac{1}{2}\right)\cos\theta}\,\mathrm{e}^{\left(N+\frac{1}{2}\right)(\delta_1\cos\theta+\delta_2|\sin\theta|)}\right),\quad s\to\infty,\ s\in\mathfrak{C},$$

where $\delta_1 := 1+\log|y|<0$ and $\delta_2 := |\theta|+\left|\arg(-y)\right|-\pi \leqslant -\delta < 0$. It follows from (2.5) that

$$\left|{}_1F_1\left[\begin{matrix}a+s\\ c\end{matrix};x\right]\right| = \mathscr{O}\left(N^{\frac{1}{4}-\frac{1}{2}\Re(c)}\mathrm{e}^{3\sqrt{N|x|}}\right),\quad s\to\infty,\ s\in\mathfrak{C}.$$

So the integral on the semi-circle $\mathfrak{C}$ tends to zero as $N\to\infty$. Finally, we get the representation (4.1) from Cauchy's residue theorem and the assertion that ([4, p. 7])

$$\operatorname*{Res}_{s=n}\Gamma(-s) = \frac{(-1)^{n-1}}{n!}.$$

(ii) Let I denote the right-hand side of (4.3a), and let $\Psi(s)$ denote the integrand given by (4.4). Choose the positive integer $M\geqslant\max\{1,\Re(-a)\}$ and shift the integration contour to the left (which is permissible on account of the exponential decay of the integrand).

Note that $s=-a-n$ ($n\in\mathbb{Z}_{\geqslant 0}$) are the only poles of $\Psi(s)$. Therefore,

$$\operatorname*{Res}_{s=-a-n}\Psi(s) = {}_1F_1\left[\begin{matrix}-n\\ c\end{matrix};x\right]\frac{\Gamma(a+n)}{\Gamma(c'-a-n)}\frac{(-1)^n}{n!}(-y)^{-a-n}.$$

By Cauchy's residue theorem, we obtain

$$\begin{aligned} I &= \frac{\Gamma(c')}{\Gamma(a)}\sum_{n=0}^{M}{}_1F_1\left[\begin{matrix}-n\\ c\end{matrix};x\right]\frac{\Gamma(a+n)}{\Gamma(c'-a-n)}\frac{(-1)^n}{n!}(-y)^{-a-n} + \underbrace{\frac{1}{2\pi\mathrm{i}}\frac{\Gamma(c')}{\Gamma(a)}\int_{C_M}\Psi(s)\mathrm{d}s}_{\text{denoted by } R_M(y)} \\ &= \frac{\Gamma(c')}{\Gamma(c'-a)}\sum_{n=0}^{M}{}_1F_1\left[\begin{matrix}-n\\ c\end{matrix};x\right]\frac{(a)_n(1+a-c')_n}{n!}(-y)^{-a-n} + R_M(y), \end{aligned}$$

where C_M denotes the vertical line $\Re(s)=\Re(-a)-M-1/2$. On account of (4.5) and (4.6), we find that the integral $R_M(y)$ is convergent and that $R_M(y)=\mathscr{O}_M(|y|^{-\Re(a)-M-\frac{1}{2}})$, which completes the proof. □

We can also obtain the asymptotics of $\Psi_2[x,y]$ as $y\to\infty$ in the right half-plane.

Theorem 4.2. *Let $\Re(a)>0$ and $c,c'\in\mathbb{C}\setminus\mathbb{Z}_{\leqslant 0}$. Fix $\delta\in\left(0,\frac{\pi}{2}\right]$ and let*

$$S_\delta := \left\{z\in\mathbb{C} : z\neq 0,\ \left|\arg(z)\right|\leqslant\frac{\pi}{2}-\delta\right\}.$$

(i) *As $t\to+\infty$, for $x,y\in S_\delta$ with $0<K_1\leqslant|x|,|y|\leqslant K_2<\infty$,*

$$\Psi_2[a;c,c';tx,ty]\sim\frac{\Gamma(c)\Gamma(c')}{2\sqrt{\pi}\,\Gamma(a)}x^{\frac14-\frac12c}y^{\frac14-\frac12c'}\left(\sqrt{x}+\sqrt{y}\right)^{2a-c-c'}t^{a-c-c'+\frac12}\mathrm{e}^{t(\sqrt{x}+\sqrt{y})^2}. \tag{4.7a}$$

(ii) *For fixed $x\in\mathbb{C}\setminus\{0\}$, as $y\to\infty$ with $y\in S_\delta$,*

$$\Psi_2[a;c,c';x,y]\sim\frac{\Gamma(c)\Gamma(c')}{2\sqrt{\pi}\,\Gamma(a)}x^{\frac14-\frac12c}y^{\frac14-\frac12c'}\left(\sqrt{x}+\sqrt{y}\right)^{2a-c-c'}\mathrm{e}^{(\sqrt{x}+\sqrt{y})^2}. \tag{4.7b}$$

Proof. Let us start with the integral representation [42, Eq. (2.4c)]

$$\Psi_2[a;c,c';x,y]=\frac{1}{\Gamma(a)}\int_0^\infty s^{a-1}\mathrm{e}^{-s}\,{}_0F_1\left[\begin{matrix}-\\c\end{matrix};xs\right]{}_0F_1\left[\begin{matrix}-\\c'\end{matrix};ys\right]\mathrm{d}s.$$

Using (3.3) and setting $s=w^2$, we obtain for $P_t(x,y):=\Psi_2[a;c,c';t^2x^2,t^2y^2]$ that

$$P_t(x,y)=\frac{2\Gamma(c)\Gamma(c')}{\Gamma(a)}x^{1-c}y^{1-c'}t^{2-c-c'}\int_0^\infty w^{2a-c-c'+1}\mathrm{e}^{-w^2}I_{c-1}(2xtw)\,I_{c'-1}\left(2ytw\right)\mathrm{d}w, \tag{4.8}$$

where we assume for the moment that t,x,y are positive.

(i) Denote $\alpha=2x(x+y)$, $\beta=2y(x+y)$ and $\tau=t(x+y)$. Set $w=\tau u$ in (4.8) and then

$$P_t(x,y)=\frac{2\Gamma(c)\Gamma(c')}{\Gamma(a)}x^{1-c}y^{1-c'}t^{2-c-c'}\tau^{2a-c-c'+2}\int_0^\infty u^{2a-c-c'+1}\mathrm{e}^{-\tau^2u^2}I_{c-1}\left(\alpha t^2u\right)I_{c'-1}\left(\beta t^2u\right)\mathrm{d}u, \tag{4.9}$$

which, by analytic continuation, is valid for $t>0$ and $x^2,y^2\in S_\delta$.

Denote by $\mathfrak{I}$ the integral on the right side of (4.9). Following the proof of Theorem 3.2, we have

$$\begin{aligned}\mathfrak{I}&=\left(\int_0^{t^{-2}}+\int_{t^{-2}}^\infty\right)u^{2a-c-c'+1}\mathrm{e}^{-\tau^2u^2}I_{c-1}\left(\alpha t^2u\right)I_{c'-1}\left(\beta t^2u\right)\mathrm{d}u\\&\sim\int_{t^{-2}}^\infty u^{2a-c-c'+1}\mathrm{e}^{-\tau^2u^2}I_{c-1}\left(\alpha t^2u\right)I_{c'-1}\left(\beta t^2u\right)\mathrm{d}u\\&\sim\frac{t^{-2}\mathrm{e}^{\tau^2}}{2\pi\sqrt{\alpha\beta}}\int_{t^{-2}}^\infty u^{2a-c-c'+1}\mathrm{e}^{-\tau^2(u-1)^2}\mathrm{d}u\sim\frac{t^{-2}\tau^{-1}\mathrm{e}^{\tau^2}}{2\sqrt{\pi\alpha\beta}},\end{aligned}$$

where in the last line we used (2.2) and Olver's Laplace method [33, Theorem I]. Hence, as $t\to+\infty$, for $x,y\in S_\delta$ with $0<K_1\leqslant|x|,|y|\leqslant K_2<\infty$,

$$\begin{aligned}P_t(x,y)&\sim\frac{\Gamma(c)\Gamma(c')}{\sqrt{\pi\alpha\beta}\,\Gamma(a)}x^{1-c}y^{1-c'}t^{-c-c'}\tau^{2a-c-c'+1}\mathrm{e}^{\tau^2}\\&=\frac{\Gamma(c)\Gamma(c')}{2\sqrt{\pi}\,\Gamma(a)}x^{\frac12-c}y^{\frac12-c'}\left(x+y\right)^{2a-c-c'}t^{2(a-c-c')+1}\mathrm{e}^{t^2(x+y)^2},\end{aligned}$$

which is consistent with the desired result (4.7a).

(ii) Denote $\lambda=x+y$. Taking $t=1$ and setting $w=\lambda v$ in (4.8), we have

$$P_1(x,y)=\frac{2\Gamma(c)\Gamma(c')}{\Gamma(a)}x^{1-c}y^{1-c'}\lambda^{2a-c-c'+2}\int_0^\infty v^{2a-c-c'+1}\mathrm{e}^{-\lambda^2v^2}I_{c-1}\left(2x\lambda v\right)I_{c'-1}\left(2y\lambda v\right)\mathrm{d}v, \tag{4.10}$$

which, by analytic continuation, is valid for fixed $x\in\mathbb{C}\setminus\{0\}$ and large y with $y^2\in S_\delta$.

Denote by $\mathfrak{T}$ the integral on the right side of (4.10). Similar to the analysis in (i), we obtain

$$\begin{aligned}\mathfrak{T}&=\left(\int_0^{|\lambda|^{-2}}+\int_{|\lambda|^{-2}}^{|\lambda|^{-1}}+\int_{|\lambda|^{-1}}^\infty\right)v^{2a-c-c'+1}\mathrm{e}^{-\lambda^2v^2}I_{c-1}\left(2x\lambda v\right)I_{c'-1}\left(2y\lambda v\right)\mathrm{d}v\\&\sim\int_{|\lambda|^{-1}}^\infty v^{2a-c-c'+1}\mathrm{e}^{-\lambda^2v^2}I_{c-1}\left(2x\lambda v\right)I_{c'-1}\left(2y\lambda v\right)\mathrm{d}v\end{aligned}$$

$$\sim \frac{1}{4\pi} x^{-\frac{1}{2}} y^{-\frac{1}{2}} \lambda^{-1} \mathrm{e}^{\lambda^2} \int_{|\lambda|^{-1}}^{\infty} v^{2a-c-c'} \mathrm{e}^{-\lambda^2 (v-1)^2} \mathrm{d}v \sim \frac{1}{4\sqrt{\pi}} x^{-\frac{1}{2}} y^{-\frac{1}{2}} \lambda^{-2} \mathrm{e}^{\lambda^2}.$$

Hence, for fixed $x \in \mathbb{C} \setminus \{0\}$, as $y \to \infty$ with $y \in S_\delta$,

$$P_1(x,y) \sim \frac{\Gamma(c)\Gamma(c')}{2\sqrt{\pi}\,\Gamma(a)} x^{\frac{1}{2}-c} y^{\frac{1}{2}-c'} \lambda^{2a-c-c'} \mathrm{e}^{\lambda^2} = \frac{\Gamma(c)\Gamma(c')}{2\sqrt{\pi}\,\Gamma(a)} x^{\frac{1}{2}-c} y^{\frac{1}{2}-c'} \big(x+y\big)^{2a-c-c'} \mathrm{e}^{(x+y)^2},$$

which yields the leading-term behavior shown in (4.7b). □

Before proceeding to the next result, we first state a useful lemma, which can be easily obtained by repeating the proof of [22, Lemma 2.3].

Lemma 4.3. *For z bounded away from the points $-b+1, \cdots, -b+n$,*

$$\big|(z+b-1)\cdots(z+b-n)\big|^{-1} \ll \lambda_n^{-1},$$

where

$$\begin{aligned} \lambda_n &:= \begin{cases} 1, & 0 \leqslant n \leqslant 2, \\ \min\limits_{1\leqslant k\leqslant n-1} (k-1)!(n-k-1)!, & n \geqslant 3 \end{cases} \\ &= \begin{cases} 1, & 0 \leqslant n \leqslant 2, \\ (m-1)!\,m!, & n = 2m+1 \geqslant 3, \\ (m-1)!^2, & n = 2m \geqslant 3. \end{cases} \end{aligned} \tag{4.11}$$

Theorem 4.4. *Assume that $a, b \in \mathbb{C}$, $c, d \in \mathbb{C} \setminus \mathbb{Z}_{\geqslant 0}$ and $a - b \in \mathbb{C} \setminus \mathbb{Z}$. Let $w > 0$ be a number such that*

$$w > \max\{\Re(a), \Re(b), \Re(d)\}$$

and that the fractional parts of $w - \Re(a)$ and $w - \Re(b)$ are both in the interval $(\varepsilon, 1)$, where $\varepsilon > 0$ is a small number. Then for any $n \in \mathbb{Z}_{\geqslant 0}$,

$${}_2F_2\left[\begin{matrix} a, b-n \\ c, \ d \end{matrix}; -z\right] = \frac{\Gamma(c)\Gamma(d)}{\Gamma(a)\Gamma(b-n)} \big\{S_n(z) + T_n(z) + R_{n,w}(z)\big\} \tag{4.12}$$

as $z \to \infty$ such that $|\arg(z)| < \pi$ and z is bounded away from the points $-b+k\,(k \in \mathbb{Z})$, where

$$S_n(z) = \sum_{k=0}^{\lfloor w-\Re(a)\rfloor} \frac{\Gamma(a+k)\Gamma(b-a-n-k)}{\Gamma(c-a-k)\Gamma(d-a-k)} \frac{(-1)^k}{k!} z^{-a-k}, \tag{4.13a}$$

$$T_n(z) = \sum_{k=0}^{\lfloor w-\Re(b)\rfloor+n} \frac{\Gamma(b-n+k)\Gamma(a-b+n-k)}{\Gamma(c-b+n-k)\Gamma(d-b+n-k)} \frac{(-1)^k}{k!} z^{n-b-k}, \tag{4.13b}$$

and

$$R_{n,w}(z) = \mathscr{O}\left(\lambda_n^{-1}\left\{|z|^{-w} + |z|^{\Re(a+b-c-d)}\,\mathrm{e}^{-\Re(z)}\right\}\right) \tag{4.13c}$$

with λ_n defined in (4.11).

Proof. The proof is much akin to that of [22, Theorem 2.4], so we show only the outline of the proof.

First of all, we use Cauchy's residue theorem to obtain expansion (4.12), where the remainder term $R_{n,w}(z)$ is given explicitly by

$$R_{n,w}(z) = \frac{1}{2\pi \mathrm{i}} \int_{\mathscr{C}} h_n(s) z^s \mathrm{d}s, \quad h_n(s) := \frac{\Gamma(a+s)\Gamma(b-n+s)}{\Gamma(c+s)\Gamma(d+s)} \Gamma(-s).$$

Here $\mathscr{C}$ is a negative-oriented loop that consists of the vertical line

$$L^v: \quad s = -w + \mathrm{i}t, \quad |t| \leqslant T$$

and the contours $L^{\pm}$ which pass to infinity in the directions $\pm\theta_0$ $(0<\theta_0<\frac{\pi}{2})$. Furthermore, $\mathscr{C}$ is taken to embrace all the poles of $h_n(s)$ in the right of L^{ν}, and $L^{\pm}$ are taken to be bounded away from the poles of $h_n(s)/\Gamma(-s)$ in the right of L^{ν}.

Then, divide $R_{n,w}(z)$ into two parts:

$$R_{n,w}(z)=\frac{1}{2\pi\mathrm{i}}\int_{L^{\nu}}h_n(s)z^s\mathrm{d}s+\frac{1}{2\pi\mathrm{i}}\int_{L^{\pm}}h_n(s)z^s\mathrm{d}s=:R_{\nu}(z)+R_{\pm}(z).$$

Note that for $s\in\mathscr{C}$,

$$|h_n(s)|=|h_0(s)|\cdot\big|(s+b-1)\cdots(s+b-n)\big|^{-1}\ll|h_0(s)|\,\lambda_n^{-1}.$$

It follows from [29, Eq. (2.4) and (2.6)] that

$$R_{\nu}(z)\ll\lambda_n^{-1}\int_{-w-\mathrm{i}\infty}^{-w+\mathrm{i}\infty}\big|h_0(s)z^s\big||\mathrm{d}s|\ll\lambda_n^{-1}|z|^{-w}.$$

According to [22, pp. 5–6], we also have

$$R_{\pm}(z)\ll\lambda_n^{-1}\int_{L^{\pm}}\big|h_0(s)z^s\big||\mathrm{d}s|\ll\lambda_n^{-1}|z|^{\Re(a+b-c-d)}\,\mathrm{e}^{-\Re(z)}.$$

Finally, combining the estimates on $R_{\nu}(z)$ and $R_{\pm}(z)$ we obtain the remainder estimate (4.13c). □

As a direct consequence of Theorem 4.4, we have the following corollary.

Corollary 4.5. *Assume that $b\in\mathbb{C}$ and $d\in\mathbb{C}\setminus\mathbb{Z}_{\leqslant 0}$. Let N be an integer such that $N>\max\{|b|,|b-d|\}$. Then for any $n\in\mathbb{Z}_{\geqslant 0}$,*

$${}_1F_1\left[\begin{matrix}b-n\\ d\end{matrix};-z\right]=\frac{\Gamma(d)}{\Gamma(b-n)}\sum_{k=0}^{n+N}\frac{\Gamma(b-n+k)}{\Gamma(d-b+n-k)}\frac{(-1)^k}{k!}z^{n-b-k}+R(n;z) \tag{4.14}$$

as $z\to\infty$ such that $\big|\arg(z)\big|<\pi$ and z is bounded away from the points $-b+k$ $(k\in\mathbb{Z})$, where

$$R(n;z)\ll(n+1)^{-\Re(b)}\,n!\,\lambda_n^{-1}\left\{|z|^{-\Re(b)-N-\frac{1}{2}}+|z|^{\Re(b-d)}\,\mathrm{e}^{-\Re(z)}\right\}.$$

Proof. It suffices to verify the error term here. Take $a=c=\frac{1}{2}$ and $w-\Re(b)=N+\frac{1}{2}$ in (4.12). Then

$$R(n;z)=\frac{\Gamma(d)}{\Gamma(b-n)}R_{n,w}(z)=(-1)^n\,n!\,\frac{\Gamma(d)}{\Gamma(b)}\frac{(1-b)_n}{n!}R_{n,w}(z).$$

The estimate on $R(n;z)$ follows from (4.13c) and [21, Lemma 2.1]. □

We are in a position to derive the asymptotics of Ψ_2 under the condition (1.3).

Theorem 4.6. *Assume that $a\in\mathbb{C}$ and $c,c'\in\mathbb{C}\setminus\mathbb{Z}_{\leqslant 0}$. Set $xy=\beta$. Then under the condition that*

$$x\to\infty,\ \big|\arg(x)\big|<\pi;\quad x\text{ is bounded away from }a-c+k\,(k\in\mathbb{Z});\quad 0<\beta_1\leqslant|\beta|\leqslant\beta_2<\infty,$$

the function $\Psi_2\equiv\Psi_2[a;c,c';x,y]$ admits the asymptotic expansion

$$\begin{aligned}\Psi_2=&\frac{\Gamma(c)}{\Gamma(c-a)}x^{-a}\sum_{m=0}^{N-1}b_1(m)x^{-m}+\frac{\Gamma(c)}{\Gamma(a)}x^{a-c}\mathrm{e}^{x}\left(\sum_{m=0}^{N-1}b_2(m)x^{-m}+\sum_{m=1}^{N-1}b_3(m)x^{-m}\right)\\&+\mathscr{O}\left(|x|^{-\Re(a)-N}+|x|^{\Re(a-c)-N-\frac{1}{2}}\,\mathrm{e}^{\Re(x)}\right),\end{aligned} \tag{4.15}$$

where N is any positive integer such that $N>\max\{1,|a|,|a-c|\}$, and

$$b_1(m)=\sum_{\substack{n,k\geqslant 0\\ 2n+k=m}}\frac{(a)_{n+k}\,(a-c+1)_{n+k}}{(c')_n\,n!\,k!}\big(-\beta\big)^n, \tag{4.16a}$$

$$b_2(m)={}_3F_4\left[\begin{matrix}1,a+m,a-c+1+m\\ a,a-c+1,1+m,c'+m\end{matrix};\beta\right]\frac{(a)_m\,(a-c+1)_m}{(c')_m\,m!^2}\beta^m, \tag{4.16b}$$

$$b_3(m)=\frac{1}{m!}\sum_{n=0}^{m-1}\frac{(1-a-n)_m\,(c-a-n)_m}{(c')_n\,n!}\beta^n. \tag{4.16c}$$

Proof. For $0 \leqslant n \leqslant N$, we get from (2.1) that

$$ {}_1F_1\left[\begin{matrix} c-a-n \\ c \end{matrix}; -x\right] = f_n(x) + g_n(x) + \mathscr{O}\left(|x|^{\Re(a-c)-N-\frac{1}{2}} + |x|^{-\Re(a)-n-N}\,\mathrm{e}^{-\Re(x)}\right), \tag{4.17} $$

where

$$ f_n(x) = \frac{\Gamma(c)}{\Gamma(a+n)} \sum_{k=0}^{n+N} \frac{(c-a-n)_k\,(1-a-n)_k}{k!}\, x^{a-c+n-k}, $$

$$ g_n(x) = \frac{\Gamma(c)}{\Gamma(c-a-n)}\mathrm{e}^{-x} \sum_{k=0}^{N-1} \frac{(a+n)_k\,(a-c+n+1)_k}{k!}\, x^{-a-n-k}. $$

For $n \geqslant N+1$, Corollary 4.5 gives that

$$ {}_1F_1\left[\begin{matrix} c-a-n \\ c \end{matrix}; -x\right] = f_n(x) + \mathscr{O}\left(n^{\Re(a-c)}\, n!\, \lambda_n^{-1} \left\{|x|^{\Re(a-c)-N-\frac{1}{2}} + |x|^{-\Re(a)}\,\mathrm{e}^{-\Re(x)}\right\}\right). \tag{4.18} $$

Write $P(x,y) = \mathrm{e}^{-x}\Psi_2[a; c, c'; x, y]$. It follows from (4.2), (4.17) and (4.18) that

$$ \begin{aligned} P(x,y) &= \left(\sum_{n=0}^{N} + \sum_{n=N+1}^{\infty}\right) \frac{(a)_n}{(c')_n}\, {}_1F_1\left[\begin{matrix} c-a-n \\ c \end{matrix}; -x\right] \frac{y^n}{n!} \\ &= \sum_{n=0}^{\infty} \frac{(a)_n}{(c')_n} f_n(x) \frac{y^n}{n!} + \sum_{n=0}^{N} \frac{(a)_n}{(c')_n} g_n(x) \frac{y^n}{n!} + R(x,y) \\ &=: F(x,y) + G(x,y) + R(x,y). \end{aligned} $$

Next, we shall first deal with the remainder $R(x,y)$ and then study the functions $F(x,y)$ and $G(x,y)$ by the series manipulation technique.

(1) *Estimate of* $R(x,y)$. The error introduced by (4.17) is clearly

$$ R_1(x,y) = \mathscr{O}\left(|x|^{\Re(a-c)-N-\frac{1}{2}} + |x|^{-\Re(a)-N}\,\mathrm{e}^{-\Re(x)}\right), $$

whereas the error introduced by (4.18) is

$$ R_2(x,y) \ll \left(|x|^{\Re(a-c)-N-\frac{1}{2}} + |x|^{-\Re(a)}\,\mathrm{e}^{-\Re(x)}\right) \sum_{n\geqslant N+1} n^{\Re(a-c)}\, n!\, \lambda_n^{-1} \left|\frac{(a)_n}{(c')_n}\right| \frac{|y|^n}{n!}. $$

Using [21, Lemma 2.1] and recalling (4.11), we have

$$ \begin{aligned} R_2(x,y) &\ll \left(|x|^{\Re(a-c)-N-\frac{1}{2}} + |x|^{-\Re(a)}\,\mathrm{e}^{-\Re(x)}\right) \sum_{n\geqslant N+1} n^{\Re(2a-c-c')}\, \lambda_n^{-1}\, \beta_2^n\, |x|^{-n} \\ &\ll |x|^{\Re(a-c)-2N-\frac{3}{2}} + |x|^{-\Re(a)-N-1}\,\mathrm{e}^{-\Re(x)}. \end{aligned} $$

Therefore, the remainder is

$$ R(x,y) = R_1(x,y) + R_2(x,y) = \mathscr{O}\left(|x|^{\Re(a-c)-N-\frac{1}{2}} + |x|^{-\Re(a)-N}\,\mathrm{e}^{-\Re(x)}\right). $$

(2) *Analysis of* $F(x,y)$ *and* $G(x,y)$. Bear in mind that $xy = \beta$. We have

$$ \begin{aligned} G(x,y) &= \frac{\Gamma(c)}{\Gamma(c-a)} x^{-a}\mathrm{e}^{-x} \sum_{n=0}^{N} \frac{(a)_n\,(a-c+1)_n}{(c')_n\, n!} (-\beta)^n x^{-2n} \sum_{k=0}^{N-1} \frac{(a+n)_k\,(a-c+n+1)_k}{k!} x^{-k} \\ &= \frac{\Gamma(c)}{\Gamma(c-a)} x^{-a}\mathrm{e}^{-x} \sum_{m=0}^{N-1} x^{-m} \sum_{\substack{n,k\geqslant 0 \\ 2n+k=m}} \frac{(a)_n\,(a-c+1)_n\,(a+n)_k\,(a-c+n+1)_k}{(c')_n\, n!\, k!} (-\beta)^n \end{aligned} $$

$$
\begin{aligned}
&+\mathcal{O}\left(|x|^{-\Re(a)-N}\,\mathrm{e}^{-\Re(x)}\right)\\
&=\frac{\Gamma(c)}{\Gamma(c-a)}x^{-a}\mathrm{e}^{-x}\sum_{m=0}^{N-1}b_1(m)x^{-m}+\mathcal{O}\left(|x|^{-\Re(a)-N}\,\mathrm{e}^{-\Re(x)}\right).
\end{aligned}
$$

For $F(x,y)$, we have

$$
\begin{aligned}
F(x,y)&=\frac{\Gamma(c)}{\Gamma(a)}x^{a-c}\sum_{n=0}^{\infty}\sum_{k=0}^{n+N}\frac{(c-a-n)_k\,(1-a-n)_k}{(c')_n\,n!\,k!}\beta^n x^{-k}\\
&=\frac{\Gamma(c)}{\Gamma(a)}x^{a-c}\left(\sum_{n=0}^{\infty}\sum_{k=0}^{n}+\sum_{n=0}^{\infty}\sum_{k=n+1}^{n+N}\right)\frac{(c-a-n)_k\,(1-a-n)_k}{(c')_n\,n!\,k!}\beta^n x^{-k}\\
&=:\frac{\Gamma(c)}{\Gamma(a)}x^{a-c}\big(F^-(x,y)+F^+(x,y)\big),
\end{aligned}
$$

where

$$
\begin{aligned}
F^-(x,y)&=\sum_{n=0}^{\infty}\sum_{k=0}^{n}\frac{(c-a-n)_k\,(1-a-n)_k}{(c')_n\,n!\,k!}\beta^n x^{-k}\\
&=\sum_{n=0}^{\infty}\sum_{k=0}^{\infty}\frac{(c-a-n-k)_k\,(1-a-n-k)_k}{(c')_{n+k}\,(n+k)!\,k!}\beta^{n+k}x^{-k}\\
&=\sum_{k=0}^{\infty}{}_3F_4\left[\begin{matrix}1,a+k,a-c+1+k\\ a,a-c+1,1+k,c'+k\end{matrix};\beta\right]\frac{(a)_k\,(a-c+1)_k}{(c')_k\,k!^2}\beta^k x^{-k}
\end{aligned}
$$

and

$$
\begin{aligned}
F^+(x,y)&=\sum_{n=0}^{\infty}\sum_{k=n+1}^{n+N}\frac{(c-a-n)_k\,(1-a-n)_k}{(c')_n\,n!\,k!}\beta^n x^{-k}\\
&=\sum_{n=0}^{N-1}\sum_{j=1}^{N-1}\frac{(c-a-n)_{n+j}\,(1-a-n)_{n+j}}{(c')_n\,n!\,(n+j)!}\beta^n x^{-n-j}+\mathcal{O}\left(|x|^{-N}\right)\\
&=\sum_{m=1}^{N-1}\frac{x^{-m}}{m!}\sum_{n=0}^{m-1}\frac{(1-a-n)_m\,(c-a-n)_m}{(c')_n\,n!}\beta^n+\mathcal{O}\left(|x|^{-N}\right)\\
&=\sum_{m=1}^{N-1}b_3(m)x^{-m}+\mathcal{O}\left(|x|^{-N}\right).
\end{aligned}
$$

It is clear that the leading terms of the expansion (4.15) are obtained from $F(x,y)$ and $G(x,y)$.

Combining the above expansions gives the asymptotic expansion (4.15) for Ψ_2. □

5. Appell function F_2

Recall that F_2 satisfies the symmetry relation $F_2[a,b,b';c,c';x,y]=F_2[a,b',b;c',c;y,x]$. So it suffices to study the asymptotic expansion of $F_2\left[\frac{x}{y},y\right]$ as $y\to 0$.

The following expression for F_2 is due to Jaeger (see, for example, [3, Eq. (9)], [26] and [39, p. 294]).

Theorem 5.1 (Jaeger). *Let $a,b,b'\in\mathbb{C}$, $c,c'\in\mathbb{C}\setminus\mathbb{Z}_{\leqslant 0}$ and $a-b\notin\mathbb{Z}$. Then for $|y|<1$ and $|x|>|y|+1$,*

$$
\begin{aligned}
F_2[a,b,b';c,c';x,y]&=\mathfrak{f}_c(b,a)\,(-x)^{-a}\sum_{n=0}^{\infty}{}_2F_1\left[\begin{matrix}-n,b'\\ c'\end{matrix};y\right]\frac{(a)_n\,(1-c+a)_n}{(1-b+a)_n\,n!}\frac{1}{x^n}\\
&\quad+\mathfrak{f}_c(a,b)\,(-x)^{-b}\sum_{n=0}^{\infty}{}_2F_1\left[\begin{matrix}a-b-n,b'\\ c'\end{matrix};y\right]\frac{(b)_n\,(1-c+b)_n}{(1-a+b)_n\,n!}\frac{1}{x^n},
\end{aligned}
\tag{5.1}
$$

where $\mathfrak{f}_\gamma(a,b)$ is given by (3.8a).

Using a similar approach to the proof of Theorem 3.6, we obtain from Jaeger's formula (5.1) the following result.

Corollary 5.2. *Let $a,b,b'\in\mathbb{C}$, $c,c'\in\mathbb{C}\setminus\mathbb{Z}_{\leqslant 0}$ and $a-b\notin\mathbb{Z}$. For fixed $x\in\mathbb{C}\setminus\{0\}$,*

$$F_2\left[a,b,b';c,c';\frac{x}{y},y\right]=\mathfrak{f}_c(b,a)\left(-\frac{x}{y}\right)^{-a}\sum_{m=0}^{N-1}c_1(m)y^m+\mathfrak{f}_c(a,b)\left(-\frac{x}{y}\right)^{-b}\sum_{m=0}^{N-1}c_2(m)y^m \\ +\mathscr{O}\left(|y|^{\Re(a)+N}+|y|^{\Re(b)+N}\right) \tag{5.2a}$$

as $y\to 0$ in $\left|\arg\left(-\frac{x}{y}\right)\right|<\pi$, where N is any positive integer, $\mathfrak{f}_\gamma(a,b)$ is given by (3.8a) and

$$c_1(m)=\sum_{\substack{k,\ell\geqslant 0\\ k+\ell=m}}\frac{(-k)_\ell\,(b')_\ell\,(a)_k\,(1-c+a)_k}{(c')_\ell\,(1-b+a)_k\,\ell!\,k!}x^{-k}, \tag{5.2b}$$

$$c_2(m)=\sum_{\substack{k,\ell\geqslant 0\\ k+\ell=m}}\frac{(b')_\ell\,(a-b-k)_\ell\,(b)_k\,(1-c+b)_k}{(c')_\ell\,(1-a+b)_k\,\ell!\,k!}x^{-k}. \tag{5.2c}$$

6. Two methods

In this section, we propose two elementary methods, which are easy to use and powerful for deducing the asymptotics of multiple hypergeometric functions.

6.1. Uniformity approach

Our proof of Theorem 4.6 is based on the approach used in [22], which is indeed effective for deriving the asymptotics of multiple hypergeometric functions. Let us give a detailed description:

- (**Uniformity approach**) *For $n\in\mathbb{Z}_{\geqslant 0}$, denote*

$$F_n(z)={}_pF_q\left[\begin{matrix}a_1+\lambda_1 n,\cdots,a_r+\lambda_r n,a_{r+1},\cdots,a_p\\ b_1+\mu_1 n,\cdots,b_s+\mu_s n,b_{s+1},\cdots,b_q\end{matrix};z\right], \tag{6.1}$$

where $\lambda_1,\cdots,\lambda_r,\mu_1,\cdots,\mu_s$ are fixed complex numbers. In view of [29, Sect. 5]*, for fixed n,*

$$F_n(z)\sim A_n(z)+E_n(z),\quad z\to\infty,$$

where $A_n(z)$ is the algebraic expansion and $E_n(z)$ is the exponential expansion.

To derive the asymptotics of the function

$$\mathscr{F}(z,w):=\sum_{n=0}^{\infty}a_nF_n(z)w^n, \tag{6.2}$$

we first deduce the estimate for $F_n(z)$, that is akin to (4.12) and valid uniformly for $n\geqslant 0$, namely,

$$F_n(z)=A_n^{(N)}(z)+E_n^{(N)}(z)+\mathscr{O}\big(c_n r(z)\big),\quad n\in\mathbb{Z}_{\geqslant 0},\,|z|\geqslant K, \tag{6.3}$$

where $A_n^{(N)}(z)$ (resp. $E_n^{(N)}(z)$) is the sum of first N terms of $A_n(z)$ (resp. $E_n(z)$). Then inserting (6.3) into (6.2) and using the series manipulation technique, we get the asymptotic expansion of $\mathscr{F}(z,w)$.

Let us present the powerful utility of the uniformity approach. Very recently, Brychkov and Savischenko systematically studied the formulas of the Horn functions $H_1,\cdots,H_7$ and the confluent Horn functions $H_1^{(c)},\cdots,H_{11}^{(c)}$. For example, in [10, 12, 13], they obtained the series expansions below:

$$H_2^{(c)}[a,b,c;d;x,y]=\sum_{n=0}^{\infty}\frac{(c)_n}{(1-a)_n}\,{}_2F_1\left[\begin{matrix}a-n,b\\ d\end{matrix};x\right]\frac{(-y)^n}{n!},$$

$$H_3^{(c)}[a,b;d;x,y]=\sum_{n=0}^{\infty}\frac{1}{(1-a)_n}\,{}_2F_1\left[\begin{matrix}a-n,b\\ d\end{matrix};x\right]\frac{(-y)^n}{n!},$$

$$H_4^{(c)}[a,c;d;x,y]=\sum_{n=0}^{\infty}\frac{(c)_n}{(1-a)_n}\,{}_1F_1\left[\begin{matrix}a-n\\ d\end{matrix};x\right]\frac{(-y)^n}{n!},$$
$$H_5^{(c)}[a;c;x,y]=\sum_{n=0}^{\infty}\frac{1}{(1-a)_n}\,{}_1F_1\left[\begin{matrix}a-n\\ c\end{matrix};x\right]\frac{(-y)^n}{n!}.$$

By repeating the proof of Theorem 4.4, one can obtain an expansion of ${}_2F_1[a-n,b;c;x]$, which is similar to (4.12). Then the use of the uniformity approach will give the full asymptotic expansions of $H_k^{(c)}\left[x,\frac{\beta}{x}\right]$ $(k=2,3,4,5)$ for large x. Furthermore, in view of the series expansions derived in [2, 11], the same apply to the Appell functions F_1,F_3 and the Humbert functions $\Phi_1,\Phi_2,\Phi_3,\Xi_1$.

6.2. Separation method

We provide a heuristic approach, called **separation method**, to reproduce (3.9) and (5.1). This method is of great benefit to physicists, although it is not rigorous in general.

Proposition 6.1. *Under the necessary conditions on a,b,c,c', for $x\to+\infty$ and fixed $y>0$, one has*

$$\begin{aligned}\Psi_1[a,b;c,c';-x,y]\sim{}&\mathfrak{f}_c(b,a)x^{-a}\sum_{k=0}^{\infty}{}_1F_1\left[\begin{matrix}-k\\ c'\end{matrix};y\right]\frac{(a)_k(1-c+a)_k}{(1-b+a)_k\,k!}\frac{1}{(-x)^k}\\&+\mathfrak{f}_c(a,b)x^{-b}\sum_{\ell=0}^{\infty}{}_1F_1\left[\begin{matrix}a-b-\ell\\ c'\end{matrix};y\right]\frac{(b)_\ell(1-c+b)_\ell}{(1-a+b)_\ell\,\ell!}\frac{1}{(-x)^\ell},\end{aligned}\tag{6.4}$$

where $\mathfrak{f}_\gamma(a,b)$ is defined by (3.8a). See also Remark 3.7 after replacing y by $\frac{y}{x}$ in (6.4).

Proof. Our starting point is the integral representation (3.1c). For large x, we split the integral into two parts:

$$\begin{aligned}\Psi_1[a,b;c,c';-x,y]&=\frac{1}{\Gamma(a)}\left(\int_0^{\eta/x}+\int_{\eta/x}^{\infty}\right)u^{a-1}\mathrm{e}^{-u}\,{}_1F_1\left[\begin{matrix}b\\ c\end{matrix};-xu\right]{}_0F_1\left[\begin{matrix}-\\ c'\end{matrix};yu\right]\mathrm{d}u\\&=:\mathfrak{T}_1^{(\eta)}+\mathfrak{T}_2^{(\eta)},\end{aligned}$$

where η is taken such that both $\eta\to+\infty$ and $\eta=o(x)$ hold as $x\to+\infty$.

We begin with the integral $\mathfrak{T}_1^{(\eta)}$. Setting $v=ux$, we obtain

$$\mathfrak{T}_1^{(\eta)}=\frac{x^{-a}}{\Gamma(a)}\int_0^{\eta}v^{a-1}\mathrm{e}^{-\frac{v}{x}}\,{}_1F_1\left[\begin{matrix}b\\ c\end{matrix};-v\right]{}_0F_1\left[\begin{matrix}-\\ c'\end{matrix};\frac{vy}{x}\right]\mathrm{d}v.$$

Since

$$\begin{aligned}\mathrm{e}^{-\frac{v}{x}}\,{}_0F_1\left[\begin{matrix}-\\ c'\end{matrix};\frac{vy}{x}\right]&=\sum_{n=0}^{\infty}\frac{(-1)^n}{n!}\left(\frac{v}{x}\right)^n\sum_{m=0}^{\infty}\frac{v^m}{(c')_m\,m!}\left(\frac{y}{x}\right)^m\\&=\sum_{k=0}^{\infty}\left(\frac{v}{x}\right)^k\sum_{m=0}^{k}\frac{(-1)^{k-m}y^m}{(c')_m\,m!(k-m)!}\\&=\sum_{k=0}^{\infty}\left(\frac{v}{x}\right)^k\frac{(-1)^k}{k!}\sum_{m=0}^{k}\frac{(-k)_m}{(c')_m}\frac{y^m}{m!}=\sum_{k=0}^{\infty}\left(\frac{v}{x}\right)^k\frac{(-1)^k}{k!}\,{}_1F_1\left[\begin{matrix}-k\\ c'\end{matrix};y\right],\end{aligned}$$

we have

$$\mathfrak{T}_1^{(\eta)}=\frac{x^{-a}}{\Gamma(a)}\sum_{k=0}^{\infty}\frac{(-x)^{-k}}{k!}\,{}_1F_1\left[\begin{matrix}-k\\ c'\end{matrix};y\right]\int_0^{\eta}v^{a-1+k}\,{}_1F_1\left[\begin{matrix}b\\ c\end{matrix};-v\right]\mathrm{d}v.$$

To proceed, we notice that [7, p. 408, Eq. (3.28.1.1)]

$$\int_0^{\infty}t^{s-1}\,{}_1F_1\left[\begin{matrix}a\\ b\end{matrix};-t\right]\mathrm{d}t=\frac{\Gamma(s)\Gamma(b)\Gamma(a-s)}{\Gamma(a)\Gamma(b-s)},\quad 0<\Re(s)<\Re(a).$$

By formally letting $x \to +\infty$, we have

$$\begin{aligned}\mathfrak{T}_1^{(\eta)} &\sim \frac{x^{-a}}{\Gamma(a)} \sum_{k=0}^{\infty} \frac{(-x)^{-k}}{k!} {}_1F_1\left[\begin{matrix}-k\\ c'\end{matrix};y\right] \int_0^{\infty} v^{a-1+k} {}_1F_1\left[\begin{matrix}b\\ c\end{matrix};-v\right] \mathrm{d}v \\ &= \frac{x^{-a}}{\Gamma(a)} \sum_{k=0}^{\infty} \frac{(-x)^{-k}}{k!} {}_1F_1\left[\begin{matrix}-k\\ c'\end{matrix};y\right] \frac{\Gamma(a+k)\Gamma(b-a-k)\Gamma(c)}{\Gamma(c-a-k)\Gamma(b)} \\ &= \frac{\Gamma(c)\Gamma(b-a)}{\Gamma(b)\Gamma(c-a)} x^{-a} \sum_{k=0}^{\infty} {}_1F_1\left[\begin{matrix}-k\\ c'\end{matrix};y\right] \frac{(a)_k (1-c+a)_k}{(1-b+a)_k\, k!} \frac{1}{(-x)^k},\end{aligned}$$

where in the last line we used the identity

$$\Gamma(z-k) = (-1)^k \frac{\Gamma(z)}{(1-z)_k}. \tag{6.5}$$

We now turn to the integral $\mathfrak{T}_2^{(\eta)}$ and use a formal derivation to obtain an expansion. Since the ${}_1F_1$ function in the integrand has a negative argument, the algebraic expansion should dominate, which inspires us to write

$$\begin{aligned}\mathfrak{T}_2^{(\eta)} &\sim \frac{1}{\Gamma(a)} \int_{\eta/x}^{\infty} u^{a-1} \mathrm{e}^{-u} {}_0F_1\left[\begin{matrix}-\\ c'\end{matrix};yu\right] \frac{\Gamma(c)}{\Gamma(c-b)} \sum_{n=0}^{\infty} \frac{(b)_n (1+b-c)_n}{n!} (xu)^{-b-n} \mathrm{d}u \\ &= \frac{\Gamma(c)\Gamma(c')}{\Gamma(a)\Gamma(c-b)} \frac{y^{\frac{1}{2}(1-c')}}{x^b} \sum_{n=0}^{\infty} \frac{(b)_n (1+b-c)_n}{n!} x^{-n} \int_{\eta/x}^{\infty} \mathrm{e}^{-u} u^{a-b-\frac{1}{2}(c'+1)-n} I_{c'-1}(2\sqrt{yu}) \mathrm{d}u,\end{aligned}$$

where we used (3.3). By formally letting $x \to +\infty$ and taking into account that $\eta = o(x)$, we get

$$\begin{aligned}\mathfrak{T}_2^{(\eta)} &\sim \frac{\Gamma(c)\Gamma(c')}{\Gamma(a)\Gamma(c-b)} \frac{y^{\frac{1}{2}(1-c')}}{x^b} \sum_{n=0}^{\infty} \frac{(b)_n (1+b-c)_n}{n!} x^{-n} \int_0^{\infty} \mathrm{e}^{-u} u^{a-b-\frac{1}{2}(c'+1)-n} I_{c'-1}(2\sqrt{yu}) \mathrm{d}u \\ &= \frac{2\Gamma(c)\Gamma(c')}{\Gamma(a)\Gamma(c-b)} \frac{y^{\frac{1}{2}(1-c')}}{x^b} \sum_{n=0}^{\infty} \frac{(b)_n (1+b-c)_n}{n!} x^{-n} \int_0^{\infty} \mathrm{e}^{-w^2} w^{2a-2b-c'-2n} I_{c'-1}(2\sqrt{y}\, w) \mathrm{d}w \\ &\sim \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-b)} x^{-b} \sum_{n=0}^{\infty} \frac{(b)_n (1+b-c)_n}{n!} \Gamma(a-b-n) {}_1F_1\left[\begin{matrix}a-b-n\\ c'\end{matrix};y\right] x^{-n},\end{aligned}$$

where we used the Mellin transform [7, Eq. (3.13.2.3)]

$$\int_0^{\infty} t^{s-1} \mathrm{e}^{-at^2} I_\nu(bt) \mathrm{d}t = \frac{2^{-\nu-1} b^\nu}{a^{\frac{1}{2}(s+\nu)}} \frac{\Gamma\left(\frac{1}{2}(s+\nu)\right)}{\Gamma(\nu+1)} {}_1F_1\left[\begin{matrix}\frac{1}{2}(s+\nu)\\ \nu+1\end{matrix};\frac{b^2}{4a}\right], \quad \Re(a)>0, \Re(s+\nu)>0.$$

Then use (6.5) to yield

$$\mathfrak{T}_2^{(\eta)} \sim \frac{\Gamma(c)\Gamma(a-b)}{\Gamma(a)\Gamma(c-b)} x^{-b} \sum_{n=0}^{\infty} {}_1F_1\left[\begin{matrix}a-b-n\\ c'\end{matrix};y\right] \frac{(b)_n (1-c+b)_n}{(1-a+b)_n\, n!} \frac{1}{(-x)^n},$$

which is the second term in (6.4). □

Remark 6.2. *Starting with* (3.1c) *and applying the separation method, one may rigorously deduce the leading asymptotics of* $\Psi_1[x,y]$ *for* $x<1$ *and* $y \to \infty$ *in* $\left|\arg(y)\right| < \frac{\pi}{2}$, *which was given in* [23, Section 3.1].

Proposition 6.3. *Under the necessary conditions on* a, b, b', c, c', *for* $x \to +\infty$ *and fixed* $y<1$, *one has*

$$\begin{aligned}F_2[a,b,b';c,c';-x,y] \sim{}& \mathfrak{f}_c(b,a) x^{-a} \sum_{n=0}^{\infty} {}_2F_1\left[\begin{matrix}-n, b'\\ c'\end{matrix};y\right] \frac{(a)_n (1-c+a)_n}{(1-b+a)_n\, n!} \frac{1}{(-x)^n} \\ &+ \mathfrak{f}_c(a,b) x^{-b} \sum_{n=0}^{\infty} {}_2F_1\left[\begin{matrix}a-b-n, b'\\ c'\end{matrix};y\right] \frac{(b)_n (1-c+b)_n}{(1-a+b)_n\, n!} \frac{1}{(-x)^n},\end{aligned} \tag{6.6}$$

where $\mathfrak{f}_\gamma(a,b)$ *is defined by* (3.8a).

Proof. Let us start with the following Laplace-type integral representation of F_2:

$$F_2[a,b,b';c,c';x,y] = \int_0^\infty t^{a-1}\mathrm{e}^{-t}\,{}_1F_1\left[\begin{matrix}b\\c\end{matrix};xt\right]{}_1F_1\left[\begin{matrix}b'\\c'\end{matrix};yt\right]\mathrm{d}t,$$

which, in view of (2.1), is valid for $\Re(a)>0$ and $\Re(x+y)<1$. As in the proof of Proposition 6.1, we split the integral into two parts:

$$\begin{aligned}F_2[a,b,b';c,c';-x,y] &= \left(\int_0^{\eta/x}+\int_{\eta/x}^\infty\right) t^{a-1}\mathrm{e}^{-t}\,{}_1F_1\left[\begin{matrix}b\\c\end{matrix};-xt\right]{}_1F_1\left[\begin{matrix}b'\\c'\end{matrix};yt\right]\mathrm{d}t\\ &=: \mathfrak{T}_1^{(\eta)}+\mathfrak{T}_2^{(\eta)},\end{aligned}$$

where $\eta>0$ is taken such that both $\eta\to+\infty$ and $\eta=o(x)$ hold as $x\to+\infty$.

Recall that [7, Eq. (3.28.2.1)]

$$\int_0^\infty t^{s-1}\mathrm{e}^{-\sigma t}\,{}_1F_1\left[\begin{matrix}a\\b\end{matrix};\omega t\right]\mathrm{d}t = \frac{\Gamma(s)}{\sigma^s}\,{}_2F_1\left[\begin{matrix}a,s\\b\end{matrix};\frac{\omega}{\sigma}\right].$$

We follow the proof of Proposition 6.1 and obtain that as $x\to+\infty$,

$$\begin{aligned}\mathfrak{T}_1^{(\eta)} &:= \frac{1}{\Gamma(a)}\int_0^{\eta/x} u^{a-1}\mathrm{e}^{-u}\,{}_1F_1\left[\begin{matrix}b\\c\end{matrix};-xu\right]{}_1F_1\left[\begin{matrix}b'\\c'\end{matrix};yu\right]\mathrm{d}u\\
&= \frac{x^{-a}}{\Gamma(a)}\sum_{k=0}^\infty\frac{(-x)^{-k}}{k!}\,{}_2F_1\left[\begin{matrix}-k,b'\\c'\end{matrix};y\right]\int_0^\eta v^{a+k-1}\,{}_1F_1\left[\begin{matrix}b\\c\end{matrix};-v\right]\mathrm{d}v\\
&\sim \frac{x^{-a}}{\Gamma(a)}\sum_{k=0}^\infty\frac{(-x)^{-k}}{k!}\,{}_2F_1\left[\begin{matrix}-k,b'\\c'\end{matrix};y\right]\int_0^\infty v^{a+k-1}\,{}_1F_1\left[\begin{matrix}b\\c\end{matrix};-v\right]\mathrm{d}v\\
&= \frac{\Gamma(c)\Gamma(b-a)}{\Gamma(b)\Gamma(c-a)}x^{-a}\sum_{n=0}^\infty{}_2F_1\left[\begin{matrix}-n,b'\\c'\end{matrix};y\right]\frac{(a)_n(1-c+a)_n}{(1-b+a)_n\,n!}\frac{1}{(-x)^n},\\
\mathfrak{T}_2^{(\eta)} &:= \frac{1}{\Gamma(a)}\int_{\eta/x}^\infty u^{a-1}\mathrm{e}^{-u}\,{}_1F_1\left[\begin{matrix}b\\c\end{matrix};-xu\right]{}_1F_1\left[\begin{matrix}b'\\c'\end{matrix};yu\right]\mathrm{d}u\\
&\sim \frac{1}{\Gamma(a)}\int_{\eta/x}^\infty u^{a-1}\mathrm{e}^{-u}\,{}_1F_1\left[\begin{matrix}b'\\c'\end{matrix};yu\right]\frac{\Gamma(c)}{\Gamma(c-b)}\sum_{n=0}^\infty\frac{(b)_n(1+b-c)_n}{n!}(xu)^{-b-n}\mathrm{d}u\\
&\sim \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-b)}x^{-b}\sum_{n=0}^\infty\frac{(b)_n(1+b-c)_n}{n!}x^{-n}\int_0^\infty\mathrm{e}^{-u}u^{a-b-n-1}\,{}_1F_1\left[\begin{matrix}b'\\c'\end{matrix};yu\right]\mathrm{d}u\\
&\sim \frac{\Gamma(c)\Gamma(a-b)}{\Gamma(a)\Gamma(c-b)}x^{-b}\sum_{n=0}^\infty{}_2F_1\left[\begin{matrix}a-b-n,b'\\c'\end{matrix};y\right]\frac{(b)_n(1-c+b)_n}{(1-a+b)_n\,n!}\frac{1}{(-x)^n}.\end{aligned}$$

Combining these results gives the desired formula. □

7. Concluding remarks

In this paper, we deduced the asymptotics of the Humbert functions Ψ_1,Ψ_2 and the Appell function F_2 under the condition (1.3). Our methods should be generalizable to find similarly the asymptotics of the Humbert functions $\Phi_1,\Phi_2,\Phi_3,\Xi_1,\Xi_2$ and the Appell functions F_1,F_3; see Remark 3.5 and Section 6.1.

We also derived the asymptotics of Ψ_2 for one or two large arguments. As an application, our result (4.7a) corrects Saran's estimate on Ψ_2 [37, Section 8, Eq. (1)]

$$\Psi_2[a,c;c';x,y]\sim C\,\mathrm{e}^{x+y+2\sqrt{xy}},\quad x,y\to\infty.$$

Recall that Saran's function F_E admits a Laplace-type integral [37, p. 134, Eq. (1)]

$$F_E[a_1,a_1,a_1,b_1,b_2,b_2;c_1,c_2,c_3;x,y,z]$$

$$= \frac{1}{\Gamma(a_1)} \int_0^\infty \mathrm{e}^{-t} t^{a_1-1} {}_1F_1\left[\begin{matrix} b_1 \\ c_1 \end{matrix}; xt\right] \Psi_2\left[b_2; c_2, c_3; yt, zt\right] \mathrm{d}t,$$

but Saran's convergence condition

$$\Re\left(x+y+z+2\sqrt{yz}\right) < 1,\ \Re(a_1) > 0$$

is wrong. Our result (4.7a) provides a sufficient condition

$$\Re(y) > 0,\ \Re(z) > 0,\ \Re\left(y+z+2\sqrt{yz}\right) < 1 + \min\left\{0, -\Re(x)\right\},\ \Re(a_1) > 0.$$

To clarify the general convergence condition, we are led to establish in later work the asymptotics of $\Psi_2[tx, ty]$ as $t \to +\infty$ with x, y in general case.

In addition, we made initial attempts on two asymptotic techniques we proposed, i.e., the uniformity approach and the separation method, both of which are currently deficient.

The uniformity approach is based on the series expansions involving the generalized hypergeometirc function $F_n(z)$ (see (6.1)), and the main difficulty is to explicitly deduce the uniform estimate (6.3). We have dealt with the simplest cases:

(i) Estimate on ${}_2F_2[a, b-n; c, d-n; -z]$: see [22, Theorem 2.4];

(ii) Estimate on ${}_2F_2[a, b-n; c, d; -z]$: see Theorem 4.4;

(iii) Estimate on ${}_2F_1[a, b-n; c; -z]$: briefly mentioned in Section 6.

Unfortunately, exponential expansions are implicit in the estimates for (i) and (ii), since nice integrals are missing for such ${}_2F_2$ functions. Please note Blaschke's conjecture [6, p. 1791]:

- (**More down conjecture**) *Denote $\mathscr{F}(\alpha, z)$ by*

$$\mathscr{F}(\alpha, z) := {}_pF_q\left[\begin{matrix} a_1 \pm \alpha, \cdots, a_k \pm \alpha, a_{k+1}, \cdots, a_p \\ b_1 \pm \alpha, \cdots, b_m \pm \alpha, b_{m+1}, \cdots, b_q \end{matrix}; z\right].$$

Regardless of the sign, when more large parameters are down than up, the resulting Taylor series defining $\mathscr{F}(\alpha, z)$ is always an asymptotic expansion for some values of z. That is, if $k < m$, then

$$\mathscr{F}(\alpha, z) \sim \sum_{n=0}^{\infty} \frac{(a_1 \pm \alpha)_n \cdots (a_k \pm \alpha)_n (a_{k+1})_n \cdots (a_p)_n}{(b_1 \pm \alpha)_n \cdots (b_m \pm \alpha)_n (b_{m+1})_n \cdots (b_q)_n} \frac{z^n}{n!}, \quad |\alpha| \to \infty.$$

The uniformity approach is indeed as difficult as the more down conjecture, since they both are somewhat ill-posed problems. It would be interesting to see whether the steepest descent method might become helpful for these problems.

The separation method is based on the Mellin convolution integrals. Another asymptotic technique for integrals of this type is the sum up and subtract (SUS) method introduced by López [30]. López and his coauthors [20] used the SUS method to derive the asymptotics of $F_2[x, y]$ for large x, whereas our separation method works as well. In addition, the SUS method fails for Ψ_1, but our separation method still works (see Proposition 6.1). These facts illustrate that the separation method is pretty powerful, though it is just heuristic. It would be desirable to make this method rigorous.

Remarkably, we could not obtain the asymptotics of $\Psi_1[x, y]$ (see [23]) and $\Psi_2[x, y]$ as $y \to \infty$ with arbitrary values of parameters, because we lack of useful expansions of ${}_2F_1$ and ${}_1F_1$, which are expanded in terms of large parameters. If such effective expansions of ${}_pF_q$ were known, one might use the Mellin-Barnes integral technique to establish the asymptotics of multiple hypergeometric functions for one large argument.

Acknowledgment

We are grateful to Petr Blaschke, Yury Brychkov and Nico Temme for their valuable advice during personal discussions with the first author. The second author was supported by the french ANR-PRME UNIOPEN (ANR-22-CE30-0004-01).

References

[1] T. Amdeberhan, O. Espinosa, I. Gonzalez, M. Harrison, V.H. Moll, A. Straub, *Ramanujan's master theorem*, Ramanujan J. **29** (1-3) (2012), pp. 103–120.

[2] L.U. Ancarani, J.A. Del Punta, G. Gasaneo, *Derivatives of Horn hypergeometric functions with respect to their parameters*, J. Math. Phys. **58** (7) (2017), 073504, 18 pages.

[3] B. Ananthanarayan, S. Bera, S. Froit, O. Marichev, T. Pathak, *On the evaluation of the Appell F_2 double hypergeometric function*, Comp. Phys. Commun. **284** (2023), 108589, 28 pages.

[4] G.E. Andrews, R. Askey, R. Roy, *Special Functions*, Cambridge Univ Press, (1999).

[5] A. Belafhal, H. Nebdi, *Generation and propagation of novel donut beams by a spiral phase plate: Humbert beams*, Opt. Quant. Electron. **46** (2014), pp. 201–208.

[6] P. Blaschke, *Asymptotic analysis via calculus of hypergeometric functions*, J. Math. Anal. Appl. **433** (2) (2016), pp. 1790–1820.

[7] Yu.A. Brychkov, O.I. Marichev, N.V. Savischenko, *Handbook of Mellin Transforms*, CRC Press, (2018).

[8] Yu.A. Brychkov, N. Saad, *On some formulas for the Appell function $F_2(a,b,b';c,c';w,z)$*, Integral Transforms Spec. Funct. **25** (2) (2014), pp. 111–123.

[9] Yu.A. Brychkov, N. Saad, *On some formulas for the Appell function $F_3(a,a',b,b';c;w,z)$*, Integral Transforms Spec. Funct. **26** (10) (2015), pp. 910–923.

[10] Yu.A. Brychkov, N.V. Savischenko, *On some formulas for the Horn function $H_2(a,b,c,c';d;w,z)$ and confluent Horn function $H_2^{(c)}(a,b,c;d;w,z)$*, Integral Transforms Spec. Funct. **32** (4) (2021), pp. 253–270.

[11] Yu.A. Brychkov, N.V. Savischenko, *On some formulas for the Horn functions $H_3(a,b;c;w,z)$, $H_6^{(c)}(a;c;w,z)$ and Humbert function $\Phi_3(b;c;w,z)$*, Integral Transforms Spec. Funct. **32** (9) (2021), pp. 661–676.

[12] Yu.A. Brychkov, N.V. Savischenko, *On some formulas for the Horn functions $H_5(a,b;c;w,z)$ and $H_5^{(c)}(a;c;w,z)$*, Integral Transforms Spec. Funct. **33** (5) (2022), pp. 373–387.

[13] Yu.A. Brychkov, N.V. Savischenko, *On some formulas for the confluent Horn functions $H_3^{(c)}(a,b,c;d;w,z)$, $H_4^{(c)}(a,c;d;w,z)$ and $H_9^{(c)}(a,b;c;w,z)$*, Integral Transforms Spec. Funct. **34** (4) (2022), pp. 275–294.

[14] S. Chib, F. Khannous, A. Belafhal, *Generation of donut Humbert beam of type-II*, Opt. Quant. Electron. **55** (2023), 936, 11 pages.

[15] J. Carvalho e Silva, H.M. Srivastava, *Asymptotic representations for a class of multiple hypergeometric functions with a dominant variable*, Integral Transforms Spec. Funct. **11** (2) (2001), pp. 137–150.

[16] A. Erdélyi, *Integration of a certain system of linear partial differential equations of hypergeometric type*, Proc. Roy. Soc. Edinburgh Sect. A **59** (1939), pp. 224–241.

[17] A. Erdélyi, *Some confluent hypergeometric functions of two variables*, Proc. Roy. Soc. Edinburgh Sect. A **60** (1940), pp. 344–361.

[18] G. Gasaneo, F.D. Colavecchia, S. Otranto, C.R. Garibotti, *The hyperbolic-hypergeometric functions*, J. Math. Phys. **42** (10) (2001), pp. 4971–4983.

[19] G. Gasaneo, F.D. Colavecchia, C.R. Garibotti, J.E. Miraglia, P. Macri, *Correlated continuum wave functions for three particles with Coulomb interactions*, Phys. Rev. A **55** (1997), pp. 2809–2820.

[20] C. Ferreira, J.L. López, E.P. Sinusía, *The second Appell function for one large variable*, Mediterr. J. Math. **10** (4) (2013), pp. 1853–1865.

[21] P.-C. Hang, M.-J. Luo, *Asymptotics of the Humbert function Ψ_1 for two large arguments*, SIGMA **20** (2024), 074, 13 pages.

[22] P.-C. Hang, M.-J. Luo, *Note on the Humbert function Ψ_1*, arXiv: 2410.21985v2.

[23] P.-C. Hang, M.-J. Luo, *Asymptotics of Saran's hypergeometric function F_K*, J. Math. Anal. Appl. **541** (2) (2025), 128707, 19 pages.

[24] M. Henkel, *Finite-size scaling in the ageing dynamics of the 1D Glauber-Ising model*, submitted to Entropy, arXiv: 2501.05912.

[25] P. Humbert, *IX.–The confluent hypergeometric functions of two variables*, Proc. R. Soc. Edinb. **41** (1922), pp. 73–96.

[26] J.C. Jaeger, *A continuation formula for Appell's function F_2*, J. Lond. Math. Soc. **13** (1938), p. 254.

[27] C.M. Joshi, J.P. Arya, *Inequalities for certain confluent hypergeometric functions of two variables*, Indian J. Pure Appl. Math. **13** (4) (1982), pp. 491–500.

[28] R. Kumar, *The generalized modified Bessel function and its connection with Voigt line profile and Humbert functions*, Adv. Appl. Math. **114** (2020), 101986, 19 pages.

[29] Y. Lin, R. Wong, *Asymptotics of generalized hypergeometric functions*, Frontiers in Orthogonal Polynomials and q-Series (2018), pp. 497–521.

[30] J.L. López, *Asymptotic expansions of Mellin convolution integrals*, SIAM Rev. **50** (2) (2008), pp. 275–293.

[31] T. Miyazaki, *On Fourier-Jacobi expansions of real analytic Eisenstein series of degree 2*, Abh. Math. Semin. Univ. Hamb. **84** (1) (2014), pp. 85–122.

[32] S. Mukai, *Confluences of Appell's F_2 system of hypergeometric differential equations*, Kyushu J. Math. **77** (2023), pp. 1–42.

[33] F.W.J. Olver, *Why steepest descents?*, SIAM Rev. **12** (2) (1970), pp. 228–247.

[34] F.W.J. Olver, A.B. Olde Daalhuis, D.W. Lozier, B.I. Schneider, R.F. Boisvert, C.W. Clark, B.R. Miller, B.V. Saunders, H.S. Cohl, M.A. McClain (Editors), *NIST Digital Library of Mathematical Functions*, Release 1.2.3 of 2024-12-15, aviable at https://dlmf.nist.gov/.

[35] A.P. Prudnikov, Yu.A. Brychkov, O.I. Marichev, *Integrals and Series, Vol. 3: More Special Functions*, Gordon and Breach Science Publishers, New York, (1990).

[36] L.E. Sánchez, D.K. Nagar, *Non-central matrix-variate Dirichlet distribution*, Taiwanese J. Math. **7** (3) (2003), pp. 477–491.

[37] S. Saran, *Integral representations of Laplace type for certain hypergeometric functions of three variables*, Riv., Mat. Univ. Parma **8** (1957), pp. 133–143.

[38] S. Shimomura, *Connection formulae for solutions of a system of partial differential equations associated with the confluent hypergeometric function* Φ_2, Proc. Japan Acad. **72** (1996), pp. 82–84.

[39] H.M. Srivastava, P.W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Ellis Horwood Ltd., Chichester, (1985).

[40] H.M. Srivastava, E.A. Miller, *A unified presentation of the Voigt functions*, Astrophys. Space Sci. **135** (1987), pp. 111–118.

[41] V.K. Tuan, M. Saigo, T.D. Dinh, *Some integral transforms in the space of entire functions of exponential type*, Proc. Japan Acad., Ser. A **72** (4) (1996), pp. 75–77.

[42] S. Wald, M. Henkel, *On integral representations and asymptotics of some hypergeometric functions in two variables*, Integral Transforms Spec. Funct. **29** (2) (2018), pp. 95–112.