arXiv:math/0602202v1 [math.CA] 9 Feb 2006

Asymptotics of Recurrence Relation Coefficients, Hankel Determinant Ratios, and Root Products Associated with Laurent Polynomials Orthogonal with Respect to Varying Exponential Weights

K. T.-R. McLaughlin*
Department of Mathematics
The University of Arizona
617 N. Santa Rita Ave.
P. O. Box 210089
Tucson, Arizona 85721–0089
U. S. A.

A. H. Vartanian†
Department of Mathematics
University of Central Florida
P. O. Box 161364
Orlando, Florida 32816–1364
U. S. A.

X. Zhou‡
Department of Mathematics
Duke University
Box 90320
Durham, North Carolina 27708–0320
U. S. A.

8 February 2006

Abstract

Let $\Lambda^{\mathbb{R}}$ denote the linear space over $\mathbb{R}$ spanned by z^k, $k \in \mathbb{Z}$. Define the real inner product (with varying exponential weights) $\langle \cdot, \cdot \rangle_{\mathcal{L}} : \Lambda^{\mathbb{R}} \times \Lambda^{\mathbb{R}} \to \mathbb{R}$, $(f,g) \mapsto \int_{\mathbb{R}} f(s)g(s)\exp(-\mathcal{N}V(s))\,\mathrm{d}s$, $\mathcal{N} \in \mathbb{N}$, where the external field V satisfies: (i) V is real analytic on $\mathbb{R}\setminus\{0\}$; (ii) $\lim_{|x|\to\infty}(V(x)/\ln(x^2+1)) = +\infty$; and (iii) $\lim_{|x|\to 0}(V(x)/\ln(x^{-2}+1)) = +\infty$. Orthogonalisation of the (ordered) base $\{1, z^{-1}, z, z^{-2}, z^2, \ldots, z^{-k}, z^k, \ldots\}$ with respect to $\langle \cdot, \cdot \rangle_{\mathcal{L}}$ yields the even degree and odd degree orthonormal Laurent polynomials (OLPs) $\{\phi_m(z)\}_{m=0}^{\infty}$: $\phi_{2n}(z) = \sum_{k=-n}^{n} \xi_k^{(2n)} z^k$, $\xi_n^{(2n)} > 0$, and $\phi_{2n+1}(z) = \sum_{k=-n-1}^{n} \xi_k^{(2n+1)} z^k$, $\xi_{-n-1}^{(2n+1)} > 0$. Associated with the even degree and odd degree OLPs are the following two pairs of recurrence relations: $z\phi_{2n}(z) = c_{2n}^{\sharp}\phi_{2n-2}(z) + b_{2n}^{\sharp}\phi_{2n-1}(z) + a_{2n}^{\sharp}\phi_{2n}(z) + b_{2n+1}^{\sharp}\phi_{2n+1}(z) + c_{2n+2}^{\sharp}\phi_{2n+2}(z)$ and $z\phi_{2n+1}(z) = b_{2n+1}^{\sharp}\phi_{2n}(z) + a_{2n+1}^{\sharp}\phi_{2n+1}(z) + b_{2n+2}^{\sharp}\phi_{2n+2}(z)$, where $c_0^{\sharp} = b_0^{\sharp} = 0$, and $c_{2k}^{\sharp} > 0$, $k \in \mathbb{N}$, and $z^{-1}\phi_{2n+1}(z) = \gamma_{2n+1}^{\sharp}\phi_{2n-1}(z) + \beta_{2n+1}^{\sharp}\phi_{2n}(z) + \alpha_{2n+1}^{\sharp}\phi_{2n+1}(z) + \beta_{2n+2}^{\sharp}\phi_{2n+2}(z) + \gamma_{2n+3}^{\sharp}\phi_{2n+3}(z)$ and $z^{-1}\phi_{2n}(z) = \beta_{2n}^{\sharp}\phi_{2n-1}(z) + \alpha_{2n}^{\sharp}\phi_{2n}(z) + \beta_{2n+1}^{\sharp}\phi_{2n+1}(z)$, where $\beta_0^{\sharp} = \gamma_1^{\sharp} = 0$, $\beta_1^{\sharp} > 0$, and $\gamma_{2l+1}^{\sharp} > 0$, $l \in \mathbb{N}$. Asymptotics in the double-scaling limit as $\mathcal{N}, n \to \infty$ such that $\mathcal{N}/n = 1 + o(1)$ of the coefficients of these two pairs of recurrence relations, Hankel determinant ratios associated with the real-valued, bi-infinite, strong moment sequence $\left\{c_k = \int_{\mathbb{R}} s^k \exp(-\mathcal{N}V(s))\,\mathrm{d}s\right\}_{k\in\mathbb{Z}}$, and the products of the (real) roots of the OLPs are obtained by formulating the even degree and odd degree OLP problems as matrix Riemann-Hilbert problems on $\mathbb{R}$, and then extracting the large-n behaviours by applying the non-linear steepest-descent method introduced in [1] and further developed in [2, 3].

*E-mail: mcl@math.arizona.edu

†E-mail: arthurv@math.ucf.edu

‡E-mail: zhou@math.duke.edu

2000 Mathematics Subject Classification. (Primary) 42C05, 30E25, 30E05, 47B36: (Secondary) 30C70, 30C15, 41A60, 31A99

Abbreviated Title. Asymptotics of Recurrence Coefficients, Hankel Determinant Ratios, and Root Products

Key Words. Asymptotics, equilibrium measures, Hankel Determinants, Laurent-Jacobi matrices, orthogonal Laurent polynomials, recurrence relations, Riemann-Hilbert problems, strong moment problems, variational problems

1 Introduction and Background

The *strong Stieltjes moment problem* (SSMP) considers the relation between a given doubly- or bi-infinite (moment) sequence $\{c_n\}_{n\in\mathbb{Z}}$ of real numbers and associated measures. There are, in fact, three general types of problems in this area:

(i) find necessary and sufficient conditions for the existence of a non-negative Borel measure $\mu := \mu_{\mathrm{MP}}^{\mathrm{SS}}$ on $[0, +\infty)$, and with infinite support, such that $c_n^{\mathrm{SSMP}} = \int_0^{+\infty} \lambda^n \, \mathrm{d}\mu_{\mathrm{MP}}^{\mathrm{SS}}(\lambda)$, $n \in \mathbb{Z}$, where the (improper) integral is to be understood in the Riemann-Stieltjes sense;

(ii) when there is a solution, in which case the SSMP is *determinate*, find conditions for the uniqueness of the solution; and

(iii) when there is more than one solution, in which case the SSMP is *indeterminate*, describe the family of all solutions.

Similarly, when the support of the non-negative Borel measure $\mu := \mu_{\mathrm{MP}}^{\mathrm{SH}}$ is only required to be contained in $(-\infty, +\infty)$, the analogous problem is referred to as the *strong Hamburger moment problem* (SHMP). (In this latter case, $c_n^{\mathrm{SHMP}} = \int_{-\infty}^{+\infty} \lambda^n \, \mathrm{d}\mu_{\mathrm{MP}}^{\mathrm{SH}}(\lambda)$, $n \in \mathbb{Z}$.) The SSMP (resp., SHMP) was introduced in 1980 (resp., 1981) by Jones *et al.* [4] (resp., Jones *et al.* [5]), and studied further in [6–10] (see, also, [11]). Unlike the moment theory for the *classical Stieltjes* (resp., *classical Hamburger*) *moment problem* (SMP) [12] (resp., (HMP) [13]), wherein the theory of orthogonal polynomials [14] played a prominent rôle (see, for example, [15]), the extension of the moment theory to the SSMP and the SHMP introduced a 'rational generalisation' of the orthogonal polynomials, namely, the *orthogonal Laurent* (or *L*-) *polynomials* [4–11, 16–20].

Orthogonal *L*-polynomials are defined in terms of certain orthogonality relations involving an associated measure μ. In this work, measures that satisfy the following conditions are considered:

⊛ $\widetilde{\mu} \in \mathcal{M}_1(\mathbb{R})$, where

$$\mathcal{M}_1(\mathbb{R}) := \left\{ \mu; \int_{\mathbb{R}} \mathrm{d}\mu(s) = 1, \int_{\mathbb{R}} s^k \mathrm{d}\mu(s) < \infty, k \in \mathbb{Z} \setminus \{0\} \right\};$$

⊛ $\mathrm{d}\widetilde{\mu}(z) = \widetilde{w}(z) \, \mathrm{d}z$, where

$$\widetilde{w}(z) := \exp(-\mathcal{N} V(z)), \quad \mathcal{N} \in \mathbb{N}.$$

Here, $\widetilde{w}(z)$ is referred to as the varying exponential weight function, and the function $V : \mathbb{R} \setminus \{0\} \to \mathbb{R}$, which is referred to as the *external field*, satisfies:

$$V \text{ is real analytic on } \mathbb{R} \setminus \{0\}; \tag{V1}$$

$$\lim_{|x|\to\infty} \left(V(x) / \ln(x^2+1) \right) = +\infty; \tag{V2}$$

$$\lim_{|x|\to 0} \left(V(x) / \ln(x^{-2}+1) \right) = +\infty. \tag{V3}$$

(For example, a rational function of the form $V(z) = \sum_{k=-2m_1}^{2m_2} \varrho_k z^k$, with $\varrho_k \in \mathbb{R}$, $k = -2m_1, \ldots, 2m_2$, $m_{1,2} \in \mathbb{N}$, and $\varrho_{-2m_1}, \varrho_{2m_2} > 0$ would satisfy conditions (V1)–(V3).)

Rational functions with poles at 0 and at ∞ are seminal for obtaining the results of this work: they are now discussed. It will be important to have the ordered sequence of (complex) monomials $\{1, z^{-1}, z, z^{-2}, z^2, \ldots, z^{-k}, z^k, \ldots\}$ corresponding to the *cyclically-repeated pole sequence* {no pole, 0, ∞, 0, ∞, ..., 0, ∞, ...}. This is a basis for the set of functions $\Lambda^{\mathbb{R}} = \cup_{m\in\mathbb{Z}_0^+} (\Lambda_{2m}^{\mathbb{R}} \cup \Lambda_{2m+1}^{\mathbb{R}})$, with $\mathbb{Z}_0^+ := \{0\} \cup \mathbb{N}$, where $\Lambda_{2m}^{\mathbb{R}} := \Lambda_{-m,m}^{\mathbb{R}}$, and $\Lambda_{2m+1}^{\mathbb{R}} := \Lambda_{-m-1,m}^{\mathbb{R}}$, where, for any pair $(p, q) \in \mathbb{Z} \times \mathbb{Z}$, with $p \leqslant q$, $\Lambda_{p,q}^{\mathbb{R}} := \left\{ f : \mathbb{C}^* \to \mathbb{C};\ f(z) = \sum_{k=p}^{q} \widetilde{\lambda}_k z^k,\ \widetilde{\lambda}_k \in \mathbb{R},\ k = p, \ldots, q \right\}$, where $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$. For each $l \in \mathbb{Z}_0^+$ and $0 \not\equiv f \in \Lambda_l^{\mathbb{R}}$, the *L-degree* of f, symbolically $LD(f)$, is defined as

$$LD(f) := l.$$

The *leading coefficient* of $0 \not\equiv f \in \Lambda_l^{\mathbb{R}}$, symbolically $LC(f)$, is defined as

$$LC(f) := \begin{cases} \widetilde{\lambda}_m, & l = 2m, \\ \widetilde{\lambda}_{-m-1}, & l = 2m+1, \end{cases}$$

and the *trailing coefficient* of $0 \not\equiv f \in \Lambda_l^{\mathbb{R}}$, symbolically $TC(f)$, is defined as

$$TC(f) := \begin{cases} \widetilde{\lambda}_{-m}, & l = 2m, \\ \widetilde{\lambda}_m, & l = 2m+1. \end{cases}$$

Thus, for $0 \not\equiv f \in \Lambda_{2m}^{\mathbb{R}}$, one writes $f(z) = TC(f)z^{-m} + \cdots + LC(f)z^m$, and, for $0 \not\equiv f \in \Lambda_{2m+1}^{\mathbb{R}}$, one writes $f(z) = LC(f)z^{-m-1} + \cdots + TC(f)z^m$. For each $l \in \mathbb{Z}_0^+$, $0 \not\equiv f \in \Lambda_l^{\mathbb{R}}$ is called *monic* if $LC(f) = 1$.

Define (uniquely) the *strong moment linear functional* $\mathcal{L}$ by its action on the basis elements of $\Lambda^{\mathbb{R}}$ as follows: $\mathcal{L}\colon \Lambda^{\mathbb{R}} \to \Lambda^{\mathbb{R}}$, $f = \sum_{k\in\mathbb{Z}} \widetilde{\lambda}_k z^k \mapsto \mathcal{L}(f) := \sum_{k\in\mathbb{Z}} \widetilde{\lambda}_k c_k$, where $c_k = \mathcal{L}(z^k) = \int_{\mathbb{R}} s^k \mathrm{e}^{-\mathcal{N}V(s)}\, \mathrm{d}s$, $(k, \mathcal{N}) \in \mathbb{Z}\times\mathbb{N}$. (Note that $\{c_k\}_{k\in\mathbb{Z}}$ is a bi-infinite, real-valued, *strong moment sequence*: c_k is called the *kth strong moment of* $\mathcal{L}$.) Associated with the above-defined bi-infinite, real-valued, strong moment sequence $\{c_k\}_{k\in\mathbb{Z}}$ are the *Hankel determinants* $H_k^{(m)}$, $(m,k) \in \mathbb{Z}\times\mathbb{N}$ (see, for example, [11]):

$$H_0^{(m)} := 1 \qquad \text{and} \qquad H_k^{(m)} := \begin{vmatrix} c_m & c_{m+1} & \cdots & c_{m+k-2} & c_{m+k-1} \\ c_{m+1} & c_{m+2} & \cdots & c_{m+k-1} & c_{m+k} \\ c_{m+2} & c_{m+3} & \cdots & c_{m+k} & c_{m+k+1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ c_{m+k-1} & c_{m+k} & \cdots & c_{m+2k-3} & c_{m+2k-2} \end{vmatrix}. \tag{1.1}$$

Define the bilinear form (with varying exponential weight) $\langle \cdot, \cdot \rangle_{\mathcal{L}}$ as follows: $\langle \cdot, \cdot \rangle_{\mathcal{L}} : \Lambda^{\mathbb{R}} \times \Lambda^{\mathbb{R}} \to \mathbb{R}$, $(f, g) \mapsto \langle f, g \rangle_{\mathcal{L}} := \mathcal{L}(f(z)g(z)) = \int_{\mathbb{R}} f(s)g(s)\exp(-\mathcal{N}V(s))\, \mathrm{d}s$, $\mathcal{N} \in \mathbb{N}$. It is a fact [6, 7, 11, 17] (see, also, the proofs of Lemmas 2.2.1 and 2.2.2 in [21] and [22], respectively) that the bilinear form $\langle \cdot, \cdot \rangle_{\mathcal{L}}$ thus defined is an inner product if and only if $H_{2m}^{(-2m)} > 0$ and $H_{2m+1}^{(-2m)} > 0$ for each $m \in \mathbb{Z}_0^+$ (see Equations (1.8) below).

Remark 1.1. These latter two (Hankel determinant) inequalities also appear when the question of the solvability of the SHMP is posed (in this case, the c_k, $k \in \mathbb{Z}$, which appear in Equations (1.1) should be replaced by c_k^{SHMP}, $k \in \mathbb{Z}$); indeed, if these two inequalities are true $\forall\ m \in \mathbb{Z}_0^+$, then there is a non-negative measure $\mu_{\mathrm{MP}}^{\mathrm{SH}}$ (on $\mathbb{R}$) with the given (real) moments. For the case of the SSMP, there are four (Hankel determinant) inequalities (in this latter case, the c_k, $k \in \mathbb{Z}$, which appear in Equations (1.1) should be replaced by c_k^{SSMP}, $k \in \mathbb{Z}$) which guarantee the existence of a non-negative measure $\mu_{\mathrm{MP}}^{\mathrm{SS}}$ (on $[0, +\infty)$) with the given moments, namely [4] (see, also, [6, 7]): for each $m \in \mathbb{Z}_0^+$, $H_{2m}^{(-2m)} > 0$, $H_{2m+1}^{(-2m)} > 0$, $H_{2m}^{(-2m+1)} > 0$, and $H_{2m+1}^{(-2m-1)} < 0$. It is interesting to note that the former solvability conditions do not automatically imply that the positive (real) moments $\{c_k^{\mathrm{SHMP}}\}_{k\in\mathbb{Z}_0^+}$ determine a measure via the HMP: a similar statement holds true for the SMP (see the latter four solvability conditions). ■

The *norm of* $f \in \Lambda^{\mathbb{R}}$ *with respect to* $\mathcal{L}$ is now defined:

$$\|f(\cdot)\|_{\mathcal{L}} := (\langle f, f \rangle_{\mathcal{L}})^{1/2}.$$

Given an inner product $\langle \cdot, \cdot \rangle_{\mathcal{L}}$, one may define the (real) orthonormal Laurent polynomial sequence $\{\phi_n^{\natural}(z)\}_{n\in\mathbb{Z}_0^+}$ with respect to $\mathcal{L}$ as follows; $\forall\ m, n \in \mathbb{Z}_0^+$:

(i) $\phi_n^{\natural}(z) \in \Lambda_n^{\mathbb{R}}$, that is, $LD(\phi_n^{\natural}) = n$;

(ii) $\langle \phi_m^{\natural}, \phi_{m'}^{\natural} \rangle_{\mathcal{L}} = 0$, $m \neq m'$, or, alternatively, $\langle f, \phi_m^{\natural} \rangle_{\mathcal{L}} = 0\ \forall\ f \in \Lambda_{m-1}^{\mathbb{R}}$;

(iii) $\langle \phi_m^{\natural}, \phi_m^{\natural} \rangle_{\mathcal{L}} =: \|\phi_m^{\natural}(\cdot)\|_{\mathcal{L}}^2 = 1$.

Hereafter, these shall be referred to as orthonormal L-polynomials. Orthonormal L-polynomials may be obtained by Gram-Schmidt orthogonalisation. To do this, however, one must have an ordered sequence of complex monomials, z^j, $j \in \mathbb{Z}$, and the orthonormal L-polynomials, themselves, will depend on how this sequence of complex monomials is ordered. In this work, orthonormalisation with respect to $\langle \cdot, \cdot \rangle_{\mathcal{L}}$ of the base sequence of $\Lambda^{\mathbb{R}}$ is considered, that is, $\{1, z^{-1}, z, z^{-2}, z^2, \ldots, z^{-k}, z^k, \ldots\}$. The result of this procedure is the orthonormal L-polynomials $\{\phi_m(z)\}_{m\in\mathbb{Z}_0^+}$:

- for $m = 2n$,

$$\phi_{2n}(z) = \sum_{k=-n}^{n} \xi_k^{(2n)} z^k, \qquad \xi_n^{(2n)} > 0; \tag{1.2}$$

- for $m=2n+1$,

$$\phi_{2n+1}(z)=\sum_{k=-n-1}^{n}\xi_k^{(2n+1)}z^k,\qquad \xi_{-n-1}^{(2n+1)}>0. \tag{1.3}$$

The ϕ_n's are normalised so that they all have real coefficients; in particular, the leading coefficients, $LC(\phi_{2n})=\xi_n^{(2n)}$ and $LC(\phi_{2n+1})=\xi_{-n-1}^{(2n+1)}$, $n\in\mathbb{Z}_0^+$, are both positive, $\xi_0^{(0)}=1$, and $\phi_0(z)\equiv 1$. Even though the leading coefficients, $\xi_n^{(2n)}$ and $\xi_{-n-1}^{(2n+1)}$, $n\in\mathbb{Z}_0^+$, are non-zero (in particular, they are positive), no such restriction applies to the trailing coefficients, $TC(\phi_{2n})=\xi_{-n}^{(2n)}$ and $TC(\phi_{2n+1})=\xi_n^{(2n+1)}$, $n\in\mathbb{Z}_0^+$ (see below). Furthermore, note that, by construction:

(i) $\langle\phi_{2n},z^j\rangle_{\mathcal{L}}=0$, $j=-n,\dots,n-1$;
(ii) $\langle\phi_{2n+1},z^j\rangle_{\mathcal{L}}=0$, $j=-n,\dots,n$;
(iii) $\langle\phi_j,\phi_k\rangle_{\mathcal{L}}=\delta_{jk}$, $j,k\in\mathbb{Z}_0^+$, where δ_{jk} is the Kronecker delta.

It is convenient to introduce the monic orthogonal Laurent (or L-) polynomials, $\boldsymbol{\pi}_j(z)$, $j\in\mathbb{Z}_0^+$: (i) for $j=2n$, with $\boldsymbol{\pi}_0(z)\equiv 1$,

$$\boldsymbol{\pi}_{2n}(z):=(\xi_n^{(2n)})^{-1}\phi_{2n}(z)=\nu_{-n}^{(2n)}z^{-n}+\cdots+z^n,\qquad \nu_{-n}^{(2n)}:=\xi_{-n}^{(2n)}/\xi_n^{(2n)}; \tag{1.4}$$

and (ii) for $j=2n+1$,

$$\boldsymbol{\pi}_{2n+1}(z):=(\xi_{-n-1}^{(2n+1)})^{-1}\phi_{2n+1}(z)=z^{-n-1}+\cdots+\nu_n^{(2n+1)}z^n,\qquad \nu_n^{(2n+1)}:=\xi_n^{(2n+1)}/\xi_{-n-1}^{(2n+1)}. \tag{1.5}$$

The monic orthogonal L-polynomials, $\boldsymbol{\pi}_j(z)$, $j\in\mathbb{Z}_0^+$, satisfy the following orthogonality and normalisation conditions:

(1) $\langle\boldsymbol{\pi}_{2n},z^j\rangle_{\mathcal{L}}=0$, $j=-n,\dots,n-1$;
(2) $\langle\boldsymbol{\pi}_{2n+1},z^j\rangle_{\mathcal{L}}=0$, $j=-n,\dots,n$;
(3) $\langle\boldsymbol{\pi}_{2n},\boldsymbol{\pi}_{2n}\rangle_{\mathcal{L}}=:\|\boldsymbol{\pi}_{2n}(\cdot)\|_{\mathcal{L}}^2=(\xi_n^{(2n)})^{-2}$;
(4) $\langle\boldsymbol{\pi}_{2n+1},\boldsymbol{\pi}_{2n+1}\rangle_{\mathcal{L}}=:\|\boldsymbol{\pi}_{2n+1}(\cdot)\|_{\mathcal{L}}^2=(\xi_{-n-1}^{(2n+1)})^{-2}$.

Conditions (3) and (4) above imply, respectively, that $\xi_n^{(2n)}=1/\|\boldsymbol{\pi}_{2n}(\cdot)\|_{\mathcal{L}}$ (>0) and $\xi_{-n-1}^{(2n+1)}=1/\|\boldsymbol{\pi}_{2n+1}(\cdot)\|_{\mathcal{L}}$ (>0).

In terms of the Hankel determinants, $H_k^{(m)}$, $(m,k)\in\mathbb{Z}\times\mathbb{N}$, associated with the real-valued, bi-infinite, strong moment sequence $\left\{c_k=\int_{\mathbb{R}}s^k\exp(-\mathcal{N}V(s))\,ds,\ \mathcal{N}\in\mathbb{N}\right\}_{k\in\mathbb{Z}}$, the monic orthogonal L-polynomials, $\boldsymbol{\pi}_j(z)$, $j\in\mathbb{Z}_0^+$, are represented via the following determinantal formulae [6,7,11,17]: for $m\in\mathbb{Z}_0^+$,

$$\boldsymbol{\pi}_{2m}(z)=\frac{1}{H_{2m}^{(-2m)}}\begin{vmatrix} c_{-2m} & c_{-2m+1} & \cdots & c_{-1} & z^{-m}\\ c_{-2m+1} & c_{-2m+2} & \cdots & c_0 & z^{-m+1}\\ \vdots & \vdots & \ddots & \vdots & \vdots\\ c_{-1} & c_0 & \cdots & c_{2m-2} & z^{m-1}\\ c_0 & c_1 & \cdots & c_{2m-1} & z^m\end{vmatrix}, \tag{1.6}$$

and

$$\boldsymbol{\pi}_{2m+1}(z)=-\frac{1}{H_{2m+1}^{(-2m)}}\begin{vmatrix} c_{-2m-1} & c_{-2m} & \cdots & c_{-1} & z^{-m-1}\\ c_{-2m} & c_{-2m+1} & \cdots & c_0 & z^{-m}\\ \vdots & \vdots & \ddots & \vdots & \vdots\\ c_{-1} & c_0 & \cdots & c_{2m-1} & z^{m-1}\\ c_0 & c_1 & \cdots & c_{2m} & z^m\end{vmatrix}; \tag{1.7}$$

moreover, it can be shown that (see, for example, [11,17]), for $n\in\mathbb{Z}_0^+$,

$$\xi_n^{(2n)}\left(=\frac{1}{\|\boldsymbol{\pi}_{2n}(\cdot)\|_{\mathcal{L}}}\right)=\sqrt{\frac{H_{2n}^{(-2n)}}{H_{2n+1}^{(-2n)}}},\qquad \xi_{-n-1}^{(2n+1)}\left(=\frac{1}{\|\boldsymbol{\pi}_{2n+1}(\cdot)\|_{\mathcal{L}}}\right)=\sqrt{\frac{H_{2n+1}^{(-2n)}}{H_{2n+2}^{(-2n-2)}}}, \tag{1.8}$$

$$\nu_{-n}^{(2n)}\left(:=\frac{\xi_{-n}^{(2n)}}{\xi_n^{(2n)}}\right)=\frac{H_{2n}^{(-2n+1)}}{H_{2n}^{(-2n)}},\qquad \nu_n^{(2n+1)}\left(:=\frac{\xi_n^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}}\right)=-\frac{H_{2n+1}^{(-2n-1)}}{H_{2n+1}^{(-2n)}}. \tag{1.9}$$

Remark 1.2. It is important to note [10] that there is a more general theory, where moments corresponding to an arbitrary, countable sequence of (fixed) points are involved, and where *orthogonal rational functions* [23] (see, also, [24]) play the rôle of orthogonal L-polynomials. Furthermore, since orthogonal L-polynomials are rational functions with (fixed) poles at the origin and at the point at infinity, the step towards a more general theory where poles are at arbitrary, but fixed, positions/locations in $\mathbb{C}\cup\{\infty\}$ is natural. ■

A basic fact about orthogonal L-polynomials is [6, 7, 17]:

⊛ for each $m\in\mathbb{Z}_0^+$, all the roots of $\boldsymbol{\pi}_{2m}(z)$ and $\boldsymbol{\pi}_{2m+1}(z)$ are real, simple, and non-zero.

In the literature (see, for example, [6, 7, 17]), there are two distinguished cases, depending on whether or not the trailing coefficients vanish. Recall that, for each $m\in\mathbb{Z}_0^+$, the trailing coefficient, $TC(\boldsymbol{\pi}_m)$, is defined as follows: $TC(\boldsymbol{\pi}_m):=\begin{cases}\nu_{-n}^{(2n)}, & m=2n,\\ \nu_n^{(2n+1)}, & m=2n+1.\end{cases}$ If, for each $m\in\mathbb{Z}_0^+$, $TC(\boldsymbol{\pi}_m)\neq 0$, then $\boldsymbol{\pi}_m(z)$ and the index m are called *non-singular*; otherwise, if $TC(\boldsymbol{\pi}_m)=0$, then $\boldsymbol{\pi}_m(z)$ and the index m are called *singular*. From Equations (1.9), it can be seen that, for each $m\in\mathbb{Z}_0^+$:

(i) $\boldsymbol{\pi}_{2m}(z)$ is non-singular (resp., singular) if $H_{2m}^{(-2m+1)}\neq 0$ (resp., $H_{2m}^{(-2m+1)}=0$);

(ii) $\boldsymbol{\pi}_{2m+1}(z)$ is non-singular (resp., singular) if $H_{2m+1}^{(-2m-1)}\neq 0$ (resp., $H_{2m+1}^{(-2m-1)}=0$).

It is an established fact [6, 7, 17] that, for $m\in\mathbb{Z}_0^+$:

(1) if $\boldsymbol{\pi}_{2m}(z)$ is non-singular, then it possesses $2m$ roots and $\boldsymbol{\pi}_{2m}(0)\neq 0$, and if $\boldsymbol{\pi}_{2m}(z)$ is singular, then it possesses $2m-1$ roots;

(2) if $\boldsymbol{\pi}_{2m+1}(z)$ is non-singular, then it possesses $2m+1$ roots and $\boldsymbol{\pi}_{2m+1}(0)\neq 0$, and if $\boldsymbol{\pi}_{2m+1}(z)$ is singular, then it possesses $2m$ roots.

For each $n\in\mathbb{Z}_0^+$, it can be shown that, via a straightforward factorisation argument and using Equations (1.6) and (1.7):

(i) if $\boldsymbol{\pi}_{2n}(z)$ is non-singular, setting $\left\{\alpha_k^{(2n)},\ k=1,\dots,2n\right\}:=\{z;\ \boldsymbol{\pi}_{2n}(z)=0\}$,

$$\prod_{k=1}^{2n}\alpha_k^{(2n)}=\nu_{-n}^{(2n)}\qquad(\in\mathbb{R});\tag{1.10}$$

(ii) if $\boldsymbol{\pi}_{2n}(z)$ is singular, setting (with abuse of notation) $\left\{\alpha_k^{(2n)},\ k=1,\dots,2n-1\right\}:=\{z;\ \boldsymbol{\pi}_{2n}(z)=0\}$,

$$\prod_{k=1}^{2n-1}\alpha_k^{(2n)}=-\nu_{-(n-1)}^{(2n)}:=-\xi_{-(n-1)}^{(2n)}(\xi_n^{(2n)})^{-1}\qquad(\in\mathbb{R});\tag{1.11}$$

(iii) if $\boldsymbol{\pi}_{2n+1}(z)$ is non-singular, setting $\left\{\alpha_k^{(2n+1)},\ k=1,\dots,2n+1\right\}:=\{z;\ \boldsymbol{\pi}_{2n+1}(z)=0\}$,

$$\left(\prod_{k=1}^{2n+1}\alpha_k^{(2n+1)}\right)^{-1}=-\nu_n^{(2n+1)}\qquad(\in\mathbb{R});\tag{1.12}$$

(iv) if $\boldsymbol{\pi}_{2n+1}(z)$ is singular, setting (with abuse of notation) $\left\{\alpha_k^{(2n+1)},\ k=1,\dots,2n\right\}:=\{z;\ \boldsymbol{\pi}_{2n+1}(z)=0\}$,

$$\left(\prod_{k=1}^{2n}\alpha_k^{(2n+1)}\right)^{-1}=\nu_{n-1}^{(2n+1)}:=\xi_{n-1}^{(2n+1)}(\xi_{-n-1}^{(2n+1)})^{-1}\qquad(\in\mathbb{R}).\tag{1.13}$$

Via Equations (1.8) and (1.9), and using the fact that $\xi_0^{(0)}=H_0^{(0)}=H_1^{(0)}=1$ (cf. Equations (1.1)), a straightforward calculation shows that, for $n\in\mathbb{Z}_0^+$,

$$H_{2n}^{(-2n)}=\frac{(\xi_n^{(2n)})^2(\xi_{-n-1}^{(2n+1)})^2}{\prod_{j=0}^{n}(\xi_j^{(2j)})^2\prod_{k=0}^{n}(\xi_{-k-1}^{(2k+1)})^2}\qquad(>0),\tag{1.14}$$

$$H_{2n+1}^{(-2n)}=\frac{(\xi_{-n-1}^{(2n+1)})^2}{\prod_{j=0}^{n}(\xi_j^{(2j)})^2\prod_{k=0}^{n}(\xi_{-k-1}^{(2k+1)})^2}\qquad(>0),\tag{1.15}$$

$$H_{2n}^{(-2n+1)} = \frac{\nu_{-n}^{(2n)}(\xi_n^{(2n)})^2(\xi_{-n-1}^{(2n+1)})^2}{\prod_{j=0}^{n}(\xi_j^{(2j)})^2 \prod_{k=0}^{n}(\xi_{-k-1}^{(2k+1)})^2} \qquad (\in \mathbb{R}), \tag{1.16}$$

$$H_{2n+1}^{(-2n-1)} = -\frac{\nu_{n}^{(2n+1)}(\xi_{-n-1}^{(2n+1)})^2}{\prod_{j=0}^{n}(\xi_j^{(2j)})^2 \prod_{k=0}^{n}(\xi_{-k-1}^{(2k+1)})^2} \qquad (\in \mathbb{R}), \tag{1.17}$$

$$\frac{H_{2n+2}^{(-2n-2)}}{H_{2n}^{(-2n)}} = \frac{1}{(\xi_n^{(2n)})^2(\xi_{-n-1}^{(2n+1)})^2} \qquad (>0), \tag{1.18}$$

$$\frac{H_{2n+3}^{(-2n-2)}}{H_{2n+1}^{(-2n)}} = \frac{1}{(\xi_{n+1}^{(2n+2)})^2(\xi_{-n-1}^{(2n+1)})^2} \qquad (>0), \tag{1.19}$$

with

$$\xi_{n+1}^{(2n+2)} := h^{+}\left[\xi_n^{(2n)}\right],$$

where the homomorphisms $h^{\pm}$ (shift-up, +, and shift-down, −, operators) are defined as follows:

$$h^{\pm}\colon \mathbb{Z}\to\mathbb{Z},\ n\mapsto h^{\pm}[n] := n\pm 1 \qquad \text{and} \qquad f(n)\mapsto h^{\pm}[f(n)] := f(n\pm 1);$$

furthermore, denoting by $\mathbf{id}_{\mathbb{Z}}$ the identity operator on $\mathbb{Z}$, that is, $\mathbf{id}_{\mathbb{Z}}\colon \mathbb{Z}\to\mathbb{Z}$, $n\mapsto \mathbf{id}_{\mathbb{Z}}[n] := n$ and $f(n)\mapsto \mathbf{id}_{\mathbb{Z}}[f(n)] := f(n)$, one arrives at (the Abelian relation) $h^{+}h^{-} = h^{-}h^{+} = \mathbf{id}_{\mathbb{Z}}$.

Remark 1.3. It turns out that, for the presentation of the asymptotic results of this work, it is convenient to re-write $\mathrm{d}\widetilde{\mu}(z) = \exp(-\mathcal{N}V(z))\,\mathrm{d}z = \exp(-n\widetilde{V}(z))\,\mathrm{d}z =: \mathrm{d}\mu(z)$, $n\in\mathbb{N}$, where

$$\widetilde{V}(z) = z_o V(z),$$

with

$$z_o\colon \mathbb{N}\times\mathbb{N}\to(0,+\infty),\ (\mathcal{N},n)\mapsto z_o := \mathcal{N}/n,$$

and where the 'scaled' external field $\widetilde{V}\colon \mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfies:

$$\widetilde{V} \text{ is real analytic on } \mathbb{R}\setminus\{0\}; \tag{1.20}$$

$$\lim_{|x|\to\infty}\left(\widetilde{V}(x)/\ln(x^2+1)\right) = +\infty; \tag{1.21}$$

$$\lim_{|x|\to 0}\left(\widetilde{V}(x)/\ln(x^{-2}+1)\right) = +\infty. \tag{1.22}$$

(For example, a rational function of the form $\widetilde{V}(z) = \sum_{k=-2m_1}^{2m_2}\widetilde{\varrho}_k z^k$, with $\widetilde{\varrho}_k\in\mathbb{R}$, $k=-2m_1,\ldots,2m_2$, $m_{1,2}\in\mathbb{N}$, and $\widetilde{\varrho}_{-2m_1}, \widetilde{\varrho}_{2m_2} > 0$ would satisfy conditions (1.20)–(1.22).) Note that, under the homomorphisms $h^{\pm}\colon \mathbb{Z}\to\mathbb{Z}$,

$$z_o\mapsto h^{\pm}[z_o] = z_o\left(1\pm\frac{1}{n}\right)^{-1}.$$

Hereafter, the double-scaling limit as $\mathcal{N}, n\to\infty$ such that $z_o = 1+o(1)$ is studied: the simplified 'notation' $n\to\infty$ will be adopted. ■

Remark 1.4. The reader is reminded that, in the more classical setting of standard Hankel determinants, they are directly related to the theory of random matrices; indeed, as is well known, the partition function in Random Matrix Theory is actually a Hankel determinant. The asymptotic (as $n\to\infty$) behaviour of Hankel determinants with respect to varying exponential weights has only been proved for the 'single-interval/one-cut' case [25]. In the present, more general, setting, the asymptotic (as $n\to\infty$) behaviour of the generalised Hankel determinants (1.14)–(1.17) with respect to the varying exponential weight $w(z) = \exp(-n\widetilde{V}(z))$, $n\in\mathbb{N}$, where $\widetilde{V}\colon \mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfies conditions (1.20)–(1.22), is a very challenging undertaking; however, with the asymptotic (as $n\to\infty$) results of this work, one can easily obtain (real-valued) asymptotics (as $n\to\infty$) for $\mathfrak{P}_{\xi}H_{2n}^{(-2n)}$, $\mathfrak{P}_{\xi}H_{2n+1}^{(-2n)}$, $\mathfrak{P}_{\xi}H_{2n}^{(-2n+1)}$, and $\mathfrak{P}_{\xi}H_{2n+1}^{(-2n-1)}$, where $\mathfrak{P}_{\xi} := \prod_{j=0}^{n}(\xi_j^{(2j)})^2\prod_{k=0}^{n}(\xi_{-k-1}^{(2k+1)})^2$. ■

Unlike orthogonal polynomials, which satisfy a system of three-term recurrence relations, orthonormal/orthogonal L-polynomials satisfy more complicated recurrence relations: there are several different versions of these recurrence relations [11, 16, 17, 26–28]. The general form of these (system of) recurrence relations is a pair of three- and five-term recurrence relations (see, for example, [26]): for $n \in \mathbb{Z}_0^+$,

$$z\phi_{2n+1}(z) = b^\sharp_{2n+1}\phi_{2n}(z) + a^\sharp_{2n+1}\phi_{2n+1}(z) + b^\sharp_{2n+2}\phi_{2n+2}(z), \tag{1.23}$$

$$z\phi_{2n}(z) = c^\sharp_{2n}\phi_{2n-2}(z) + b^\sharp_{2n}\phi_{2n-1}(z) + a^\sharp_{2n}\phi_{2n}(z) + b^\sharp_{2n+1}\phi_{2n+1}(z) + c^\sharp_{2n+2}\phi_{2n+2}(z), \tag{1.24}$$

where $c^\sharp_0 = b^\sharp_0 = 0$, and $c^\sharp_{2k} > 0$, $k \in \mathbb{N}$, and

$$z^{-1}\phi_{2n}(z) = \beta^\sharp_{2n}\phi_{2n-1}(z) + \alpha^\sharp_{2n}\phi_{2n}(z) + \beta^\sharp_{2n+1}\phi_{2n+1}(z), \tag{1.25}$$

$$z^{-1}\phi_{2n+1}(z) = \gamma^\sharp_{2n+1}\phi_{2n-1}(z) + \beta^\sharp_{2n+1}\phi_{2n}(z) + \alpha^\sharp_{2n+1}\phi_{2n+1}(z) + \beta^\sharp_{2n+2}\phi_{2n+2}(z) + \gamma^\sharp_{2n+3}\phi_{2n+3}(z), \tag{1.26}$$

where $\beta^\sharp_0 = \gamma^\sharp_1 = 0$, $\beta^\sharp_1 > 0$, and $\gamma^\sharp_{2k+1} > 0$, $k \in \mathbb{N}$, leading, respectively, to the *real-symmetric*, *tri-penta-diagonal-type Laurent-Jacobi matrices*, $\mathcal{F}$ and $\mathcal{G}$, for the mappings $\mathcal{F}\colon \Lambda^{\mathbb{R}} \to \Lambda^{\mathbb{R}}$, $\mathfrak{f}(z) \mapsto z\mathfrak{f}(z)$, and $\mathcal{G}\colon \Lambda^{\mathbb{R}} \to \Lambda^{\mathbb{R}}$, $\mathfrak{g}(z) \mapsto z^{-1}\mathfrak{g}(z)$,

$$\mathcal{F} = \begin{pmatrix}
a^\sharp_0 & b^\sharp_1 & c^\sharp_2 & & & & & & & & & & \\
b^\sharp_1 & a^\sharp_1 & b^\sharp_2 & & & & & & & & & & \\
c^\sharp_2 & b^\sharp_2 & a^\sharp_2 & b^\sharp_3 & c^\sharp_4 & & & & & & & & \\
 & & b^\sharp_3 & a^\sharp_3 & b^\sharp_4 & & & & & & & & \\
 & & c^\sharp_4 & b^\sharp_4 & a^\sharp_4 & b^\sharp_5 & c^\sharp_6 & & & & & & \\
 & & & & b^\sharp_5 & a^\sharp_5 & b^\sharp_6 & & & & & & \\
 & & & & c^\sharp_6 & b^\sharp_6 & a^\sharp_6 & b^\sharp_7 & c^\sharp_8 & & & & \\
 & & & & & & b^\sharp_7 & a^\sharp_7 & b^\sharp_8 & & & & \\
 & & & & & & c^\sharp_8 & b^\sharp_8 & a^\sharp_8 & b^\sharp_9 & c^\sharp_{10} & & \\
 & & & & & & & & \ddots & \ddots & \ddots & & \\
 & & & & & & & & b^\sharp_{2k+1} & a^\sharp_{2k+1} & b^\sharp_{2k+2} & & \\
 & & & & & & & & c^\sharp_{2k+2} & b^\sharp_{2k+2} & a^\sharp_{2k+2} & b^\sharp_{2k+3} & c^\sharp_{2k+4} \\
 & & & & & & & & & & \ddots & \ddots & \ddots
\end{pmatrix}, \tag{1.27}$$

and

$$\mathcal{G} = \begin{pmatrix}
\alpha^\sharp_0 & \beta^\sharp_1 & & & & & & & & & & & \\
\beta^\sharp_1 & \alpha^\sharp_1 & \beta^\sharp_2 & \gamma^\sharp_3 & & & & & & & & & \\
 & \beta^\sharp_2 & \alpha^\sharp_2 & \beta^\sharp_3 & & & & & & & & & \\
 & \gamma^\sharp_3 & \beta^\sharp_3 & \alpha^\sharp_3 & \beta^\sharp_4 & \gamma^\sharp_5 & & & & & & & \\
 & & & \beta^\sharp_4 & \alpha^\sharp_4 & \beta^\sharp_5 & & & & & & & \\
 & & & \gamma^\sharp_5 & \beta^\sharp_5 & \alpha^\sharp_5 & \beta^\sharp_6 & \gamma^\sharp_7 & & & & & \\
 & & & & & \beta^\sharp_6 & \alpha^\sharp_6 & \beta^\sharp_7 & & & & & \\
 & & & & & \gamma^\sharp_7 & \beta^\sharp_7 & \alpha^\sharp_7 & \beta^\sharp_8 & \gamma^\sharp_9 & & & \\
 & & & & & & \beta^\sharp_8 & \alpha^\sharp_8 & \beta^\sharp_9 & & & & \\
 & & & & & & \ddots & \ddots & \ddots & & & & \\
 & & & & & & & & \gamma^\sharp_{2k+1} & \beta^\sharp_{2k+1} & \alpha^\sharp_{2k+1} & \beta^\sharp_{2k+2} & \gamma^\sharp_{2k+3} \\
 & & & & & & & & & & \beta^\sharp_{2k+2} & \alpha^\sharp_{2k+2} & \beta^\sharp_{2k+3} \\
 & & & & & & & & & & \ddots & \ddots & \ddots
\end{pmatrix}, \tag{1.28}$$

with zeros outside the indicated diagonals; furthermore, as shown in [26], $\mathcal{F}$ and $\mathcal{G}$ are formal inverses, that is, $\mathcal{F}\mathcal{G} = \mathcal{G}\mathcal{F} = \mathrm{diag}(1, 1, \ldots, 1, \ldots)$.

Remark 1.5. There are multitudinous applications of orthogonal L-polynomials and their associated Laurent-Jacobi matrices. To list a few: (i) numerical analysis (quadrature formulae) and trigonometric moment problems [29]; (ii) the spectral theory of one-to-one self-adjoint operators in infinite-dimensional (necessarily separable) Hilbert spaces [26, 30]; (iii) complex approximation theory (two-point Padé approximants) [31]; (iv) the direct/inverse scattering theory for the finite relativistic Toda lattice [32, 33] (see, also, the recent work [34], related to an extension of the Toda lattice); and (v) the classical Pick-Nevanlinna problem [35]. Some further comments on these connections can be found in Section 1 of [21] (see, also, [23], and the many references therein). It turns out that $n \to \infty$ asymptotics of orthogonal L-polynomials are an essential calculational ingredient in analyses related to the above-mentioned, seemingly disparate, topics. ■

It is, by now, a well-known, if not established, mathematical fact that variational conditions for minimisation problems in logarithmic potential theory, via the *equilibrium measure* [36–39], play a crucial rôle in the asymptotic analysis of (matrix) Riemann-Hilbert problems (RHPs) associated with (continuous and discrete) orthogonal polynomials, their roots, and corresponding recurrence relation coefficients (see, for example, [40–47]). The situation with respect to the large-n asymptotic analysis for the monic orthogonal L-polynomials, $\boldsymbol{\pi}_n(z)$, is analogous; however, unlike the asymptotic analysis for the orthogonal polynomials case, the asymptotic analysis for $\boldsymbol{\pi}_n(z)$ requires the consideration of two different families of RHPs, one for even degree (see **RHP1** below) and one for odd degree (see **RHP2** below). Thus, one must consider two sets of variational conditions for two (suitably posed) minimisation problems. Recall that the orthonormal L-polynomials, $\{\phi_m(z)\}_{m\in\mathbb{Z}_0^+}$ (cf. Equations (1.2) and (1.3)), are related to the monic orthogonal L-polynomials, $\{\boldsymbol{\pi}_m(z)\}_{m\in\mathbb{Z}_0^+}$, via Equations (1.4) and (1.5). The even degree and odd degree monic orthogonal L-polynomial problems for $\boldsymbol{\pi}_{2n}(z)$ ($m = 2n$) and $\boldsymbol{\pi}_{2n+1}(z)$ ($m = 2n+1$), respectively, are formulated, *à la* Fokas-Its-Kitaev [48, 49], as the following matrix RHPs on $\mathbb{R}$.

Before stating them, however, the notations/nomenclaturae used throughout the text are now summarised.

Notational Conventions

(1) $\mathrm{I} = \left(\begin{smallmatrix} 1 & 0 \\ 0 & 1 \end{smallmatrix}\right)$ is the 2×2 identity matrix, $\sigma_1 = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$, $\sigma_2 = \left(\begin{smallmatrix} 0 & -\mathrm{i} \\ \mathrm{i} & 0 \end{smallmatrix}\right)$, and $\sigma_3 = \left(\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix}\right)$ are the Pauli matrices, $\sigma_\pm := \frac{1}{2}(\sigma_1 \pm \mathrm{i}\sigma_2)$, $\mathbb{R}_\pm := \{x \in \mathbb{R};\ \pm x > 0\}$, and $\mathbb{C}_\pm := \{z \in \mathbb{C};\ \pm \mathrm{Im}(z) > 0\}$;

(2) a contour $\mathcal{D}$ which is the finite union of piecewise-smooth, simple curves (as closed sets) is said to be *orientable* if its complement, $\mathbb{C} \setminus \mathcal{D}$, can always be divided into two, possibly disconnected, disjoint open sets $\boldsymbol{\mho^+}$ and $\boldsymbol{\mho^-}$, either of which has finitely many components, such that $\mathcal{D}$ admits an orientation so that it can either be viewed as a positively oriented boundary $\mathcal{D}^+$ for $\boldsymbol{\mho^+}$ or as a negatively oriented boundary $\mathcal{D}^-$ for $\boldsymbol{\mho^-}$ [50], that is, the (possibly disconnected) components of $\mathbb{C} \setminus \mathcal{D}$ can be coloured by + or − in such a way that the + regions do not share boundary with the − regions, except, possibly, at finitely many points [51];

(3) for each segment of an oriented contour $\mathcal{D}$, according to the given orientation, the "+" side is to the left and the "-" side is to the right as one traverses the contour in the direction of orientation; so, for a matrix-valued function $\mathcal{A}(z)$, $\mathcal{A}_\pm(z)$ denote the non-tangential limits $\mathcal{A}_\pm(z) := \lim_{\substack{z' \to z \\ z' \in \pm \text{ side of } \mathcal{D}}} \mathcal{A}(z')$;

(4) unless stated otherwise, exponents such as z^ν shall be interpreted as principal branches;

(5) for some point set $\mathcal{D} \subset \mathcal{X}$, with $\mathcal{X} = \mathbb{C}$ or $\mathbb{R}$, $\overline{\mathcal{D}} := \mathcal{D} \cup \partial\mathcal{D}$, and $\mathcal{D}^c := \mathcal{X} \setminus \overline{\mathcal{D}}$;

(6) unless stated otherwise, $\mathbb{R}$ is oriented from $-\infty$ to $+\infty$;

(7) for a 2×2 matrix-valued function $\mathcal{A}(z)$, $\mathcal{A}_{ij}(z)$, $i, j = 1, 2$, denotes the $(i\,j)$-element of $\mathcal{A}(z)$;

(8) for a 2×2 matrix-valued function $\mathcal{Y}(z)$, the notation $\mathcal{Y}(z) =_{z \to z_0} O(*)$ means $\mathcal{Y}_{ij}(z) =_{z \to z_0} O(*_{ij})$, $i, j = 1, 2$ (*mutatis mutandis* for $o(1)$).

The RHP which characterises the even degree monic orthogonal L-polynomials $\boldsymbol{\pi}_{2n}(z)$, $n \in \mathbb{Z}_0^+$, is now stated.

RHP1 ([21])**.** Let $V:\mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfy conditions (V1)–(V3). Find $\overset{e}{\mathrm{Y}}:\mathbb{C}\setminus\mathbb{R}\to \mathrm{SL}_2(\mathbb{C})$ solving: (i) $\overset{e}{\mathrm{Y}}(z)$ is holomorphic for $z\in\mathbb{C}\setminus\mathbb{R}$; (ii) for $z\in\mathbb{R}$, the boundary values $\overset{e}{\mathrm{Y}}_{\pm}(z)$ satisfy the jump condition

$$\overset{e}{\mathrm{Y}}_{+}(z) = \overset{e}{\mathrm{Y}}_{-}(z)(\mathrm{I}+\exp(-\mathcal{N}V(z))\sigma_{+});$$

(iii) $\overset{e}{\mathrm{Y}}(z)z^{-n\sigma_3} =_{\substack{z\to\infty\\ z\in\mathbb{C}\setminus\mathbb{R}}} \mathrm{I}+O(z^{-1})$; and (iv) $\overset{e}{\mathrm{Y}}(z)z^{n\sigma_3} =_{\substack{z\to 0\\ z\in\mathbb{C}\setminus\mathbb{R}}} O(1)$.

Lemma 1.1 ([21])**.** *Let* $\overset{e}{\mathrm{Y}}:\mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$ *solve* **RHP1**. **RHP1** *possesses a unique solution given by:* (i) *for* $n=0$,

$$\overset{e}{\mathrm{Y}}(z)=\begin{pmatrix}1 & \int_{\mathbb{R}}\frac{\exp(-\mathcal{N}V(s))}{s-z}\frac{\mathrm{d}s}{2\pi\mathrm{i}}\\ 0 & 1\end{pmatrix},\quad z\in\mathbb{C}\setminus\mathbb{R},$$

where $\boldsymbol{\pi}_0(z):=\overset{e}{\mathrm{Y}}_{11}(z)\equiv 1$; *and* (ii) *for* $n\in\mathbb{N}$,

$$\overset{e}{\mathrm{Y}}(z)=\begin{pmatrix}\boldsymbol{\pi}_{2n}(z) & \int_{\mathbb{R}}\frac{\boldsymbol{\pi}_{2n}(s)\exp(-\mathcal{N}V(s))}{s-z}\frac{\mathrm{d}s}{2\pi\mathrm{i}}\\ \overset{e}{\mathrm{Y}}_{21}(z) & \int_{\mathbb{R}}\frac{\overset{e}{\mathrm{Y}}_{21}(s)\exp(-\mathcal{N}V(s))}{s-z}\frac{\mathrm{d}s}{2\pi\mathrm{i}}\end{pmatrix},\quad z\in\mathbb{C}\setminus\mathbb{R},\tag{1.29}$$

where $\overset{e}{\mathrm{Y}}_{21}:\mathbb{C}^*\to\mathbb{C}$, *and* $\boldsymbol{\pi}_{2n}(z)$ *is the even degree monic* OLP *defined in Equation* (1.4).

The RHP which characterises the odd degree monic orthogonal L-polynomials $\boldsymbol{\pi}_{2n+1}(z)$, $n\in\mathbb{Z}_0^+$, is now stated.

RHP2 ([22])**.** Let $V:\mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfy conditions (V1)–(V3). Find $\overset{o}{\mathrm{Y}}:\mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$ solving: (i) $\overset{o}{\mathrm{Y}}(z)$ is holomorphic for $z\in\mathbb{C}\setminus\mathbb{R}$; (ii) for $z\in\mathbb{R}$, the boundary values $\overset{o}{\mathrm{Y}}_{\pm}(z)$ satisfy the jump condition

$$\overset{o}{\mathrm{Y}}_{+}(z) = \overset{o}{\mathrm{Y}}_{-}(z)(\mathrm{I}+\exp(-\mathcal{N}V(z))\sigma_{+});$$

(iii) $\overset{o}{\mathrm{Y}}(z)z^{n\sigma_3} =_{\substack{z\to 0\\ z\in\mathbb{C}\setminus\mathbb{R}}} \mathrm{I}+O(z)$; and (iv) $\overset{o}{\mathrm{Y}}(z)z^{-(n+1)\sigma_3} =_{\substack{z\to\infty\\ z\in\mathbb{C}\setminus\mathbb{R}}} O(1)$.

Lemma 1.2 ([22])**.** *Let* $\overset{o}{\mathrm{Y}}:\mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$ *solve* **RHP2**. **RHP2** *possesses a unique solution given by:* (i) *for* $n=0$,

$$\overset{o}{\mathrm{Y}}(z)=\begin{pmatrix}z\boldsymbol{\pi}_1(z) & z\int_{\mathbb{R}}\frac{(s\boldsymbol{\pi}_1(s))\exp(-\mathcal{N}V(s))}{s(s-z)}\frac{\mathrm{d}s}{2\pi\mathrm{i}}\\ 2\pi\mathrm{i}z & 1+z\int_{\mathbb{R}}\frac{\exp(-\mathcal{N}V(s))}{s-z}\mathrm{d}s\end{pmatrix},\quad z\in\mathbb{C}\setminus\mathbb{R},$$

where $\boldsymbol{\pi}_1(z)=\frac{1}{z}+\frac{\xi_0^{(1)}}{\xi_{-1}^{(1)}}$, *with* $\frac{\xi_0^{(1)}}{\xi_{-1}^{(1)}}=-\int_{\mathbb{R}}s^{-1}\exp(-\mathcal{N}V(s))\,\mathrm{d}s$, $\mathcal{N}\in\mathbb{N}$; *and* (ii) *for* $n\in\mathbb{N}$,

$$\overset{o}{\mathrm{Y}}(z)=\begin{pmatrix}z\boldsymbol{\pi}_{2n+1}(z) & z\int_{\mathbb{R}}\frac{(s\boldsymbol{\pi}_{2n+1}(s))\exp(-\mathcal{N}V(s))}{s(s-z)}\frac{\mathrm{d}s}{2\pi\mathrm{i}}\\ \overset{o}{\mathrm{Y}}_{21}(z) & z\int_{\mathbb{R}}\frac{\overset{o}{\mathrm{Y}}_{21}(s)\exp(-\mathcal{N}V(s))}{s(s-z)}\frac{\mathrm{d}s}{2\pi\mathrm{i}}\end{pmatrix},\quad z\in\mathbb{C}\setminus\mathbb{R},\tag{1.30}$$

where $\overset{o}{\mathrm{Y}}_{21}:\mathbb{C}^*\to\mathbb{C}$, *and* $\boldsymbol{\pi}_{2n+1}(z)$ *is the odd degree monic* OLP *defined in Equation* (1.5).

This paper, which is not completely self-contained (see Remark 1.6 below), is a continuation, proper, of [21, 22], where, for example, asymptotics in the double-scaling limit as $\mathcal{N},n\to\infty$ such that $z_o=1+o(1)$ of the monic orthogonal L-polynomials, $\{\boldsymbol{\pi}_n(z)\}_{n\in\mathbb{Z}_0^+}$, and orthonormal L-polynomials, $\{\phi_n(z)\}_{n\in\mathbb{Z}_0^+}$, in the entire complex plane, were obtained. In [21, 22], the L-polynomials were taken to be orthogonal with respect to the varying exponential measure (cf. Remark 1.3)[1] $\mathrm{d}\mu(z)=\mathrm{e}^{-n\widetilde{V}(z)}\,\mathrm{d}z$, where

[1]Note that $LD(\boldsymbol{\pi}_m)=LD(\phi_m)=\begin{cases}2n, & m=\text{even},\\ 2n+1, & m=\text{odd},\end{cases}$ coincides with the parameter in the measure of orthogonality: the large parameter, n, enters simultaneously into the L-degree of the L-polynomials and the (varying exponential) weight; thus, asymptotics of the L-polynomials are studied along a 'diagonal strip' of a doubly-indexed sequence.

the external field[2] $\widetilde{V}\colon \mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfies conditions (1.20)–(1.22). Furthermore, asymptotics of the 'even' and 'odd' norming constants, $\xi_n^{(2n)}$ and $\xi_{-n-1}^{(2n+1)}$, respectively, as well as, subsequently, those of the ('even' and 'odd') norms $\|\boldsymbol{\pi}_{2n}(\cdot)\|_{\mathcal{L}}$, $\|\boldsymbol{\pi}_{2n+1}(\cdot)\|_{\mathcal{L}}$, and the Hankel determinant ratios $H_{2n}^{(-2n)}/H_{2n+1}^{(-2n)}$, $H_{2n+1}^{(-2n)}/H_{2n+2}^{(-2n-2)}$, were also obtained. The aforementioned asymptotics were derived by extracting the large-n behaviours of the respective solutions of **RHP1** and **RHP2**. The paradigm for the asymptotic analysis of **RHP1** and **RHP2** is a union of the Deift-Zhou (DZ) non-linear steepest-descent method [1, 2], used for the asymptotic analysis of undulatory RHPs, and the extension of Deift-Venakides-Zhou [3], incorporating into the DZ method a non-linear analogue of the WKB method, making the asymptotic analysis of fully non-linear problems tractable. It should be re-emphasized that, in this context, the equilibrium measure plays an absolutely crucial rôle in the analysis [36–39]; multitudinous extensions and applications of the DZ method can be found in [40–47, 53–67].

In this paper, asymptotics for the following quantities are obtained:

- the (real) root products of the L-polynomials, for both the non-singular and singular cases,

$$\prod_{k=1}^{2n}\alpha_k^{(2n)},\qquad \prod_{k=1}^{2n+1}\alpha_k^{(2n+1)},\qquad \prod_{k=1}^{2n-1}\alpha_k^{(2n)},\qquad \prod_{k=1}^{2n}\alpha_k^{(2n+1)};$$

- Hankel determinant ratios (all real valued) associated with the bi-infinite, real-valued, strong moment sequence $\left\{c_k=\int_{\mathbb{R}}s^k\exp(-n\widetilde{V}(s))\,\mathrm{d}s,\ n\in\mathbb{N}\right\}_{k\in\mathbb{Z}}$,

$$\frac{H_{2n+2}^{(-2n-2)}}{H_{2n}^{(-2n)}},\qquad \frac{H_{2n+3}^{(-2n-2)}}{H_{2n+1}^{(-2n)}},\qquad \frac{H_{2n}^{(-2n+1)}}{H_{2n}^{(-2n)}},\qquad \frac{H_{2n+1}^{(-2n-1)}}{H_{2n+1}^{(-2n)}};$$

- recurrence relation coefficients (all real valued),

$$a_{2n}^{\sharp},\ b_{2n}^{\sharp},\ c_{2n}^{\sharp},\ a_{2n+1}^{\sharp},\ b_{2n+1}^{\sharp},\ c_{2n+2}^{\sharp},\ b_{2n+2}^{\sharp},\ \alpha_{2n}^{\sharp},\ \beta_{2n}^{\sharp},\ \gamma_{2n+1}^{\sharp},\ \alpha_{2n+1}^{\sharp},\ \beta_{2n+1}^{\sharp},\ \gamma_{2n+3}^{\sharp},\ \beta_{2n+2}^{\sharp},$$

  and subsequently those of the (real-symmetric, tri-penta-diagonal-type) Laurent-Jacobi matrices $\mathcal{F}$ and $\mathcal{G}$.

In order to accomplish this, the asymptotic results of [21, 22], in particular, asymptotics of $\overset{e}{\mathrm{Y}}(z)$ and $\overset{o}{\mathrm{Y}}(z)$ as $z\to 0$ and $z\to\infty$, where $z\in\mathbb{C}\setminus\mathbb{R}$, with n-dependent coefficients, are used.

Given the results of [21, 22], carrying through with the above-mentioned programme involves lengthy algebraic calculations. A plethora of notations remains to be introduced and defined; in particular, basic elements associated with the construction of hyperelliptic and finite genus (compact) Riemann surfaces and the corresponding Riemann theta functions, and the construction of the respective 'even' and 'odd' equilibrium measures (see Section 2).

Remark 1.6. Strictly speaking, this paper is a hybrid continuation of [21, 22], and is not completely self-contained: concepts, definitions, formulas, and theorems proven in [21, 22] are used. ■

Remark 1.7. The superscripts $^{\pm}$, and sometimes the subscripts $_{\pm}$, used throughout the paper should not be confused with the subscripts $_{\pm}$ appearing in the various RHPs. ■

Remark 1.8. Although $\overline{\mathbb{C}}$ (or $\mathbb{CP}^1$) $:=\mathbb{C}\cup\{\infty\}$ (resp., $\overline{\mathbb{R}}:=\mathbb{R}\cup\{-\infty\}\cup\{+\infty\}$) is the standard definition for the (closed) Riemann sphere (resp., closed real line), the simplified, and somewhat abusive, notation $\mathbb{C}$ (resp., $\mathbb{R}$) is used to denote both the (closed) Riemann sphere, $\overline{\mathbb{C}}$ (resp., closed real line, $\overline{\mathbb{R}}$), and the (open) complex field, $\mathbb{C}$ (resp., open real line, $\mathbb{R}$), and the context(s) should make clear which object(s) the notation $\mathbb{C}$ (resp., $\mathbb{R}$) represents. ■

2 Preliminary Considerations and Auxiliary Functions

In this section, auxiliary information which is necessary in order to obtain the asymptotic (as $n\to\infty$) results presented in Section 3 is gathered. Towards this end, a terse presentation of all the relevant particulars follows (see [21, 22] for complete details and proofs).

[2] For real non-analytic external fields, see the recent work [52].

It turns out that, for the asymptotic analysis of this work, the following multi-valued functions, and their associated hyperelliptic Riemman surfaces of *genus* N ($\in\mathbb{N}$), assumed finite, are essential; for further details, and proofs, see, for example, [68, 69]. There are two cases to consider: the case of even degree, and the case of odd degree. The even degree case is treated first.

$\sqrt{R_e(z)}$ Given the set of real points $-\infty < a_0^e < b_0^e < a_1^e < b_1^e < \cdots < a_N^e < b_N^e < +\infty$, where $a_{N+1}^e \equiv a_0^e$ (as points on the Riemann sphere, $\overline{\mathbb{C}}$), define the multi-valued function $(R_e(z))^{1/2}$ as follows:

$$(R_e(z))^{1/2} := \left(\prod_{k=0}^{N}\left(z-b_k^e\right)\left(z-a_{k+1}^e\right)\right)^{1/2} \tag{2.1}$$

(see [21], Figure 1). It must be noted that the set of points $\{b_{j-1}^e, a_j^e\}_{j=1}^{N+1}$, which form the end-points of the support of the associated 'even' equilibrium measure (see below), are not arbitrary; rather, they satisfy the (real) n-dependent and locally solvable system of $2(N+1)$ *moment conditions* (2.12). The function $R_e(z)$ ($\in\mathbb{R}[z]$: the algebra of polynomials in z with coefficients in $\mathbb{R}$) is a unital polynomial of even degree ($\deg(R_e(z)) = 2(N+1)$) whose simple roots/zeros are $\{b_{j-1}^e, a_j^e\}_{j=1}^{N+1}$.

Let $\mathcal{Y}_e := \left\{(y,z);\, y^2 = R_e(z)\right\}$ denote the two-sheeted Riemann surface of genus N associated with $y^2 = R_e(z)$: the first sheet of $\mathcal{Y}_e$ is denoted by $\mathcal{Y}_e^+$, and points on the first sheet are represented as $z^+ := (z, +(R_e(z))^{1/2})$; similarly, the second sheet of $\mathcal{Y}_e$ is denoted by $\mathcal{Y}_e^-$, and points on the second sheet are represented as $z^- := (z, -(R_e(z))^{1/2})$. As points on the plane $\mathbb{C}$, $z^+ = z^- := z$. The single-valued branch for the square root of $(R_e(z))^{1/2}$ is chosen so that $z^{-(N+1)}(R_e(z))^{1/2} \underset{\substack{z\to\infty\\ z\in\mathcal{Y}_e^{\pm}}}{\sim} \pm 1$. $\mathcal{Y}_e$ is realised as a (two-sheeted) branched/ramified covering of the Riemann sphere such that its two sheets are two identical copies of $\mathbb{C}$ with branch cuts (slits) along the intervals $(a_0^e, b_0^e), (a_1^e, b_1^e), \ldots, (a_N^e, b_N^e)$ and glued together 'cross-wise' in the canonical manner along $\cup_{j=1}^{N+1}(a_{j-1}^e, b_{j-1}^e)$. The *canonical* **1**-*homology basis* for $\mathcal{Y}_e$ are the cycles (closed contours) $\boldsymbol{\alpha}_0^e$ and $\{\boldsymbol{\alpha}_j^e, \boldsymbol{\beta}_j^e\}$, $j = 1, \ldots, N$, which are characterised by the fact that the cycles $\boldsymbol{\alpha}_j^e$, $j = 0, \ldots, N$, lie on $\mathcal{Y}_e^+$, and the cycles $\boldsymbol{\beta}_j^e$, $j = 1, \ldots, N$, pass from $\mathcal{Y}_e^+$ (starting from the slit (a_j^e, b_j^e)), through the slit (a_0^e, b_0^e) to $\mathcal{Y}_e^-$, and back again to $\mathcal{Y}_e^+$ through the slit (a_j^e, b_j^e) (see [21], Figure 3). (Note: in this work, $\mathcal{Y}_e^{\pm} = \mathbb{C}_{\pm}$.)

The canonical **1**-homology basis $\{\boldsymbol{\alpha}_j^e, \boldsymbol{\beta}_j^e\}$, $j = 1, \ldots, N$, generates, on $\mathcal{Y}_e$, the (corresponding) $\boldsymbol{\alpha}^e$-normalised basis of holomorphic Abelian differentials (one-forms) $\{\omega_1^e, \omega_2^e, \ldots, \omega_N^e\}$, where $\omega_j^e := \sum_{k=1}^{N} \frac{c_{jk}^e z^{N-k}}{\sqrt{R_e(z)}}\,\mathrm{d}z$, $c_{jk}^e \in \mathbb{C}$, $j = 1, \ldots, N$, and $\oint_{\boldsymbol{\alpha}_k^e} \omega_j^e = \delta_{kj}$, $k, j = 1, \ldots, N$: ω_l^e, $l = 1, \ldots, N$, is real valued on $\cup_{j=1}^{N+1}(a_{j-1}^e, b_{j-1}^e)$, and has exactly one (real) root in any (open) interval (a_{j-1}^e, b_{j-1}^e), $j = 1, \ldots, N+1$; furthermore, in the intervals (b_{j-1}^e, a_j^e), $j = 1, \ldots, N$, ω_l^e, $l = 1, \ldots, N$, take non-zero, pure imaginary values. Let $\boldsymbol{\omega}^e := (\omega_1^e, \omega_2^e, \ldots, \omega_N^e)$ denote the basis of holomorphic one-forms on $\mathcal{Y}_e$ as normalised above with the associated $N \times N$ *Riemann matrix* of $\boldsymbol{\beta}^e$-periods, $\tau^e = (\tau^e)_{i,j=1,\ldots,N} := (\oint_{\boldsymbol{\beta}_j^e} \omega_i^e)_{i,j=1,\ldots,N}$. The Riemann matrix, τ^e, is symmetric and pure imaginary, $-\mathrm{i}\tau^e$ is positive definite ($\mathrm{Im}(\tau_{ij}^e) > 0$), and $\det(\tau^e) \neq 0$. For the holomorphic Abelian differential (one-form) $\boldsymbol{\omega}^e$ defined above, choose a_{N+1}^e as the *base point*, and set $\boldsymbol{u}^e : \mathcal{Y}_e \to \mathrm{Jac}(\mathcal{Y}_e), z \mapsto \boldsymbol{u}^e(z) := \int_{a_{N+1}^e}^{z} \boldsymbol{\omega}^e$, where $\mathrm{Jac}(\mathcal{Y}_e) := \mathbb{C}^N / \{N + \tau^e M\}$, $(N, M) \in \mathbb{Z}^N \times \mathbb{Z}^N$, is the *Jacobi variety* of $\mathcal{Y}_e$, and the integration from a_{N+1}^e to z ($\in \mathcal{Y}_e$) is taken along any path on $\mathcal{Y}_e^+$.

Remark 2.1. From the representation $\omega_j^e = \sum_{k=1}^{N} \frac{c_{jk}^e z^{N-k}}{\sqrt{R_e(z)}}\,\mathrm{d}z$, $j = 1, \ldots, N$, and the normalisation condition $\oint_{\boldsymbol{\alpha}_k^e} \omega_j^e = \delta_{kj}$, $k, j = 1, \ldots, N$, one shows that c_{jk}^e, $k, j = 1, \ldots, N$, are obtained from

$$\begin{pmatrix} c_{11}^e & c_{12}^e & \cdots & c_{1N}^e \\ c_{21}^e & c_{22}^e & \cdots & c_{2N}^e \\ \vdots & \vdots & \ddots & \vdots \\ c_{N1}^e & c_{N2}^e & \cdots & c_{NN}^e \end{pmatrix} = \widetilde{\mathfrak{S}}_e^{-1}, \tag{E1}$$

where

$$\widetilde{\Xi}_e := \begin{pmatrix} \oint_{\boldsymbol{\alpha}_1^e} \frac{\mathrm{d}s_1}{\sqrt{R_e(s_1)}} & \oint_{\boldsymbol{\alpha}_2^e} \frac{\mathrm{d}s_2}{\sqrt{R_e(s_2)}} & \cdots & \oint_{\boldsymbol{\alpha}_N^e} \frac{\mathrm{d}s_N}{\sqrt{R_e(s_N)}} \\ \oint_{\boldsymbol{\alpha}_1^e} \frac{s_1\mathrm{d}s_1}{\sqrt{R_e(s_1)}} & \oint_{\boldsymbol{\alpha}_2^e} \frac{s_2\mathrm{d}s_2}{\sqrt{R_e(s_2)}} & \cdots & \oint_{\boldsymbol{\alpha}_N^e} \frac{s_N\mathrm{d}s_N}{\sqrt{R_e(s_N)}} \\ \vdots & \vdots & \ddots & \vdots \\ \oint_{\boldsymbol{\alpha}_1^e} \frac{s_1^{N-1}\mathrm{d}s_1}{\sqrt{R_e(s_1)}} & \oint_{\boldsymbol{\alpha}_2^e} \frac{s_2^{N-1}\mathrm{d}s_2}{\sqrt{R_e(s_2)}} & \cdots & \oint_{\boldsymbol{\alpha}_N^e} \frac{s_N^{N-1}\mathrm{d}s_N}{\sqrt{R_e(s_N)}} \end{pmatrix}. \tag{E2}$$

For a (representation-independent) proof of the fact that $\det(\widetilde{\Xi}_e) \neq 0$, see, for example, Chapter 10, Section 10–2, of [68]. ■

Set [21]

$$\gamma^e(z) := \begin{cases} \left(\prod_{k=1}^{N+1} (z - b_{k-1}^e)(z - a_k^e)^{-1} \right)^{1/4}, & z \in \mathbb{C}_+, \\ -\mathrm{i} \left(\prod_{k=1}^{N+1} (z - b_{k-1}^e)(z - a_k^e)^{-1} \right)^{1/4}, & z \in \mathbb{C}_-. \end{cases} \tag{2.2}$$

It is shown in Section 4 of [21] that $\gamma^e(z) =_{\substack{z\to\infty \\ z\in \mathcal{Y}_e^{\pm}}} (-\mathrm{i})^{(1\mp 1)/2}(1 + O(z^{-1}))$, and that

$$\left\{ z_j^{e,\pm} \right\}_{j=1}^N = \left\{ z^{\pm} \in \mathcal{Y}_e^{\pm};\ (\gamma^e(z) \mp (\gamma^e(z))^{-1})|_{z=z^{\pm}} = 0 \right\}, \tag{2.3}$$

with $z_j^{e,\pm} \in (a_j^e, b_j^e)^{\pm}$ $(\subset \mathcal{Y}_e^{\pm})$, $j = 1, \ldots, N$, where, as points on the plane, $z_j^{e,+} = z_j^{e,-} := z_j^e$, $j = 1, \ldots, N$ (on the plane, $z_j^e \in (a_j^e, b_j^e)$, $j = 1, \ldots, N$).

Corresponding to $\mathcal{Y}_e$, define $\boldsymbol{d}_e := -\boldsymbol{K}_e - \sum_{j=1}^N \int_{a_{N+1}^e}^{z_j^{e,-}} \boldsymbol{\omega}^e$ $(\in \mathbb{C}^N)$, where $\boldsymbol{K}_e$ is the associated 'even' *vector of Riemann constants*, and the integration from a_{N+1}^e to $z_j^{e,-}$, $j = 1, \ldots, N$, is taken along a fixed path in $\mathcal{Y}_e^-$. It is shown in Chapter VII of [69] that $\boldsymbol{K}_e = \sum_{j=1}^N \int_{a_j^e}^{a_{N+1}^e} \boldsymbol{\omega}^e$; furthermore, $\boldsymbol{K}_e$ is a point of order 2, that is, $2\boldsymbol{K}_e = 0$ and $s\boldsymbol{K}_e \neq 0$ for $0 < s < 2$. Recalling the definition of $\boldsymbol{\omega}^e$ and that $z^{-(N+1)}(R_e(z))^{1/2} \sim_{\substack{z\to\infty \\ z\in\mathbb{C}_{\pm}}} \pm 1$, using the fact that $\boldsymbol{K}_e$ is a point of order 2, one arrives at

$$\begin{aligned} \boldsymbol{d}_e &= -\boldsymbol{K}_e - \sum_{j=1}^N \int_{a_{N+1}^e}^{z_j^{e,-}} \boldsymbol{\omega}^e = \boldsymbol{K}_e - \sum_{j=1}^N \int_{a_{N+1}^e}^{z_j^{e,-}} \boldsymbol{\omega}^e = -\boldsymbol{K}_e + \sum_{j=1}^N \int_{a_{N+1}^e}^{z_j^{e,+}} \boldsymbol{\omega}^e = \boldsymbol{K}_e + \sum_{j=1}^N \int_{a_{N+1}^e}^{z_j^{e,+}} \boldsymbol{\omega}^e \\ &= -\sum_{j=1}^N \int_{a_j^e}^{z_j^{e,-}} \boldsymbol{\omega}^e = \sum_{j=1}^N \int_{a_j^e}^{z_j^{e,+}} \boldsymbol{\omega}^e. \end{aligned} \tag{2.4}$$

Associated with the Riemann matrix of $\boldsymbol{\beta}^e$-periods, τ^e, is the 'even' *Riemann theta function*:

$$\boldsymbol{\theta}(z; \tau^e) =: \boldsymbol{\theta}^e(z) = \sum_{m\in\mathbb{Z}^N} \mathrm{e}^{2\pi\mathrm{i}(m,z) + \pi\mathrm{i}(m, \tau^e m)}, \quad z \in \mathbb{C}^N, \tag{2.5}$$

where $(\cdot, \cdot)$ denotes the real Euclidean inner/scalar product[3]; $\boldsymbol{\theta}^e(z)$ has the following evenness and (quasi-) periodicity properties,

$$\boldsymbol{\theta}^e(-z) = \boldsymbol{\theta}^e(z), \qquad \boldsymbol{\theta}^e(z + e_j) = \boldsymbol{\theta}^e(z), \qquad \text{and} \qquad \boldsymbol{\theta}^e(z \pm \tau_j^e) = \mathrm{e}^{\mp 2\pi\mathrm{i}z_j - \mathrm{i}\pi\tau_{jj}^e} \boldsymbol{\theta}^e(z),$$

where e_j is the standard (basis) column vector in $\mathbb{C}^N$ with 1 in the jth entry and 0 elsewhere, and $\tau_j^e := \tau^e e_j$ $(\in \mathbb{C}^N)$, $j = 1, \ldots, N$.

The odd degree case is now treated.

[3] For $A = (A_1, A_2, \ldots, A_N) \in \mathbb{E}^N$ and $B = (B_1, B_2, \ldots, B_N) \in \mathbb{E}^N$, $(A, B) := \sum_{k=1}^N A_k B_k$.

$\sqrt{R_o(z)}$ Given the set of real points $-\infty < a_0^o < b_0^o < a_1^o < b_1^o < \cdots < a_N^o < b_N^o < +\infty$, where $a_{N+1}^o \equiv a_0^o$ (as points on the Riemann sphere, $\overline{\mathbb{C}}$), define the multi-valued function $(R_o(z))^{1/2}$ as follows:

$$(R_o(z))^{1/2} := \left(\prod_{k=0}^{N} \left(z - b_k^o\right)\left(z - a_{k+1}^o\right) \right)^{1/2} \tag{2.6}$$

(see [22], Figure 2). It must be noted that the set of points $\{b_{j-1}^o, a_j^o\}_{j=1}^{N+1}$, which form the endpoints of the support of the associated 'odd' equilibrium measure (see below), are not arbitrary; rather, they satisfy the (real) n-dependent and locally solvable system of $2(N+1)$ moment conditions (2.18). The function $R_o(z)$ ($\in \mathbb{R}[z]$) is a unital polynomial of even degree ($\deg(R_o(z)) = 2(N+1)$) whose simple roots/zeros are $\{b_{j-1}^o, a_j^o\}_{j=1}^{N+1}$.

Let $\mathcal{Y}_o := \left\{(y,z);\, y^2 = R_o(z)\right\}$ denote the two-sheeted Riemann surface of genus N associated with $y^2 = R_o(z)$: the first sheet of $\mathcal{Y}_o$ is denoted by $\mathcal{Y}_o^+$, and points on the first sheet are represented as $z^+ := (z, +(R_o(z))^{1/2})$; similarly, the second sheet of $\mathcal{Y}_o$ is denoted by $\mathcal{Y}_o^-$, and points on the second sheet are represented as $z^- := (z, -(R_o(z))^{1/2})$. As points on the plane $\mathbb{C}$, $z^+ = z^- := z$. The single-valued branch for the square root of $(R_o(z))^{1/2}$ is chosen so that $z^{-(N+1)}(R_o(z))^{1/2} \sim_{\substack{z\to\infty \\ z\in\mathcal{Y}_o^{\pm}}} \pm 1$. $\mathcal{Y}_o$ is realised as a (two-sheeted) branched/ramified covering of the Riemann sphere such that its two sheets are two identical copies of $\mathbb{C}$ with branch cuts (slits) along the intervals $(a_0^o, b_0^o), (a_1^o, b_1^o), \ldots, (a_N^o, b_N^o)$ and glued together 'cross-wise' in the canonical manner along $\cup_{j=1}^{N+1}(a_{j-1}^o, b_{j-1}^o)$. The canonical **1**-homology basis for $\mathcal{Y}_o$ are the cycles (closed contours) $\boldsymbol{\alpha}_0^o$ and $\{\boldsymbol{\alpha}_j^o, \boldsymbol{\beta}_j^o\}$, $j = 1, \ldots, N$, which are characterised by the fact that the cycles $\boldsymbol{\alpha}_j^o$, $j = 0, \ldots, N$, lie on $\mathcal{Y}_o^+$, and the cycles $\boldsymbol{\beta}_j^o$, $j = 1, \ldots, N$, pass from $\mathcal{Y}_o^+$ (starting from the slit (a_j^o, b_j^o)), through the slit (a_0^o, b_0^o) to $\mathcal{Y}_o^-$, and back again to $\mathcal{Y}_o^+$ through the slit (a_j^o, b_j^o) (see [22], Figure 4). (Note: in this work, $\mathcal{Y}_o^{\pm} = \mathbb{C}_{\pm}$.)

The canonical **1**-homology basis $\{\boldsymbol{\alpha}_j^o, \boldsymbol{\beta}_j^o\}$, $j = 1, \ldots, N$, generates, on $\mathcal{Y}_o$, the (corresponding) $\boldsymbol{\alpha}^o$-normalised basis of holomorphic Abelian differentials (one-forms) $\{\omega_1^o, \omega_2^o, \ldots, \omega_N^o\}$, where $\omega_j^o := \sum_{k=1}^{N} \frac{c_{jk}^o z^{N-k}}{\sqrt{R_o(z)}}\,\mathrm{d}z$, $c_{jk}^o \in \mathbb{C}$, $j = 1, \ldots, N$, and $\oint_{\boldsymbol{\alpha}_k^o} \omega_j^o = \delta_{kj}$, $k, j = 1, \ldots, N$: ω_l^o, $l = 1, \ldots, N$, is real valued on $\cup_{j=1}^{N+1}(a_{j-1}^o, b_{j-1}^o)$, and has exactly one (real) root in any (open) interval (a_{j-1}^o, b_{j-1}^o), $j = 1, \ldots, N{+}1$; furthermore, in the intervals (b_{j-1}^o, a_j^o), $j = 1, \ldots, N$, ω_l^o, $l = 1, \ldots, N$, take non-zero, pure imaginary values. Let $\boldsymbol{\omega}^o := (\omega_1^o, \omega_2^o, \ldots, \omega_N^o)$ denote the basis of holomorphic one-forms on $\mathcal{Y}_o$ as normalised above with the associated $N \times N$ Riemann matrix of $\boldsymbol{\beta}^o$-periods, $\tau^o = (\tau^o)_{i,j=1,\ldots,N} := (\oint_{\boldsymbol{\beta}_j^o} \omega_i^o)_{i,j=1,\ldots,N}$. The Riemann matrix, τ^o, is symmetric and pure imaginary, $-\mathrm{i}\tau^o$ is positive definite ($\mathrm{Im}(\tau_{ij}^o) > 0$), and $\det(\tau^o) \neq 0$. For the holomorphic Abelian differential (one-form) $\boldsymbol{\omega}^o$ defined above, choose a_{N+1}^o as the base point, and set $\boldsymbol{u}^o : \mathcal{Y}_o \to \mathrm{Jac}(\mathcal{Y}_o), z \mapsto \boldsymbol{u}^o(z) := \int_{a_{N+1}^o}^{z} \boldsymbol{\omega}^o$, where $\mathrm{Jac}(\mathcal{Y}_o) := \mathbb{C}^N / \{N + \tau^o M\}$, $(N, M) \in \mathbb{Z}^N \times \mathbb{Z}^N$, is the Jacobi variety of $\mathcal{Y}_o$, and the integration from a_{N+1}^o to z ($\in \mathcal{Y}_o$) is taken along any path on $\mathcal{Y}_o^+$.

Remark 2.2. From the representation $\omega_j^o = \sum_{k=1}^{N} \frac{c_{jk}^o z^{N-k}}{\sqrt{R_o(z)}}\,\mathrm{d}z$, $j = 1, \ldots, N$, and the normalisation condition $\oint_{\boldsymbol{\alpha}_k^o} \omega_j^o = \delta_{kj}$, $k, j = 1, \ldots, N$, one shows that c_{jk}^o, $k, j = 1, \ldots, N$, are obtained from

$$\begin{pmatrix} c_{11}^o & c_{12}^o & \cdots & c_{1N}^o \\ c_{21}^o & c_{22}^o & \cdots & c_{2N}^o \\ \vdots & \vdots & \ddots & \vdots \\ c_{N1}^o & c_{N2}^o & \cdots & c_{NN}^o \end{pmatrix} = \widetilde{\Xi}_o^{-1}, \tag{O1}$$

where

$$\widetilde{\Xi}_o := \begin{pmatrix} \oint_{\boldsymbol{\alpha}_1^o} \frac{\mathrm{d}s_1}{\sqrt{R_o(s_1)}} & \oint_{\boldsymbol{\alpha}_2^o} \frac{\mathrm{d}s_2}{\sqrt{R_o(s_2)}} & \cdots & \oint_{\boldsymbol{\alpha}_N^o} \frac{\mathrm{d}s_N}{\sqrt{R_o(s_N)}} \\ \oint_{\boldsymbol{\alpha}_1^o} \frac{s_1\mathrm{d}s_1}{\sqrt{R_o(s_1)}} & \oint_{\boldsymbol{\alpha}_2^o} \frac{s_2\mathrm{d}s_2}{\sqrt{R_o(s_2)}} & \cdots & \oint_{\boldsymbol{\alpha}_N^o} \frac{s_N\mathrm{d}s_N}{\sqrt{R_o(s_N)}} \\ \vdots & \vdots & \ddots & \vdots \\ \oint_{\boldsymbol{\alpha}_1^o} \frac{s_1^{N-1}\mathrm{d}s_1}{\sqrt{R_o(s_1)}} & \oint_{\boldsymbol{\alpha}_2^o} \frac{s_2^{N-1}\mathrm{d}s_2}{\sqrt{R_o(s_2)}} & \cdots & \oint_{\boldsymbol{\alpha}_N^o} \frac{s_N^{N-1}\mathrm{d}s_N}{\sqrt{R_o(s_N)}} \end{pmatrix}. \tag{O2}$$

For a (representation-independent) proof of the fact that $\det(\widetilde{\Xi}_o)\neq 0$, see, for example, Chapter 10, Section 10–2, of [68]. ■

Set [22]

$$\gamma^o(z):=\begin{cases}\left(\displaystyle\prod_{k=1}^{N+1}(z-b_{k-1}^o)(z-a_k^o)^{-1}\right)^{1/4}, & z\in\mathbb{C}_+,\\ -\mathrm{i}\left(\displaystyle\prod_{k=1}^{N+1}(z-b_{k-1}^o)(z-a_k^o)^{-1}\right)^{1/4}, & z\in\mathbb{C}_-.\end{cases}\tag{2.7}$$

It is shown in Section 4 of [22] that $\gamma^o(z)=_{\substack{z\to 0\\ z\in\mathcal{Y}_o^\pm}}(-\mathrm{i})^{(1\mp 1)/2}\gamma^o(0)(1+\mathcal{O}(z))$, where $\gamma^o(0):=(\prod_{k=1}^{N+1}b_{k-1}^o(a_k^o)^{-1})^{1/4}$ (>0), and that a set of N upper-edge and lower-edge finite-length-gap roots/zeros are

$$\left\{z_j^{o,\pm}\right\}_{j=1}^N=\left\{z^\pm\in\mathcal{Y}_o^\pm;\ ((\gamma^o(0))^{-1}\gamma^o(z)\mp\gamma^o(0)(\gamma^o(z))^{-1})|_{z=z^\pm}=0\right\},\tag{2.8}$$

with $z_j^{o,\pm}\in(a_j^o,b_j^o)^\pm$ $(\subset\mathcal{Y}_o^\pm)$, $j=1,\dots,N$, where, as points on the plane, $z_j^{o,+}=z_j^{o,-}:=z_j^o$, $j=1,\dots,N$ (on the plane, $z_j^o\in(a_j^o,b_j^o)$, $j=1,\dots,N$).

Corresponding to $\mathcal{Y}_o$, define $\boldsymbol{d}_o:=-\boldsymbol{K}_o-\sum_{j=1}^N\int_{a_{N+1}^o}^{z_j^{o,-}}\boldsymbol{\omega}^o$ $(\in\mathbb{C}^N)$, where $\boldsymbol{K}_o$ is the associated 'odd' vector of Riemann constants, and the integration from a_{N+1}^o to $z_j^{o,-}$, $j=1,\dots,N$, is taken along a fixed path in $\mathcal{Y}_o^-$. It is shown in Chapter VII of [69] that $\boldsymbol{K}_o=\sum_{j=1}^N\int_{a_j^o}^{a_{N+1}^o}\boldsymbol{\omega}^o$; furthermore, $\boldsymbol{K}_o$ is a point of order 2, that is, $2\boldsymbol{K}_o=0$ and $s\boldsymbol{K}_o\neq 0$ for $0<s<2$. Recalling the definition of $\boldsymbol{\omega}^o$ and that $z^{-(N+1)}(R_o(z))^{1/2}\sim_{\substack{z\to\infty\\ z\in\mathbb{C}_\pm}}\pm 1$, using the fact that $\boldsymbol{K}_o$ is a point of order 2, one arrives at

$$\begin{aligned}\boldsymbol{d}_o&=-\boldsymbol{K}_o-\sum_{j=1}^N\int_{a_{N+1}^o}^{z_j^{o,-}}\boldsymbol{\omega}^o=\boldsymbol{K}_o-\sum_{j=1}^N\int_{a_{N+1}^o}^{z_j^{o,-}}\boldsymbol{\omega}^o=-\boldsymbol{K}_o+\sum_{j=1}^N\int_{a_{N+1}^o}^{z_j^{o,+}}\boldsymbol{\omega}^o=\boldsymbol{K}_o+\sum_{j=1}^N\int_{a_{N+1}^o}^{z_j^{o,+}}\boldsymbol{\omega}^o\\ &=-\sum_{j=1}^N\int_{a_j^o}^{z_j^{o,-}}\boldsymbol{\omega}^o=\sum_{j=1}^N\int_{a_j^o}^{z_j^{o,+}}\boldsymbol{\omega}^o.\end{aligned}\tag{2.9}$$

Associated with the Riemann matrix of $\boldsymbol{\beta}^o$-periods, τ^o, is the 'odd' Riemann theta function:

$$\boldsymbol{\theta}(z;\tau^o)=:\boldsymbol{\theta}^o(z)=\sum_{m\in\mathbb{Z}^N}\mathrm{e}^{2\pi\mathrm{i}(m,z)+\pi\mathrm{i}(m,\tau^o m)},\quad z\in\mathbb{C}^N;\tag{2.10}$$

$\boldsymbol{\theta}^o(z)$ has the following evenness and (quasi-) periodicity properties,

$$\boldsymbol{\theta}^o(-z)=\boldsymbol{\theta}^o(z),\qquad \boldsymbol{\theta}^o(z+e_j)=\boldsymbol{\theta}^o(z),\qquad\text{and}\qquad \boldsymbol{\theta}^o(z\pm\tau_j^o)=\mathrm{e}^{\mp 2\pi\mathrm{i}z_j-\mathrm{i}\pi\tau_{jj}^o}\boldsymbol{\theta}^o(z),$$

where $\tau_j^o:=\tau^o e_j$ $(\in\mathbb{C}^N)$, $j=1,\dots,N$.

Some variational theory from [21] and [22] is now summarised. The following discussion is decomposed into two parts: one part corresponding to **RHP1** for $\overset{e}{Y}\colon\mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$, denoted as **P₁**; and the other part corresponding to **RHP2** for $\overset{o}{Y}\colon\mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$, denoted as **P₂**.

P₁ Let $\widetilde{V}\colon\mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfy conditions (1.20)–(1.22). Let $\mathrm{I}_V^e[\mu^e]\colon\mathcal{M}_1(\mathbb{R})\to\mathbb{R}$ denote the functional

$$\mathrm{I}_V^e[\mu^e]=\iint_{\mathbb{R}^2}\ln\left(\frac{|st|}{|s-t|^2}\right)\mathrm{d}\mu^e(s)\,\mathrm{d}\mu^e(t)+2\int_{\mathbb{R}}\widetilde{V}(s)\,\mathrm{d}\mu^e(s),\tag{2.11}$$

and consider the associated minimisation problem,

$$E_V^e=\inf\left\{\mathrm{I}_V^e[\mu^e];\ \mu^e\in\mathcal{M}_1(\mathbb{R})\right\}.$$

In [21], it is proven that the infimum is finite, and there is a unique measure μ_V^e $(\in\mathcal{M}_1(\mathbb{R}))$ achieving the infimum. The measure μ_V^e is referred to as the 'even' equilibrium measure. Furthermore, the following 'regularity' (see below) properties of μ_V^e are also established in [21]:

- the 'even' equilibrium measure has compact support which consists of the disjoint union of a finite number of bounded real intervals; in fact, as shown in [21], $\operatorname{supp}(\mu_V^e) =: J_e = \cup_{j=1}^{N+1}(b_{j-1}^e, a_j^e)$ $(\subset \mathbb{R}\setminus\{0\})$, where $\{b_{j-1}^e, a_j^e\}_{j=1}^{N+1}$, as described in item $\boxed{\sqrt{R_e(z)}}$ above, with $b_0^e := \min\{\operatorname{supp}(\mu_V^e)\} \notin \{-\infty, 0\}$, $a_{N+1}^e := \max\{\operatorname{supp}(\mu_V^e)\} \notin \{0, +\infty\}$, and $-\infty < b_0^e < a_1^e < b_1^e < a_2^e < \cdots < b_N^e < a_{N+1}^e < +\infty$, constitute the end-points of the support of the 'even' equilibrium measure;
- the end-points $\{b_{j-1}^e, a_j^e\}_{j=1}^{N+1}$ satisfy the following n-dependent and (locally) solvable system of $2(N+1)$ real moment conditions [21]:

$$\begin{aligned}
&\int_{J_e} \frac{(2s^{-1}+\widetilde{V}'(s))s^j}{(R_e(s))_+^{1/2}}\,\mathrm{d}s = 0, \quad j=0,\dots,N, \qquad \int_{J_e} \frac{(2s^{-1}+\widetilde{V}'(s))s^{N+1}}{(R_e(s))_+^{1/2}}\,\mathrm{d}s = -4\pi\mathrm{i},\\
&\int_{a_j^e}^{b_j^e}\left(\frac{\mathrm{i}(R_e(s))^{1/2}}{2\pi}\int_{J_e}\frac{(2\xi^{-1}+\widetilde{V}'(\xi))}{(R_e(\xi))_+^{1/2}(\xi-s)}\,\mathrm{d}\xi\right)\mathrm{d}s = \ln\left|\frac{a_j^e}{b_j^e}\right| + \frac{1}{2}\left(\widetilde{V}(a_j^e)-\widetilde{V}(b_j^e)\right), \quad j=1,\dots,N,
\end{aligned} \tag{2.12}$$

  where $'$ denotes differentiation with respect to the argument,

$$(R_e(z))^{1/2} := \left(\prod_{k=1}^{N+1}(z-b_{k-1}^e)(z-a_k^e)\right)^{1/2},$$

  $(R_e(x))_\pm^{1/2} := \lim_{\varepsilon\downarrow 0}(R_e(x\pm\mathrm{i}\varepsilon))^{1/2}$, and the branch of the square root is chosen so that $z^{-(N+1)}(R_e(z))^{1/2} \sim_{\substack{z\to\infty\\ z\in\mathbb{C}_\pm}} \pm 1$;
- the 'even' equilibrium measure is absolutely continuous with respect to Lebesgue measure. The *density* is given by

$$\mathrm{d}\mu_V^e(x) := \psi_V^e(x)\,\mathrm{d}x = \frac{1}{2\pi\mathrm{i}}(R_e(x))_+^{1/2}h_V^e(x)\mathbf{1}_{J_e}(x)\,\mathrm{d}x, \tag{2.13}$$

  where

$$h_V^e(z) = \frac{1}{2}\oint_{C_{\mathrm{R}}^e}\frac{(\frac{\mathrm{i}}{\pi s}+\frac{\mathrm{i}\widetilde{V}'(s)}{2\pi})}{\sqrt{R_e(s)}\,(s-z)}\,\mathrm{d}s \tag{2.14}$$

  (real analytic for $z\in\mathbb{R}\setminus\{0\}$), C_{R}^e $(\subset\mathbb{C}^*)$ is the union of two circular contours, one outer one of large radius $R^\natural$ traversed clockwise and one inner one of small radius $r^\natural$ traversed counter-clockwise, with the numbers $0 < r^\natural < R^\natural < +\infty$ chosen such that, for (any) non-real z in the domain of analyticity of $\widetilde{V}$ (that is, $\mathbb{C}^*$), $\operatorname{int}(C_{\mathrm{R}}^e) \supset J_e \cup \{z\}$, and $\mathbf{1}_{J_e}(x)$ denotes the indicator (characteristic) function of the set J_e. (Note that $\psi_V^e(x) \geqslant 0$ $\forall$ $x\in\overline{J_e} := \cup_{j=1}^{N+1}[b_{j-1}^e, a_j^e]$: it vanishes like a square root at the end-points of the support of the 'even' equilibrium measure, that is, $\psi_V^e(s) =_{s\downarrow b_{j-1}^e} O((s-b_{j-1}^e)^{1/2})$ and $\psi_V^e(s) =_{s\uparrow a_j^e} O((a_j^e-s)^{1/2})$, $j=1,\dots,N+1$.);
- the 'even' equilibrium measure and its (compact) support are (uniquely) characterised by the following Euler-Lagrange variational equations: there exists $\ell_e\in\mathbb{R}$, the 'even' Lagrange multiplier, and $\mu^e\in\mathcal{M}_1(\mathbb{R})$ such that

$$4\int_{J_e}\ln(|x-s|)\,\mathrm{d}\mu^e(s) - 2\ln|x| - \widetilde{V}(x) - \ell_e = 0, \quad x\in\overline{J_e}, \tag{$\mathbf{P}_1^{(a)}$}$$

$$4\int_{J_e}\ln(|x-s|)\,\mathrm{d}\mu^e(s) - 2\ln|x| - \widetilde{V}(x) - \ell_e \leqslant 0, \quad x\in\mathbb{R}\setminus\overline{J_e}; \tag{$\mathbf{P}_1^{(b)}$}$$

- the Euler-Lagrange variational equations can be conveniently recast in terms of the complex potential $g^e(z)$ of μ_V^e:

$$g^e(z) := \int_{J_e}\ln\left((z-s)^2(zs)^{-1}\right)\mathrm{d}\mu_V^e(s), \quad z\in\mathbb{C}\setminus(-\infty,\max\{0,a_{N+1}^e\}). \tag{2.15}$$

  The function $g^e:\mathbb{C}\setminus(-\infty,\max\{0,a_{N+1}^e\})\to\mathbb{C}$ so defined satisfies:

($\mathbf{P_1^{(1)}}$) $g^e(z)$ is analytic for $z\in\mathbb{C}\setminus(-\infty,\max\{0,a^e_{N+1}\})$;

($\mathbf{P_1^{(2)}}$) $g^e(z)=_{\substack{z\to\infty\\ z\in\mathbb{C}\setminus\mathbb{R}}}\ln(z)+O(1)$;

($\mathbf{P_1^{(3)}}$) $g^e_+(z)+g^e_-(z)-\widetilde{V}(z)-\ell_e+2Q_e=0$, $z\in\overline{J_e}$, where $g^e_\pm(z):=\lim_{\varepsilon\downarrow 0}g^e(z\pm\mathrm{i}\varepsilon)$, and

$$Q_e:=\int_{J_e}\ln(s)\,\mathrm{d}\mu^e_V(s)=\int_{J_e}\ln(|s|)\,\mathrm{d}\mu^e_V(s)+\mathrm{i}\pi\int_{J_e\cap\mathbb{R}_-}\mathrm{d}\mu^e_V(s); \tag{2.16}$$

($\mathbf{P_1^{(4)}}$) $g^e_+(z)+g^e_-(z)-\widetilde{V}(z)-\ell_e+2Q_e\leqslant 0$, $z\in\mathbb{R}\setminus\overline{J_e}$, where equality holds for at most a finite number of points;

($\mathbf{P_1^{(5)}}$) $g^e_+(z)-g^e_-(z)=\mathrm{i}f^{\mathbb{R}}_{g^e}(z)$, $z\in\mathbb{R}$, where $f^{\mathbb{R}}_{g^e}\colon\mathbb{R}\to\mathbb{R}$, and, in particular, $g^e_+(z)-g^e_-(z)=\mathrm{i}\,\mathrm{const.}$, $z\in\mathbb{R}\setminus\overline{J_e}$, with $\mathrm{const.}\in\mathbb{R}$;

($\mathbf{P_1^{(6)}}$) $\mathrm{i}(g^e_+(z)-g^e_-(z))'\geqslant 0$, $z\in J_e$, where equality holds for at most a finite number of points.

$\mathbf{P_2}$ Let $\widetilde{V}\colon\mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfy conditions (1.20)–(1.22). Let $\mathrm{I}^o_V[\mu^o]\colon\mathcal{M}_1(\mathbb{R})\to\mathbb{R}$ denote the functional

$$\mathrm{I}^o_V[\mu^o]=\iint_{\mathbb{R}^2}\ln\left(\frac{|st|}{|s-t|^{2+\frac{1}{n}}}\right)\mathrm{d}\mu^o(s)\,\mathrm{d}\mu^o(t)+2\int_{\mathbb{R}}\widetilde{V}(s)\,\mathrm{d}\mu^o(s),\quad n\in\mathbb{N}, \tag{2.17}$$

and consider the associated minimisation problem,

$$E^o_V=\inf\left\{\mathrm{I}^o_V[\mu^o];\ \mu^o\in\mathcal{M}_1(\mathbb{R})\right\}.$$

In [22], it is proven that the infimum is finite, and there is a unique measure μ^o_V ($\in\mathcal{M}_1(\mathbb{R})$) achieving the infimum. The measure μ^o_V is referred to as the 'odd' equilibrium measure. Furthermore, the following 'regularity' (see below) properties of μ^o_V are also established in [22]:

- the 'odd' equilibrium measure has compact support which consists of the disjoint union of a finite number of bounded real intervals; in fact, as shown in [22], $\mathrm{supp}(\mu^o_V)=:J_o=\cup_{j=1}^{N+1}(b^o_{j-1},a^o_j)$ ($\subset\mathbb{R}\setminus\{0\}$), where $\{b^o_{j-1},a^o_j\}_{j=1}^{N+1}$, as described in item $\boxed{\sqrt{R_o(z)}}$ above, with $b^o_0:=\min\{\mathrm{supp}(\mu^o_V)\}\notin\{-\infty,0\}$, $a^o_{N+1}:=\max\{\mathrm{supp}(\mu^o_V)\}\notin\{0,+\infty\}$, and $-\infty<b^o_0<a^o_1<b^o_1<a^o_2<\cdots<b^o_N<a^o_{N+1}<+\infty$, constitute the end-points of the support of the 'odd' equilibrium measure; (The number of intervals, $N+1$, is the same in the 'odd' case as in the 'even' case, which can be established by a lengthy analysis similar to that contained in [38].)
- the end-points $\{b^o_{j-1},a^o_j\}_{j=1}^{N+1}$ satisfy the following n-dependent and (locally) solvable system of $2(N+1)$ real moment conditions [22]:

$$\int_{J_o}\frac{(2s^{-1}+\widetilde{V}'(s))s^j}{(R_o(s))^{1/2}_+}\,\mathrm{d}s=0,\quad j=0,\dots,N,\qquad \int_{J_o}\frac{(2s^{-1}+\widetilde{V}'(s))s^{N+1}}{(R_o(s))^{1/2}_+}\,\frac{\mathrm{d}s}{2\pi\mathrm{i}}=-\left(2+\frac{1}{n}\right),$$
$$\int_{a^o_j}^{b^o_j}\left(\frac{\mathrm{i}(R_o(s))^{1/2}}{2\pi}\int_{J_o}\frac{(2\xi^{-1}+\widetilde{V}'(\xi))}{(R_o(\xi))^{1/2}_+(\xi-s)}\,\mathrm{d}\xi\right)\mathrm{d}s=\ln\left|\frac{a^o_j}{b^o_j}\right|+\frac{1}{2}\left(\widetilde{V}(a^o_j)-\widetilde{V}(b^o_j)\right),\quad j=1,\dots,N, \tag{2.18}$$

where

$$(R_o(z))^{1/2}:=\left(\prod_{k=1}^{N+1}(z-b^o_{k-1})(z-a^o_k)\right)^{1/2},$$

$(R_o(x))^{1/2}_\pm:=\lim_{\varepsilon\downarrow 0}(R_o(x\pm\mathrm{i}\varepsilon))^{1/2}$, and the branch of the square root is chosen so that $z^{-(N+1)}(R_o(z))^{1/2}\sim_{\substack{z\to\infty\\ z\in\mathbb{C}_\pm}}\pm 1$;

- the 'odd' equilibrium measure is absolutely continuous with respect to Lebesgue measure. The density is given by

$$\mathrm{d}\mu^o_V(x):=\psi^o_V(x)\,\mathrm{d}x=\frac{1}{2\pi\mathrm{i}}(R_o(x))^{1/2}_+h^o_V(x)\mathbf{1}_{J_o}(x)\,\mathrm{d}x, \tag{2.19}$$

where

$$h_V^o(z)=\left(2+\frac{1}{n}\right)^{-1}\oint_{C_{\mathrm{R}}^o}\frac{(\frac{\mathrm{i}}{\pi s}+\frac{\mathrm{i}\widetilde{V}'(s)}{2\pi})}{\sqrt{R_o(s)}\,(s-z)}\,\mathrm{d}s \tag{2.20}$$

(real analytic for $z\in\mathbb{R}\setminus\{0\}$), C_{R}^o ($\subset\mathbb{C}^*$) is the union of two circular contours, one outer one of large radius $R^\flat$ traversed clockwise and one inner one of small radius $r^\flat$ traversed counter-clockwise, with the numbers $0<r^\flat<R^\flat<+\infty$ chosen such that, for (any) non-real z in the domain of analyticity of $\widetilde{V}$ (that is, $\mathbb{C}^*$), $\mathrm{int}(C_{\mathrm{R}}^o)\supset J_o\cup\{z\}$, and $\mathbf{1}_{J_o}(x)$ denotes the indicator (characteristic) function of the set J_o. (Note that $\psi_V^o(x)\geqslant 0\ \forall\ x\in\overline{J_o}:=\cup_{j=1}^{N+1}[b_{j-1}^o,a_j^o]$: it vanishes like a square root at the end-points of the support of the 'odd' equilibrium measure, that is, $\psi_V^o(s)=_{s\downarrow b_{j-1}^o}O((s-b_{j-1}^o)^{1/2})$ and $\psi_V^o(s)=_{s\uparrow a_j^o}O((a_j^o-s)^{1/2})$, $j=1,\dots,N+1$.);

- the 'odd' equilibrium measure and its (compact) support are (uniquely) characterised by the following Euler-Lagrange variational equations: there exists $\ell_o\in\mathbb{R}$, the 'odd' Lagrange multiplier, and $\mu^o\in\mathcal{M}_1(\mathbb{R})$ such that

$$2\left(2+\frac{1}{n}\right)\int_{J_o}\ln(|x-s|)\,\mathrm{d}\mu^o(s)-2\ln|x|-\widetilde{V}(x)-\ell_o-2\left(2+\frac{1}{n}\right)\widetilde{Q}_o=0,\quad x\in\overline{J_o},\qquad(\mathbf{P_2^{(a)}})$$

$$2\left(2+\frac{1}{n}\right)\int_{J_o}\ln(|x-s|)\,\mathrm{d}\mu^o(s)-2\ln|x|-\widetilde{V}(x)-\ell_o-2\left(2+\frac{1}{n}\right)\widetilde{Q}_o\leqslant 0,\quad x\in\mathbb{R}\setminus\overline{J_o},\qquad(\mathbf{P_2^{(b)}})$$

where $\widetilde{Q}_o:=\int_{J_o}\ln(|s|)\,\mathrm{d}\mu^o(s)$;

- the Euler-Lagrange variational equations can be conveniently recast in terms of the complex potential $g^o(z)$ of μ_V^o:

$$g^o(z):=\int_{J_o}\ln\left((z-s)^{2+\frac{1}{n}}(zs)^{-1}\right)\mathrm{d}\mu_V^o(s),\quad z\in\mathbb{C}\setminus(-\infty,\max\{0,a_{N+1}^o\}).\tag{2.21}$$

The function $g^o:\mathbb{C}\setminus(-\infty,\max\{0,a_{N+1}^o\})\to\mathbb{C}$ so defined satisfies:

$(\mathbf{P_2^{(1)}})$ $g^o(z)$ is analytic for $z\in\mathbb{C}\setminus(-\infty,\max\{0,a_{N+1}^o\})$;

$(\mathbf{P_2^{(2)}})$ $g^o(z)=_{\substack{z\to 0\\ z\in\mathbb{C}\setminus\mathbb{R}}}-\ln(z)+O(1)$;

$(\mathbf{P_2^{(3)}})$ $g_+^o(z)+g_-^o(z)-\widetilde{V}(z)-\ell_o-\mathfrak{Q}_{\mathcal{A}}^+-\mathfrak{Q}_{\mathcal{A}}^-=0$, $z\in\overline{J_o}$, where $g_\pm^o(z):=\lim_{\varepsilon\downarrow 0}g^o(z\pm\mathrm{i}\varepsilon)$, and

$$\mathfrak{Q}_{\mathcal{A}}^\pm:=\left(1+\frac{1}{n}\right)\int_{J_o}\ln(|s|)\,\mathrm{d}\mu_V^o(s)-\mathrm{i}\pi\int_{J_o\cap\mathbb{R}_-}\mathrm{d}\mu_V^o(s)\pm\mathrm{i}\pi\left(2+\frac{1}{n}\right)\int_{J_o\cap\mathbb{R}_+}\mathrm{d}\mu_V^o(s);\tag{2.22}$$

$(\mathbf{P_2^{(4)}})$ $g_+^o(z)+g_-^o(z)-\widetilde{V}(z)-\ell_o-\mathfrak{Q}_{\mathcal{A}}^+-\mathfrak{Q}_{\mathcal{A}}^-\leqslant 0$, $z\in\mathbb{R}\setminus\overline{J_o}$, where equality holds for at most a finite number of points;

$(\mathbf{P_2^{(5)}})$ $g_+^o(z)-g_-^o(z)-\mathfrak{Q}_{\mathcal{A}}^++\mathfrak{Q}_{\mathcal{A}}^-=\mathrm{i}f_{g^o}^{\mathbb{R}}(z)$, $z\in\mathbb{R}$, where $f_{g^o}^{\mathbb{R}}:\mathbb{R}\to\mathbb{R}$, and, in particular, $g_+^o(z)-g_-^o(z)-\mathfrak{Q}_{\mathcal{A}}^++\mathfrak{Q}_{\mathcal{A}}^-=\mathrm{i}\,\mathrm{const.}$, $z\in\mathbb{R}\setminus\overline{J_o}$, with $\mathrm{const.}\in\mathbb{R}$;

$(\mathbf{P_2^{(6)}})$ $\mathrm{i}(g_+^o(z)-g_-^o(z)-\mathfrak{Q}_{\mathcal{A}}^++\mathfrak{Q}_{\mathcal{A}}^-)'\geqslant 0$, $z\in J_o$, where equality holds for at most a finite number of points.

In this work, as in [21, 22], the so-called 'regular case' is considered, namely:

- $\mathrm{d}\mu_V^e$, or $\widetilde{V}:\mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfying conditions (1.20)–(1.22), is *regular* if

$$h_V^e(x)\not\equiv 0\quad\text{on}\quad\overline{J_e},$$

$$4\int_{J_e}\ln(|x-s|)\,\mathrm{d}\mu_V^e(s)-2\ln|x|-\widetilde{V}(x)-\ell_e<0,\quad x\in\mathbb{R}\setminus\overline{J_e},$$

and inequalities **$(\mathbf{P_1^{(4)}})$** and **$(\mathbf{P_1^{(6)}})$** in **$\mathbf{P_1}$** are strict, that is, $\leqslant$ (resp., $\geqslant$) is replaced by $<$ (resp., $>$);

- $\mathrm{d}\mu_V^o$, or $\widetilde{V}\colon \mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfying conditions (1.20)–(1.22), is regular if

$$h_V^o(x)\not\equiv 0 \quad \text{on} \quad \overline{J_o},$$
$$2\left(2+\frac{1}{n}\right)\int_{J_o}\ln(|x-s|)\,\mathrm{d}\mu_V^o(s)-2\ln|x|-\widetilde{V}(x)-\ell_o-2\left(2+\frac{1}{n}\right)Q_o<0, \quad x\in\mathbb{R}\setminus\overline{J_o},$$

  where $Q_o:=\int_{J_o}\ln(|s|)\,\mathrm{d}\mu_V^o(s)$, and inequalities $\mathbf{(P_2^{(4)})}$ and $\mathbf{(P_2^{(6)})}$ in $\boxed{\mathbf{P_2}}$ are strict, that is, $\leqslant$ (resp., $\geqslant$) is replaced by $<$ (resp., $>$)[4].

The (density of the) 'even' and 'odd' equilibrium measures, $\mathrm{d}\mu_V^e$ and $\mathrm{d}\mu_V^o$, respectively, together with the variational problems, emerge naturally in the asymptotic analysis of **RHP1** and **RHP2**.

Remark 2.3. The following correspondences should be noted:

- $g^e\colon \mathbb{C}\setminus(-\infty,\max\{0,a_{N+1}^e\})\to\mathbb{C}$ solves the phase conditions $\mathbf{(P_1^{(1)})}$—$\mathbf{(P_1^{(6)})}\Leftrightarrow \mu_V^e\in\mathcal{M}_1(\mathbb{R})$ solves the variational conditions $(\mathbf{P_1^{(a)}})$ and $(\mathbf{P_1^{(b)}})$;
- $g^o\colon \mathbb{C}\setminus(-\infty,\max\{0,a_{N+1}^o\})\to\mathbb{C}$ solves the phase conditions $\mathbf{(P_2^{(1)})}$—$\mathbf{(P_2^{(6)})}\Leftrightarrow \mu_V^o\in\mathcal{M}_1(\mathbb{R})$ solves the variational conditions $(\mathbf{P_2^{(a)}})$ and $(\mathbf{P_2^{(b)}})$. ■

3 Asymptotics for the Root Products, Recurrence Relation Coefficients, and Hankel Determinant Ratios of OLPs

In this section, asymptotics in the double-scaling limit as $\mathcal{N},n\to\infty$ such that $z_o=1+o(1)$, denoted, hereafter, via the simplified 'notation' $n\to\infty$, for the following quantities are obtained:

- the products of the (real) roots of the L-polynomials, for both the non-singular and singular cases (cf. Equations (1.10)–(1.13));
- the Hankel determinant ratios associated with the bi-infinite, real-valued, strong moment sequence $\left\{c_k=\int_{\mathbb{R}}s^k\exp(-n\widetilde{V}(s))\,\mathrm{d}s,\ n\in\mathbb{N}\right\}_{k\in\mathbb{Z}}$ (cf. Equations (1.9), (1.18), and (1.19));
- the (real-valued) recurrence relation coefficients (cf. Equations (1.23)–(1.26)), and subsequently those of the (real-symmetric, tri-penta-diagonal-type) Laurent-Jacobi matrices $\mathcal{F}$ and $\mathcal{G}$ (cf. Equations (1.27) and (1.28)).

Now that the relevant auxiliary information, functions, and results have been given in Section 2, one turns to the large-n asymptotic description of the solutions to **RHP1** and **RHP2**. As mentioned in Sections 1 and 2, this asymptotic analysis was the subject of [21] (for **RHP1**) and [22] (for **RHP2**). In this manuscript, all that is necessary from the results of [21, 22] are the large-n asymptotic behaviours of the n-dependent coefficients of the following asymptotic expansions for $\overset{e}{\mathrm{Y}}(z)$ and $\overset{o}{\mathrm{Y}}(z)$:

$$\overset{e}{\mathrm{Y}}(z)z^{-n\sigma_3}\underset{\substack{z\to\infty\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=}\mathrm{I}+\frac{1}{z}\mathrm{Y}_1^{e,\infty}+\frac{1}{z^2}\mathrm{Y}_2^{e,\infty}+O\left(\frac{1}{z^3}\right), \tag{3.1}$$

$$\overset{e}{\mathrm{Y}}(z)z^{n\sigma_3}\underset{\substack{z\to 0\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=}\mathrm{Y}_0^{e,0}+z\mathrm{Y}_1^{e,0}+z^2\mathrm{Y}_2^{e,0}+O(z^3), \tag{3.2}$$

[4]There are three distinct situations in which these conditions may fail: (i) for at least one $\widetilde{x}_e\in\mathbb{R}\setminus\overline{J_e}$ (resp., $\widetilde{x}_o\in\mathbb{R}\setminus\overline{J_o}$), $4\int_{J_e}\ln(|\widetilde{x}_e-s|)\,\mathrm{d}\mu_V^e(s)-2\ln|\widetilde{x}_e|-\widetilde{V}(\widetilde{x}_e)-\ell_e=0$ (resp., $2(2+\frac{1}{n})\int_{J_o}\ln(|\widetilde{x}_o-s|)\,\mathrm{d}\mu_V^o(s)-2\ln|\widetilde{x}_o|-\widetilde{V}(\widetilde{x}_o)-\ell_o-2(2+\frac{1}{n})Q_o=0$), that is, for n even (resp., n odd) equality is attained for at least one point $\widetilde{x}_e$ (resp., $\widetilde{x}_o$) in the complement of the closure of the support of the 'even' (resp., 'odd') equilibrium measure μ_V^e (resp., μ_V^o), which corresponds to the situation in which a 'band' has just closed, or is about to open, about $\widetilde{x}_e$ (resp., $\widetilde{x}_o$); (ii) for at least one $\widehat{x}_e$ (resp., $\widehat{x}_o$), $h_V^e(\widehat{x}_e)=0$ (resp., $h_V^o(\widehat{x}_o)=0$), that is, for n even (resp., n odd) the function h_V^e (resp., h_V^o) vanishes for at least one point $\widehat{x}_e$ (resp., $\widehat{x}_o$) within the support of the 'even' (resp., 'odd') equilibrium measure μ_V^e (resp., μ_V^o), which corresponds to the situation in which a 'gap' is about to open, or close, about $\widehat{x}_e$ (resp., $\widehat{x}_o$); and (iii) there exists at least one $j\in\{1,\dots,N+1\}$, denoted j_e (resp., j_o), such that $h_V^e(b_{j_e-1}^e)=0$ and/or $h_V^e(a_{j_e}^e)=0$ (resp., $h_V^o(b_{j_o-1}^o)=0$ and/or $h_V^o(a_{j_o}^o)=0$). Each of these three cases can occur only a finite number of times due to the fact that $\widetilde{V}\colon\mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfies conditions (1.20)–(1.22) [38, 40].

$$\overset{o}{\mathrm{Y}}(z)z^{n\sigma_3} \underset{\substack{z\to 0\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=} \mathrm{I}+z\mathrm{Y}_1^{o,0}+z^2\mathrm{Y}_2^{o,0}+O(z^3), \tag{3.3}$$

$$\overset{o}{\mathrm{Y}}(z)z^{-(n+1)\sigma_3} \underset{\substack{z\to \infty\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=} \mathrm{Y}_0^{o,\infty}+\frac{1}{z}\mathrm{Y}_1^{o,\infty}+\frac{1}{z^2}\mathrm{Y}_2^{o,\infty}+O\left(\frac{1}{z^3}\right), \tag{3.4}$$

where, for the convenience of the presentation which follows, as well as that of the reader's, large-n asymptotics of $\mathrm{Y}_j^{e,\infty}$, $j=1,2$, and $\mathrm{Y}_k^{e,0}$, $k=0,1,2$, are given in Appendix A (Theorems A.1 and A.2, respectively), and large-n asymptotics of $\mathrm{Y}_j^{o,0}$, $j=1,2$, and $\mathrm{Y}_k^{o,\infty}$, $k=0,1,2$, are given in Appendix B (Theorems B.1 and B.2, respectively). Furthermore, large-n asymptotics of the 'even' and 'odd' leading coefficients $\xi_n^{(2n)}$ and $\xi_{-n-1}^{(2n+1)}$, respectively, are given in Appendix A (Theorem A.3) and Appendix B (Theorem B.3).

The underlying idea of the following analysis is relatively straightforward: the quantities of interest, namely, the coefficients in the (system of) three- and five-term recurrence relations, Hankel determinant ratios, etc., are all expressible in terms of the n-dependent coefficients of asymptotic expansions (3.1)–(3.4), for which rather complete large-n asymptotics are available.

Remark 3.1. It turns out that (cf. Equations (1.29) and (1.30)) only the (1 1)- and (1 2)-elements of asymptotics (3.1)–(3.4) are necessary for obtaining the results of this work. In Lemma 3.1 (see below), and Propositions 3.1–3.3 (see below), a series of formulae are presented which provide explicit relations between the quantities of interest and the (n-dependent) coefficients in the asymptotic expansions (3.1)–(3.4). The statements, and proofs, are somewhat lengthy; but the important point should not be missed, namely: all quantities of interest are expressed, via straightforward algebraic calculations, in terms of the above asymptotic expansions. ■

Lemma 3.1. *Let the external field* $\widetilde{V}\colon \mathbb{R}\setminus\{0\}\to\mathbb{R}$ *be regular and satisfy conditions* (1.20)–(1.22). *Let the orthonormal L-polynomials (resp., monic orthogonal L-polynomials),* $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ *(resp.,* $\{\boldsymbol{\pi}_k(z)\}_{k\in\mathbb{Z}_0^+}$*) be as defined in Equations* (1.2) *and* (1.3) *(resp., Equations* (1.4) *and* (1.5)), *and let* $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ *satisfy the system of recurrence relations* (1.23)–(1.26). *Let* $\overset{e}{\mathrm{Y}}\colon \mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$ *(resp.,* $\overset{o}{\mathrm{Y}}\colon \mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$*) be the (unique) solution of* **RHP1** *(resp.,* **RHP2**) *with integral representation* (1.29) *(resp.,* (1.30)). *Then,*

$$\begin{aligned} z^n\overset{e}{\mathrm{Y}}_{12}(z) \underset{\substack{z\to\infty\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=} \frac{1}{z}\Bigg(-\frac{(\xi_n^{(2n)})^{-2}}{2\pi\mathrm{i}}\Bigg)+\frac{1}{z^2}\Bigg(-\frac{a_{2n}^\sharp(\xi_n^{(2n)})^{-2}}{2\pi\mathrm{i}}+\frac{(\xi_n^{(2n)})^{-1}}{2\pi\mathrm{i}}\Bigg(\frac{b_{2n}^\sharp\nu_{-n}^{(2n)}}{\xi_{-n}^{(2n-1)}} \\ +\frac{c_{2n}^\sharp}{\xi_{n-1}^{(2n-2)}}\Bigg(\frac{\xi_{n-1}^{(2n)}}{\xi_n^{(2n)}}-\frac{\nu_{-n}^{(2n)}\xi_{n-1}^{(2n-1)}}{\xi_{-n}^{(2n-1)}}\Bigg)\Bigg)\Bigg)+O\left(\frac{1}{z^3}\right), \end{aligned} \tag{3.5}$$

$$\begin{aligned} z^{-n}\overset{e}{\mathrm{Y}}_{12}(z) \underset{\substack{z\to 0\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=} \frac{1}{2\pi\mathrm{i}}\frac{\beta_{2n}^\sharp}{\xi_n^{(2n)}\xi_{-n}^{(2n-1)}}+z\Bigg(\frac{(\xi_n^{(2n)})^{-1}}{2\pi\mathrm{i}}\Bigg(\frac{\beta_{2n+1}^\sharp}{\xi_{-n-1}^{(2n+1)}}-\frac{\alpha_{2n}^\sharp\nu_n^{(2n+1)}}{\xi_n^{(2n)}} \\ +\frac{\beta_{2n}^\sharp}{\xi_{-n}^{(2n-1)}}\Bigg(\nu_n^{(2n+1)}\nu_{-n}^{(2n)}-\frac{\xi_{-n}^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}}\Bigg)\Bigg)\Bigg)+O(z^2), \end{aligned} \tag{3.6}$$

and

$$\begin{aligned} z^{-n}\overset{o}{\mathrm{Y}}_{12}(z) \underset{\substack{z\to 0\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=} z\Bigg(\frac{(\xi_{-n-1}^{(2n+1)})^{-2}}{2\pi\mathrm{i}}\Bigg)+z^2\Bigg(\frac{\alpha_{2n+1}^\sharp(\xi_{-n-1}^{(2n+1)})^{-2}}{2\pi\mathrm{i}}-\frac{(\xi_{-n-1}^{(2n+1)})^{-1}}{2\pi\mathrm{i}}\Bigg(\frac{\beta_{2n+1}^\sharp\nu_n^{(2n+1)}}{\xi_n^{(2n)}} \\ +\frac{\gamma_{2n+1}^\sharp}{\xi_{-n}^{(2n-1)}}\Bigg(\frac{\xi_{-n}^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}}-\nu_n^{(2n+1)}\nu_{-n}^{(2n)}\Bigg)\Bigg)\Bigg)+O(z^3), \end{aligned} \tag{3.7}$$

$$\begin{aligned} z^{n+1}\overset{o}{\mathrm{Y}}_{12}(z) \underset{\substack{z\to\infty\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=} -\frac{1}{2\pi\mathrm{i}}\frac{b_{2n+1}^\sharp}{\xi_{-n-1}^{(2n+1)}\xi_n^{(2n)}}+\frac{1}{z}\Bigg(\frac{(\xi_{-n-1}^{(2n+1)})^{-1}}{2\pi\mathrm{i}}\Bigg(\frac{a_{2n+1}^\sharp}{\xi_{-n-1}^{(2n+1)}}\frac{\xi_{-(n+1)}^{(2n+2)}}{\xi_{n+1}^{(2n+2)}}-\frac{b_{2n+2}^\sharp}{\xi_{n+1}^{(2n+2)}} \\ +\frac{b_{2n+1}^\sharp}{\xi_n^{(2n)}}\Bigg(\frac{\xi_n^{(2n+2)}}{\xi_{n+1}^{(2n+2)}}-\nu_n^{(2n+1)}\frac{\xi_{-(n+1)}^{(2n+2)}}{\xi_{n+1}^{(2n+2)}}\Bigg)\Bigg)\Bigg)+O\left(\frac{1}{z^2}\right). \end{aligned} \tag{3.8}$$

Proof. Noting from Lemma 1.1, Equation (1.29) that

$$\boldsymbol{\pi}_{2n}(z) := \overset{e}{\mathrm{Y}}_{11}(z) \qquad \text{and} \qquad \overset{e}{\mathrm{Y}}_{12}(z) = \int_{\mathbb{R}} \frac{\boldsymbol{\pi}_{2n}(s)\exp(-n\widetilde{V}(s))}{s-z}\,\frac{\mathrm{d}s}{2\pi\mathrm{i}}, \quad z \in \mathbb{C}\setminus\mathbb{R},$$

and recalling that, for $\widetilde{V}\colon \mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfying conditions (1.20)–(1.22) and the regularity assumption, $\int_{\mathbb{R}} s^k \exp(-n\widetilde{V}(s))\,\mathrm{d}s < \infty$, $(k,n)\in\mathbb{Z}\times\mathbb{N}$, one uses the expansion $\frac{1}{s-z} = -\sum_{j=0}^{m}\frac{s^j}{z^{j+1}} + \frac{s^{m+1}}{z^{m+1}(s-z)}$, $m\in\mathbb{Z}_0^+$, for $(\mathbb{C}\setminus\mathbb{R}\ni)\ z\to\infty$, and $\frac{1}{z-s} = -\sum_{j=0}^{m}\frac{z^j}{s^{j+1}} + \frac{z^{m+1}}{s^{m+1}(z-s)}$ for $(\mathbb{C}\setminus\mathbb{R}\ni)\ z\to 0$, and proceeds exactly as in the proof of Proposition 5.4 of [21]. The proof of expansions (3.5) and (3.6) is now completed upon using, repeatedly, the orthogonality relations $\langle \boldsymbol{\pi}_{2n}, z^j\rangle_{\mathcal{L}} = 0$, $j = -n,\dots,n-1$, $\langle \boldsymbol{\pi}_{2n+1}, z^i\rangle_{\mathcal{L}} = 0$, $i = -n,\dots,n$, $\langle \boldsymbol{\pi}_{2n}, \boldsymbol{\pi}_{2n}\rangle_{\mathcal{L}} = (\xi_n^{(2n)})^{-2}$, and $\langle \boldsymbol{\pi}_{2n+1}, \boldsymbol{\pi}_{2n+1}\rangle_{\mathcal{L}} = (\xi_{-n-1}^{(2n+1)})^{-2}$, as well as Equations (1.2)–(1.5), the recurrence relations (1.23)–(1.26), and (the (1 2)-elements of) asymptotic expansions (3.1) and (3.2).

Mutatis mutandis for expansions (3.7) and (3.8); but, in the latter case, one begins with Lemma 1.2, Equation (1.30), that is,

$$z\boldsymbol{\pi}_{2n+1}(z) := \overset{o}{\mathrm{Y}}_{11}(z) \qquad \text{and} \qquad \overset{o}{\mathrm{Y}}_{12}(z) = z\int_{\mathbb{R}} \frac{(s\boldsymbol{\pi}_{2n+1}(s))\exp(-n\widetilde{V}(s))}{s(s-z)}\,\frac{\mathrm{d}s}{2\pi\mathrm{i}}, \quad z \in \mathbb{C}\setminus\mathbb{R},$$

uses (the (1 2)-elements of) asymptotic expansions (3.3) and (3.4), and proceeds exactly as in the proof of Proposition 5.4 of [22]. □

Proposition 3.1. *Let the external field $\widetilde{V}\colon \mathbb{R}\setminus\{0\}\to\mathbb{R}$ be regular and satisfy conditions* (1.20)–(1.22). *Let the orthonormal L-polynomials (resp., monic orthogonal L-polynomials), $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ (resp., $\{\boldsymbol{\pi}_k(z)\}_{k\in\mathbb{Z}_0^+}$) be as defined in Equations* (1.2) *and* (1.3) *(resp., Equations* (1.4) *and* (1.5)), *and let $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ satisfy the system of recurrence relations* (1.23)–(1.26). *Let $\overset{e}{\mathrm{Y}}\colon \mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$ (resp., $\overset{o}{\mathrm{Y}}\colon \mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$) be the (unique) solution of* **RHP1** *(resp.,* **RHP2**) *with integral representation* (1.29) *(resp.,* (1.30)). *Then a set of relations between the coefficients of asymptotic expansions* (3.1)–(3.4), *the 'even' and 'odd' norming constants, and the recurrence coefficients is:*

$$\xi_n^{(2n)} = \frac{1}{(-2\pi\mathrm{i}(\mathrm{Y}_1^{e,\infty})_{12})^{1/2}} \qquad (>0), \tag{3.9}$$

$$2\pi\mathrm{i}(\mathrm{Y}_0^{e,0})_{12} = \frac{\beta_{2n}^{\sharp}}{\xi_n^{(2n)}\xi_{-n}^{(2n-1)}}, \tag{3.10}$$

$$\xi_{-n-1}^{(2n+1)} = \frac{1}{(2\pi\mathrm{i}(\mathrm{Y}_1^{o,0})_{12})^{1/2}} \qquad (>0), \tag{3.11}$$

$$-2\pi\mathrm{i}(\mathrm{Y}_0^{o,\infty})_{12} = \frac{b_{2n+1}^{\sharp}}{\xi_{-n-1}^{(2n+1)}\xi_n^{(2n)}}, \tag{3.12}$$

$$-2\pi\mathrm{i}(\mathrm{Y}_2^{e,\infty})_{12} = \frac{a_{2n}^{\sharp}}{(\xi_n^{(2n)})^2} - \frac{1}{\xi_n^{(2n)}}\left(\frac{b_{2n}^{\sharp}\nu_{-n}^{(2n)}}{\xi_{-n}^{(2n-1)}} + \frac{c_{2n}^{\sharp}}{\xi_{n-1}^{(2n-2)}}\left(\frac{\xi_{n-1}^{(2n)}}{\xi_n^{(2n)}} - \nu_{-n}^{(2n)}\frac{\xi_{n-1}^{(2n-1)}}{\xi_{-n}^{(2n-1)}}\right)\right), \tag{3.13}$$

$$2\pi\mathrm{i}(\mathrm{Y}_1^{e,0})_{12} = \frac{1}{\xi_n^{(2n)}}\left(\frac{\beta_{2n+1}^{\sharp}}{\xi_{-n-1}^{(2n+1)}} - \frac{\alpha_{2n}^{\sharp}\nu_n^{(2n+1)}}{\xi_n^{(2n)}} + \frac{\beta_{2n}^{\sharp}}{\xi_{-n}^{(2n-1)}}\left(\nu_n^{(2n+1)}\nu_{-n}^{(2n)} - \frac{\xi_{-n}^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}}\right)\right), \tag{3.14}$$

$$2\pi\mathrm{i}(\mathrm{Y}_2^{o,0})_{12} = \frac{\alpha_{2n+1}^{\sharp}}{(\xi_{-n-1}^{(2n+1)})^2} - \frac{1}{\xi_{-n-1}^{(2n+1)}}\left(\frac{\beta_{2n+1}^{\sharp}\nu_n^{(2n+1)}}{\xi_n^{(2n)}} + \frac{\gamma_{2n+1}^{\sharp}}{\xi_{-n}^{(2n-1)}}\left(\frac{\xi_{-n}^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}} - \nu_n^{(2n+1)}\nu_{-n}^{(2n)}\right)\right), \tag{3.15}$$

$$-2\pi\mathrm{i}(\mathrm{Y}_1^{o,\infty})_{12} = \frac{1}{\xi_{-n-1}^{(2n+1)}}\left(\frac{b_{2n+2}^{\sharp}}{\xi_{n+1}^{(2n+2)}} - \frac{a_{2n+1}^{\sharp}}{\xi_{-n-1}^{(2n+1)}}\frac{\xi_{-(n+1)}^{(2n+2)}}{\xi_{n+1}^{(2n+2)}} + \frac{b_{2n+1}^{\sharp}}{\xi_n^{(2n)}}\left(\nu_n^{(2n+1)}\frac{\xi_{-(n+1)}^{(2n+2)}}{\xi_{n+1}^{(2n+2)}} - \frac{\xi_n^{(2n+2)}}{\xi_{n+1}^{(2n+2)}}\right)\right). \tag{3.16}$$

Proof. Equate coefficients of like powers of z^{-k} (resp., z^k), $k = 1,2$, and z^i (resp., z^{-i}), $i = 0,1$, respectively, in—the (1 2)-elements of—asymptotics (3.1) and (3.5) (resp., asymptotics (3.3) and (3.7)) and asymptotics (3.2) and (3.6) (resp., asymptotics (3.4) and (3.8)). □

Remark 3.2. Large-n asymptotics for $\xi_n^{(2n)}$ are given in Appendix A, Theorem A.3; similarly, asymptotics for $\xi_{-n-1}^{(2n+1)}$ are given in Appendix B, Theorem B.3. These are obtained by using the $n \to \infty$ asymptotics for $(Y_1^{e,\infty})_{12}$ and $(Y_1^{o,0})_{12}$, and Equations (3.9) and (3.11), respectively (see, also, [21, 22]).

It turns out that the following recurrence relation coefficients are the essential building blocks from which the remaining ones can be expressed:

$$a_{2n+1}^{\sharp}, \qquad b_{2n+1}^{\sharp}, \qquad b_{2n+2}^{\sharp}, \qquad c_{2n+2}^{\sharp}, \qquad \alpha_{2n}^{\sharp}, \qquad \beta_{2n+1}^{\sharp}, \qquad \beta_{2n+2}^{\sharp}, \qquad \text{and} \qquad \gamma_{2n+3}^{\sharp};$$

for example, formulae for $a_{2n}^{\sharp}$ and $\alpha_{2n+1}^{\sharp}$ may be derived: they have the functional form $a_{2n}^{\sharp} = a_{2n}^{\sharp}(b_{2n}^{\sharp}, b_{2n+1}^{\sharp}, \alpha_{2n}^{\sharp}, \beta_{2n}^{\sharp}, \beta_{2n+1}^{\sharp})$ and $\alpha_{2n+1}^{\sharp} = \alpha_{2n+1}^{\sharp}(a_{2n}^{\sharp}, b_{2n}^{\sharp}, b_{2n+1}^{\sharp}, c_{2n+2}^{\sharp}, \beta_{2n+1}^{\sharp}, \beta_{2n+2}^{\sharp}, \gamma_{2n+1}^{\sharp})$. Furthermore, even though Equations (3.13)–(3.16) are not used explicitly hereafter, they give rise to several very interesting identities/relations between the 'even' and 'odd' Riemann theta functions, $\boldsymbol{\theta}^e(z)$ and $\boldsymbol{\theta}^o(z)$, respectively, and the end-points of the supports of the 'even' and 'odd' equilibrium measures, $\{b_{j-1}^e, a_j^e\}_{j=1}^{N+1}$ and $\{b_{j-1}^o, a_j^o\}_{j=1}^{N+1}$, respectively: the details are left to the reader. ■

Proposition 3.2. *Let the external field* $\widetilde{V}\colon \mathbb{R} \setminus \{0\} \to \mathbb{R}$ *be regular and satisfy conditions* (1.20)–(1.22). *Let the orthonormal L-polynomials (resp., monic orthogonal L-polynomials),* $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ *(resp.,* $\{\boldsymbol{\pi}_k(z)\}_{k\in\mathbb{Z}_0^+}$*) be as defined in Equations* (1.2) *and* (1.3) *(resp., Equations* (1.4) *and* (1.5)), *and let* $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ *satisfy the system of recurrence relations* (1.23)–(1.26). *Let* $\overset{e}{Y}\colon \mathbb{C} \setminus \mathbb{R} \to \mathrm{SL}_2(\mathbb{C})$ *(resp.,* $\overset{o}{Y}\colon \mathbb{C} \setminus \mathbb{R} \to \mathrm{SL}_2(\mathbb{C})$*) be the (unique) solution of* **RHP1** *(resp.,* **RHP2**) *with integral representation* (1.29) *(resp.,* (1.30)). *Then,*

$$\begin{gathered}
\frac{\xi_{-n}^{(2n)}}{\xi_n^{(2n)}} =: \nu_{-n}^{(2n)} = (Y_0^{e,0})_{11}, \qquad \frac{\xi_{-(n-1)}^{(2n)}}{\xi_n^{(2n)}} =: \nu_{-(n-1)}^{(2n)} = (Y_1^{e,0})_{11}, \qquad \frac{\xi_{-(n-2)}^{(2n)}}{\xi_n^{(2n)}} = (Y_2^{e,0})_{11}, \\
\frac{\xi_{n-1}^{(2n)}}{\xi_n^{(2n)}} = (Y_1^{e,\infty})_{11}, \qquad \frac{\xi_{n-2}^{(2n)}}{\xi_n^{(2n)}} = (Y_2^{e,\infty})_{11},
\end{gathered} \tag{3.17}$$

$$\begin{gathered}
\frac{\xi_n^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}} =: \nu_n^{(2n+1)} = (Y_0^{o,\infty})_{11}, \qquad \frac{\xi_{n-1}^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}} =: \nu_{n-1}^{(2n+1)} = (Y_1^{o,\infty})_{11}, \qquad \frac{\xi_{n-2}^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}} = (Y_2^{o,\infty})_{11}, \\
\frac{\xi_{-n}^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}} = (Y_1^{o,0})_{11}, \qquad \frac{\xi_{-(n-1)}^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}} = (Y_2^{o,0})_{11},
\end{gathered} \tag{3.18}$$

and

$$\xi_n^{(2n+1)} = b_{2n+2}^{\sharp}\xi_{n+1}^{(2n+2)}, \tag{3.19}$$

$$\xi_{n-1}^{(2n+1)} = b_{2n+1}^{\sharp}\xi_n^{(2n)} + a_{2n+1}^{\sharp}\xi_n^{(2n+1)} + b_{2n+2}^{\sharp}\xi_n^{(2n+2)}, \tag{3.20}$$

$$\xi_n^{(2n)} = c_{2n+2}^{\sharp}\xi_{n+1}^{(2n+2)}, \tag{3.21}$$

$$\xi_{n-1}^{(2n)} = a_{2n}^{\sharp}\xi_n^{(2n)} + b_{2n+1}^{\sharp}\xi_n^{(2n+1)} + c_{2n+2}^{\sharp}\xi_n^{(2n+2)}, \tag{3.22}$$

$$b_{2n}^{\sharp}\xi_{-n}^{(2n-1)} + a_{2n}^{\sharp}\xi_{-n}^{(2n)} + b_{2n+1}^{\sharp}\xi_{-n}^{(2n+1)} + c_{2n+2}^{\sharp}\xi_{-n}^{(2n+2)} = 0, \tag{3.23}$$

$$b_{2n+1}^{\sharp}\xi_{-n-1}^{(2n+1)} + c_{2n+2}^{\sharp}\xi_{-(n+1)}^{(2n+2)} = 0, \tag{3.24}$$

$$\xi_{-n-1}^{(2n+1)} = b_{2n+1}^{\sharp}\xi_{-n}^{(2n)} + a_{2n+1}^{\sharp}\xi_{-n}^{(2n+1)} + b_{2n+2}^{\sharp}\xi_{-n}^{(2n+2)}, \tag{3.25}$$

$$a_{2n+1}^{\sharp}\xi_{-n-1}^{(2n+1)} + b_{2n+2}^{\sharp}\xi_{-(n+1)}^{(2n+2)} = 0, \tag{3.26}$$

where Equations (3.19), (3.21), (3.24), *and* (3.26) *are used to determine* $b_{2n+2}^{\sharp}$, $c_{2n+2}^{\sharp}$, $b_{2n+1}^{\sharp}$, *and* $a_{2n+1}^{\sharp}$, *respectively (see Proposition* 3.3 *below),*

$$\xi_{-n-1}^{(2n+1)} = \gamma_{2n+3}^{\sharp}\xi_{-(n+2)}^{(2n+3)}, \tag{3.27}$$

$$\xi_{-n}^{(2n+1)} = \alpha_{2n+1}^{\sharp}\xi_{-n-1}^{(2n+1)} + \beta_{2n+2}^{\sharp}\xi_{-(n+1)}^{(2n+2)} + \gamma_{2n+3}^{\sharp}\xi_{-(n+1)}^{(2n+3)}, \tag{3.28}$$

$$\beta^{\sharp}_{2n+2}\xi^{(2n+2)}_{n+1}+\gamma^{\sharp}_{2n+3}\xi^{(2n+3)}_{n+1}=0, \tag{3.29}$$

$$\beta^{\sharp}_{2n+1}\xi^{(2n)}_{n}+\alpha^{\sharp}_{2n+1}\xi^{(2n+1)}_{n}+\beta^{\sharp}_{2n+2}\xi^{(2n+2)}_{n}+\gamma^{\sharp}_{2n+3}\xi^{(2n+3)}_{n}=0, \tag{3.30}$$

$$\xi^{(2n)}_{-n}=\beta^{\sharp}_{2n+1}\xi^{(2n+1)}_{-n-1}, \tag{3.31}$$

$$\xi^{(2n)}_{-(n-1)}=\beta^{\sharp}_{2n}\xi^{(2n-1)}_{-n}+\alpha^{\sharp}_{2n}\xi^{(2n)}_{-n}+\beta^{\sharp}_{2n+1}\xi^{(2n+1)}_{-n}, \tag{3.32}$$

$$\alpha^{\sharp}_{2n}\xi^{(2n)}_{n}+\beta^{\sharp}_{2n+1}\xi^{(2n+1)}_{n}=0, \tag{3.33}$$

$$\xi^{(2n)}_{n}=\beta^{\sharp}_{2n}\xi^{(2n-1)}_{n-1}+\alpha^{\sharp}_{2n}\xi^{(2n)}_{n-1}+\beta^{\sharp}_{2n+1}\xi^{(2n+1)}_{n-1}, \tag{3.34}$$

where Equations (3.27), (3.29), (3.31), *and* (3.33) *are used to determine* $\gamma^{\sharp}_{2n+3}$, $\beta^{\sharp}_{2n+2}$, $\beta^{\sharp}_{2n+1}$, *and* $\alpha^{\sharp}_{2n}$, *respectively (see Proposition* 3.3 *below), with*

$$a^{\sharp}_{2n}=\frac{1}{\alpha^{\sharp}_{2n}}-\frac{1}{\alpha^{\sharp}_{2n}}\left(b^{\sharp}_{2n}\beta^{\sharp}_{2n}+b^{\sharp}_{2n+1}\beta^{\sharp}_{2n+1}\right), \tag{3.35}$$

$$\alpha^{\sharp}_{2n+1}=-\frac{1}{b^{\sharp}_{2n+1}}\left(b^{\sharp}_{2n}\gamma^{\sharp}_{2n+1}+a^{\sharp}_{2n}\beta^{\sharp}_{2n+1}+c^{\sharp}_{2n+2}\beta^{\sharp}_{2n+2}\right). \tag{3.36}$$

Proof. One shows from Equation (1.4) that

$$z^{-n}\boldsymbol{\pi}_{2n}(z)\underset{\substack{z\to\infty\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=}1+\frac{1}{z}\frac{\xi^{(2n)}_{n-1}}{\xi^{(2n)}_{n}}+\frac{1}{z^2}\frac{\xi^{(2n)}_{n-2}}{\xi^{(2n)}_{n}}+O\left(\frac{1}{z^3}\right),$$

$$z^{n}\boldsymbol{\pi}_{2n}(z)\underset{\substack{z\to 0\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=}\nu^{(2n)}_{-n}+z\frac{\xi^{(2n)}_{-(n-1)}}{\xi^{(2n)}_{n}}+z^2\frac{\xi^{(2n)}_{-(n-2)}}{\xi^{(2n)}_{n}}+O(z^3),$$

where $\nu^{(2n)}_{-n}$ is defined in Equations (1.9), and, from (the (1 1)-elements of) asymptotics (3.1) and (3.2), that

$$z^{-n}\overset{e}{\mathrm{Y}}_{11}(z)\underset{\substack{z\to\infty\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=}1+\frac{1}{z}(\mathrm{Y}^{e,\infty}_{1})_{11}+\frac{1}{z^2}(\mathrm{Y}^{e,\infty}_{2})_{11}+O\left(\frac{1}{z^3}\right),$$

$$z^{n}\overset{e}{\mathrm{Y}}_{11}(z)\underset{\substack{z\to 0\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=}(\mathrm{Y}^{e,0}_{0})_{11}+z(\mathrm{Y}^{e,0}_{1})_{11}+z^2(\mathrm{Y}^{e,0}_{2})_{11}+O(z^3);$$

via Lemma 1.1 (Equation (1.29)), upon equating coefficients of like powers of z^{-k}, $k=1,2$, and z^{i}, $i=0,1,2$, in the above asymptotic expansions, one arrives at Equations (3.17). *Mutatis mutandis* for Equations (3.18); but in the latter case, via Lemma 1.2 (Equation (1.30)), one equates coefficients of like powers of z^{k}, $k=1,2$, and z^{-i}, $i=0,1,2$, in the asymptotics (cf. Equation (1.5))

$$z^{n+1}\boldsymbol{\pi}_{2n+1}(z)\underset{\substack{z\to 0\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=}1+z\frac{\xi^{(2n+1)}_{-n}}{\xi^{(2n+1)}_{-n-1}}+z^2\frac{\xi^{(2n+1)}_{-(n-1)}}{\xi^{(2n+1)}_{-n-1}}+O(z^3),$$

$$z^{-n}\boldsymbol{\pi}_{2n+1}(z)\underset{\substack{z\to\infty\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=}\nu^{(2n+1)}_{n}+\frac{1}{z}\frac{\xi^{(2n+1)}_{n-1}}{\xi^{(2n+1)}_{-n-1}}+\frac{1}{z^2}\frac{\xi^{(2n+1)}_{n-2}}{\xi^{(2n+1)}_{-n-1}}+O\left(\frac{1}{z^3}\right),$$

where $\nu^{(2n+1)}_{n}$ is defined in Equations (1.9), and (cf. the (1 1)-elements of asymptotic expansions (3.3) and (3.4))

$$z^{n+1}\left(z^{-1}\overset{o}{\mathrm{Y}}_{11}(z)\right)\underset{\substack{z\to 0\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=}1+z(\mathrm{Y}^{o,0}_{1})_{11}+z^2(\mathrm{Y}^{o,0}_{2})_{11}+O(z^3),$$

$$z^{-n}\left(z^{-1}\overset{o}{\mathrm{Y}}_{11}(z)\right)\underset{\substack{z\to\infty\\ z\in\mathbb{C}\setminus\mathbb{R}}}{=}(\mathrm{Y}^{o,\infty}_{0})_{11}+\frac{1}{z}(\mathrm{Y}^{o,\infty}_{1})_{11}+\frac{1}{z^2}(\mathrm{Y}^{o,\infty}_{2})_{11}+O\left(\frac{1}{z^3}\right).$$

Equations (3.19)–(3.34) are derived by substituting Equations (1.2) and (1.3) into recurrence relations (1.23)–(1.26) and equating coefficients of like powers of $z^{\pm 1}$ as $(\mathbb{C}\setminus\mathbb{R}\ni)\ z\to\infty$ and $z\to 0$. Recall

that (cf. Equations (1.27) and (1.28)) $\mathcal{F}\mathcal{G} = \mathcal{G}\mathcal{F} = \mathrm{diag}(1, 1, \ldots, 1, \ldots)$; forming, for example, the product $\mathcal{F}\mathcal{G}$ (the result remains unchanged upon forming the product $\mathcal{G}\mathcal{F}$), using Equations (3.19)–(3.34), and recalling that (due to the chosen normalisation for the orthonormal L-polynomials) $\xi_0^{(0)} = 1$, one shows, after a tedious, but otherwise straightforward, algebraic calculation that the only non-trivial relations between the recurrence relation coefficients are Equations (3.35) and (3.36). □

Proposition 3.3. *Let the external field* $\widetilde{V}\colon \mathbb{R} \setminus \{0\} \to \mathbb{R}$ *be regular and satisfy conditions* (1.20)–(1.22). *Let the orthonormal L-polynomials (resp., monic orthogonal L-polynomials),* $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ *(resp.,* $\{\boldsymbol{\pi}_k(z)\}_{k\in\mathbb{Z}_0^+}$*) be as defined in Equations* (1.2) *and* (1.3) *(resp., Equations* (1.4) *and* (1.5)), *and let* $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ *satisfy the system of recurrence relations* (1.23)–(1.26). *Let* $\overset{e}{\mathrm{Y}}\colon \mathbb{C} \setminus \mathbb{R} \to \mathrm{SL}_2(\mathbb{C})$ *(resp.,* $\overset{o}{\mathrm{Y}}\colon \mathbb{C} \setminus \mathbb{R} \to \mathrm{SL}_2(\mathbb{C})$*) be the (unique) solution of* **RHP1** *(resp.,* **RHP2**) *with integral representation* (1.29) *(resp.,* (1.30)) *and asymptotics* (3.1) *and* (3.2) *(resp., asymptotics* (3.3) *and* (3.4)). *Then,*

$$a_{2n+1}^{\sharp} = -\nu_n^{(2n+1)} \frac{(\mathrm{Y}_0^{o,\infty})_{12}}{(\mathrm{Y}_1^{o,0})_{12}} \qquad (\in \mathbb{R}), \tag{3.37}$$

$$b_{2n+1}^{\sharp} = -2\pi\mathrm{i}\,\xi_n^{(2n)} \xi_{-n-1}^{(2n+1)} (\mathrm{Y}_0^{o,\infty})_{12} \qquad (\in \mathbb{R}), \tag{3.38}$$

$$b_{2n+2}^{\sharp} = \nu_n^{(2n+1)} \sqrt{\frac{-2\pi\mathrm{i}h^+[(\mathrm{Y}_1^{e,\infty})_{12}]}{2\pi\mathrm{i}(\mathrm{Y}_1^{o,0})_{12}}} \qquad (\in \mathbb{R}), \tag{3.39}$$

$$c_{2n+2}^{\sharp} = \sqrt{\frac{-2\pi\mathrm{i}h^+[(\mathrm{Y}_1^{e,\infty})_{12}]}{-2\pi\mathrm{i}(\mathrm{Y}_1^{e,\infty})_{12}}} \qquad (\in \mathbb{R}_+), \tag{3.40}$$

$$\alpha_{2n}^{\sharp} = -\nu_{-n}^{(2n)} \nu_n^{(2n+1)} \qquad (\in \mathbb{R}), \tag{3.41}$$

$$\beta_{2n+1}^{\sharp} = \nu_{-n}^{(2n)} \frac{\xi_n^{(2n)}}{\xi_{-n-1}^{(2n+1)}} \qquad (\in \mathbb{R}), \tag{3.42}$$

$$\beta_{2n+2}^{\sharp} = -\left(h^+\left[\nu_n^{(2n+1)}\right]\right) \sqrt{\frac{-2\pi\mathrm{i}h^+[(\mathrm{Y}_1^{e,\infty})_{12}]}{2\pi\mathrm{i}(\mathrm{Y}_1^{o,0})_{12}}} \qquad (\in \mathbb{R}), \tag{3.43}$$

$$\gamma_{2n+3}^{\sharp} = \sqrt{\frac{2\pi\mathrm{i}h^+[(\mathrm{Y}_1^{o,0})_{12}]}{2\pi\mathrm{i}(\mathrm{Y}_1^{o,0})_{12}}} \qquad (\in \mathbb{R}_+), \tag{3.44}$$

where all the square roots are positive.

Proof. One commences by establishing the following identity:

$$h^+\left[\nu_{-n}^{(2n)}\right] = \frac{(\mathrm{Y}_0^{o,\infty})_{12}}{(\mathrm{Y}_1^{o,0})_{12}}, \quad n \in \mathbb{Z}_0^+. \tag{3.45}$$

From Equations (3.9), (3.11), (3.21), and (3.24), one shows that

$$\begin{aligned} b_{2n+1}^{\sharp} &= -c_{2n+2}^{\sharp} \frac{\xi_{-(n+1)}^{(2n+2)}}{\xi_{-n-1}^{(2n+1)}} = -\frac{\xi_n^{(2n)}}{\xi_{-n-1}^{(2n+1)}} \frac{\xi_{-(n+1)}^{(2n+2)}}{\xi_{n+1}^{(2n+2)}} = -\frac{\xi_n^{(2n)}}{\xi_{-n-1}^{(2n+1)}} h^+\left[\xi_{-n}^{(2n)} / \xi_n^{(2n)}\right] \\ &= -\frac{\xi_n^{(2n)}}{\xi_{-n-1}^{(2n+1)}} h^+\left[\nu_{-n}^{(2n)}\right] = -\left(h^+\left[\nu_{-n}^{(2n)}\right]\right) \frac{(2\pi\mathrm{i}(\mathrm{Y}_1^{o,0})_{12})^{1/2}}{(-2\pi\mathrm{i}(\mathrm{Y}_1^{e,\infty})_{12})^{1/2}}; \end{aligned}$$

but, from Equation (3.12), which, in fact, leads to Equation (3.38) for $b_{2n+1}^{\sharp}$, via Equations (3.9) and (3.11), one has that

$$b_{2n+1}^{\sharp} = -\frac{2\pi\mathrm{i}(\mathrm{Y}_0^{o,\infty})_{12}}{(-2\pi\mathrm{i}(\mathrm{Y}_1^{e,\infty})_{12})^{1/2} (2\pi\mathrm{i}(\mathrm{Y}_1^{o,0})_{12})^{1/2}} :$$

equating the above two expressions for $b_{2n+1}^{\sharp}$, for the **same** choices of branches, one arrives at Identity (3.45). From Equations (3.19) and (3.26), one has that

$$a_{2n+1}^{\sharp} = -b_{2n+2}^{\sharp} \frac{\xi_{-(n+1)}^{(2n+2)}}{\xi_{-n-1}^{(2n+1)}} = -\frac{\xi_n^{(2n+1)}}{\xi_{n+1}^{(2n+2)}} \frac{\xi_{-(n+1)}^{(2n+2)}}{\xi_{-n-1}^{(2n+1)}} = -\frac{\xi_n^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}} \frac{\xi_{-(n+1)}^{(2n+2)}}{\xi_{n+1}^{(2n+2)}}$$

$$= -\nu_n^{(2n+1)} h^+\left[\xi_{-n}^{(2n)}/\xi_n^{(2n)}\right] = -\nu_n^{(2n+1)} h^+\left[\nu_{-n}^{(2n)}\right],$$

which, via Identity (3.45), leads to Equation (3.37). From Equation (3.19), one has that

$$b_{2n+2}^\sharp = \frac{\xi_n^{(2n+1)}}{\xi_{n+1}^{(2n+2)}} = \frac{\xi_n^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}} \frac{\xi_{-n-1}^{(2n+1)}}{\xi_{n+1}^{(2n+2)}} = \nu_n^{(2n+1)} \frac{\xi_{-n-1}^{(2n+1)}}{h^+\left[\xi_n^{(2n)}\right]},$$

which, via Equations (3.9) and (3.11), lead to Equation (3.39). From Equation (3.21), one has that

$$c_{2n+2}^\sharp = \frac{\xi_n^{(2n)}}{\xi_{n+1}^{(2n+2)}} = \frac{\xi_n^{(2n)}}{h^+\left[\xi_n^{(2n)}\right]},$$

which, via Equation (3.9), leads to Equation (3.40). From Equations (3.31) and (3.33), one shows that

$$\alpha_{2n}^\sharp = -\beta_{2n+1}^\sharp \frac{\xi_n^{(2n+1)}}{\xi_n^{(2n)}} = -\frac{\xi_{-n}^{(2n)}}{\xi_{-n-1}^{(2n+1)}} \frac{\xi_n^{(2n+1)}}{\xi_n^{(2n)}} = -\frac{\xi_{-n}^{(2n)}}{\xi_n^{(2n)}} \frac{\xi_n^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}} = -\nu_{-n}^{(2n)} \nu_n^{(2n+1)},$$

which is Equation (3.41). From Equation (3.31), one has that

$$\beta_{2n+1}^\sharp = \frac{\xi_{-n}^{(2n)}}{\xi_{-n-1}^{(2n+1)}} = \frac{\xi_{-n}^{(2n)}}{\xi_n^{(2n)}} \frac{\xi_n^{(2n)}}{\xi_{-n-1}^{(2n+1)}} = \nu_{-n}^{(2n)} \frac{\xi_n^{(2n)}}{\xi_{-n-1}^{(2n+1)}},$$

which is Equation (3.42). From Equations (3.27) and (3.29), one has that

$$\begin{aligned}\beta_{2n+2}^\sharp &= -\gamma_{2n+3}^\sharp \frac{\xi_{n+1}^{(2n+3)}}{\xi_{n+1}^{(2n+2)}} = -\frac{\xi_{-n-1}^{(2n+1)}}{\xi_{-(n+2)}^{(2n+3)}} \frac{\xi_{n+1}^{(2n+3)}}{\xi_{n+1}^{(2n+2)}} = -\frac{\xi_{n+1}^{(2n+3)}}{\xi_{-(n+2)}^{(2n+3)}} \frac{\xi_{-n-1}^{(2n+1)}}{\xi_{n+1}^{(2n+2)}} \\ &= -\left(h^+\left[\xi_n^{(2n+1)}/\xi_{-n-1}^{(2n+1)}\right]\right) \frac{\xi_{-n-1}^{(2n+1)}}{h^+\left[\xi_n^{(2n)}\right]} = -\left(h^+\left[\nu_n^{(2n+1)}\right]\right) \frac{\xi_{-n-1}^{(2n+1)}}{h^+\left[\xi_n^{(2n)}\right]},\end{aligned}$$

which, via Equations (3.9) and (3.11), lead to Equation (3.43). Finally, from Equation (3.27), one has that

$$\gamma_{2n+3}^\sharp = \frac{\xi_{-n-1}^{(2n+1)}}{\xi_{-(n+2)}^{(2n+3)}} = \frac{\xi_{-n-1}^{(2n+1)}}{h^+\left[\xi_{-n-1}^{(2n+1)}\right]},$$

which, via Equation (3.11), leads to Equation (3.44). □

Remark 3.3. Now, explicit formulae are at hand relating all quantities of interest (such as recurrence relation coefficients, real root products, and leading coefficients) to the n-dependent coefficients of asymptotic expansions (3.1)–(3.4). Using these, formulae for certain ratios of Hankel determinants are also obtained. ■

Recalling Equations (1.9)–(1.13), via relations (3.17) and (3.18), one arrives at: (i) for the non-singular case,

$$\prod_{k=1}^{2n} \alpha_k^{(2n)} = \frac{H_{2n}^{(-2n+1)}}{H_{2n}^{(-2n)}} = \frac{\xi_{-n}^{(2n)}}{\xi_n^{(2n)}} =: \nu_{-n}^{(2n)} = (\mathrm{Y}_0^{e,0})_{11},$$

where (for $\xi_{-n}^{(2n)} \neq 0$) $\left\{\alpha_k^{(2n)},\ k=1,\ldots,2n\right\} = \{z;\ \boldsymbol{\pi}_{2n}(z)=0\}$, with $\alpha_k^{(2n)}$ real and simple (see Theorem 3.1, Equation (3.48)), and

$$\left(\prod_{k=1}^{2n+1} \alpha_k^{(2n+1)}\right)^{-1} = \frac{H_{2n+1}^{(-2n-1)}}{H_{2n+1}^{(-2n)}} = -\frac{\xi_n^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}} =: -\nu_n^{(2n+1)} = -(\mathrm{Y}_0^{o,\infty})_{11},$$

where (for $\xi_n^{(2n+1)} \neq 0$) $\left\{\alpha_k^{(2n+1)},\ k=1,\ldots,2n+1\right\} = \{z;\ \boldsymbol{\pi}_{2n+1}(z)=0\}$, with $\alpha_k^{(2n+1)}$ real and simple (see Theorem 3.1, Equation (3.49)); and (ii) for the singular case,

$$\prod_{k=1}^{2n-1} \alpha_k^{(2n)} = -\frac{\xi_{-(n-1)}^{(2n)}}{\xi_n^{(2n)}} =: -\nu_{-(n-1)}^{(2n)} = -(\mathrm{Y}_1^{e,0})_{11},$$

where (for $\xi_{-n}^{(2n)}=0$) $\left\{\alpha_k^{(2n)},\ k=1,\dots,2n-1\right\}=\{z;\ \boldsymbol{\pi}_{2n}(z)=0\}$, with $\alpha_k^{(2n)}$ real and simple (see Theorem 3.1, Equation (3.50)), and

$$\left(\prod_{k=1}^{2n}\alpha_k^{(2n+1)}\right)^{-1}=\frac{\xi_{n-1}^{(2n+1)}}{\xi_{-n-1}^{(2n+1)}}=:\nu_{n-1}^{(2n+1)}=(\mathrm{Y}_1^{o,\infty})_{11},$$

where (for $\xi_n^{(2n+1)}=0$) $\left\{\alpha_k^{(2n+1)},\ k=1,\dots,2n\right\}=\{z;\ \boldsymbol{\pi}_{2n+1}(z)=0\}$, with $\alpha_k^{(2n+1)}$ real and simple (see Theorem 3.1, Equation (3.51)).

Remark 3.4. In order to eschew a flood of superfluous notation, the 'symbol' $c(n)$ appears hereafter in the various error estimations (as $n\to\infty$) of the form $O(c(n)n^{-2})$ and $O(c(n)(n+1/2)^{-2})$, where $c(n)=_{n\to\infty}O(1)$. Even though, *sensus strictu*, the symbol $c(n)$ should properly be denoted as $c_1(n),c_2(n),\dots,c_k(n),\dots$, where, in general, $c_i(n)\neq c_j(n)$, $i\neq j\in\mathbb{N}$, and $c_j(n)=_{n\to\infty}O(1)$, $j\in\mathbb{N}$, with a certain abuse of notation, and concomitant with the above theme, the simplified 'notation' $c(n)$ will be retained. ■

Remark 3.5. The notations $\infty^{\pm}$ (resp., $0^{\pm}$), which appear in the theorems below, are to be understood as follows: let $\mathcal{P}:=(y,z)$ denote an arbitrary point on the (hyperelliptic) Riemann surface $\mathcal{Y}_e$ or $\mathcal{Y}_o$, where, for $q\in\{e,o\}$, $\mathcal{Y}_q:=\left\{(y,z);\ y^2=R_q(z)\right\}$, with $R_e(z)$ (resp., $R_o(z)$) defined in $\boxed{\mathbf{P_1}}$ (resp., $\boxed{\mathbf{P_2}}$). Then: (i) $\mathcal{P}\to\infty^{\pm}\Leftrightarrow z\to\infty$, $y\sim\pm z^{N+1}$; and (ii) $\mathcal{P}\to 0^{\pm}\Leftrightarrow z\to 0$, $y\sim\pm(-1)^{\mathcal{N}_+^q}(\prod_{k=1}^{N+1}|b_{k-1}^q a_k^q|)^{1/2}$ for $q\in\{e,o\}$, where $\mathcal{N}_+^q\in\{0,\dots,N+1\}$ is the number of bands to the right of $z=0$. Furthermore, since in this manuscript $\mathcal{Y}_q^{\pm}=\mathbb{C}_{\pm}$ for $q\in\{e,o\}$, $\infty^{\pm}$ and $0^{\pm}\in\mathbb{C}_{\pm}$. ■

At this point, one is ready to use the preceding formulae and relations subsumed in Propositions 3.1–3.3 to establish large-n asymptotic expansions for the various quantities of interest.

Using, now, $n\to\infty$ asymptotics for the (1 1)-elements of $\mathrm{Y}_0^{e,0}$ and $\mathrm{Y}_1^{e,0}$ (resp., $\mathrm{Y}_0^{o,\infty}$ and $\mathrm{Y}_1^{o,\infty}$) given in Appendix A, Theorem A.2 (resp., Appendix B, Theorem B.2), one proves the following theorem concerning the products of the (real) roots of the L-polynomials: the asymptotic results are contained in Equations (3.48)–(3.51).

Theorem 3.1. *Let the external field* $\widetilde{V}\colon\mathbb{R}\setminus\{0\}\to\mathbb{R}$ *satisfy conditions* (1.20)–(1.22). *Let the orthonormal L-polynomials (resp., monic orthogonal L-polynomials)* $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ *(resp.,* $\{\boldsymbol{\pi}_k(z)\}_{k\in\mathbb{Z}_0^+}$*) be as defined in Equations* (1.2) *and* (1.3) *(resp., Equations* (1.4) *and* (1.5)). *Let* $\overset{e}{\mathrm{Y}}\colon\mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$ *(resp.,* $\overset{o}{\mathrm{Y}}\colon\mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$*) be the (unique) solution of* **RHP1** *(resp.,* **RHP2**) *with integral representation* (1.29) *(resp.,* (1.30)). *Define the density of the 'even' (resp., 'odd') equilibrium measure,* $\mathrm{d}\mu_V^e(x)$ *(resp.,* $\mathrm{d}\mu_V^o(x)$), *as in Equation* (2.13) *(resp., Equation* (2.19)), *and set* $J_q:=\mathrm{supp}(\mu_V^q)=\cup_{j=1}^{N+1}(b_{j-1}^q,a_j^q)$, $q\in\{e,o\}$, *where* $\{b_{j-1}^e,a_j^e\}_{j=1}^{N+1}$ *(resp.,* $\{b_{j-1}^o,a_j^o\}_{j=1}^{N+1}$*) satisfy the (real) n-dependent and locally solvable system of* $2(N+1)$ *moment conditions* (2.12) *(resp.,* (2.18)).

Suppose, furthermore, that $\widetilde{V}\colon\mathbb{R}\setminus\{0\}\to\mathbb{R}$ *is regular, namely:* (i) $h_V^q(z)\not\equiv 0$ *on* $\overline{J_q}:=\cup_{k=1}^{N+1}[b_{k-1}^q,a_k^q]$, $q\in\{e,o\}$; *and* (ii) *the variational inequalities* $(\mathbf{P_1^{(b)}})$ *and* $(\mathbf{P_2^{(b)}})$ *are strict, and take the following form,*

$$4\int_{J_e}\ln(|x-s|)\,\mathrm{d}\mu_V^e(s)-2\ln|x|-\widetilde{V}(x)-\ell_e<0,\quad x\in\mathbb{R}\setminus\overline{J_e},\tag{3.46}$$

$$2\left(2+\frac{1}{n}\right)\int_{J_o}\ln(|x-s|)\,\mathrm{d}\mu_V^o(s)-2\ln|x|-\widetilde{V}(x)-\ell_o-2\left(2+\frac{1}{n}\right)Q_o<0,\quad x\in\mathbb{R}\setminus\overline{J_o},\tag{3.47}$$

where the 'even' and 'odd' variational constants, ℓ_e $(\in\mathbb{R})$ *and* ℓ_o $(\in\mathbb{R})$, *respectively, are defined by Equations* $(\mathbf{P_1^{(a)}})$ *and* $(\mathbf{P_2^{(a)}})$, *and* $Q_o:=\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s$. *Then,*

$$\begin{aligned}\nu_{-n}^{(2n)}=\prod_{k=1}^{2n}\alpha_k^{(2n)}\underset{n\to\infty}{=}&\frac{\omega_+^e}{2}\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}\\&\times\left\{1+\frac{1}{n}\left((\mathcal{R}_0^{e,0})_{11}+(\mathcal{R}_0^{e,0})_{12}\frac{\omega_-^e}{\mathrm{i}\omega_+^e}\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}\right.\right.\\&\left.\left.\times\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\boldsymbol{d}_e)}\right)+O\left(\frac{c(n)}{n^2}\right)\right\}\end{aligned}$$

$$\times \exp\left(2n\left(\int_{J_e} \ln(|s|)\psi_V^e(s)\,\mathrm{d}s + \mathrm{i}\pi \int_{J_e\cap\mathbb{R}_+} \psi_V^e(s)\,\mathrm{d}s\right)\right), \tag{3.48}$$

$$\begin{aligned}
-\nu_n^{(2n+1)} = \left(\prod_{k=1}^{2n+1} \alpha_k^{(2n+1)}\right)^{-1} &\underset{n\to\infty}{=} -\frac{1}{2}\frac{\omega_+^o}{\mathbb{E}}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)} \\
&\times \left\{1+\frac{1}{(n+\frac{1}{2})}\left((\mathcal{R}_0^{o,\infty})_{11}-(\mathcal{R}_0^{o,\infty})_{12}\frac{\omega_-^o\mathbb{E}^2}{\mathrm{i}\omega_+^o}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}\right.\right. \\
&\left.\left.\times\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}\right)+O\left(\frac{c(n)}{(n+\frac{1}{2})^2}\right)\right\} \\
&\times \exp\left(-2\left(n+\frac{1}{2}\right)\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s\right),
\end{aligned} \tag{3.49}$$

$$\begin{aligned}
-\nu_{-(n-1)}^{(2n)} = \prod_{k=1}^{2n-1}\alpha_k^{(2n)} &\underset{n\to\infty}{=} n\omega_+^e\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)} \\
&\times\left\{1+\frac{1}{n}\left((\mathcal{R}_0^{e,0})_{11}+(\mathcal{R}_0^{e,0})_{12}\frac{\omega_-^e}{\mathrm{i}\omega_+^e}\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}\right.\right. \\
&\times\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}+\frac{1}{2}\left(\int_{J_e}s^{-1}\psi_V^e(s)\,\mathrm{d}s\right)^{-1}\left[\frac{\widetilde{\alpha}_0^e(1,1,\mathbf{0})}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)+\boldsymbol{d}_e)}\right. \\
&\left.\left.-\frac{\widetilde{\alpha}_0^e(1,1,\boldsymbol{\Omega}^e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}+\frac{\omega_-^e\eta_-^e}{4\omega_+^e}\right]\right)+O\left(\frac{c(n)}{n^2}\right)\left.\right\}\left(\int_{J_e}s^{-1}\psi_V^e(s)\,\mathrm{d}s\right) \\
&\times\exp\left(2n\left(\int_{J_e}\ln(|s|)\psi_V^e(s)\,\mathrm{d}s+\mathrm{i}\pi\int_{J_e\cap\mathbb{R}_+}\psi_V^e(s)\,\mathrm{d}s\right)\right),
\end{aligned} \tag{3.50}$$

$$\begin{aligned}
\nu_{n-1}^{(2n+1)} = \left(\prod_{k=1}^{2n}\alpha_k^{(2n+1)}\right)^{-1} &\underset{n\to\infty}{=} -\left(n+\frac{1}{2}\right)\frac{\omega_+^o}{\mathbb{E}}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)} \\
&\times\left\{1+\frac{1}{(n+\frac{1}{2})}\left((\mathcal{R}_0^{o,\infty})_{11}-(\mathcal{R}_0^{o,\infty})_{12}\frac{\omega_-^o\mathbb{E}^2}{\mathrm{i}\omega_+^o}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}\right.\right. \\
&\times\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\boldsymbol{d}_o)}+\frac{1}{2}\left(\int_{J_o}s\psi_V^o(s)\,\mathrm{d}s\right)^{-1} \\
&\times\left[\frac{\widetilde{\alpha}_\infty^o(1,1,\boldsymbol{\Omega}^o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}-\frac{\widetilde{\alpha}_\infty^o(1,1,\mathbf{0})}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)+\boldsymbol{d}_o)}+\frac{\omega_-^o\eta_+^o}{4\omega_+^o}\right]\left.\right)+O\left(\frac{c(n)}{(n+\frac{1}{2})^2}\right)\left.\right\} \\
&\times\left(\int_{J_o}s\psi_V^o(s)\,\mathrm{d}s\right)\exp\left(-2\left(n+\frac{1}{2}\right)\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s\right).
\end{aligned} \tag{3.51}$$

In the above asymptotic expansions (3.48)–(3.51)*, there are a number of new parameters, which are now defined:*

$$\omega_\pm^q := \gamma_0^q \pm (\gamma_0^q)^{-1}, \quad q\in\{e,o\}, \tag{3.52}$$

$$\gamma_0^q := \gamma^q(0) = \left(\prod_{k=1}^{N+1} b_{k-1}^q (a_k^q)^{-1}\right)^{1/4} \quad (>0), \tag{3.53}$$

$$\eta_+^q := \sum_{k=1}^{N+1}\left(a_k^q - b_{k-1}^q\right), \qquad \eta_-^q := \sum_{k=1}^{N+1}\left((b_{k-1}^q)^{-1} - (a_k^q)^{-1}\right), \tag{3.54}$$

$\gamma^e(z)$ (resp., $\gamma^o(z)$) is defined in Equation (2.2) *(resp., Equation* (2.7)*), $\boldsymbol{\theta}^e(z)$ (resp., $\boldsymbol{\theta}^o(z)$) is defined in Equation* (2.5) *(resp., Equation* (2.10)*), $\boldsymbol{u}^q(z) := \int_{a_{N+1}^q}^z \boldsymbol{\omega}^q$, $q\in\{e,o\}$, $\boldsymbol{u}_+^q(\infty) := \int_{a_{N+1}^q}^{\infty^+}\boldsymbol{\omega}^q$, $\boldsymbol{u}_+^q(0) := \int_{a_{N+1}^q}^{0^+}\boldsymbol{\omega}^q$, $\boldsymbol{d}_e$ (resp.,*

$\boldsymbol{d}_o$) *is given in Equations* (2.4) *(resp., Equations* (2.9)*), with* $z_j^{e,\pm} \in (a_j^e, b_j^e)^{\pm} \subset \mathbb{C}_{\pm}$ *(resp.,* $z_j^{o,\pm} \in (a_j^o, b_j^o)^{\pm} \subset \mathbb{C}_{\pm}$*),* $j = 1, \dots, N$*, given in Equation* (2.3) *(resp., Equation* (2.8)*),* $\boldsymbol{\Omega}^q := (\Omega_1^q, \Omega_2^q, \dots, \Omega_N^q)^{\mathrm{T}}$ $(\in \mathbb{R}^N)$*, with* $\Omega_j^q = 4\pi \int_{b_j^q}^{a_{N+1}^q} \psi_V^q(s)\, \mathrm{d}s$*,* $j = 1, \dots, N$*,*

$$\mathcal{R}_0^{q,p} := \operatorname{sgn}(q) \sum_{j=1}^{N+1} \left(\frac{\left(\mathcal{A}^q(b_{j-1}^q)\left(\widehat{\alpha}_1^q(b_{j-1}^q) + (b_{j-1}^q)^{-1}\widehat{\alpha}_0^q(b_{j-1}^q)\right) - \mathcal{B}^q(b_{j-1}^q)\widehat{\alpha}_0^q(b_{j-1}^q)\right)}{(\widehat{\alpha}_0^q(b_{j-1}^q))^2 b_{j-1}^q} + \frac{\left(\mathcal{A}^q(a_j^q)\left(\widehat{\alpha}_1^q(a_j^q) + (a_j^q)^{-1}\widehat{\alpha}_0^q(a_j^q)\right) - \mathcal{B}^q(a_j^q)\widehat{\alpha}_0^q(a_j^q)\right)}{(\widehat{\alpha}_0^q(a_j^q))^2 a_j^q} \right), \qquad q \in \{e, o\}, \quad p \in \{0, \infty\}, \tag{3.55}$$

$$\operatorname{sgn}(q) := \begin{cases} +1, & q = e, \\ -1, & q = o, \end{cases} \tag{3.56}$$

where, for $j = 1, \dots, N+1$*,*

$$\mathcal{A}^e(b_{j-1}^e) = -s_1 (Q_0^e(b_{j-1}^e))^{-1} \mathrm{e}^{\mathrm{i}n\mho_{j-1}^e} \begin{pmatrix} \varkappa_1^e(b_{j-1}^e)\varkappa_2^e(b_{j-1}^e) & \mathrm{i}(\varkappa_1^e(b_{j-1}^e))^2 \\ \mathrm{i}(\varkappa_2^e(b_{j-1}^e))^2 & -\varkappa_1^e(b_{j-1}^e)\varkappa_2^e(b_{j-1}^e) \end{pmatrix}, \tag{3.57}$$

$$\mathcal{A}^e(a_j^e) = s_1 Q_0^e(a_j^e) \mathrm{e}^{\mathrm{i}n\mho_j^e} \begin{pmatrix} -\varkappa_1^e(a_j^e)\varkappa_2^e(a_j^e) & \mathrm{i}(\varkappa_1^e(a_j^e))^2 \\ \mathrm{i}(\varkappa_2^e(a_j^e))^2 & \varkappa_1^e(a_j^e)\varkappa_2^e(a_j^e) \end{pmatrix}, \tag{3.58}$$

$$\mathcal{A}^o(b_{j-1}^o) = -\frac{s_1(\gamma^o(0))^2 \mathrm{e}^{\mathrm{i}(n+\frac{1}{2})\mho_{j-1}^o}}{Q_0^o(b_{j-1}^o)} \begin{pmatrix} \varkappa_1^o(b_{j-1}^o)\varkappa_2^o(b_{j-1}^o) & \mathrm{i}(\varkappa_1^o(b_{j-1}^o))^2 \\ \mathrm{i}(\varkappa_2^o(b_{j-1}^o))^2 & -\varkappa_1^o(b_{j-1}^o)\varkappa_2^o(b_{j-1}^o) \end{pmatrix}, \tag{3.59}$$

$$\mathcal{A}^o(a_j^o) = \frac{s_1 Q_0^o(a_j^o) \mathrm{e}^{\mathrm{i}(n+\frac{1}{2})\mho_j^o}}{(\gamma^o(0))^2} \begin{pmatrix} -\varkappa_1^o(a_j^o)\varkappa_2^o(a_j^o) & \mathrm{i}(\varkappa_1^o(a_j^o))^2 \\ \mathrm{i}(\varkappa_2^o(a_j^o))^2 & \varkappa_1^o(a_j^o)\varkappa_2^o(a_j^o) \end{pmatrix}, \tag{3.60}$$

$$\frac{\mathcal{B}^e(b_{j-1}^e)}{\mathrm{e}^{\mathrm{i}n\mho_{j-1}^e}} = \begin{pmatrix} \varkappa_1^e(b_{j-1}^e)\varkappa_2^e(b_{j-1}^e)\left(-s_1(Q_0^e(b_{j-1}^e))^{-1}\right. \\ \times \left\{ \overset{e}{\urcorner}_1^1(b_{j-1}^e) + \overset{e}{\urcorner}_{-1}^1(b_{j-1}^e) - Q_1^e(b_{j-1}^e) \right. \\ \left. \times (Q_0^e(b_{j-1}^e))^{-1} \right\} - t_1 \left\{ Q_0^e(b_{j-1}^e) \right. \\ \left. + (Q_0^e(b_{j-1}^e))^{-1} \overset{e}{\aleph}_1^1(b_{j-1}^e) \overset{e}{\aleph}_{-1}^1(b_{j-1}^e) \right\} \\ \left. + \mathrm{i}(s_1 + t_1) \left\{ \overset{e}{\aleph}_{-1}^1(b_{j-1}^e) - \overset{e}{\aleph}_1^1(b_{j-1}^e) \right\} \right) \\ (\varkappa_1^e(b_{j-1}^e))^2 \left(-\mathrm{i}s_1 (Q_0^e(b_{j-1}^e))^{-1} \left\{ 2\overset{e}{\urcorner}_1^1(b_{j-1}^e) \right. \right. \\ \left. - Q_1^e(b_{j-1}^e)(Q_0^e(b_{j-1}^e))^{-1} \right\} + \mathrm{i}t_1 \left\{ Q_0^e(b_{j-1}^e) \right. \\ \left. \left. - (Q_0^e(b_{j-1}^e))^{-1} (\overset{e}{\aleph}_1^1(b_{j-1}^e))^2 \right\} \right. \\ \left. - 2(s_1 - t_1) \overset{e}{\aleph}_1^1(b_{j-1}^e) \right) \end{pmatrix} \begin{pmatrix} (\varkappa_1^e(b_{j-1}^e))^2 \left(-\mathrm{i}s_1 (Q_0^e(b_{j-1}^e))^{-1} \left\{ 2\overset{e}{\urcorner}_1^1(b_{j-1}^e) \right. \right. \\ \left. - Q_1^e(b_{j-1}^e)(Q_0^e(b_{j-1}^e))^{-1} \right\} + \mathrm{i}t_1 \left\{ Q_0^e(b_{j-1}^e) \right. \\ \left. \left. - (Q_0^e(b_{j-1}^e))^{-1} (\overset{e}{\aleph}_1^1(b_{j-1}^e))^2 \right\} \right. \\ \left. + 2(s_1 - t_1) \overset{e}{\aleph}_1^1(b_{j-1}^e) \right) \\ \varkappa_1^e(b_{j-1}^e)\varkappa_2^e(b_{j-1}^e)\left(s_1(Q_0^e(b_{j-1}^e))^{-1}\right. \\ \times \left\{ \overset{e}{\urcorner}_1^1(b_{j-1}^e) + \overset{e}{\urcorner}_{-1}^1(b_{j-1}^e) - Q_1^e(b_{j-1}^e) \right. \\ \left. \times (Q_0^e(b_{j-1}^e))^{-1} \right\} + t_1 \left\{ Q_0^e(b_{j-1}^e) \right. \\ \left. + (Q_0^e(b_{j-1}^e))^{-1} \overset{e}{\aleph}_1^1(b_{j-1}^e) \overset{e}{\aleph}_{-1}^1(b_{j-1}^e) \right\} \\ \left. + \mathrm{i}(s_1 + t_1) \left\{ \overset{e}{\aleph}_1^1(b_{j-1}^e) - \overset{e}{\aleph}_{-1}^1(b_{j-1}^e) \right\} \right) \end{pmatrix}, \tag{3.61}$$

$$\frac{\mathcal{B}^e(a_j^e)}{\mathrm{e}^{\mathrm{i}n\mathcal{O}_j^e}}=\begin{pmatrix}\kappa_1^e(a_j^e)\kappa_2^e(a_j^e)\Big(-s_1\Big\{Q_1^e(a_j^e)+Q_0^e(a_j^e)\Big[\overset{e}{\urcorner}{}_1^1(a_j^e)+\overset{e}{\urcorner}{}_{-1}^1(a_j^e)\Big]\Big\}-t_1\times\Big\{(Q_0^e(a_j^e))^{-1}+Q_0^e(a_j^e)\overset{e}{\aleph}{}_1^1(a_j^e)\overset{e}{\aleph}{}_{-1}^1(a_j^e)\Big\}+\mathrm{i}(s_1+t_1)\Big\{\overset{e}{\aleph}{}_{-1}^1(a_j^e)-\overset{e}{\aleph}{}_1^1(a_j^e)\Big\}\Big) & (\kappa_1^e(a_j^e))^2\Big(\mathrm{i}s_1\Big\{Q_1^e(a_j^e)+2Q_0^e(a_j^e)\times\overset{e}{\urcorner}{}_1^1(a_j^e)\Big\}+\mathrm{i}t_1\Big\{Q_0^e(a_j^e)(\overset{e}{\aleph}{}_1^1(a_j^e))^2-(Q_0^e(a_j^e))^{-1}\Big\}-2(s_1-t_1)\overset{e}{\aleph}{}_1^1(a_j^e)\Big)\\ (\kappa_2^e(a_j^e))^2\Big(\mathrm{i}s_1\Big\{Q_1^e(a_j^e)+2Q_0^e(a_j^e)\times\overset{e}{\urcorner}{}_{-1}^1(a_j^e)\Big\}+\mathrm{i}t_1\Big\{Q_0^e(a_j^e)(\overset{e}{\aleph}{}_{-1}^1(a_j^e))^2-(Q_0^e(a_j^e))^{-1}\Big\}+2(s_1-t_1)\overset{e}{\aleph}{}_{-1}^1(a_j^e)\Big) & \kappa_1^e(a_j^e)\kappa_2^e(a_j^e)\Big(s_1\Big\{Q_1^e(a_j^e)+Q_0^e(a_j^e)\Big[\overset{e}{\urcorner}{}_1^1(a_j^e)+\overset{e}{\urcorner}{}_{-1}^1(a_j^e)\Big]\Big\}+t_1\times\Big\{(Q_0^e(a_j^e))^{-1}+Q_0^e(a_j^e)\overset{e}{\aleph}{}_1^1(a_j^e)\overset{e}{\aleph}{}_{-1}^1(a_j^e)\Big\}+\mathrm{i}(s_1+t_1)\Big\{\overset{e}{\aleph}{}_1^1(a_j^e)-\overset{e}{\aleph}{}_{-1}^1(a_j^e)\Big\}\Big)\end{pmatrix},\tag{3.62}$$

$$\frac{\mathcal{B}^o(b_{j-1}^o)}{\mathrm{e}^{\mathrm{i}(n+\frac{1}{2})\mathcal{O}_{j-1}^o}}=\begin{pmatrix}\kappa_1^o(b_{j-1}^o)\kappa_2^o(b_{j-1}^o)\Big(-\frac{s_1(\gamma^o(0))^2}{Q_0^o(b_{j-1}^o)}\times\Big\{\overset{o}{\urcorner}{}_1^1(b_{j-1}^o)+\overset{o}{\urcorner}{}_{-1}^1(b_{j-1}^o)-Q_1^o(b_{j-1}^o)\times(Q_0^o(b_{j-1}^o))^{-1}\Big\}-t_1(\gamma^o(0))^2\Big\{Q_0^o(b_{j-1}^o)+(Q_0^o(b_{j-1}^o))^{-1}\overset{o}{\aleph}{}_1^1(b_{j-1}^o)\overset{o}{\aleph}{}_{-1}^1(b_{j-1}^o)\Big\}+\mathrm{i}(s_1+t_1)\Big\{\overset{o}{\aleph}{}_{-1}^1(b_{j-1}^o)-\overset{o}{\aleph}{}_1^1(b_{j-1}^o)\Big\}\Big) & (\kappa_1^o(b_{j-1}^o))^2\Big(-\frac{\mathrm{i}s_1(\gamma^o(0))^2}{Q_0^o(b_{j-1}^o)}\Big\{2\overset{o}{\urcorner}{}_1^1(b_{j-1}^o)-Q_1^o(b_{j-1}^o)(Q_0^o(b_{j-1}^o))^{-1}\Big\}+\mathrm{i}t_1\Big\{Q_0^o(b_{j-1}^o)\times(\gamma^o(0))^{-2}-(Q_0^o(b_{j-1}^o))^{-1}(\gamma^o(0))^2\times(\overset{o}{\aleph}{}_1^1(b_{j-1}^o))^2\Big\}+2(s_1-t_1)\overset{o}{\aleph}{}_1^1(b_{j-1}^o)\Big)\\ (\kappa_2^o(b_{j-1}^o))^2\Big(-\frac{\mathrm{i}s_1(\gamma^o(0))^2}{Q_0^o(b_{j-1}^o)}\Big\{2\overset{o}{\urcorner}{}_{-1}^1(b_{j-1}^o)-Q_1^o(b_{j-1}^o)(Q_0^o(b_{j-1}^o))^{-1}\Big\}+\mathrm{i}t_1\Big\{Q_0^o(b_{j-1}^o)\times(\gamma^o(0))^{-2}-(Q_0^o(b_{j-1}^o))^{-1}(\gamma^o(0))^2\times(\overset{o}{\aleph}{}_{-1}^1(b_{j-1}^o))^2\Big\}-2(s_1-t_1)\overset{o}{\aleph}{}_{-1}^1(b_{j-1}^o)\Big) & \kappa_1^o(b_{j-1}^o)\kappa_2^o(b_{j-1}^o)\Big(\frac{s_1(\gamma^o(0))^2}{Q_0^o(b_{j-1}^o)}\times\Big\{\overset{o}{\urcorner}{}_1^1(b_{j-1}^o)+\overset{o}{\urcorner}{}_{-1}^1(b_{j-1}^o)-Q_1^o(b_{j-1}^o)\times(Q_0^o(b_{j-1}^o))^{-1}\Big\}+t_1(\gamma^o(0))^2\Big\{Q_0^o(b_{j-1}^o)+(Q_0^o(b_{j-1}^o))^{-1}\overset{o}{\aleph}{}_1^1(b_{j-1}^o)\overset{o}{\aleph}{}_{-1}^1(b_{j-1}^o)\Big\}+\mathrm{i}(s_1+t_1)\Big\{\overset{o}{\aleph}{}_1^1(b_{j-1}^o)-\overset{o}{\aleph}{}_{-1}^1(b_{j-1}^o)\Big\}\Big)\end{pmatrix},\tag{3.63}$$

$$\frac{\mathcal{B}^o(a_j^o)}{\mathrm{e}^{\mathrm{i}(n+\frac{1}{2})\mathcal{O}_j^o}}=\begin{pmatrix}\kappa_1^o(a_j^o)\kappa_2^o(a_j^o)\Big(-\frac{s_1}{(\gamma^o(0))^2}\Big\{Q_1^o(a_j^o)+Q_0^o(a_j^o)\Big[\overset{o}{\urcorner}{}_1^1(a_j^o)+\overset{o}{\urcorner}{}_{-1}^1(a_j^o)\Big]\Big\}-t_1\times\Big\{(\gamma^o(0))^{-2}Q_0^o(a_j^o)\overset{o}{\aleph}{}_1^1(a_j^o)\overset{o}{\aleph}{}_{-1}^1(a_j^o)+(\gamma^o(0))^2(Q_0^o(a_j^o))^{-1}\Big\}+\mathrm{i}(s_1+t_1)\Big\{\overset{o}{\aleph}{}_{-1}^1(a_j^o)-\overset{o}{\aleph}{}_1^1(a_j^o)\Big\}\Big) & (\kappa_1^o(a_j^o))^2\Big(\frac{\mathrm{i}s_1}{(\gamma^o(0))^2}\Big\{Q_1^o(a_j^o)+2Q_0^o(a_j^o)\times\overset{o}{\urcorner}{}_1^1(a_j^o)\Big\}+\mathrm{i}t_1\Big\{Q_0^o(a_j^o)(\overset{o}{\aleph}{}_1^1(a_j^o))^2\times(\gamma^o(0))^{-2}-(Q_0^o(a_j^o))^{-1}(\gamma^o(0))^2\Big\}-2(s_1-t_1)\overset{o}{\aleph}{}_1^1(a_j^o)\Big)\\ (\kappa_2^o(a_j^o))^2\Big(\frac{\mathrm{i}s_1}{(\gamma^o(0))^2}\Big\{Q_1^o(a_j^o)+2Q_0^o(a_j^o)\times\overset{o}{\urcorner}{}_{-1}^1(a_j^o)\Big\}+\mathrm{i}t_1\Big\{Q_0^o(a_j^o)(\overset{o}{\aleph}{}_{-1}^1(a_j^o))^2\times(\gamma^o(0))^{-2}-(Q_0^o(a_j^o))^{-1}(\gamma^o(0))^2\Big\}+2(s_1-t_1)\overset{o}{\aleph}{}_{-1}^1(a_j^o)\Big) & \kappa_1^o(a_j^o)\kappa_2^o(a_j^o)\Big(\frac{s_1}{(\gamma^o(0))^2}\Big\{Q_1^o(a_j^o)+Q_0^o(a_j^o)\Big[\overset{o}{\urcorner}{}_1^1(a_j^o)+\overset{o}{\urcorner}{}_{-1}^1(a_j^o)\Big]\Big\}+t_1\times\Big\{(\gamma^o(0))^{-2}Q_0^o(a_j^o)\overset{o}{\aleph}{}_1^1(a_j^o)\overset{o}{\aleph}{}_{-1}^1(a_j^o)+(\gamma^o(0))^2(Q_0^o(a_j^o))^{-1}\Big\}+\mathrm{i}(s_1+t_1)\Big\{\overset{o}{\aleph}{}_1^1(a_j^o)-\overset{o}{\aleph}{}_{-1}^1(a_j^o)\Big\}\Big)\end{pmatrix},\tag{3.64}$$

$$\mathbb{E}:=\exp\Big(\mathrm{i}2\pi\Big(n+\frac{1}{2}\Big)\int_{J_o\cap\mathbb{R}_+}\psi_V^o(s)\,\mathrm{d}s\Big),\tag{3.65}$$

and, for $q\in\{e,o\}$,

$$s_1=\frac{5}{72},\qquad t_1=-\frac{7}{72},\qquad \mathcal{O}_k^q=\begin{cases}\Omega_k^q, & k=1,\dots,N,\\ 0, & k=0,N+1,\end{cases}\tag{3.66}$$

$$Q_0^q(b_0^q) = -\mathrm{i}\left((a_{N+1}^q - b_0^q)^{-1}\prod_{k=1}^{N}\left(\frac{b_k^q - b_0^q}{a_k^q - b_0^q}\right)\right)^{1/2}, \tag{3.67}$$

$$Q_1^q(b_0^q) = \frac{1}{2}Q_0^q(b_0^q)\left(\sum_{k=1}^{N}\left(\frac{1}{b_0^q - b_k^q} - \frac{1}{b_0^q - a_k^q}\right) - \frac{1}{b_0^q - a_{N+1}^q}\right), \tag{3.68}$$

$$Q_0^q(a_{N+1}^q) = \left((a_{N+1}^q - b_0^q)\prod_{k=1}^{N}\left(\frac{a_{N+1}^q - b_k^q}{a_{N+1}^q - a_k^q}\right)\right)^{1/2}, \tag{3.69}$$

$$Q_1^q(a_{N+1}^q) = \frac{1}{2}Q_0^q(a_{N+1}^q)\left(\sum_{k=1}^{N}\left(\frac{1}{a_{N+1}^q - b_k^q} - \frac{1}{a_{N+1}^q - a_k^q}\right) + \frac{1}{a_{N+1}^q - b_0^q}\right), \tag{3.70}$$

$$Q_0^q(b_j^q) = -\mathrm{i}\left(\frac{(b_j^q - b_0^q)}{(a_{N+1}^q - b_j^q)(b_j^q - a_j^q)}\prod_{k=1}^{j-1}\left(\frac{b_j^q - b_k^q}{b_j^q - a_k^q}\right)\prod_{l=j+1}^{N}\left(\frac{b_l^q - b_j^q}{a_l^q - b_j^q}\right)\right)^{1/2}, \tag{3.71}$$

$$Q_1^q(b_j^q) = \frac{1}{2}Q_0^q(b_j^q)\left(\sum_{\substack{k=1\\k\neq j}}^{N}\left(\frac{1}{b_j^q - b_k^q} - \frac{1}{b_j^q - a_k^q}\right) + \frac{1}{b_j^q - b_0^q} - \frac{1}{b_j^q - a_{N+1}^q} - \frac{1}{b_j^q - a_j^q}\right), \tag{3.72}$$

$$Q_0^q(a_j^q) = \left(\frac{(a_j^q - b_0^q)(b_j^q - a_j^q)}{(a_{N+1}^q - a_j^q)}\prod_{k=1}^{j-1}\left(\frac{a_j^q - b_k^q}{a_j^q - a_k^q}\right)\prod_{l=j+1}^{N}\left(\frac{b_l^q - a_j^q}{a_l^q - a_j^q}\right)\right)^{1/2}, \tag{3.73}$$

$$Q_1^q(a_j^q) = \frac{1}{2}Q_0^q(a_j^q)\left(\sum_{\substack{k=1\\k\neq j}}^{N}\left(\frac{1}{a_j^q - b_k^q} - \frac{1}{a_j^q - a_k^q}\right) + \frac{1}{a_j^q - b_0^q} - \frac{1}{a_j^q - a_{N+1}^q} + \frac{1}{a_j^q - b_j^q}\right), \tag{3.74}$$

with $\mathrm{i}Q_0^q(b_{j-1}^q), Q_0^q(a_j^q) > 0$, $j = 1, \dots, N+1$,

$$\varkappa_1^e(\xi) = \frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty) + \boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\xi) - \frac{n}{2\pi}\boldsymbol{\Omega}^e + \boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty) - \frac{n}{2\pi}\boldsymbol{\Omega}^e + \boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\xi) + \boldsymbol{d}_e)}, \tag{3.75}$$

$$\varkappa_2^e(\xi) = \frac{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty) - \boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\xi) - \frac{n}{2\pi}\boldsymbol{\Omega}^e - \boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty) - \frac{n}{2\pi}\boldsymbol{\Omega}^e - \boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\xi) - \boldsymbol{d}_e)}, \tag{3.76}$$

$$\varkappa_1^o(\xi) = \frac{1}{\mathbb{E}}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0) + \boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\xi) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\xi) + \boldsymbol{d}_o)}, \tag{3.77}$$

$$\varkappa_2^o(\xi) = \mathbb{E}\frac{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0) - \boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\xi) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o - \boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o - \boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\xi) - \boldsymbol{d}_o)}, \tag{3.78}$$

$$\overset{e}{\aleph}{}_{\varepsilon_2}^{\varepsilon_1}(\xi) = -\frac{\mathfrak{u}^e(\varepsilon_1, \varepsilon_2, \mathbf{0}; \xi)}{\boldsymbol{\theta}^e(\varepsilon_1\boldsymbol{u}_+^e(\xi) + \varepsilon_2\boldsymbol{d}_e)} + \frac{\mathfrak{u}^e(\varepsilon_1, \varepsilon_2, \boldsymbol{\Omega}^e; \xi)}{\boldsymbol{\theta}^e(\varepsilon_1\boldsymbol{u}_+^e(\xi) - \frac{n}{2\pi}\boldsymbol{\Omega}^e + \varepsilon_2\boldsymbol{d}_e)}, \quad \varepsilon_1, \varepsilon_2 = \pm 1, \tag{3.79}$$

$$\overset{o}{\aleph}{}_{\varepsilon_2}^{\varepsilon_1}(\xi) = -\frac{\mathfrak{u}^o(\varepsilon_1, \varepsilon_2, \mathbf{0}; \xi)}{\boldsymbol{\theta}^o(\varepsilon_1\boldsymbol{u}_+^o(\xi) + \varepsilon_2\boldsymbol{d}_o)} + \frac{\mathfrak{u}^o(\varepsilon_1, \varepsilon_2, \boldsymbol{\Omega}^o; \xi)}{\boldsymbol{\theta}^o(\varepsilon_1\boldsymbol{u}_+^o(\xi) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \varepsilon_2\boldsymbol{d}_o)}, \tag{3.80}$$

$$\begin{aligned}\overset{e}{\daleth}{}_{\varepsilon_2}^{\varepsilon_1}(\xi) = {} & -\frac{\mathfrak{v}^e(\varepsilon_1, \varepsilon_2, \mathbf{0}; \xi)}{\boldsymbol{\theta}^e(\varepsilon_1\boldsymbol{u}_+^e(\xi) + \varepsilon_2\boldsymbol{d}_e)} + \frac{\mathfrak{v}^e(\varepsilon_1, \varepsilon_2, \boldsymbol{\Omega}^e; \xi)}{\boldsymbol{\theta}^e(\varepsilon_1\boldsymbol{u}_+^e(\xi) - \frac{n}{2\pi}\boldsymbol{\Omega}^e + \varepsilon_2\boldsymbol{d}_e)} - \left(\frac{\mathfrak{u}^e(\varepsilon_1, \varepsilon_2, \mathbf{0}; \xi)}{\boldsymbol{\theta}^e(\varepsilon_1\boldsymbol{u}_+^e(\xi) + \varepsilon_2\boldsymbol{d}_e)}\right)^2 \\ & + \frac{\mathfrak{u}^e(\varepsilon_1, \varepsilon_2, \mathbf{0}; \xi)\mathfrak{u}^e(\varepsilon_1, \varepsilon_2, \boldsymbol{\Omega}^e; \xi)}{\boldsymbol{\theta}^e(\varepsilon_1\boldsymbol{u}_+^e(\xi) + \varepsilon_2\boldsymbol{d}_e)\boldsymbol{\theta}^e(\varepsilon_1\boldsymbol{u}_+^e(\xi) - \frac{n}{2\pi}\boldsymbol{\Omega}^e + \varepsilon_2\boldsymbol{d}_e)},\end{aligned} \tag{3.81}$$

$$\begin{aligned}\overset{o}{\daleth}{}_{\varepsilon_2}^{\varepsilon_1}(\xi) = {} & -\frac{\mathfrak{v}^o(\varepsilon_1, \varepsilon_2, \mathbf{0}; \xi)}{\boldsymbol{\theta}^o(\varepsilon_1\boldsymbol{u}_+^o(\xi) + \varepsilon_2\boldsymbol{d}_o)} + \frac{\mathfrak{v}^o(\varepsilon_1, \varepsilon_2, \boldsymbol{\Omega}^o; \xi)}{\boldsymbol{\theta}^o(\varepsilon_1\boldsymbol{u}_+^o(\xi) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \varepsilon_2\boldsymbol{d}_o)} - \left(\frac{\mathfrak{u}^o(\varepsilon_1, \varepsilon_2, \mathbf{0}; \xi)}{\boldsymbol{\theta}^o(\varepsilon_1\boldsymbol{u}_+^o(\xi) + \varepsilon_2\boldsymbol{d}_o)}\right)^2 \\ & + \frac{\mathfrak{u}^o(\varepsilon_1, \varepsilon_2, \mathbf{0}; \xi)\mathfrak{u}^o(\varepsilon_1, \varepsilon_2, \boldsymbol{\Omega}^o; \xi)}{\boldsymbol{\theta}^o(\varepsilon_1\boldsymbol{u}_+^o(\xi) + \varepsilon_2\boldsymbol{d}_o)\boldsymbol{\theta}^o(\varepsilon_1\boldsymbol{u}_+^o(\xi) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \varepsilon_2\boldsymbol{d}_o)},\end{aligned} \tag{3.82}$$

$$\mathfrak{u}^q(\varepsilon_1, \varepsilon_2, \boldsymbol{\Omega}^q, \xi) := 2\pi\Lambda_q^1(\varepsilon_1, \varepsilon_2, \boldsymbol{\Omega}^q, \xi), \qquad \mathfrak{v}^q(\varepsilon_1, \varepsilon_2, \boldsymbol{\Omega}^q, \xi) := -2\pi^2\Lambda_q^2(\varepsilon_1, \varepsilon_2, \boldsymbol{\Omega}^q, \xi), \tag{3.83}$$

$$\Lambda_e^{j'}(\varepsilon_1,\varepsilon_2,\boldsymbol{\Omega}^e,\xi)=\sum_{m\in\mathbb{Z}^N}(\mathrm{r}_e(\xi))^{j'}\,\mathrm{e}^{2\pi\mathrm{i}(m,\varepsilon_1\boldsymbol{u}_+^e(\xi)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\varepsilon_2\boldsymbol{d}_e)+\pi\mathrm{i}(m,\boldsymbol{\tau}^e m)},\quad j'=1,2,\tag{3.84}$$

$$\Lambda_o^{j'}(\varepsilon_1,\varepsilon_2,\boldsymbol{\Omega}^o,\xi)=\sum_{m\in\mathbb{Z}^N}(\mathrm{r}_o(\xi))^{j'}\,\mathrm{e}^{2\pi\mathrm{i}(m,\varepsilon_1\boldsymbol{u}_+^o(\xi)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\varepsilon_2\boldsymbol{d}_o)+\pi\mathrm{i}(m,\boldsymbol{\tau}^o m)},\tag{3.85}$$

$$\mathrm{r}_q(\xi):=\frac{2(m,\vec{\curlywedge}_q(\xi))}{\nwarrow^q(\xi)},\qquad \vec{\curlywedge}_q(\xi)=\left(\curlywedge_1^q(\xi),\curlywedge_2^q(\xi),\dots,\curlywedge_N^q(\xi)\right),\tag{3.86}$$

$$\curlywedge_{j'}^q(\xi):=\sum_{k=1}^N c_{j'k}^q\xi^{N-k},\quad j'=1,\dots,N,\tag{3.87}$$

$$\nwarrow^q(b_0^q)=\mathrm{i}(-1)^N\eta_{b_0^q},\quad \nwarrow^q(a_{N+1}^q)=\eta_{a_{N+1}^q},\quad \nwarrow^q(b_j^q)=\mathrm{i}(-1)^{N-j}\eta_{b_j^q},\quad \nwarrow^q(a_j^q)=(-1)^{N-j+1}\eta_{a_j^q},\tag{3.88}$$

$$\eta_{b_0^q}:=\left((a_{N+1}^q-b_0^q)\prod_{k=1}^N(b_k^q-b_0^q)(a_k^q-b_0^q)\right)^{1/2},\tag{3.89}$$

$$\eta_{a_{N+1}^q}:=\left((a_{N+1}^q-b_0^q)\prod_{k=1}^N(a_{N+1}^q-b_k^q)(a_{N+1}^q-a_k^q)\right)^{1/2},\tag{3.90}$$

$$\eta_{b_j^q}:=\left((b_j^q-a_j^q)(a_{N+1}^q-b_j^q)(b_j^q-b_0^q)\prod_{k=1}^{j-1}(b_j^q-b_k^q)(b_j^q-a_k^q)\prod_{l=j+1}^N(b_l^q-b_j^q)(a_l^q-b_j^q)\right)^{1/2},\tag{3.91}$$

$$\eta_{a_j^q}:=\left((b_j^q-a_j^q)(a_{N+1}^q-a_j^q)(a_j^q-b_0^q)\prod_{k=1}^{j-1}(a_j^q-b_k^q)(a_j^q-a_k^q)\prod_{l=j+1}^N(b_l^q-a_j^q)(a_l^q-a_j^q)\right)^{1/2},\tag{3.92}$$

where $c_{j'k'}^e$ *(resp.,* $c_{j'k'}^o$*),* $j',k'=1,\dots,N$*, are obtained from Equations* (E1) *and* (E2) *(resp., Equations* (O1) *and* (O2)*),* $\eta_{b_{j-1}^q},\eta_{a_j^q}>0$*,* $j=1,\dots,N+1$*, and*

$$\widehat{\alpha}_0^q(b_0^q)=\frac{4}{3}\mathrm{i}(-1)^N h_V^q(b_0^q)\eta_{b_0^q},\tag{3.93}$$

$$\widehat{\alpha}_1^q(b_0^q)=\mathrm{i}(-1)^N\left(\frac{2}{5}h_V^q(b_0^q)\eta_{b_0^q}\left(\sum_{l=1}^N\left(\frac{1}{b_0^q-b_l^q}+\frac{1}{b_0^q-a_l^q}\right)+\frac{1}{b_0^q-a_{N+1}^q}\right)+\frac{4}{5}(h_V^q(b_0^q))'\eta_{b_0^q}\right),\tag{3.94}$$

$$\widehat{\alpha}_0^q(a_{N+1}^q)=\frac{4}{3}h_V^q(a_{N+1}^q)\eta_{a_{N+1}^q},\tag{3.95}$$

$$\widehat{\alpha}_1^q(a_{N+1}^q)=\frac{2}{5}h_V^q(a_{N+1}^q)\eta_{a_{N+1}^q}\left(\sum_{l=1}^N\left(\frac{1}{a_{N+1}^q-b_l^q}+\frac{1}{a_{N+1}^q-a_l^q}\right)+\frac{1}{a_{N+1}^q-b_0^q}\right)+\frac{4}{5}(h_V^q(a_{N+1}^q))'\eta_{a_{N+1}^q},\tag{3.96}$$

$$\widehat{\alpha}_0^q(b_j^q)=\frac{4}{3}\mathrm{i}(-1)^{N-j}h_V^q(b_j^q)\eta_{b_j^q},\tag{3.97}$$

$$\widehat{\alpha}_1^q(b_j^q)=\mathrm{i}(-1)^{N-j}\left(\frac{2}{5}h_V^q(b_j^q)\eta_{b_j^q}\left(\sum_{\substack{k=1\\k\neq j}}^N\left(\frac{1}{b_j^q-b_k^q}+\frac{1}{b_j^q-a_k^q}\right)+\frac{1}{b_j^q-a_j^q}+\frac{1}{b_j^q-b_0^q}+\frac{1}{b_j^q-a_{N+1}^q}\right)+\frac{4}{5}(h_V^q(b_j^q))'\eta_{b_j^q}\right),\tag{3.98}$$

$$\widehat{\alpha}_0^q(a_j^q)=\frac{4}{3}(-1)^{N-j+1}h_V^q(a_j^q)\eta_{a_j^q},\tag{3.99}$$

$$\widehat{\alpha}_1^q(a_j^q)=(-1)^{N-j+1}\left(\frac{2}{5}h_V^q(a_j^q)\eta_{a_j^q}\left(\sum_{\substack{k=1\\k\neq j}}^N\left(\frac{1}{a_j^q-b_k^q}+\frac{1}{a_j^q-a_k^q}\right)+\frac{1}{a_j^q-b_j^q}+\frac{1}{a_j^q-a_{N+1}^q}+\frac{1}{a_j^q-b_0^q}\right)+\frac{4}{5}(h_V^q(a_j^q))'\eta_{a_j^q}\right),\tag{3.100}$$

and

$$\widetilde{\alpha}_0^e(1,1,\mathfrak{Z}):=2\pi\mathrm{i}\sum_{m\in\mathbb{Z}^N}(m,\widehat{\boldsymbol{\alpha}}_0^e)\mathrm{e}^{2\pi\mathrm{i}(m,\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\mathfrak{Z}+\boldsymbol{d}_e)+\pi\mathrm{i}(m,\tau^e m)},\tag{3.101}$$

$$\widetilde{\alpha}_\infty^o(1,1,\mathfrak{Z}):=2\pi\mathrm{i}\sum_{m\in\mathbb{Z}^N}(m,\widehat{\boldsymbol{\alpha}}_\infty^o)\mathrm{e}^{2\pi\mathrm{i}(m,\boldsymbol{u}_+^o(\infty)-\frac{1}{2\pi}(n+\frac{1}{2})\mathfrak{Z}+\boldsymbol{d}_o)+\pi\mathrm{i}(m,\tau^o m)},\tag{3.102}$$

where $\widehat{\boldsymbol{\alpha}}_p^q = (\widehat{\alpha}_{p,1}^q, \widehat{\alpha}_{p,2}^q, \dots, \widehat{\alpha}_{p,N}^q)$, $q \in \{e,o\}$, $p \in \{0,\infty\}$, *with*

$$\widehat{\alpha}_{0,j}^e := (-1)^{\mathcal{N}_+^e} \left(\prod_{i=1}^{N+1} |b_{i-1}^e a_i^e| \right)^{-1/2} c_{jN}^e, \qquad \widehat{\alpha}_{\infty,j}^o := c_{j1}^o, \quad j = 1, \dots, N, \tag{3.103}$$

and $\mathbf{0} := (0, 0, \dots, 0)^{\mathrm{T}}$ $(\in \mathbb{R}^N)$.

Via Proposition 3.3, the formulae

$$b_{2n}^\sharp := h^-\left[b_{2n+2}^\sharp\right], \qquad c_{2n}^\sharp := h^-\left[c_{2n+2}^\sharp\right], \qquad \beta_{2n}^\sharp := h^-\left[\beta_{2n+2}^\sharp\right], \qquad \gamma_{2n+1}^\sharp := h^-\left[\gamma_{2n+3}^\sharp\right],$$

and Equations (3.35) and (3.36), one proves the following theorem for asymptotics (as $n \to \infty$) of the coefficients of the recurrence relations (1.23)–(1.26). Theorem 3.2 (see below) presents asymptotics for $a_{2n+1}^\sharp$ (see Equation (3.104)), $b_{2n+1}^\sharp$ (see Equation (3.106)), $b_{2n+2}^\sharp$ (see Equation (3.109)), $c_{2n+2}^\sharp$ (see Equation (3.114)), $\alpha_{2n}^\sharp$ (see Equation (3.115)), $\beta_{2n+1}^\sharp$ (see Equation (3.116)), $\beta_{2n+2}^\sharp$ (see Equation (3.117)), $\gamma_{2n+3}^\sharp$ (see Equation (3.118)), $b_{2n}^\sharp$ (see Equation (3.119)), $c_{2n}^\sharp$ (see Equation (3.120)), $\beta_{2n}^\sharp$ (see Equation (3.121)), and $\gamma_{2n+1}^\sharp$ (see Equation (3.122)). Furthermore, asymptotics for $\xi_{n-1}^{(2n)}/\xi_n^{(2n)}$ (cf. Equations (3.17)) and $\xi_{-n}^{(2n+1)}/\xi_{-n-1}^{(2n+1)}$ (cf. Equations (3.18)) are presented in Equations (3.123) and (3.124), respectively.

Theorem 3.2. *Let the external field* $\widetilde{V}\colon \mathbb{R} \setminus \{0\} \to \mathbb{R}$ *satisfy conditions* (1.20)–(1.22). *Let the orthonormal L-polynomials (resp., monic orthogonal L-polynomials)* $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ *(resp.,* $\{\boldsymbol{\pi}_k(z)\}_{k\in\mathbb{Z}_0^+}$*) be as defined in Equations* (1.2) *and* (1.3) *(resp., Equations* (1.4) *and* (1.5)*), and let* $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ *satisfy the system of recurrence relations* (1.23)–(1.26). *Let* $\overset{e}{\mathrm{Y}}\colon \mathbb{C} \setminus \mathbb{R} \to \mathrm{SL}_2(\mathbb{C})$ *(resp.,* $\overset{o}{\mathrm{Y}}\colon \mathbb{C} \setminus \mathbb{R} \to \mathrm{SL}_2(\mathbb{C})$*) be the (unique) solution of* **RHP1** *(resp.,* **RHP2***) with integral representation* (1.29) *(resp.,* (1.30)*). Define the density of the 'even' (resp., 'odd') equilibrium measure,* $\mathrm{d}\mu_V^e(x)$ *(resp.,* $\mathrm{d}\mu_V^o(x)$*), as in Equation* (2.13) *(resp., Equation* (2.19)*), and set* $J_q := \operatorname{supp}(\mu_V^q) = \cup_{j=1}^{N+1}(b_{j-1}^q, a_j^q)$, $q \in \{e,o\}$, *where* $\{b_{j-1}^e, a_j^e\}_{j=1}^{N+1}$ *(resp.,* $\{b_{j-1}^o, a_j^o\}_{j=1}^{N+1}$*) satisfy the (real) n-dependent and locally solvable system of* $2(N+1)$ *moment conditions* (2.12) *(resp.,* (2.18)*).*

Suppose, furthermore, that $\widetilde{V}\colon \mathbb{R}\setminus\{0\} \to \mathbb{R}$ *is regular, namely:* (i) $h_V^q(z) \not\equiv 0$ *on* $\overline{J_q} := \cup_{k=1}^{N+1}[b_{k-1}^q, a_k^q]$, $q \in \{e,o\}$; *and* (ii) *the strict variational inequalities of Theorem* 3.1, *Equations* (3.46) *and* (3.47)*, are valid. Then,*

$$\begin{aligned}
a_{2n+1}^\sharp \underset{n\to\infty}{=} & -\frac{\omega_+^o\omega_-^o}{\eta_-^o} \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(\infty) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0) - \frac{1}{2}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)} \\
& \times \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0) + \boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0) + \boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty) + \boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(\infty) + \boldsymbol{d}_o)} \Bigg\{ 1 + \frac{1}{(n+\frac{1}{2})} \Bigg((\mathcal{R}_0^{o,\infty})_{11} - (\mathcal{R}_0^{o,\infty})_{12} \frac{\omega_-^o \mathbb{E}^2}{\mathrm{i}\omega_+^o} \\
& \times \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty) + \boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o - \boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty) - \boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o - \boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)} \\
& + (\mathcal{R}_0^{o,\infty})_{11} - (\mathcal{R}_0^{o,\infty})_{12} \frac{\omega_+^o \mathbb{E}^2}{\mathrm{i}\omega_-^o} \frac{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(\infty) + \boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(\infty) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o - \boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(\infty) - \boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(\infty) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)} \\
& \times \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o - \boldsymbol{d}_o)} - \frac{4\mathrm{i}(\mathcal{R}_1^{o,0})_{12}\mathbb{E}^2}{\eta_-^o} \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)} \\
& \times \frac{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0) + \boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0) + \boldsymbol{d}_o)} \Bigg) + O\left(\frac{c(n)}{(n+\frac{1}{2})^2} \right) \Bigg\},
\end{aligned} \tag{3.104}$$

where

$$\begin{aligned}
\mathcal{R}_1^{o,0} := \sum_{j=1}^{N+1} \Bigg(& \frac{\left(\mathcal{A}^o(b_{j-1}^o)\left(\widehat{\alpha}_1^o(b_{j-1}^o) + 2(b_{j-1}^o)^{-1}\widehat{\alpha}_0^o(b_{j-1}^o)\right) - \mathcal{B}^o(b_{j-1}^o)\widehat{\alpha}_0^o(b_{j-1}^o)\right)}{(b_{j-1}^o)^2(\widehat{\alpha}_0^o(b_{j-1}^o))^2} \\
& + \frac{\left(\mathcal{A}^o(a_j^o)\left(\widehat{\alpha}_1^o(a_j^o) + 2(a_j^o)^{-1}\widehat{\alpha}_0^o(a_j^o)\right) - \mathcal{B}^o(a_j^o)\widehat{\alpha}_0^o(a_j^o)\right)}{(a_j^o)^2(\widehat{\alpha}_0^o(a_j^o))^2} \Bigg),
\end{aligned} \tag{3.105}$$

with all parameters defined in Theorem 3.1, *Equations* (3.52)–(3.100),

$$
\begin{aligned}
b^{\sharp}_{2n+1} \underset{n\to\infty}{=} & -\frac{2\varpi^o_-}{\mathbb{E}}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(\infty)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(\infty)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)} \\
& \times\sqrt{\frac{1}{\eta^e_+}\frac{\boldsymbol{\theta}^e(\boldsymbol{u}^e_+(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(-\boldsymbol{u}^e_+(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}^e_+(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}^e_+(\infty)+\boldsymbol{d}_e)}} \\
& \times\sqrt{\frac{\mathbb{E}^2}{\eta^o_-}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(0)+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)+\boldsymbol{d}_o)}} \\
& \times\left\{1+\frac1n\left(\frac{2\mathfrak{Q}^{\flat}_{12}}{\eta^e_+}\frac{\boldsymbol{\theta}^e(\boldsymbol{u}^e_+(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(-\boldsymbol{u}^e_+(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}^e_+(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}^e_+(\infty)+\boldsymbol{d}_e)}\right)\right. \\
& +\frac{1}{(n+\frac12)}\left(\frac{2\mathbb{E}^2\mathfrak{Q}^{\sharp}_{12}}{\eta^o_-}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(0)+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)+\boldsymbol{d}_o)}\right. \\
& +(\mathcal{R}^{o,\infty}_0)_{11}+(\mathcal{R}^{o,\infty}_0)_{12}\frac{\mathrm{i}\varpi^o_+\mathbb{E}^2}{\varpi^o_-}\frac{\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(\infty)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(\infty)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)} \\
& \left.\left.\times\frac{\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(\infty)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(\infty)-\boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}\right)+O\left(\frac{c(n)}{n^2}\right)\right\} \\
& \times\exp\left(2\left(n+\frac12\right)\int_{J_o}\ln(|s|)\psi^o_V(s)\,\mathrm{d}s+\frac{n}{2}(\ell_o-\ell_e)\right),
\end{aligned}
\tag{3.106}
$$

where

$$
\mathfrak{Q}^{\flat}:=\mathrm{i}\mathcal{R}^{e,\infty}_1,\qquad\qquad \mathfrak{Q}^{\sharp}:=-\mathrm{i}\mathcal{R}^{o,0}_1,
\tag{3.107}
$$

with

$$
\mathcal{R}^{e,\infty}_1:=\sum_{j=1}^{N+1}\left(\frac{\left(\mathcal{B}^e(a^e_j)\widehat{\alpha}^e_0(a^e_j)-\mathcal{A}^e(a^e_j)\widehat{\alpha}^e_1(a^e_j)\right)}{(\widehat{\alpha}^e_0(a^e_j))^2}+\frac{\left(\mathcal{B}^e(b^e_{j-1})\widehat{\alpha}^e_0(b^e_{j-1})-\mathcal{A}^e(b^e_{j-1})\widehat{\alpha}^e_1(b^e_{j-1})\right)}{(\widehat{\alpha}^e_0(b^e_{j-1}))^2}\right),
\tag{3.108}
$$

and all parameters defined in Theorem 3.1, *Equations* (3.52)–(3.100),

$$
b^{\sharp}_{2n+2}\underset{n\to\infty}{=}\nu^{(2n+1)}_n\sqrt{\frac{h^+\left[\mathrm{e}^{n\ell_e}\eta^e_+\mathfrak{a}^{\uparrow}(n)\left(1+\frac1n\mathfrak{A}^{\uparrow}(n)+O\left(\frac{c(n)}{n^2}\right)\right)\right]}{\mathrm{e}^{n\ell_o}\eta^o_-\mathfrak{b}^{\downarrow}(n)\left(1+\frac{1}{(n+\frac12)}\mathfrak{B}^{\downarrow}(n)+O\left(\frac{c(n)}{(n+\frac12)^2}\right)\right)}},
\tag{3.109}
$$

where asymptotics (as $n\to\infty$*) of* $\nu^{(2n+1)}_n$ *is given in Theorem* 3.1, *Equation* (3.49), *with*

$$
\mathfrak{a}^{\uparrow}(n):=\frac{\boldsymbol{\theta}^e(\boldsymbol{u}^e_+(\infty)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(-\boldsymbol{u}^e_+(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}^e_+(\infty)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}^e_+(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)},
\tag{3.110}
$$

$$
\mathfrak{b}^{\downarrow}(n):=\frac{1}{\mathbb{E}^2}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(0)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)},
\tag{3.111}
$$

$$
\mathfrak{A}^{\uparrow}(n):=-\frac{4\mathrm{i}(\mathcal{R}^{e,\infty}_1)_{12}}{\eta^e_+}\frac{\boldsymbol{\theta}^e(-\boldsymbol{u}^e_+(\infty)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}^e_+(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}^e_+(\infty)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(-\boldsymbol{u}^e_+(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)},
\tag{3.112}
$$

$$
\mathfrak{B}^{\downarrow}(n):=\frac{4\mathrm{i}(\mathcal{R}^{o,0}_1)_{12}\mathbb{E}^2}{\eta^o_-}\frac{\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(0)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)},
\tag{3.113}
$$

$$
c^{\sharp}_{2n+2}\underset{n\to\infty}{=}\sqrt{\frac{h^+\left[\mathrm{e}^{n\ell_e}\eta^e_+\mathfrak{a}^{\uparrow}(n)\left(1+\frac1n\mathfrak{A}^{\uparrow}(n)+O\left(\frac{c(n)}{n^2}\right)\right)\right]}{\mathrm{e}^{n\ell_e}\eta^e_+\mathfrak{a}^{\uparrow}(n)\left(1+\frac1n\mathfrak{A}^{\uparrow}(n)+O\left(\frac{c(n)}{n^2}\right)\right)}},
\tag{3.114}
$$

$$\alpha_{2n}^{\sharp} \underset{n\to\infty}{=} -\frac{\omega_+^e\omega_+^o}{4\mathbb{E}}\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)+\boldsymbol{d}_e)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)+\boldsymbol{d}_o)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}$$
$$\times\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}\Bigg\{1+\frac1n\Bigg((\mathcal{R}_0^{e,0})_{11}+(\mathcal{R}_0^{e,0})_{12}\frac{\omega_-^e}{\mathrm{i}\omega_+^e}$$
$$\times\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}\Bigg)$$
$$+\frac{1}{(n+\frac12)}\Bigg((\mathcal{R}_0^{o,\infty})_{11}-(\mathcal{R}_0^{o,\infty})_{12}\frac{\omega_-^o\mathbb{E}^2}{\mathrm{i}\omega_+^o}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}$$
$$\times\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(\infty)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}\Bigg)+O\Bigg(\frac{c(n)}{n^2}\Bigg)\Bigg\}$$
$$\times\exp\Bigg(2n\Bigg(\int_{J_e}\ln(|s|)\psi_V^e(s)\,\mathrm{d}s+\mathrm{i}\pi\int_{J_e\cap\mathbb{R}_+}\psi_V^e(s)\,\mathrm{d}s\Bigg)-2\Bigg(n+\frac12\Bigg)\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s\Bigg), \tag{3.115}$$

$$\beta_{2n+1}^{\sharp} \underset{n\to\infty}{=} \frac{\omega_+^e}{2}\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}$$
$$\times\sqrt{\frac{1}{\eta_+^e}\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}}$$
$$\times\left(\frac{\mathbb{E}^2}{\eta_-^o}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}\right)^{-1/2}$$
$$\times\Bigg\{1+\frac{1}{(n+\frac12)}\Bigg(-\frac{2\mathbb{E}^2\mathfrak{Q}_{12}^{\natural}}{\eta_-^o}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}\Bigg)$$
$$+\frac1n\Bigg(\frac{2\mathfrak{Q}_{12}^{\flat}}{\eta_+^e}\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}+(\mathcal{R}_0^{e,0})_{11}+(\mathcal{R}_0^{e,0})_{12}\frac{\omega_-^e}{\mathrm{i}\omega_+^e}$$
$$\times\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(0)-\boldsymbol{d}_e)\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}\Bigg)+O\Bigg(\frac{c(n)}{n^2}\Bigg)\Bigg\}$$
$$\times\exp\Bigg(2n\Bigg(\int_{J_e}\ln(|s|)\psi_V^e(s)\,\mathrm{d}s+\mathrm{i}\pi\int_{J_e\cap\mathbb{R}_+}\psi_V^e(s)\,\mathrm{d}s\Bigg)+\frac{n}{2}(\ell_o-\ell_e)\Bigg), \tag{3.116}$$

$$\beta_{2n+2}^{\sharp} \underset{n\to\infty}{=} -\Big(h^+\Big[\nu_n^{(2n+1)}\Big]\Big)\sqrt{\frac{h^+\Big[\mathrm{e}^{n\ell_e}\eta_+^e\mathfrak{a}^{\uparrow}(n)\Big(1+\frac1n\mathfrak{A}^{\uparrow}(n)+O\Big(\frac{c(n)}{n^2}\Big)\Big)\Big]}{\mathrm{e}^{n\ell_o}\eta_-^o\mathfrak{b}^{\downarrow}(n)\Big(1+\frac{1}{(n+\frac12)}\mathfrak{B}^{\downarrow}(n)+O\Big(\frac{c(n)}{(n+\frac12)^2}\Big)\Big)}}, \tag{3.117}$$

$$\gamma_{2n+3}^{\sharp} \underset{n\to\infty}{=} \sqrt{\frac{h^+\Big[\mathrm{e}^{n\ell_o}\eta_-^o\mathfrak{b}^{\downarrow}(n)\Big(1+\frac{1}{(n+\frac12)}\mathfrak{B}^{\downarrow}(n)+O\Big(\frac{c(n)}{(n+\frac12)^2}\Big)\Big)\Big]}{\mathrm{e}^{n\ell_o}\eta_-^o\mathfrak{b}^{\downarrow}(n)\Big(1+\frac{1}{(n+\frac12)}\mathfrak{B}^{\downarrow}(n)+O\Big(\frac{c(n)}{(n+\frac12)^2}\Big)\Big)}}, \tag{3.118}$$

$$b_{2n}^{\sharp} \underset{n\to\infty}{=} h^-\Big[\nu_n^{(2n+1)}\Big]\sqrt{\frac{\mathrm{e}^{n\ell_e}\eta_+^e\mathfrak{a}^{\uparrow}(n)\Big(1+\frac1n\mathfrak{A}^{\uparrow}(n)+O\Big(\frac{c(n)}{n^2}\Big)\Big)}{h^-\Big[\mathrm{e}^{n\ell_o}\eta_-^o\mathfrak{b}^{\downarrow}(n)\Big(1+\frac{1}{(n+\frac12)}\mathfrak{B}^{\downarrow}(n)+O\Big(\frac{c(n)}{(n+\frac12)^2}\Big)\Big)\Big]}}, \tag{3.119}$$

$$c_{2n}^{\sharp} \underset{n\to\infty}{=} \sqrt{\frac{\mathrm{e}^{n\ell_e}\eta_+^e\mathfrak{a}^{\uparrow}(n)\Big(1+\frac1n\mathfrak{A}^{\uparrow}(n)+O\Big(\frac{c(n)}{n^2}\Big)\Big)}{h^-\Big[\mathrm{e}^{n\ell_e}\eta_+^e\mathfrak{a}^{\uparrow}(n)\Big(1+\frac1n\mathfrak{A}^{\uparrow}(n)+O\Big(\frac{c(n)}{n^2}\Big)\Big)\Big]}}, \tag{3.120}$$

$$\beta^{\sharp}_{2n} \underset{n\to\infty}{=} -\nu_n^{(2n+1)} \sqrt{\frac{\mathrm{e}^{n\ell_e}\eta^e_+ \mathfrak{a}^{\uparrow}(n)\left(1+\frac{1}{n}\mathfrak{A}^{\uparrow}(n)+O\left(\frac{c(n)}{n^2}\right)\right)}{h^-\left[\mathrm{e}^{n\ell_o}\eta^o_- \mathfrak{b}^{\downarrow}(n)\left(1+\frac{1}{(n+\frac12)}\mathfrak{B}^{\downarrow}(n)+O\left(\frac{c(n)}{(n+\frac12)^2}\right)\right)\right]}}, \tag{3.121}$$

$$\gamma^{\sharp}_{2n+1} \underset{n\to\infty}{=} \sqrt{\frac{\mathrm{e}^{n\ell_o}\eta^o_- \mathfrak{b}^{\downarrow}(n)\left(1+\frac{1}{(n+\frac12)}\mathfrak{B}^{\downarrow}(n)+O\left(\frac{c(n)}{(n+\frac12)^2}\right)\right)}{h^-\left[\mathrm{e}^{n\ell_o}\eta^o_- \mathfrak{b}^{\downarrow}(n)\left(1+\frac{1}{(n+\frac12)}\mathfrak{B}^{\downarrow}(n)+O\left(\frac{c(n)}{(n+\frac12)^2}\right)\right)\right]}}, \tag{3.122}$$

and $n\to\infty$ asymptotics for $a^{\sharp}_{2n}$ and $\alpha^{\sharp}_{2n+1}$ are obtained by substituting, formally, asymptotics (3.106), (3.114)–(3.117), (3.119), (3.121), *and* (3.122) *into Equations* (3.35) *and* (3.56). *Furthermore,*

$$\frac{\xi^{(2n)}_{n-1}}{\xi^{(2n)}_{n}} \underset{n\to\infty}{=} -2n\int_{J_e} s\psi^e_V(s)\,\mathrm{d}s + \frac{\alpha^e_\infty(1,1,\mathbf{0})}{\boldsymbol{\theta}^e(\boldsymbol{u}^e_+(\infty)+\boldsymbol{d}_e)} - \frac{\alpha^e_\infty(1,1,\boldsymbol{\Omega}^e)}{\boldsymbol{\theta}^e(\boldsymbol{u}^e_+(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)} + \frac{1}{n}(\mathcal{R}^{e,\infty}_1)_{11} + O\left(\frac{c(n)}{n^2}\right), \tag{3.123}$$

$$\begin{aligned}\frac{\xi^{(2n+1)}_{-n}}{\xi^{(2n+1)}_{-n-1}} \underset{n\to\infty}{=}& -2\left(n+\frac12\right)\int_{J_o} s^{-1}\psi^o_V(s)\,\mathrm{d}s - \frac{\alpha^o_0(1,1,\mathbf{0})}{\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)+\boldsymbol{d}_o)} + \frac{\alpha^o_0(1,1,\boldsymbol{\Omega}^o)}{\boldsymbol{\theta}^o(\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)} + \frac{1}{(n+\frac12)}(\mathcal{R}^{o,0}_1)_{11}\\ &+ O\left(\frac{c(n)}{(n+\frac12)^2}\right),\end{aligned} \tag{3.124}$$

where $\mathcal{R}^{o,0}_1$ (resp., $\mathcal{R}^{e,\infty}_1$) is defined in Equation (3.105) *(resp., Equation* (3.108)),

$$\alpha^e_\infty(1,1,\mathfrak{Z}) := 2\pi\mathrm{i}\sum_{m\in\mathbb{Z}^N}(m,\widehat{\boldsymbol{\alpha}}^e_\infty)\mathrm{e}^{2\pi\mathrm{i}(m,\boldsymbol{u}^e_+(\infty)-\frac{n}{2\pi}\mathfrak{Z}+\boldsymbol{d}_e)+\pi\mathrm{i}(m,\tau^e m)}, \tag{3.125}$$

$$\alpha^o_0(1,1,\mathfrak{Z}) := 2\pi\mathrm{i}\sum_{m\in\mathbb{Z}^N}(m,\widehat{\boldsymbol{\alpha}}^o_0)\mathrm{e}^{2\pi\mathrm{i}(m,\boldsymbol{u}^o_+(0)-\frac{1}{2\pi}(n+\frac12)\mathfrak{Z}+\boldsymbol{d}_o)+\pi\mathrm{i}(m,\tau^o m)}, \tag{3.126}$$

where $\widehat{\boldsymbol{\alpha}}^q_p = (\widehat{\alpha}^q_{p,1}, \widehat{\alpha}^q_{p,2}, \ldots, \widehat{\alpha}^q_{p,N})$, $q\in\{e,o\}$, $p\in\{0,\infty\}$, with

$$\widehat{\alpha}^e_{\infty,j} := c^e_{j1}, \qquad \widehat{\alpha}^o_{0,j} := (-1)^{N^o_+}\left(\prod_{i=1}^{N+1}|b^o_{i-1}a^o_i|\right)^{-1/2} c^o_{jN}, \quad j=1,\ldots,N, \tag{3.127}$$

where c^e_{j1} (resp., c^o_{jN}), $j=1,\ldots,N$, are obtained from Equations (E1) *and* (E2) *(resp., Equations* (O1) *and* (O2)), *and* $\mathbf{0} := (0,0,\ldots,0)^{\mathrm{T}}$ $(\in\mathbb{R}^N)$.

Recalling Equations (1.9), asymptotics (as $n\to\infty$) for the Hankel determinant ratios

$$H^{(-2n+1)}_{2n}/H^{(-2n)}_{2n} \qquad \text{and} \qquad H^{(-2n-1)}_{2n+1}/H^{(-2n)}_{2n+1}$$

have already been presented in Theorem 3.1, Equations (3.48) and (3.49), respectively. Via Equations (1.18) and (1.19), and asymptotics for $\xi^{(2n)}_n$ (resp., $\xi^{(2n+1)}_{-n-1}$) given in Appendix A, Theorem A.3 (resp., Appendix B, Theorem B.3), one proves the following theorem concerning asymptotics of the remaining Hankel determinant ratios $H^{(-2n-2)}_{2n+2}/H^{(-2n)}_{2n}$ (cf. Equation (1.18)) and $H^{(-2n-2)}_{2n+3}/H^{(-2n)}_{2n+1}$ (cf. Equation (1.19)); see, in particular, Equations (3.128) and (3.129).

Theorem 3.3. *Let the external field $\widetilde{V}\colon \mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfy conditions* (1.20)–(1.22). *Let $H^{(m)}_k$, $(m,k)\in\mathbb{Z}\times\mathbb{N}$, be the Hankel determinants associated with the bi-infinite, real-valued, strong moment sequence $\left\{c_j = \int_{\mathbb{R}} s^j \exp(-n\widetilde{V}(s))\,\mathrm{d}s,\ n\in\mathbb{N}\right\}_{j\in\mathbb{Z}}$ defined in Equations* (1.1). *Let the orthonormal L-polynomials (resp., monic orthogonal L-polynomials) $\{\phi_k(z)\}_{k\in\mathbb{Z}^+_0}$ (resp., $\{\boldsymbol{\pi}_k(z)\}_{k\in\mathbb{Z}^+_0}$) be as defined in Equations* (1.2) *and* (1.3) *(resp., Equations* (1.4) *and* (1.5)). *Let $\overset{e}{\mathrm{Y}}\colon \mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$ (resp., $\overset{o}{\mathrm{Y}}\colon \mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$) be the (unique) solution of* **RHP1** *(resp.,* **RHP2**) *with integral representation* (1.29) *(resp.,* (1.30)). *Define the density of the 'even' (resp., 'odd') equilibrium measure, $\mathrm{d}\mu^e_V(x)$ (resp., $\mathrm{d}\mu^o_V(x)$), as in Equation* (2.13) *(resp., Equation* (2.19)), *and set $J_q := \mathrm{supp}(\mu^q_V) = \cup_{j=1}^{N+1}(b^q_{j-1}, a^q_j)$, $q\in\{e,o\}$, where $\{b^e_{j-1}, a^e_j\}_{j=1}^{N+1}$ (resp., $\{b^o_{j-1}, a^o_j\}_{j=1}^{N+1}$) satisfy the (real) n-dependent and locally solvable system of $2(N+1)$ moment conditions* (2.12) *(resp.,* (2.18)).

Suppose, furthermore, that $\widetilde{V}\colon \mathbb{R}\setminus\{0\}\to\mathbb{R}$ *is regular, namely:* (i) $h_V^q(z)\not\equiv 0$ *on* $\overline{J_q}:=\cup_{k=1}^{N+1}[b_{k-1}^q,a_k^q]$, $q\in\{e,o\}$; *and* (ii) *the strict variational inequalities of Theorem* 3.1, *Equations* (3.46) *and* (3.47), *are valid. Then,*

$$\begin{aligned}\frac{H_{2n+2}^{(-2n-2)}}{H_{2n}^{(-2n)}} \underset{n\to\infty}{=} & \frac{\pi^2\eta_+^e\eta_-^o}{4\mathbb{E}^2}\frac{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}\\ &\times\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}\left\{1+\frac1n\left(-\frac{4\mathfrak{Q}_{12}^\flat}{\eta_+^e}\frac{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}\right.\right.\\ &\times\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}\Big)+\frac{1}{(n+\frac12)}\left(-\frac{4\mathbb{E}^2\mathfrak{Q}_{12}^\natural}{\eta_-^o}\frac{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}\right.\\ &\left.\left.\times\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}\right)+O\left(\frac{c(n)}{n^2}\right)\right\}\exp(n(\ell_e+\ell_o))\,,\end{aligned}\tag{3.128}$$

and

$$\begin{aligned}\frac{H_{2n+3}^{(-2n-2)}}{H_{2n+1}^{(-2n)}} \underset{n\to\infty}{=} & \frac{\pi^2\mathrm{e}^{n\ell_o}\eta_-^o}{4\mathbb{E}^2}\frac{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}\\ &\times\left\{1+\frac{1}{(n+\frac12)}\left(-\frac{4\mathbb{E}^2\mathfrak{Q}_{12}^\natural}{\eta_-^o}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac12)\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}\right)\right.\\ &\left.+O\left(\frac{c(n)}{(n+\frac12)^2}\right)\right\}\left\{h^+\left[\frac{\mathrm{e}^{-n\ell_e}}{\eta_+^e}\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}\right.\right.\\ &\left.\left.\times\left(1+\frac1n\left(\frac{4\mathfrak{Q}_{12}^\flat}{\eta_+^e}\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}\right)+O\left(\frac{c(n)}{n^2}\right)\right)\right]\right\}^{-1},\end{aligned}\tag{3.129}$$

with all parameters defined in Theorem 3.1, *Equations* (3.52)–(3.100), *and Theorem* 3.2, *Equations* (3.105), (3.107), *and* (3.108).

Acknowledgements

K. T.-R. McLaughlin was supported, in part, by National Science Foundation Grant Nos. DMS–0200749 and DMS–0451495; he also acknowledges the partial support of NATO Collaborative Linkage Grant 'Orthogonal Polynomials: Theory, Applications and Generalizations', Ref. No. PST.CLG.979738. X. Zhou was supported, in part, by National Science Foundation Grant No. DMS–0300844.

Appendix A. Asymptotics for $\overset{e}{\mathrm{Y}}(z)$

In this appendix, large-n asymptotics for the coefficients of the expansions (3.1) and (3.2) for the solution of **RHP1**, with integral representation (1.29), are presented (for complete details and proofs, see [21]).

Remark A.1. In [21], the following formula was established for complex z away from an open neighbourhood around $\overline{J_e} := \cup_{j=1}^{N+1}[b_{j-1}^e, a_j^e]$:

$$\overset{e}{\mathrm{Y}}(z) = \mathrm{e}^{\frac{n\ell_e}{2}\sigma_3}\mathcal{R}^e(z)\overset{e}{m}^{\infty}(z)\mathrm{e}^{n(g^e(z)-\frac{\ell_e}{2}+\int_{J_e}\ln(s)\psi_V^e(s)\,\mathrm{d}s)\sigma_3},$$

where ℓ_e, $g^e(z)$, and $\psi_V^e(z)$ have been defined heretofore, the matrix-valued function $\overset{e}{m}^{\infty}(z)$ is determined explicitly, and the matrix-valued function $\mathcal{R}^e(z)$ solves a "small norm" RHP for which a complete asymptotic expansion as $n \to \infty$ for $z \in \mathbb{C}$ is computed. The above explicit formula for the solution of **RHP1** yields an explicit asymptotic description for the even degree OLPs in question (for complete details and proofs, see [21]). ■

Remark A.2. In Theorems A.1 and A.2 below, only $n \to \infty$ asymptotics for $\mathcal{R}_j^{e,\infty}(n)$, $j = 1, 2$, and $\mathcal{R}_{k-1}^{e,0}(n)$, $k = 1, 2, 3$, are given, as they are all that are actually necessary in order to obtain the results of Theorems 3.1–3.3: finite-n formulae for $\mathcal{R}_j^{e,\infty}(n)$, $j = 1, 2$, and $\mathcal{R}_{k-1}^{e,0}(n)$, $k = 1, 2, 3$, are given in [21]; furthermore, the 'symbol' $c(n)$ appearing in the asymptotic (as $n \to \infty$) expressions for $\mathcal{R}_j^{e,\infty}(n)$, $j = 1, 2$, and $\mathcal{R}_{k-1}^{e,0}(n)$, $k = 1, 2, 3$, is to be understood as a uniformly bounded ($O(1)$), $\mathrm{M}_2(\mathbb{C})$-valued, n-dependent function, that is, $\mathrm{M}_2(\mathbb{C}) \ni c(n) =_{n\to\infty} O(1)$. ■

Theorem A.1 ([21]). *Let the external field $\widetilde{V}: \mathbb{R} \setminus \{0\} \to \mathbb{R}$ satisfy conditions (1.20)–(1.22). Let the orthonormal L-polynomials (resp., monic orthogonal L-polynomials) $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ (resp., $\{\boldsymbol{\pi}_k(z)\}_{k\in\mathbb{Z}_0^+}$) be as defined in Equations (1.2) and (1.3) (resp., Equations (1.4) and (1.5)). Let $\overset{e}{\mathrm{Y}}: \mathbb{C} \setminus \mathbb{R} \to \mathrm{SL}_2(\mathbb{C})$ be the unique solution of* **RHP1** *with integral representation (1.29). Define the density of the 'even' equilibrium measure, $\mathrm{d}\mu_V^e(x)$, as in Equation (2.13), and set $J_e := \mathrm{supp}(\mu_V^e) = \cup_{j=1}^{N+1}(b_{j-1}^e, a_j^e)$, where $\{b_{j-1}^e, a_j^e\}_{j=1}^{N+1}$ satisfy the (real) n-dependent and locally solvable system of $2(N+1)$ moment conditions (2.12).*

Suppose, furthermore, that $\widetilde{V}: \mathbb{R} \setminus \{0\} \to \mathbb{R}$ is regular, namely: (i) $h_V^e(z) \not\equiv 0$ *on* $\overline{J_e} := \cup_{j=1}^{N+1}[b_{j-1}^e, a_j^e]$; (ii)

$$4\int_{J_e}\ln(|x-s|)\,\mathrm{d}\mu_V^e(s) - 2\ln|x| - \widetilde{V}(x) - \ell_e = 0, \quad x \in \overline{J_e},$$

which defines the 'even' variational constant, ℓ_e ($\in \mathbb{R}$) (the same on each compact interval $[b_{j-1}^e, a_j^e]$, $j = 1, \ldots, N+1$), and

$$4\int_{J_e}\ln(|x-s|)\,\mathrm{d}\mu_V^e(s) - 2\ln|x| - \widetilde{V}(x) - \ell_e < 0, \quad x \in \mathbb{R} \setminus \overline{J_e};$$

(iii)

$$g_+^e(z) + g_-^e(z) - \widetilde{V}(z) - \ell_e + 2Q_e < 0, \quad z \in \mathbb{R} \setminus \overline{J_e},$$

where $g^e(z)$ is defined by Equation (2.15), $g_\pm^e(z) := \lim_{\varepsilon\downarrow 0} g^e(z \pm \mathrm{i}\varepsilon)$, and Q_e is defined in Equation (2.16); and (iv)

$$\mathrm{i}(g_+^e(z) - g_-^e(z))' > 0, \quad z \in J_e.$$

Then,

$$\overset{e}{\mathrm{Y}}(z)z^{-n\sigma_3} \underset{z\to\infty}{=} \mathrm{I} + \frac{1}{z}\mathrm{Y}_1^{e,\infty} + \frac{1}{z^2}\mathrm{Y}_2^{e,\infty} + O\left(\frac{1}{z^3}\right),$$

where

$$(\mathrm{Y}_1^{e,\infty})_{11} = -2n\int_{J_e} s\psi_V^e(s)\,\mathrm{d}s + (\overset{e}{m}_1^{\infty})_{11} + (\mathcal{R}_1^{e,\infty}(n))_{11},$$
$$(\mathrm{Y}_1^{e,\infty})_{12} = \mathrm{e}^{n\ell_e}\left((\overset{e}{m}_1^{\infty})_{12} + (\mathcal{R}_1^{e,\infty}(n))_{12}\right),$$

$$
\begin{aligned}
(\mathrm{Y}_1^{e,\infty})_{21} &= \mathrm{e}^{-n\ell_e}\Big((\overset{e}{m}{}_1^{\infty})_{21}+(\mathcal{R}_1^{e,\infty}(n))_{21}\Big),\\
(\mathrm{Y}_1^{e,\infty})_{22} &= 2n\int_{J_e} s\psi_V^e(s)\,\mathrm{d}s+(\overset{e}{m}{}_1^{\infty})_{22}+(\mathcal{R}_1^{e,\infty}(n))_{22},\\
(\mathrm{Y}_2^{e,\infty})_{11} &= 2n^2\left(\int_{J_e} s\psi_V^e(s)\,\mathrm{d}s\right)^2-n\int_{J_e} s^2\psi_V^e(s)\,\mathrm{d}s-2n\Big((\overset{e}{m}{}_1^{\infty})_{11}+(\mathcal{R}_1^{e,\infty}(n))_{11}\Big)\int_{J_e} s\psi_V^e(s)\,\mathrm{d}s\\
&\quad+(\overset{e}{m}{}_2^{\infty})_{11}+(\mathcal{R}_2^{e,\infty}(n))_{11}+(\mathcal{R}_1^{e,\infty}(n))_{11}(\overset{e}{m}{}_1^{\infty})_{11}+(\mathcal{R}_1^{e,\infty}(n))_{12}(\overset{e}{m}{}_1^{\infty})_{21},\\
(\mathrm{Y}_2^{e,\infty})_{12} &= \mathrm{e}^{n\ell_e}\left(2n\Big((\overset{e}{m}{}_1^{\infty})_{12}+(\mathcal{R}_1^{e,\infty}(n))_{12}\Big)\int_{J_e} s\psi_V^e(s)\,\mathrm{d}s+(\overset{e}{m}{}_2^{\infty})_{12}+(\mathcal{R}_2^{e,\infty}(n))_{12}\right.\\
&\quad\left.+(\mathcal{R}_1^{e,\infty}(n))_{11}(\overset{e}{m}{}_1^{\infty})_{12}+(\mathcal{R}_1^{e,\infty}(n))_{12}(\overset{e}{m}{}_1^{\infty})_{22}\right),\\
(\mathrm{Y}_2^{e,\infty})_{21} &= \mathrm{e}^{-n\ell_e}\left(-2n\Big((\overset{e}{m}{}_1^{\infty})_{21}+(\mathcal{R}_1^{e,\infty}(n))_{21}\Big)\int_{J_e} s\psi_V^e(s)\,\mathrm{d}s+(\overset{e}{m}{}_2^{\infty})_{21}+(\mathcal{R}_2^{e,\infty}(n))_{21}\right.\\
&\quad\left.+(\mathcal{R}_1^{e,\infty}(n))_{21}(\overset{e}{m}{}_1^{\infty})_{11}+(\mathcal{R}_1^{e,\infty}(n))_{22}(\overset{e}{m}{}_1^{\infty})_{21}\right),\\
(\mathrm{Y}_2^{e,\infty})_{22} &= 2n^2\left(\int_{J_e} s\psi_V^e(s)\,\mathrm{d}s\right)^2+n\int_{J_e} s^2\psi_V^e(s)\,\mathrm{d}s+2n\Big((\overset{e}{m}{}_1^{\infty})_{22}+(\mathcal{R}_1^{e,\infty}(n))_{22}\Big)\int_{J_e} s\psi_V^e(s)\,\mathrm{d}s\\
&\quad+(\overset{e}{m}{}_2^{\infty})_{22}+(\mathcal{R}_2^{e,\infty}(n))_{22}+(\mathcal{R}_1^{e,\infty}(n))_{21}(\overset{e}{m}{}_1^{\infty})_{12}+(\mathcal{R}_1^{e,\infty}(n))_{22}(\overset{e}{m}{}_1^{\infty})_{22},
\end{aligned}
$$

with

$$
\begin{aligned}
(\overset{e}{m}{}_1^{\infty})_{11} &= \frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}\left(\frac{\theta_\infty^e(1,1,\boldsymbol{\Omega}^e)\alpha_\infty^e(1,1,\mathbf{0})-\alpha_\infty^e(1,1,\boldsymbol{\Omega}^e)\theta_\infty^e(1,1,\mathbf{0})}{(\theta_\infty^e(1,1,\mathbf{0}))^2}\right),\\
(\overset{e}{m}{}_1^{\infty})_{12} &= \frac{1}{4\mathrm{i}}\left(\sum_{k=1}^{N+1}(b_{k-1}^e-a_k^e)\right)\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)\theta_\infty^e(-1,1,\boldsymbol{\Omega}^e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)\theta_\infty^e(-1,1,\mathbf{0})},\\
(\overset{e}{m}{}_1^{\infty})_{21} &= -\frac{1}{4\mathrm{i}}\left(\sum_{k=1}^{N+1}(b_{k-1}^e-a_k^e)\right)\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)\theta_\infty^e(1,-1,\boldsymbol{\Omega}^e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)\theta_\infty^e(1,-1,\mathbf{0})},\\
(\overset{e}{m}{}_1^{\infty})_{22} &= \frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}\left(\frac{\theta_\infty^e(-1,-1,\boldsymbol{\Omega}^e)\alpha_\infty^e(-1,-1,\mathbf{0})-\alpha_\infty^e(-1,-1,\boldsymbol{\Omega}^e)\theta_\infty^e(-1,-1,\mathbf{0})}{(\theta_\infty^e(-1,-1,\mathbf{0}))^2}\right),\\
(\overset{e}{m}{}_2^{\infty})_{11} &= \Big(\theta_\infty^e(1,1,\boldsymbol{\Omega}^e)\Big(\beta_\infty^e(1,1,\mathbf{0})\theta_\infty^e(1,1,\mathbf{0})+(\alpha_\infty^e(1,1,\mathbf{0}))^2\Big)-\alpha_\infty^e(1,1,\boldsymbol{\Omega}^e)\alpha_\infty^e(1,1,\mathbf{0})\theta_\infty^e(1,1,\mathbf{0})\\
&\quad-\beta_\infty^e(1,1,\boldsymbol{\Omega}^e)(\theta_\infty^e(1,1,\mathbf{0}))^2\Big)\frac{(\theta_\infty^e(1,1,\mathbf{0}))^{-3}\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}+\frac{1}{32}\left(\sum_{k=1}^{N+1}(b_{k-1}^e-a_k^e)\right)^2,\\
(\overset{e}{m}{}_2^{\infty})_{12} &= \frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}\left(\left(\frac{\theta_\infty^e(-1,1,\boldsymbol{\Omega}^e)\alpha_\infty^e(-1,1,\mathbf{0})-\alpha_\infty^e(-1,1,\boldsymbol{\Omega}^e)\theta_\infty^e(-1,1,\mathbf{0})}{(\theta_\infty^e(-1,1,\mathbf{0}))^2}\right)\right.\\
&\quad\left.\times\frac{1}{4\mathrm{i}}\left(\sum_{k=1}^{N+1}(b_{k-1}^e-a_k^e)\right)+\frac{1}{8\mathrm{i}}\left(\sum_{k=1}^{N+1}((b_{k-1}^e)^2-(a_k^e)^2)\right)\frac{\theta_\infty^e(-1,1,\boldsymbol{\Omega}^e)}{\theta_\infty^e(-1,1,\mathbf{0})}\right),\\
(\overset{e}{m}{}_2^{\infty})_{21} &= -\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}\left(\left(\frac{\theta_\infty^e(1,-1,\boldsymbol{\Omega}^e)\alpha_\infty^e(1,-1,\mathbf{0})-\alpha_\infty^e(1,-1,\boldsymbol{\Omega}^e)\theta_\infty^e(1,-1,\mathbf{0})}{(\theta_\infty^e(1,-1,\mathbf{0}))^2}\right)\right.\\
&\quad\left.\times\frac{1}{4\mathrm{i}}\left(\sum_{k=1}^{N+1}(b_{k-1}^e-a_k^e)\right)+\frac{1}{8\mathrm{i}}\left(\sum_{k=1}^{N+1}((b_{k-1}^e)^2-(a_k^e)^2)\right)\frac{\theta_\infty^e(1,-1,\boldsymbol{\Omega}^e)}{\theta_\infty^e(1,-1,\mathbf{0})}\right),\\
(\overset{e}{m}{}_2^{\infty})_{22} &= \Big(\theta_\infty^e(-1,-1,\boldsymbol{\Omega}^e)\Big(\beta_\infty^e(-1,-1,\mathbf{0})\theta_\infty^e(-1,-1,\mathbf{0})+(\alpha_\infty^e(-1,-1,\mathbf{0}))^2\Big)-\alpha_\infty^e(-1,-1,\boldsymbol{\Omega}^e)\\
&\quad\times\alpha_\infty^e(-1,-1,\mathbf{0})\theta_\infty^e(-1,-1,\mathbf{0})-\beta_\infty^e(-1,-1,\boldsymbol{\Omega}^e)(\theta_\infty^e(-1,-1,\mathbf{0}))^2\Big)(\theta_\infty^e(-1,-1,\mathbf{0}))^{-3}\\
&\quad\times\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}+\frac{1}{32}\left(\sum_{k=1}^{N+1}(b_{k-1}^e-a_k^e)\right)^2,
\end{aligned}
$$

and $(\star)_{ij}$, $i,j=1,2$, *denoting the* $(i\,j)$*-element of* $\star$, *where, for* $\varepsilon_1,\varepsilon_2=\pm1$,

$$\theta^e_\infty(\varepsilon_1,\varepsilon_2,\mathcal{Z}):=\boldsymbol{\theta}^e(\varepsilon_1\boldsymbol{u}^e_+(\infty)-\tfrac{n}{2\pi}\mathcal{Z}+\varepsilon_2\boldsymbol{d}_e),$$
$$\alpha^e_\infty(\varepsilon_1,\varepsilon_2,\mathcal{Z}):=2\pi\mathrm{i}\varepsilon_1\sum_{m\in\mathbb{Z}^N}(m,\widehat{\boldsymbol{\alpha}}^e_\infty)\mathrm{e}^{2\pi\mathrm{i}(m,\varepsilon_1\boldsymbol{u}^e_+(\infty)-\frac{n}{2\pi}\mathcal{Z}+\varepsilon_2\boldsymbol{d}_e)+\pi\mathrm{i}(m,\tau^e m)},$$

where $\boldsymbol{u}^e_+(\infty)=\int_{a^e_{N+1}}^{\infty^+}\boldsymbol{\omega}^e$, $\widehat{\boldsymbol{\alpha}}^e_\infty=(\widehat{\alpha}^e_{\infty,1},\widehat{\alpha}^e_{\infty,2},\dots,\widehat{\alpha}^e_{\infty,N})$, *with* $\widehat{\alpha}^e_{\infty,j}:=c^e_{j1}$, $j=1,\dots,N$, *and*

$$\beta^e_\infty(\varepsilon_1,\varepsilon_2,\mathcal{Z}):=2\pi\sum_{m\in\mathbb{Z}^N}\left(\pi(m,\widehat{\boldsymbol{\alpha}}^e_\infty)^2+\mathrm{i}\varepsilon_1(m,\widehat{\boldsymbol{\beta}}^e_\infty)\right)\mathrm{e}^{2\pi\mathrm{i}(m,\varepsilon_1\boldsymbol{u}^e_+(\infty)-\frac{n}{2\pi}\mathcal{Z}+\varepsilon_2\boldsymbol{d}_e)+\pi\mathrm{i}(m,\tau^e m)},$$

where $\widehat{\boldsymbol{\beta}}^e_\infty=(\widehat{\beta}^e_{\infty,1},\widehat{\beta}^e_{\infty,2},\dots,\widehat{\beta}^e_{\infty,N})$, *with* $\widehat{\beta}^e_{\infty,j}:=\frac12(c^e_{j2}+\frac12c^e_{j1}\sum_{i=1}^{N+1}(b^e_{i-1}+a^e_i))$, $j=1,\dots,N$, *where* c^e_{j1},c^e_{j2}, $j=1,\dots,N$, *are obtained from Equations* (E1) *and* (E2), *and*

$$\mathcal{R}^{e,\infty}_1(n)\underset{n\to\infty}{=}\frac1n\sum_{j=1}^{N+1}\left(\frac{\left(\mathcal{B}^e(a^e_j)\widehat{\alpha}^e_0(a^e_j)-\mathcal{A}^e(a^e_j)\widehat{\alpha}^e_1(a^e_j)\right)}{(\widehat{\alpha}^e_0(a^e_j))^2}+\frac{\left(\mathcal{B}^e(b^e_{j-1})\widehat{\alpha}^e_0(b^e_{j-1})-\mathcal{A}^e(b^e_{j-1})\widehat{\alpha}^e_1(b^e_{j-1})\right)}{(\widehat{\alpha}^e_0(b^e_{j-1}))^2}\right)+O\left(\frac{c(n)}{n^2}\right),$$

$$\mathcal{R}^{e,\infty}_2(n)\underset{n\to\infty}{=}\frac1n\sum_{j=1}^{N+1}\left(\frac{\left(\widehat{\alpha}^e_0(b^e_{j-1})\mathcal{A}^e(b^e_{j-1})+b^e_{j-1}\left(\mathcal{B}^e(b^e_{j-1})\widehat{\alpha}^e_0(b^e_{j-1})-\mathcal{A}^e(b^e_{j-1})\widehat{\alpha}^e_1(b^e_{j-1})\right)\right)}{(\widehat{\alpha}^e_0(b^e_{j-1}))^2}\right.$$
$$\left.+\frac{\left(\widehat{\alpha}^e_0(a^e_j)\mathcal{A}^e(a^e_j)+a^e_j\left(\mathcal{B}^e(a^e_j)\widehat{\alpha}^e_0(a^e_j)-\mathcal{A}^e(a^e_j)\widehat{\alpha}^e_1(a^e_j)\right)\right)}{(\widehat{\alpha}^e_0(a^e_j))^2}\right)+O\left(\frac{c(n)}{n^2}\right),$$

with all parameters defined in Theorem 3.1, *Equations* (3.52)–(3.100).

Theorem A.2 ([21]**).** *Let all the conditions stated in Theorem* A.1 *be valid, and let all parameters be as defined therein. Let* $\overset{e}{\mathrm{Y}}\colon\mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$ *be the unique solution of* **RHP1** *with integral representation* (1.29). *Then,*

$$\overset{e}{\mathrm{Y}}(z)z^{n\sigma_3}\underset{z\to0}{=}\mathrm{Y}^{e,0}_0+z\mathrm{Y}^{e,0}_1+z^2\mathrm{Y}^{e,0}_2+O(z^3),$$

where

$$(\mathrm{Y}^{e,0}_0)_{11}=(\widehat{\mathrm{Q}}^e_0)_{11}\mathrm{e}^{2n(\int_{J_e}\ln(|s|)\psi^e_V(s)\,\mathrm{d}s+\mathrm{i}\pi\int_{J_e\cap\mathbb{R}_+}\psi^e_V(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}^{e,0}_0)_{12}=(\widehat{\mathrm{Q}}^e_0)_{12}\mathrm{e}^{n(\ell_e-2\int_{J_e}\ln(|s|)\psi^e_V(s)\,\mathrm{d}s-\mathrm{i}2\pi\int_{J_e\cap\mathbb{R}_+}\psi^e_V(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}^{e,0}_0)_{21}=(\widehat{\mathrm{Q}}^e_0)_{21}\mathrm{e}^{-n(\ell_e-2\int_{J_e}\ln(|s|)\psi^e_V(s)\,\mathrm{d}s-\mathrm{i}2\pi\int_{J_e\cap\mathbb{R}_+}\psi^e_V(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}^{e,0}_0)_{22}=(\widehat{\mathrm{Q}}^e_0)_{22}\mathrm{e}^{-2n(\int_{J_e}\ln(|s|)\psi^e_V(s)\,\mathrm{d}s+\mathrm{i}\pi\int_{J_e\cap\mathbb{R}_+}\psi^e_V(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}^{e,0}_1)_{11}=\left((\widehat{\mathrm{Q}}^e_1)_{11}-2n(\widehat{\mathrm{Q}}^e_0)_{11}\int_{J_e}s^{-1}\psi^e_V(s)\,\mathrm{d}s\right)\mathrm{e}^{2n(\int_{J_e}\ln(|s|)\psi^e_V(s)\,\mathrm{d}s+\mathrm{i}\pi\int_{J_e\cap\mathbb{R}_+}\psi^e_V(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}^{e,0}_1)_{12}=\left((\widehat{\mathrm{Q}}^e_1)_{12}+2n(\widehat{\mathrm{Q}}^e_0)_{12}\int_{J_e}s^{-1}\psi^e_V(s)\,\mathrm{d}s\right)\mathrm{e}^{n(\ell_e-2\int_{J_e}\ln(|s|)\psi^e_V(s)\,\mathrm{d}s-\mathrm{i}2\pi\int_{J_e\cap\mathbb{R}_+}\psi^e_V(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}^{e,0}_1)_{21}=\left((\widehat{\mathrm{Q}}^e_1)_{21}-2n(\widehat{\mathrm{Q}}^e_0)_{21}\int_{J_e}s^{-1}\psi^e_V(s)\,\mathrm{d}s\right)\mathrm{e}^{-n(\ell_e-2\int_{J_e}\ln(|s|)\psi^e_V(s)\,\mathrm{d}s-\mathrm{i}2\pi\int_{J_e\cap\mathbb{R}_+}\psi^e_V(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}^{e,0}_1)_{22}=\left((\widehat{\mathrm{Q}}^e_1)_{22}+2n(\widehat{\mathrm{Q}}^e_0)_{22}\int_{J_e}s^{-1}\psi^e_V(s)\,\mathrm{d}s\right)\mathrm{e}^{-2n(\int_{J_e}\ln(|s|)\psi^e_V(s)\,\mathrm{d}s+\mathrm{i}\pi\int_{J_e\cap\mathbb{R}_+}\psi^e_V(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}^{e,0}_2)_{11}=\left((\widehat{\mathrm{Q}}^e_2)_{11}-2n(\widehat{\mathrm{Q}}^e_1)_{11}\int_{J_e}s^{-1}\psi^e_V(s)\,\mathrm{d}s+(\widehat{\mathrm{Q}}^e_0)_{11}\left(2n^2\left(\int_{J_e}s^{-1}\psi^e_V(s)\,\mathrm{d}s\right)^2\right.\right.$$
$$\left.\left.-\,n\int_{J_e}s^{-2}\psi^e_V(s)\,\mathrm{d}s\right)\right)\mathrm{e}^{2n(\int_{J_e}\ln(|s|)\psi^e_V(s)\,\mathrm{d}s+\mathrm{i}\pi\int_{J_e\cap\mathbb{R}_+}\psi^e_V(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}^{e,0}_2)_{12}=\left((\widehat{\mathrm{Q}}^e_2)_{12}+2n(\widehat{\mathrm{Q}}^e_1)_{12}\int_{J_e}s^{-1}\psi^e_V(s)\,\mathrm{d}s+(\widehat{\mathrm{Q}}^e_0)_{12}\left(2n^2\left(\int_{J_e}s^{-1}\psi^e_V(s)\,\mathrm{d}s\right)^2\right.\right.$$

$$+\ n\int_{J_e} s^{-2}\psi_V^e(s)\,\mathrm{d}s\Bigg)\Bigg)\mathrm{e}^{n(\ell_e-2\int_{J_e}\ln(|s|)\psi_V^e(s)\,\mathrm{d}s-\mathrm{i}2\pi\int_{J_e\cap\mathbb{R}_+}\psi_V^e(s)\,\mathrm{d}s)},$$

$$(\mathrm{Y}_2^{e,0})_{21}=\Bigg((\widehat{Q}_2^e)_{21}-2n(\widehat{Q}_1^e)_{21}\int_{J_e} s^{-1}\psi_V^e(s)\,\mathrm{d}s+(\widehat{Q}_0^e)_{21}\Bigg(2n^2\Bigg(\int_{J_e} s^{-1}\psi_V^e(s)\,\mathrm{d}s\Bigg)^2$$
$$-\ n\int_{J_e} s^{-2}\psi_V^e(s)\,\mathrm{d}s\Bigg)\Bigg)\mathrm{e}^{-n(\ell_e-2\int_{J_e}\ln(|s|)\psi_V^e(s)\,\mathrm{d}s-\mathrm{i}2\pi\int_{J_e\cap\mathbb{R}_+}\psi_V^e(s)\,\mathrm{d}s)},$$

$$(\mathrm{Y}_2^{e,0})_{22}=\Bigg((\widehat{Q}_2^e)_{22}+2n(\widehat{Q}_1^e)_{22}\int_{J_e} s^{-1}\psi_V^e(s)\,\mathrm{d}s+(\widehat{Q}_0^e)_{22}\Bigg(2n^2\Bigg(\int_{J_e} s^{-1}\psi_V^e(s)\,\mathrm{d}s\Bigg)^2$$
$$+\ n\int_{J_e} s^{-2}\psi_V^e(s)\,\mathrm{d}s\Bigg)\Bigg)\mathrm{e}^{-2n(\int_{J_e}\ln(|s|)\psi_V^e(s)\,\mathrm{d}s+\mathrm{i}\pi\int_{J_e\cap\mathbb{R}_+}\psi_V^e(s)\,\mathrm{d}s)},$$

with

$$(\widehat{Q}_0^e)_{11}:=(\overset{e}{m}{}_0^0)_{11}\Big(1+(\mathcal{R}_0^{e,0}(n))_{11}\Big)+(\mathcal{R}_0^{e,0}(n))_{12}(\overset{e}{m}{}_0^0)_{21},$$
$$(\widehat{Q}_0^e)_{12}:=(\overset{e}{m}{}_0^0)_{12}\Big(1+(\mathcal{R}_0^{e,0}(n))_{11}\Big)+(\mathcal{R}_0^{e,0}(n))_{12}(\overset{e}{m}{}_0^0)_{22},$$
$$(\widehat{Q}_0^e)_{21}:=(\overset{e}{m}{}_0^0)_{21}\Big(1+(\mathcal{R}_0^{e,0}(n))_{22}\Big)+(\mathcal{R}_0^{e,0}(n))_{21}(\overset{e}{m}{}_0^0)_{11},$$
$$(\widehat{Q}_0^e)_{22}:=(\overset{e}{m}{}_0^0)_{22}\Big(1+(\mathcal{R}_0^{e,0}(n))_{22}\Big)+(\mathcal{R}_0^{e,0}(n))_{21}(\overset{e}{m}{}_0^0)_{12},$$
$$(\widehat{Q}_1^e)_{11}:=(\overset{e}{m}{}_1^0)_{11}\Big(1+(\mathcal{R}_0^{e,0}(n))_{11}\Big)+(\mathcal{R}_0^{e,0}(n))_{12}(\overset{e}{m}{}_1^0)_{21}+(\mathcal{R}_1^{e,0}(n))_{11}(\overset{e}{m}{}_0^0)_{11}$$
$$+(\mathcal{R}_1^{e,0}(n))_{12}(\overset{e}{m}{}_0^0)_{21},$$
$$(\widehat{Q}_1^e)_{12}:=(\overset{e}{m}{}_1^0)_{12}\Big(1+(\mathcal{R}_0^{e,0}(n))_{11}\Big)+(\mathcal{R}_0^{e,0}(n))_{12}(\overset{e}{m}{}_1^0)_{22}+(\mathcal{R}_1^{e,0}(n))_{11}(\overset{e}{m}{}_0^0)_{12}$$
$$+(\mathcal{R}_1^{e,0}(n))_{12}(\overset{e}{m}{}_0^0)_{22},$$
$$(\widehat{Q}_1^e)_{21}:=(\overset{e}{m}{}_1^0)_{21}\Big(1+(\mathcal{R}_0^{e,0}(n))_{22}\Big)+(\mathcal{R}_0^{e,0}(n))_{21}(\overset{e}{m}{}_1^0)_{11}+(\mathcal{R}_1^{e,0}(n))_{21}(\overset{e}{m}{}_0^0)_{11}$$
$$+(\mathcal{R}_1^{e,0}(n))_{22}(\overset{e}{m}{}_0^0)_{21},$$
$$(\widehat{Q}_1^e)_{22}:=(\overset{e}{m}{}_1^0)_{22}\Big(1+(\mathcal{R}_0^{e,0}(n))_{22}\Big)+(\mathcal{R}_0^{e,0}(n))_{21}(\overset{e}{m}{}_1^0)_{12}+(\mathcal{R}_1^{e,0}(n))_{21}(\overset{e}{m}{}_0^0)_{12}$$
$$+(\mathcal{R}_1^{e,0}(n))_{22}(\overset{e}{m}{}_0^0)_{22},$$
$$(\widehat{Q}_2^e)_{11}:=(\overset{e}{m}{}_2^0)_{11}\Big(1+(\mathcal{R}_0^{e,0}(n))_{11}\Big)+(\mathcal{R}_0^{e,0}(n))_{12}(\overset{e}{m}{}_2^0)_{21}+(\mathcal{R}_1^{e,0}(n))_{11}(\overset{e}{m}{}_1^0)_{11}$$
$$+(\mathcal{R}_1^{e,0}(n))_{12}(\overset{e}{m}{}_1^0)_{21}+(\mathcal{R}_2^{e,0}(n))_{11}(\overset{e}{m}{}_0^0)_{11}+(\mathcal{R}_2^{e,0}(n))_{12}(\overset{e}{m}{}_0^0)_{21},$$
$$(\widehat{Q}_2^e)_{12}:=(\overset{e}{m}{}_2^0)_{12}\Big(1+(\mathcal{R}_0^{e,0}(n))_{11}\Big)+(\mathcal{R}_0^{e,0}(n))_{12}(\overset{e}{m}{}_2^0)_{22}+(\mathcal{R}_1^{e,0}(n))_{11}(\overset{e}{m}{}_1^0)_{12}$$
$$+(\mathcal{R}_1^{e,0}(n))_{12}(\overset{e}{m}{}_1^0)_{22}+(\mathcal{R}_2^{e,0}(n))_{11}(\overset{e}{m}{}_0^0)_{12}+(\mathcal{R}_2^{e,0}(n))_{12}(\overset{e}{m}{}_0^0)_{22},$$
$$(\widehat{Q}_2^e)_{21}:=(\overset{e}{m}{}_2^0)_{21}\Big(1+(\mathcal{R}_0^{e,0}(n))_{22}\Big)+(\mathcal{R}_0^{e,0}(n))_{21}(\overset{e}{m}{}_2^0)_{11}+(\mathcal{R}_1^{e,0}(n))_{21}(\overset{e}{m}{}_1^0)_{11}$$
$$+(\mathcal{R}_1^{e,0}(n))_{22}(\overset{e}{m}{}_1^0)_{21}+(\mathcal{R}_2^{e,0}(n))_{21}(\overset{e}{m}{}_0^0)_{11}+(\mathcal{R}_2^{e,0}(n))_{22}(\overset{e}{m}{}_0^0)_{21},$$
$$(\widehat{Q}_2^e)_{22}:=(\overset{e}{m}{}_2^0)_{22}\Big(1+(\mathcal{R}_0^{e,0}(n))_{22}\Big)+(\mathcal{R}_0^{e,0}(n))_{21}(\overset{e}{m}{}_2^0)_{12}+(\mathcal{R}_1^{e,0}(n))_{21}(\overset{e}{m}{}_1^0)_{12}$$
$$+(\mathcal{R}_1^{e,0}(n))_{22}(\overset{e}{m}{}_1^0)_{22}+(\mathcal{R}_2^{e,0}(n))_{21}(\overset{e}{m}{}_0^0)_{12}+(\mathcal{R}_2^{e,0}(n))_{22}(\overset{e}{m}{}_0^0)_{22},$$

and

$$(\overset{e}{m}{}_0^0)_{11}=\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{2}\right)\frac{\theta_0^e(1,1,\boldsymbol{\Omega}^e)}{\theta_0^e(1,1,\mathbf{0})},$$
$$(\overset{e}{m}{}_0^0)_{12}=-\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{2\mathrm{i}}\right)\frac{\theta_0^e(-1,1,\boldsymbol{\Omega}^e)}{\theta_0^e(-1,1,\mathbf{0})},$$
$$(\overset{e}{m}{}_0^0)_{21}=\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{2\mathrm{i}}\right)\frac{\theta_0^e(1,-1,\boldsymbol{\Omega}^e)}{\theta_0^e(1,-1,\mathbf{0})},$$

$$(\overset{e}{m}{}_0^0)_{22} = \frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{2}\right)\frac{\theta_0^e(-1,-1,\boldsymbol{\Omega}^e)}{\theta_0^e(-1,-1,\mathbf{0})},$$

$$(\overset{e}{m}{}_1^0)_{11} = \frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}\left(\left(\frac{\widetilde{\alpha}_0^e(1,1,\boldsymbol{\Omega}^e)\theta_0^e(1,1,\mathbf{0})-\widetilde{\alpha}_0^e(1,1,\mathbf{0})\theta_0^e(1,1,\boldsymbol{\Omega}^e)}{(\theta_0^e(1,1,\mathbf{0}))^2}\right)\right.$$
$$\left.\times\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{2}\right)+\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{8}\right)\left(\sum_{k=1}^{N+1}\left(\frac{1}{a_k^e}-\frac{1}{b_{k-1}^e}\right)\right)\frac{\theta_0^e(1,1,\boldsymbol{\Omega}^e)}{\theta_0^e(1,1,\mathbf{0})}\right),$$

$$(\overset{e}{m}{}_1^0)_{12} = -\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}\left(\left(\frac{\widetilde{\alpha}_0^e(-1,1,\boldsymbol{\Omega}^e)\theta_0^e(-1,1,\mathbf{0})-\widetilde{\alpha}_0^e(-1,1,\mathbf{0})\theta_0^e(-1,1,\boldsymbol{\Omega}^e)}{(\theta_0^e(-1,1,\mathbf{0}))^2}\right)\right.$$
$$\left.\times\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{2\mathrm{i}}\right)+\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{8\mathrm{i}}\right)\left(\sum_{k=1}^{N+1}\left(\frac{1}{a_k^e}-\frac{1}{b_{k-1}^e}\right)\right)\frac{\theta_0^e(-1,1,\boldsymbol{\Omega}^e)}{\theta_0^e(-1,1,\mathbf{0})}\right),$$

$$(\overset{e}{m}{}_1^0)_{21} = \frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}\left(\left(\frac{\widetilde{\alpha}_0^e(1,-1,\boldsymbol{\Omega}^e)\theta_0^e(1,-1,\mathbf{0})-\widetilde{\alpha}_0^e(1,-1,\mathbf{0})\theta_0^e(1,-1,\boldsymbol{\Omega}^e)}{(\theta_0^e(1,-1,\mathbf{0}))^2}\right)\right.$$
$$\left.\times\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{2\mathrm{i}}\right)+\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{8\mathrm{i}}\right)\left(\sum_{k=1}^{N+1}\left(\frac{1}{a_k^e}-\frac{1}{b_{k-1}^e}\right)\right)\frac{\theta_0^e(1,-1,\boldsymbol{\Omega}^e)}{\theta_0^e(1,-1,\mathbf{0})}\right),$$

$$(\overset{e}{m}{}_1^0)_{22} = \frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}\left(\left(\frac{\widetilde{\alpha}_0^e(-1,-1,\boldsymbol{\Omega}^e)\theta_0^e(-1,-1,\mathbf{0})-\widetilde{\alpha}_0^e(-1,-1,\mathbf{0})\theta_0^e(-1,-1,\boldsymbol{\Omega}^e)}{(\theta_0^e(-1,-1,\mathbf{0}))^2}\right)\right.$$
$$\left.\times\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{2}\right)+\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{8}\right)\left(\sum_{k=1}^{N+1}\left(\frac{1}{a_k^e}-\frac{1}{b_{k-1}^e}\right)\right)\frac{\theta_0^e(-1,-1,\boldsymbol{\Omega}^e)}{\theta_0^e(-1,-1,\mathbf{0})}\right),$$

$$(\overset{e}{m}{}_2^0)_{11} = \frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}\left(\left(\theta_0^e(1,1,\boldsymbol{\Omega}^e)\left((\widetilde{\alpha}_0^e(1,1,\mathbf{0}))^2-\beta_0^e(1,1,\mathbf{0})\theta_0^e(1,1,\mathbf{0})\right)\right.\right.$$
$$\left.-\widetilde{\alpha}_0^e(1,1,\boldsymbol{\Omega}^e)\widetilde{\alpha}_0^e(1,1,\mathbf{0})\theta_0^e(1,1,\mathbf{0})+\beta_0^e(1,1,\boldsymbol{\Omega}^e)(\theta_0^e(1,1,\mathbf{0}))^2\right)\frac{1}{(\theta_0^e(1,1,\mathbf{0}))^3}$$
$$\times\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{2}\right)+\left(\frac{\widetilde{\alpha}_0^e(1,1,\boldsymbol{\Omega}^e)\theta_0^e(1,1,\mathbf{0})-\widetilde{\alpha}_0^e(1,1,\mathbf{0})\theta_0^e(1,1,\boldsymbol{\Omega}^e)}{(\theta_0^e(1,1,\mathbf{0}))^2}\right)$$
$$\times\left(\sum_{k=1}^{N+1}\left(\frac{1}{a_k^e}-\frac{1}{b_{k-1}^e}\right)\right)\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{8}\right)+\left(\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{16}\right)\sum_{k=1}^{N+1}\left(\frac{1}{(a_k^e)^2}-\frac{1}{(b_{k-1}^e)^2}\right)\right.$$
$$\left.\left.+\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{64}\right)\left(\sum_{k=1}^{N+1}\left(\frac{1}{a_k^e}-\frac{1}{b_{k-1}^e}\right)\right)^2\right)\frac{\theta_0^e(1,1,\boldsymbol{\Omega}^e)}{\theta_0^e(1,1,\mathbf{0})}\right),$$

$$(\overset{e}{m}{}_2^0)_{12} = -\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e+\boldsymbol{d}_e)}\left(\left(\theta_0^e(-1,1,\boldsymbol{\Omega}^e)\left((\widetilde{\alpha}_0^e(-1,1,\mathbf{0}))^2-\beta_0^e(-1,1,\mathbf{0})\theta_0^e(-1,1,\mathbf{0})\right)\right.\right.$$
$$\left.-\widetilde{\alpha}_0^e(-1,1,\boldsymbol{\Omega}^e)\widetilde{\alpha}_0^e(-1,1,\mathbf{0})\theta_0^e(-1,1,\mathbf{0})+\beta_0^e(-1,1,\boldsymbol{\Omega}^e)(\theta_0^e(-1,1,\mathbf{0}))^2\right)\frac{1}{(\theta_0^e(-1,1,\mathbf{0}))^3}$$
$$\times\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{2\mathrm{i}}\right)+\left(\frac{\widetilde{\alpha}_0^e(-1,1,\boldsymbol{\Omega}^e)\theta_0^e(-1,1,\mathbf{0})-\widetilde{\alpha}_0^e(-1,1,\mathbf{0})\theta_0^e(-1,1,\boldsymbol{\Omega}^e)}{(\theta_0^e(-1,1,\mathbf{0}))^2}\right)$$
$$\times\left(\sum_{k=1}^{N+1}\left(\frac{1}{a_k^e}-\frac{1}{b_{k-1}^e}\right)\right)\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{8\mathrm{i}}\right)+\left(\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{16\mathrm{i}}\right)\sum_{k=1}^{N+1}\left(\frac{1}{(a_k^e)^2}-\frac{1}{(b_{k-1}^e)^2}\right)\right.$$
$$\left.\left.+\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{64\mathrm{i}}\right)\left(\sum_{k=1}^{N+1}\left(\frac{1}{a_k^e}-\frac{1}{b_{k-1}^e}\right)\right)^2\right)\frac{\theta_0^e(-1,1,\boldsymbol{\Omega}^e)}{\theta_0^e(-1,1,\mathbf{0})}\right),$$

$$(\overset{e}{m}{}_2^0)_{21} = \frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}\left(\left(\theta_0^e(1,-1,\boldsymbol{\Omega}^e)\left((\widetilde{\alpha}_0^e(1,-1,\mathbf{0}))^2-\beta_0^e(1,-1,\mathbf{0})\theta_0^e(1,-1,\mathbf{0})\right)\right.\right.$$
$$\left.-\widetilde{\alpha}_0^e(1,-1,\boldsymbol{\Omega}^e)\widetilde{\alpha}_0^e(1,-1,\mathbf{0})\theta_0^e(1,-1,\mathbf{0})+\beta_0^e(1,-1,\boldsymbol{\Omega}^e)(\theta_0^e(1,-1,\mathbf{0}))^2\right)\frac{1}{(\theta_0^e(1,-1,\mathbf{0}))^3}$$

$$\times\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{2\mathrm{i}}\right)+\left(\frac{\widetilde{\alpha}_0^e(1,-1,\boldsymbol{\Omega}^e)\theta_0^e(1,-1,\mathbf{0})-\widetilde{\alpha}_0^e(1,-1,\mathbf{0})\theta_0^e(1,-1,\boldsymbol{\Omega}^e)}{(\theta_0^e(1,-1,\mathbf{0}))^2}\right)$$
$$\times\left(\sum_{k=1}^{N+1}\left(\frac{1}{a_k^e}-\frac{1}{b_{k-1}^e}\right)\right)\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{8\mathrm{i}}\right)+\left(\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{16\mathrm{i}}\right)\sum_{k=1}^{N+1}\left(\frac{1}{(a_k^e)^2}-\frac{1}{(b_{k-1}^e)^2}\right)\right.$$
$$\left.+\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{64\mathrm{i}}\right)\left(\sum_{k=1}^{N+1}\left(\frac{1}{a_k^e}-\frac{1}{b_{k-1}^e}\right)\right)^2\right)\frac{\theta_0^e(1,-1,\boldsymbol{\Omega}^e)}{\theta_0^e(1,-1,\mathbf{0})}\Bigg),$$

$$(\overset{e}{m}{}_2^0)_{22}=\frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty)+\boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty)-\frac{n}{2\pi}\boldsymbol{\Omega}^e-\boldsymbol{d}_e)}\Big(\big(\theta_0^e(-1,-1,\boldsymbol{\Omega}^e)\big((\widetilde{\alpha}_0^e(-1,-1,\mathbf{0}))^2-\beta_0^e(-1,-1,\mathbf{0})\theta_0^e(-1,-1,\mathbf{0})\big)$$
$$-\widetilde{\alpha}_0^e(-1,-1,\boldsymbol{\Omega}^e)\widetilde{\alpha}_0^e(-1,-1,\mathbf{0})\theta_0^e(-1,-1,\mathbf{0})+\beta_0^e(-1,-1,\boldsymbol{\Omega}^e)(\theta_0^e(-1,-1,\mathbf{0}))^2\big)\tfrac{1}{(\theta_0^e(-1,-1,\mathbf{0}))^3}$$
$$\times\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{2}\right)+\left(\frac{\widetilde{\alpha}_0^e(-1,-1,\boldsymbol{\Omega}^e)\theta_0^e(-1,-1,\mathbf{0})-\widetilde{\alpha}_0^e(-1,-1,\mathbf{0})\theta_0^e(-1,-1,\boldsymbol{\Omega}^e)}{(\theta_0^e(-1,-1,\mathbf{0}))^2}\right)$$
$$\times\left(\sum_{k=1}^{N+1}\left(\frac{1}{a_k^e}-\frac{1}{b_{k-1}^e}\right)\right)\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{8}\right)+\left(\left(\frac{\gamma_0^e-(\gamma_0^e)^{-1}}{16}\right)\sum_{k=1}^{N+1}\left(\frac{1}{(a_k^e)^2}-\frac{1}{(b_{k-1}^e)^2}\right)\right.$$
$$\left.+\left(\frac{\gamma_0^e+(\gamma_0^e)^{-1}}{64}\right)\left(\sum_{k=1}^{N+1}\left(\frac{1}{a_k^e}-\frac{1}{b_{k-1}^e}\right)\right)^2\right)\frac{\theta_0^e(-1,-1,\boldsymbol{\Omega}^e)}{\theta_0^e(-1,-1,\mathbf{0})}\Bigg),$$

where, for $\varepsilon_1,\varepsilon_2=\pm1$,

$$\theta_0^e(\varepsilon_1,\varepsilon_2,\mathfrak{Z}):=\boldsymbol{\theta}^e(\varepsilon_1\boldsymbol{u}_+^e(0)-\tfrac{n}{2\pi}\mathfrak{Z}+\varepsilon_2\boldsymbol{d}_e),$$
$$\widetilde{\alpha}_0^e(\varepsilon_1,\varepsilon_2,\mathfrak{Z}):=2\pi\mathrm{i}\varepsilon_1\sum_{m\in\mathbb{Z}^N}(m,\widehat{\boldsymbol{\alpha}}_0^e)\mathrm{e}^{2\pi\mathrm{i}(m,\varepsilon_1\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\mathfrak{Z}+\varepsilon_2\boldsymbol{d}_e)+\pi\mathrm{i}(m,\tau^e m)},$$

with $\boldsymbol{u}_+^e(0)=\int_{a_{N+1}^e}^{0^+}\boldsymbol{\omega}^e$, $\widehat{\boldsymbol{\alpha}}_0^e=(\widehat{\alpha}_{0,1}^e,\widehat{\alpha}_{0,2}^e,\dots,\widehat{\alpha}_{0,N}^e)$, *and* $\widehat{\alpha}_{0,j}^e:=(-1)^{\mathcal{N}_+^e}(\prod_{i=1}^{N+1}|b_{i-1}^e a_i^e|)^{-1/2}c_{jN}^e$, $j=1,\dots,N$, *where* $\mathcal{N}_+^e\in\{0,\dots,N+1\}$ *is the number of bands to the right of* $z=0$,

$$\beta_0^e(\varepsilon_1,\varepsilon_2,\mathfrak{Z}):=2\pi\sum_{m\in\mathbb{Z}^N}\left(\mathrm{i}\varepsilon_1(m,\widehat{\boldsymbol{\beta}}_0^e)-\pi(m,\widehat{\boldsymbol{\alpha}}_0^e)^2\right)\mathrm{e}^{2\pi\mathrm{i}(m,\varepsilon_1\boldsymbol{u}_+^e(0)-\frac{n}{2\pi}\mathfrak{Z}+\varepsilon_2\boldsymbol{d}_e)+\pi\mathrm{i}(m,\tau^e m)},$$

where $\widehat{\boldsymbol{\beta}}_0^e=(\widehat{\beta}_{0,1}^e,\widehat{\beta}_{0,2}^e,\dots,\widehat{\beta}_{0,N}^e)$, *with* $\widehat{\beta}_{0,j}^e:=\frac{1}{2}(-1)^{\mathcal{N}_+^e}(\prod_{i=1}^{N+1}|b_{i-1}^e a_i^e|)^{-1/2}(c_{jN-1}^e+\frac{1}{2}c_{jN}^e\sum_{k=1}^{N+1}((a_k^e)^{-1}+(b_{k-1}^e)^{-1}))$, $j=1,\dots,N$, *where* c_{jN}^e, c_{jN-1}^e, $j=1,\dots,N$, *are obtained from Equations* (E1) *and* (E2), *and* γ_0^e *is defined in Theorem* 3.1, *Equations* (3.53), *and, for* $k=1,2,3$,

$$\mathcal{R}_{k-1}^{e,0}(n)\underset{n\to\infty}{=}\frac{1}{n}\sum_{j=1}^{N+1}\Bigg(\frac{\left(\mathcal{A}^e(b_{j-1}^e)\left(\widehat{\alpha}_1^e(b_{j-1}^e)+k(b_{j-1}^e)^{-1}\widehat{\alpha}_0^e(b_{j-1}^e)\right)-\mathcal{B}^e(b_{j-1}^e)\widehat{\alpha}_0^e(b_{j-1}^e)\right)}{(b_{j-1}^e)^k(\widehat{\alpha}_0^e(b_{j-1}^e))^2}$$
$$+\frac{\left(\mathcal{A}^e(a_j^e)\left(\widehat{\alpha}_1^e(a_j^e)+k(a_j^e)^{-1}\widehat{\alpha}_0^e(a_j^e)\right)-\mathcal{B}^e(a_j^e)\widehat{\alpha}_0^e(a_j^e)\right)}{(a_j^e)^k(\widehat{\alpha}_0^e(a_j^e))^2}\Bigg)+O\left(\frac{c(n)}{n^2}\right),$$

with all parameters defined in Theorem A.1.

Theorem A.3 ([21]**).** *Let all the conditions stated in Theorem* A.1 *be valid, and let all parameters be as defined therein. Let* $\overset{e}{Y}:\mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$ *be the unique solution of* **RHP1** *with integral representation* (1.29). *Let* $\boldsymbol{\pi}_{2n}(z)$ *be the even degree monic orthogonal L-polynomial defined in Equations* (1.4) *with* $n\to\infty$ *asymptotics (in the entire complex plane) given by Theorem* 2.3.1 *of* [21]. *Then,*

$$\left(\xi_n^{(2n)}\right)^2=\frac{1}{\|\boldsymbol{\pi}_{2n}(\cdot)\|_{\mathcal{L}}^2}=\frac{H_{2n}^{(-2n)}}{H_{2n+1}^{(-2n)}}\underset{n\to\infty}{=}\frac{\mathrm{e}^{-n\ell_e}}{\pi}\Xi^{\flat}\left(1+\frac{2}{n}\Xi^{\flat}(\mathfrak{Q}^{\flat})_{12}+O\left(\frac{c^{\flat}(n)}{n^2}\right)\right),\tag{A.1}$$

where

$$\Xi^{\flat} := 2\left(\sum_{k=1}^{N+1}\left(a_k^e - b_{k-1}^e\right)\right)^{-1} \frac{\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty) - \frac{n}{2\pi}\boldsymbol{\Omega}^e + \boldsymbol{d}_e)\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty) + \boldsymbol{d}_e)}{\boldsymbol{\theta}^e(-\boldsymbol{u}_+^e(\infty) - \frac{n}{2\pi}\boldsymbol{\Omega}^e + \boldsymbol{d}_e)\boldsymbol{\theta}^e(\boldsymbol{u}_+^e(\infty) + \boldsymbol{d}_e)} \quad (>0),$$

$\mathfrak{Q}^{\flat}$ *is defined in Theorem* 3.2, *Equations* (3.107), $(\mathfrak{Q}^{\flat})_{12}$ *denotes the* (1 2)*-element of* $\mathfrak{Q}^{\flat}$, $c^{\flat}(n) =_{n\to\infty} O(1)$, *and all the relevant parameters are defined in Theorem* A.1. *Asymptotics for* $\xi_n^{(2n)}$ *are obtained by taking the positive square root of both sides of Equation* (A.1).

Appendix B. Asymptotics of $\overset{o}{\mathrm{Y}}(z)$

In this appendix, large-n asymptotics for the coefficients of the expansions (3.3) and (3.4) for the solution of **RHP2**, with integral representation (1.30), are presented (for complete details and proofs, see [22]).

Remark B.1. In [22], the following formula was established for complex z away from an open neighbourhood around $\overline{J_o} := \cup_{j=1}^{N+1}[b_{j-1}^o, a_j^o]$:

$$\overset{o}{\mathrm{Y}}(z) = \begin{cases} \mathrm{e}^{\frac{n\ell_o}{2}\sigma_3}\mathcal{R}^o(z)\overset{o}{m}^{\infty}(z)\mathbb{E}^{\sigma_3}\mathrm{e}^{n(g^o(z)-\frac{\ell_o}{2}-\mathfrak{Q}_{\mathcal{A}}^+)\sigma_3}, & z\in\mathbb{C}_+, \\ \mathrm{e}^{\frac{n\ell_o}{2}\sigma_3}\mathcal{R}^o(z)\overset{o}{m}^{\infty}(z)\mathbb{E}^{-\sigma_3}\mathrm{e}^{n(g^o(z)-\frac{\ell_o}{2}-\mathfrak{Q}_{\mathcal{A}}^-)\sigma_3}, & z\in\mathbb{C}_-, \end{cases}$$

where ℓ_o, $g^o(z)$, $\mathfrak{Q}_{\mathcal{A}}^{\pm}$, and $\mathbb{E}$ have been defined heretofore, the matrix-valued function $\overset{o}{m}^{\infty}(z)$ is determined explicitly, and the matrix-valued function $\mathcal{R}^o(z)$ solves a "small norm" RHP for which a complete asymptotic expansion as $n\to\infty$ for $z\in\mathbb{C}$ is computed. The above explicit formula for the solution of **RHP2** yields an explicit asymptotic description for the odd degree OLPs in question (for complete details and proofs, see [22]). ■

Remark B.2. In Theorems B.1 and B.2 below, only $n\to\infty$ asymptotics for $\mathcal{R}_{k-1}^{o,0}(n)$, $k=2,3$, and $\mathcal{R}_j^{o,\infty}(n)$, $j=0,1,2$, are given, as they are all that are actually necessary in order to obtain the results of Theorems 3.1–3.3: finite-n formulae for $\mathcal{R}_{k-1}^{o,0}(n)$, $k=2,3$, and $\mathcal{R}_j^{o,\infty}(n)$, $j=0,1,2$, are given in [22]; furthermore, the 'symbol' $c(n)$ appearing in the asymptotic (as $n\to\infty$) expressions for $\mathcal{R}_{k-1}^{o,0}(n)$, $k=2,3$, and $\mathcal{R}_j^{o,\infty}(n)$, $j=0,1,2$, is to be understood as a uniformly bounded ($O(1)$), $\mathrm{M}_2(\mathbb{C})$-valued, n-dependent function, that is, $\mathrm{M}_2(\mathbb{C})\ni c(n)=_{n\to\infty}O(1)$. ■

Theorem B.1 ([22]**).** *Let the external field $\widetilde{V}\colon \mathbb{R}\setminus\{0\}\to\mathbb{R}$ satisfy conditions* (1.20)–(1.22). *Let the orthonormal L-polynomials (resp., monic orthogonal L-polynomials) $\{\phi_k(z)\}_{k\in\mathbb{Z}_0^+}$ (resp., $\{\boldsymbol{\pi}_k(z)\}_{k\in\mathbb{Z}_0^+}$) be as defined in Equations* (1.2) *and* (1.3) *(resp., Equations* (1.4) *and* (1.5)). *Let $\overset{o}{\mathrm{Y}}\colon \mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$ be the unique solution of* **RHP2** *with integral representation* (1.30). *Define the density of the 'odd' equilibrium measure, $\mathrm{d}\mu_V^o(x)$, as in Equation* (2.19), *and set $J_o := \mathrm{supp}(\mu_V^o) = \cup_{j=1}^{N+1}(b_{j-1}^o, a_j^o)$, where $\{b_{j-1}^o, a_j^o\}_{j=1}^{N+1}$ satisfy the (real) n-dependent and locally solvable system of $2(N+1)$ moment conditions* (2.18).

Suppose, furthermore, that $\widetilde{V}\colon \mathbb{R}\setminus\{0\}\to\mathbb{R}$ is regular, namely: (i) $h_V^o(z)\not\equiv 0$ *on* $\overline{J_o} := \cup_{j=1}^{N+1}[b_{j-1}^o, a_j^o]$; (ii)

$$2\left(2+\frac{1}{n}\right)\int_{J_o}\ln(|x-s|)\,\mathrm{d}\mu_V^o(s) - 2\ln|x| - \widetilde{V}(x) - \ell_o - 2\left(2+\frac{1}{n}\right)Q_o = 0, \quad x\in\overline{J_o},$$

which defines the 'odd' variational constant, $l_o\,(\in\mathbb{R})$ (the same on each compact interval $[b_{j-1}^o, a_j^o]$, $j=1,\dots,N+1$), where $Q_o := \int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s$, and

$$2\left(2+\frac{1}{n}\right)\int_{J_o}\ln(|x-s|)\,\mathrm{d}\mu_V^o(s) - 2\ln|x| - \widetilde{V}(x) - \ell_o - 2\left(2+\frac{1}{n}\right)Q_o < 0, \quad x\in\mathbb{R}\setminus\overline{J_o};$$

(iii)

$$g_+^o(z) + g_-^o(z) - \widetilde{V}(z) - \ell_o - \mathfrak{Q}_{\mathcal{A}}^+ - \mathfrak{Q}_{\mathcal{A}}^- < 0, \quad z\in\mathbb{R}\setminus\overline{J_o},$$

where $g^o(z)$ is defined by Equation (2.21), *$g_{\pm}^o(z) := \lim_{\varepsilon\downarrow 0} g^o(z\pm\mathrm{i}\varepsilon)$, and $\mathfrak{Q}_{\mathcal{A}}^{\pm}$ are defined in Equations* (2.22); *and* (iv)

$$\mathrm{i}(g_+^o(z) - g_-^o(z) - \mathfrak{Q}_{\mathcal{A}}^+ + \mathfrak{Q}_{\mathcal{A}}^-)' > 0, \quad z\in J_o.$$

Then,

$$\overset{o}{\mathrm{Y}}(z)z^{n\sigma_3} \underset{z\to 0}{=} \mathrm{I} + z\mathrm{Y}_1^{o,0} + z^2\mathrm{Y}_2^{o,0} + O(z^3),$$

where

$$(\mathrm{Y}_1^{o,0})_{11} = -(2n+1)\int_{J_o} s^{-1}\psi_V^o(s)\,\mathrm{d}s + (\overset{o}{m}_1^0)_{11}\mathbb{E} + (\mathcal{R}_1^{o,0}(n))_{11},$$

$$
\begin{aligned}
(\Upsilon_1^{o,0})_{12} &= \mathrm{e}^{n\ell_o}\Big((\overset{o}{m}{}_1^0)_{12}\mathbb{E}^{-1}+(\mathcal{R}_1^{o,0}(n))_{12}\Big),\\
(\Upsilon_1^{o,0})_{21} &= \mathrm{e}^{-n\ell_o}\Big((\overset{o}{m}{}_1^0)_{21}\mathbb{E}+(\mathcal{R}_1^{o,0}(n))_{21}\Big),\\
(\Upsilon_1^{o,0})_{22} &= (2n+1)\int_{J_o} s^{-1}\psi_V^o(s)\,\mathrm{d}s+(\overset{o}{m}{}_1^0)_{22}\mathbb{E}^{-1}+(\mathcal{R}_1^{o,0}(n))_{22},\\
(\Upsilon_2^{o,0})_{11} &= \tfrac{1}{2}(2n+1)^2\left(\int_{J_o} s^{-1}\psi_V^o(s)\,\mathrm{d}s\right)^2-\tfrac{1}{2}(2n+1)\int_{J_o} s^{-2}\psi_V^o(s)\,\mathrm{d}s-(2n+1)\Big((\overset{o}{m}{}_1^0)_{11}\mathbb{E}+(\mathcal{R}_1^{o,0}(n))_{11}\Big)\\
&\quad\times\int_{J_o} s^{-1}\psi_V^o(s)\,\mathrm{d}s+(\overset{o}{m}{}_2^0)_{11}\mathbb{E}+(\mathcal{R}_2^{o,0}(n))_{11}+\Big((\mathcal{R}_1^{o,0}(n))_{11}(\overset{o}{m}{}_1^0)_{11}+(\mathcal{R}_1^{o,0}(n))_{12}(\overset{o}{m}{}_1^0)_{21}\Big)\mathbb{E},\\
(\Upsilon_2^{o,0})_{12} &= \mathrm{e}^{n\ell_o}\bigg((2n+1)\Big((\overset{o}{m}{}_1^0)_{12}\mathbb{E}^{-1}+(\mathcal{R}_1^{o,0}(n))_{12}\Big)\int_{J_o} s^{-1}\psi_V^o(s)\,\mathrm{d}s+(\overset{o}{m}{}_2^0)_{12}\mathbb{E}^{-1}+(\mathcal{R}_2^{o,0}(n))_{12}\\
&\quad+\Big((\mathcal{R}_1^{o,0}(n))_{11}(\overset{o}{m}{}_1^0)_{12}+(\mathcal{R}_1^{o,0}(n))_{12}(\overset{o}{m}{}_1^0)_{22}\Big)\mathbb{E}^{-1}\bigg),\\
(\Upsilon_2^{o,0})_{21} &= \mathrm{e}^{-n\ell_o}\bigg(-(2n+1)\Big((\overset{o}{m}{}_1^0)_{21}\mathbb{E}+(\mathcal{R}_1^{o,0}(n))_{21}\Big)\int_{J_o} s^{-1}\psi_V^o(s)\,\mathrm{d}s+(\overset{o}{m}{}_2^0)_{21}\mathbb{E}+(\mathcal{R}_2^{o,0}(n))_{21}\\
&\quad+\Big((\mathcal{R}_1^{o,0}(n))_{21}(\overset{o}{m}{}_1^0)_{11}+(\mathcal{R}_1^{o,0}(n))_{22}(\overset{o}{m}{}_1^0)_{21}\Big)\mathbb{E}\bigg),\\
(\Upsilon_2^{o,0})_{22} &= \tfrac{1}{2}(2n+1)^2\left(\int_{J_o} s^{-1}\psi_V^o(s)\,\mathrm{d}s\right)^2+\tfrac{1}{2}(2n+1)\int_{J_o} s^{-2}\psi_V^o(s)\,\mathrm{d}s+(2n+1)\Big((\overset{o}{m}{}_1^0)_{22}\mathbb{E}^{-1}+(\mathcal{R}_1^{o,0}(n))_{22}\Big)\\
&\quad\times\int_{J_o} s^{-1}\psi_V^o(s)\,\mathrm{d}s+(\overset{o}{m}{}_2^0)_{22}\mathbb{E}^{-1}+(\mathcal{R}_2^{o,0}(n))_{22}+\Big((\mathcal{R}_1^{o,0}(n))_{21}(\overset{o}{m}{}_1^0)_{12}+(\mathcal{R}_1^{o,0}(n))_{22}(\overset{o}{m}{}_1^0)_{22}\Big)\mathbb{E}^{-1},
\end{aligned}
$$

with

$$
\begin{aligned}
(\overset{o}{m}{}_1^0)_{11} &= -\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}^{-1}}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}\left(\frac{\theta_0^o(1,1,\boldsymbol{\Omega}^o)\alpha_0^o(1,1,\mathbf{0})-\alpha_0^o(1,1,\boldsymbol{\Omega}^o)\theta_0^o(1,1,\mathbf{0})}{(\theta_0^o(1,1,\mathbf{0}))^2}\right),\\
(\overset{o}{m}{}_1^0)_{12} &= \frac{1}{4\mathrm{i}}\left(\sum_{k=1}^{N+1}\left(\frac{1}{b_{k-1}^o}-\frac{1}{a_k^o}\right)\right)\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\theta_0^o(-1,1,\boldsymbol{\Omega}^o)\mathbb{E}^{-1}}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\theta_0^o(-1,1,\mathbf{0})},\\
(\overset{o}{m}{}_1^0)_{21} &= -\frac{1}{4\mathrm{i}}\left(\sum_{k=1}^{N+1}\left(\frac{1}{b_{k-1}^o}-\frac{1}{a_k^o}\right)\right)\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\theta_0^o(1,-1,\boldsymbol{\Omega}^o)\mathbb{E}}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)\theta_0^o(1,-1,\mathbf{0})},\\
(\overset{o}{m}{}_1^0)_{22} &= -\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}\left(\frac{\theta_0^o(-1,-1,\boldsymbol{\Omega}^o)\alpha_0^o(-1,-1,\mathbf{0})-\alpha_0^o(-1,-1,\boldsymbol{\Omega}^o)\theta_0^o(-1,-1,\mathbf{0})}{(\theta_0^o(-1,-1,\mathbf{0}))^2}\right),\\
(\overset{o}{m}{}_2^0)_{11} &= \Big(\theta_0^o(1,1,\boldsymbol{\Omega}^o)\Big(-\beta_0^0(1,1,\mathbf{0})\theta_0^o(1,1,\mathbf{0})+(\alpha_0^o(1,1,\mathbf{0}))^2\Big)-\alpha_0^o(1,1,\boldsymbol{\Omega}^o)\alpha_0^o(1,1,\mathbf{0})\theta_0^o(1,1,\mathbf{0})\\
&\quad+\beta_0^o(1,1,\boldsymbol{\Omega}^o)(\theta_0^o(1,1,\mathbf{0}))^2\Big)\frac{(\theta_0^o(1,1,\mathbf{0}))^{-3}\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}^{-1}}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}+\frac{1}{32}\left(\sum_{k=1}^{N+1}\left(\frac{1}{b_{k-1}^o}-\frac{1}{a_k^o}\right)\right)^2\mathbb{E}^{-1},\\
(\overset{o}{m}{}_2^0)_{12} &= -\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}^{-1}}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}\Bigg(\left(\frac{\theta_0^o(-1,1,\boldsymbol{\Omega}^o)\alpha_0^o(-1,1,\mathbf{0})-\alpha_0^o(-1,1,\boldsymbol{\Omega}^o)\theta_0^o(-1,1,\mathbf{0})}{(\theta_0^o(-1,1,\mathbf{0}))^2}\right)\\
&\quad\times\frac{1}{4\mathrm{i}}\left(\sum_{k=1}^{N+1}\left(\frac{1}{b_{k-1}^o}-\frac{1}{a_k^o}\right)\right)-\frac{1}{8\mathrm{i}}\left(\sum_{k=1}^{N+1}\left(\frac{1}{(b_{k-1}^o)^2}-\frac{1}{(a_k^o)^2}\right)\right)\frac{\theta_0^o(-1,1,\boldsymbol{\Omega}^o)}{\theta_0^o(-1,1,\mathbf{0})}\Bigg),\\
(\overset{o}{m}{}_2^0)_{21} &= \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}\Bigg(\left(\frac{\theta_0^o(1,-1,\boldsymbol{\Omega}^o)\alpha_0^o(1,-1,\mathbf{0})-\alpha_0^o(1,-1,\boldsymbol{\Omega}^o)\theta_0^o(1,-1,\mathbf{0})}{(\theta_0^o(1,-1,\mathbf{0}))^2}\right)\\
&\quad\times\frac{1}{4\mathrm{i}}\left(\sum_{k=1}^{N+1}\left(\frac{1}{b_{k-1}^o}-\frac{1}{a_k^o}\right)\right)-\frac{1}{8\mathrm{i}}\left(\sum_{k=1}^{N+1}\left(\frac{1}{(b_{k-1}^o)^2}-\frac{1}{(a_k^o)^2}\right)\right)\frac{\theta_0^o(1,-1,\boldsymbol{\Omega}^o)}{\theta_0^o(1,-1,\mathbf{0})}\Bigg),\\
(\overset{o}{m}{}_2^0)_{22} &= \Big(\theta_0^o(-1,-1,\boldsymbol{\Omega}^o)\Big(-\beta_0^o(-1,-1,\mathbf{0})\theta_0^o(-1,-1,\mathbf{0})+(\alpha_0^o(-1,-1,\mathbf{0}))^2\Big)-\alpha_0^o(-1,-1,\boldsymbol{\Omega}^o)\\
&\quad\times\alpha_0^o(-1,-1,\mathbf{0})\theta_0^o(-1,-1,\mathbf{0})+\beta_0^o(-1,-1,\boldsymbol{\Omega}^o)(\theta_0^o(-1,-1,\mathbf{0}))^2\Big)(\theta_0^o(-1,-1,\mathbf{0}))^{-3}
\end{aligned}
$$

$$\times \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}+\frac{1}{32}\left(\sum_{k=1}^{N+1}\left(\frac{1}{b_{k-1}^o}-\frac{1}{a_k^o}\right)\right)^2\mathbb{E},$$

and $(\star)_{ij}$, $i,j=1,2$, *denoting the* $(i\,j)$*-element of* $\star$, *where, for* $\varepsilon_1,\varepsilon_2=\pm1$,

$$\theta_0^o(\varepsilon_1,\varepsilon_2,\mathfrak{Z}):=\boldsymbol{\theta}^o(\varepsilon_1\boldsymbol{u}_+^o(0)-\tfrac{1}{2\pi}(n+\tfrac{1}{2})\mathfrak{Z}+\varepsilon_2\boldsymbol{d}_o),$$
$$\alpha_0^o(\varepsilon_1,\varepsilon_2,\mathfrak{Z}):=2\pi\mathrm{i}\varepsilon_1\sum_{m\in\mathbb{Z}^N}(m,\widehat{\boldsymbol{\alpha}}_0^o)\mathrm{e}^{2\pi\mathrm{i}(m,\varepsilon_1\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\mathfrak{Z}+\varepsilon_2\boldsymbol{d}_o)+\pi\mathrm{i}(m,\tau^o m)},$$

where $\boldsymbol{u}_+^o(0)=\int_{a_{N+1}^o}^{0^+}\boldsymbol{\omega}^o$, $\widehat{\boldsymbol{\alpha}}_0^o=(\widehat{\alpha}_{0,1}^o,\widehat{\alpha}_{0,2}^o,\dots,\widehat{\alpha}_{0,N}^o)$, *with* $\widehat{\alpha}_{0,j}^o:=(-1)^{\mathcal{N}_+^o}(\prod_{i=1}^{N+1}|b_{i-1}^o a_i^o|)^{-1/2}c_{jN}^o$, $j=1,\dots,N$, *where* $\mathcal{N}_+^o\in\{0,\dots,N+1\}$ *is the number of bands to the right of* $z=0$, *and*

$$\beta_0^o(\varepsilon_1,\varepsilon_2,\mathfrak{Z}):=2\pi\sum_{m\in\mathbb{Z}^N}\left(\mathrm{i}\varepsilon_1(m,\widehat{\boldsymbol{\beta}}_0^o)-\pi(m,\widehat{\boldsymbol{\alpha}}_0^o)^2\right)\mathrm{e}^{2\pi\mathrm{i}(m,\varepsilon_1\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\mathfrak{Z}+\varepsilon_2\boldsymbol{d}_o)+\pi\mathrm{i}(m,\tau^o m)},$$

where $\widehat{\boldsymbol{\beta}}_0^o=(\widehat{\beta}_{0,1}^o,\widehat{\beta}_{0,2}^o,\dots,\widehat{\beta}_{0,N}^o)$, *with* $\widehat{\beta}_{0,j}^o:=\frac{1}{2}(-1)^{\mathcal{N}_+^o}(\prod_{i=1}^{N+1}|b_{i-1}^o a_i^o|)^{-1/2}(c_{jN-1}^o+\frac{1}{2}c_{jN}^o\sum_{k=1}^{N+1}((a_k^o)^{-1}+(b_{k-1}^o)^{-1}))$, $j=1,\dots,N$, *where* c_{jN}^o, c_{jN-1}^o, $j=1,\dots,N$, *are obtained from Equations* (O1) *and* (O2), *and, for* $k=2,3$,

$$\mathcal{R}_{k-1}^{o,0}(n)\underset{n\to\infty}{=}\frac{1}{(n+\frac{1}{2})}\sum_{j=1}^{N+1}\left(\frac{\left(\mathcal{A}^o(b_{j-1}^o)\left(\widehat{\alpha}_1^o(b_{j-1}^o)+k(b_{j-1}^o)^{-1}\widehat{\alpha}_0^o(b_{j-1}^o)\right)-\mathcal{B}^o(b_{j-1}^o)\widehat{\alpha}_0^o(b_{j-1}^o)\right)}{(b_{j-1}^o)^k(\widehat{\alpha}_0^o(b_{j-1}^o))^2}\right.$$
$$\left.+\frac{\left(\mathcal{A}^o(a_j^o)\left(\widehat{\alpha}_1^o(a_j^o)+k(a_j^o)^{-1}\widehat{\alpha}_0^o(a_j^o)\right)-\mathcal{B}^o(a_j^o)\widehat{\alpha}_0^o(a_j^o)\right)}{(a_j^o)^k(\widehat{\alpha}_0^o(a_j^o))^2}\right)+O\left(\frac{c(n)}{(n+\frac{1}{2})^2}\right),$$

with all parameters defined in Theorem 3.1, *Equations* (3.52)–(3.100).

Theorem B.2 ([22]**).** *Let all the conditions stated in Theorem* B.1 *be valid, and let all parameters be as defined therein. Let* $\overset{o}{\mathrm{Y}}:\mathbb{C}\setminus\mathbb{R}\to\mathrm{SL}_2(\mathbb{C})$ *be the unique solution of* **RHP2** *with integral representation* (1.30). *Then,*

$$\overset{o}{\mathrm{Y}}(z)z^{-(n+1)\sigma_3}\underset{z\to\infty}{=}\mathrm{Y}_0^{o,\infty}+\frac{1}{z}\mathrm{Y}_1^{o,\infty}+\frac{1}{z^2}\mathrm{Y}_2^{o,\infty}+O\left(\frac{1}{z^3}\right),$$

where

$$(\mathrm{Y}_0^{o,\infty})_{11}=(\widehat{Q}_0^o)_{11}\mathrm{e}^{-2(n+\frac{1}{2})\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s},$$
$$(\mathrm{Y}_0^{o,\infty})_{12}=(\widehat{Q}_0^o)_{12}\mathrm{e}^{n(\ell_o+2(1+\frac{1}{2n})\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}_0^{o,\infty})_{21}=(\widehat{Q}_0^o)_{21}\mathrm{e}^{-n(\ell_o+2(1+\frac{1}{2n})\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}_0^{o,\infty})_{22}=(\widehat{Q}_0^o)_{22}\mathrm{e}^{2(n+\frac{1}{2})\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s},$$
$$(\mathrm{Y}_1^{o,\infty})_{11}=\left((\widehat{Q}_1^o)_{11}-(2n+1)(\widehat{Q}_0^o)_{11}\int_{J_o}s\psi_V^o(s)\,\mathrm{d}s\right)\mathrm{e}^{-2(n+\frac{1}{2})\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s},$$
$$(\mathrm{Y}_1^{o,\infty})_{12}=\left((\widehat{Q}_1^o)_{12}+(2n+1)(\widehat{Q}_0^o)_{12}\int_{J_o}s\psi_V^o(s)\,\mathrm{d}s\right)\mathrm{e}^{n(\ell_o+2(1+\frac{1}{2n})\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}_1^{o,\infty})_{21}=\left((\widehat{Q}_1^o)_{21}-(2n+1)(\widehat{Q}_0^o)_{21}\int_{J_o}s\psi_V^o(s)\,\mathrm{d}s\right)\mathrm{e}^{-n(\ell_o+2(1+\frac{1}{2n})\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s)},$$
$$(\mathrm{Y}_1^{o,\infty})_{22}=\left((\widehat{Q}_1^o)_{22}+(2n+1)(\widehat{Q}_0^o)_{22}\int_{J_o}s\psi_V^o(s)\,\mathrm{d}s\right)\mathrm{e}^{2(n+\frac{1}{2})\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s},$$
$$(\mathrm{Y}_2^{o,\infty})_{11}=\left((\widehat{Q}_2^o)_{11}-(2n+1)(\widehat{Q}_1^o)_{11}\int_{J_o}s\psi_V^o(s)\,\mathrm{d}s+(\widehat{Q}_0^o)_{11}\left(\tfrac{1}{2}(2n+1)^2\left(\int_{J_o}s\psi_V^o(s)\,\mathrm{d}s\right)^2\right.\right.$$
$$\left.\left.-\tfrac{1}{2}(2n+1)\int_{J_o}s^2\psi_V^o(s)\,\mathrm{d}s\right)\right)\mathrm{e}^{-2(n+\frac{1}{2})\int_{J_o}\ln(|s|)\psi_V^o(s)\,\mathrm{d}s},$$

$$(\Upsilon_2^{o,\infty})_{12} = \bigg((\widehat{Q}_2^o)_{12} + (2n+1)(\widehat{Q}_1^o)_{12} \int_{J_o} s\psi_V^o(s)\,\mathrm{d}s + (\widehat{Q}_0^o)_{12} \bigg(\tfrac{1}{2}(2n+1)^2 \bigg(\int_{J_o} s\psi_V^o(s)\,\mathrm{d}s \bigg)^2 + \tfrac{1}{2}(2n+1) \int_{J_o} s^2 \psi_V^o(s)\,\mathrm{d}s \bigg) \bigg) \mathrm{e}^{n(\ell_o + 2(1+\frac{1}{2n}) \int_{J_o} \ln(|s|)\psi_V^o(s)\,\mathrm{d}s)},$$

$$(\Upsilon_2^{o,\infty})_{21} = \bigg((\widehat{Q}_2^o)_{21} - (2n+1)(\widehat{Q}_1^o)_{21} \int_{J_o} s\psi_V^o(s)\,\mathrm{d}s + (\widehat{Q}_0^o)_{21} \bigg(\tfrac{1}{2}(2n+1)^2 \bigg(\int_{J_o} s\psi_V^o(s)\,\mathrm{d}s \bigg)^2 - \tfrac{1}{2}(2n+1) \int_{J_o} s^2 \psi_V^o(s)\,\mathrm{d}s \bigg) \bigg) \mathrm{e}^{-n(\ell_o + 2(1+\frac{1}{2n}) \int_{J_o} \ln(|s|)\psi_V^o(s)\,\mathrm{d}s)},$$

$$(\Upsilon_2^{o,\infty})_{22} = \bigg((\widehat{Q}_2^o)_{22} + (2n+1)(\widehat{Q}_1^o)_{22} \int_{J_o} s\psi_V^o(s)\,\mathrm{d}s + (\widehat{Q}_0^o)_{22} \bigg(\tfrac{1}{2}(2n+1)^2 \bigg(\int_{J_o} s\psi_V^o(s)\,\mathrm{d}s \bigg)^2 + \tfrac{1}{2}(2n+1) \int_{J_o} s^2 \psi_V^o(s)\,\mathrm{d}s \bigg) \bigg) \mathrm{e}^{2(n+\frac{1}{2}) \int_{J_o} \ln(|s|)\psi_V^o(s)\,\mathrm{d}s},$$

with

$$\begin{aligned}
(\widehat{Q}_0^o)_{11} &:= (\overset{o}{m}{}_0^{\infty})_{11}\big(1 + (\mathcal{R}_0^{o,\infty}(n))_{11}\big) + (\mathcal{R}_0^{o,\infty}(n))_{12}(\overset{o}{m}{}_0^{\infty})_{21}, \\
(\widehat{Q}_0^o)_{12} &:= (\overset{o}{m}{}_0^{\infty})_{12}\big(1 + (\mathcal{R}_0^{o,\infty}(n))_{11}\big) + (\mathcal{R}_0^{o,\infty}(n))_{12}(\overset{o}{m}{}_0^{\infty})_{22}, \\
(\widehat{Q}_0^o)_{21} &:= (\overset{o}{m}{}_0^{\infty})_{21}\big(1 + (\mathcal{R}_0^{o,\infty}(n))_{22}\big) + (\mathcal{R}_0^{o,\infty}(n))_{21}(\overset{o}{m}{}_0^{\infty})_{11}, \\
(\widehat{Q}_0^o)_{22} &:= (\overset{o}{m}{}_0^{\infty})_{22}\big(1 + (\mathcal{R}_0^{o,\infty}(n))_{22}\big) + (\mathcal{R}_0^{o,\infty}(n))_{21}(\overset{o}{m}{}_0^{\infty})_{12}, \\
(\widehat{Q}_1^o)_{11} &:= (\overset{o}{m}{}_1^{\infty})_{11}\big(1 + (\mathcal{R}_0^{o,\infty}(n))_{11}\big) + (\mathcal{R}_0^{o,\infty}(n))_{12}(\overset{o}{m}{}_1^{\infty})_{21} + (\mathcal{R}_1^{o,\infty}(n))_{11}(\overset{o}{m}{}_0^{\infty})_{11} \\
&\quad + (\mathcal{R}_1^{o,\infty}(n))_{12}(\overset{o}{m}{}_0^{\infty})_{21}, \\
(\widehat{Q}_1^o)_{12} &:= (\overset{o}{m}{}_1^{\infty})_{12}\big(1 + (\mathcal{R}_0^{o,\infty}(n))_{11}\big) + (\mathcal{R}_0^{o,\infty}(n))_{12}(\overset{o}{m}{}_1^{\infty})_{22} + (\mathcal{R}_1^{o,\infty}(n))_{11}(\overset{o}{m}{}_0^{\infty})_{12} \\
&\quad + (\mathcal{R}_1^{o,\infty}(n))_{12}(\overset{o}{m}{}_0^{\infty})_{22}, \\
(\widehat{Q}_1^o)_{21} &:= (\overset{o}{m}{}_1^{\infty})_{21}\big(1 + (\mathcal{R}_0^{o,\infty}(n))_{22}\big) + (\mathcal{R}_0^{o,\infty}(n))_{21}(\overset{o}{m}{}_1^{\infty})_{11} + (\mathcal{R}_1^{o,\infty}(n))_{21}(\overset{o}{m}{}_0^{\infty})_{11} \\
&\quad + (\mathcal{R}_1^{o,\infty}(n))_{22}(\overset{o}{m}{}_0^{\infty})_{21}, \\
(\widehat{Q}_1^o)_{22} &:= (\overset{o}{m}{}_1^{\infty})_{22}\big(1 + (\mathcal{R}_0^{o,\infty}(n))_{22}\big) + (\mathcal{R}_0^{o,\infty}(n))_{21}(\overset{o}{m}{}_1^{\infty})_{12} + (\mathcal{R}_1^{o,\infty}(n))_{21}(\overset{o}{m}{}_0^{\infty})_{12} \\
&\quad + (\mathcal{R}_1^{o,\infty}(n))_{22}(\overset{o}{m}{}_0^{\infty})_{22}, \\
(\widehat{Q}_2^o)_{11} &:= (\overset{o}{m}{}_2^{\infty})_{11}\big(1 + (\mathcal{R}_0^{o,\infty}(n))_{11}\big) + (\mathcal{R}_0^{o,\infty}(n))_{12}(\overset{o}{m}{}_2^{\infty})_{21} + (\mathcal{R}_1^{o,\infty}(n))_{11}(\overset{o}{m}{}_1^{\infty})_{11} \\
&\quad + (\mathcal{R}_1^{o,\infty}(n))_{12}(\overset{o}{m}{}_1^{\infty})_{21} + (\mathcal{R}_2^{o,\infty}(n))_{11}(\overset{o}{m}{}_0^{\infty})_{11} + (\mathcal{R}_2^{o,\infty}(n))_{12}(\overset{o}{m}{}_0^{\infty})_{21}, \\
(\widehat{Q}_2^o)_{12} &:= (\overset{o}{m}{}_2^{\infty})_{12}\big(1 + (\mathcal{R}_0^{o,\infty}(n))_{11}\big) + (\mathcal{R}_0^{o,\infty}(n))_{12}(\overset{o}{m}{}_2^{\infty})_{22} + (\mathcal{R}_1^{o,\infty}(n))_{11}(\overset{o}{m}{}_1^{\infty})_{12} \\
&\quad + (\mathcal{R}_1^{o,\infty}(n))_{12}(\overset{o}{m}{}_1^{\infty})_{22} + (\mathcal{R}_2^{o,\infty}(n))_{11}(\overset{o}{m}{}_0^{\infty})_{12} + (\mathcal{R}_2^{o,\infty}(n))_{12}(\overset{o}{m}{}_0^{\infty})_{22}, \\
(\widehat{Q}_2^o)_{21} &:= (\overset{o}{m}{}_2^{\infty})_{21}\big(1 + (\mathcal{R}_0^{o,\infty}(n))_{22}\big) + (\mathcal{R}_0^{o,\infty}(n))_{21}(\overset{o}{m}{}_2^{\infty})_{11} + (\mathcal{R}_1^{o,\infty}(n))_{21}(\overset{o}{m}{}_1^{\infty})_{11} \\
&\quad + (\mathcal{R}_1^{o,\infty}(n))_{22}(\overset{o}{m}{}_1^{\infty})_{21} + (\mathcal{R}_2^{o,\infty}(n))_{21}(\overset{o}{m}{}_0^{\infty})_{11} + (\mathcal{R}_2^{o,\infty}(n))_{22}(\overset{o}{m}{}_0^{\infty})_{21}, \\
(\widehat{Q}_2^o)_{22} &:= (\overset{o}{m}{}_2^{\infty})_{22}\big(1 + (\mathcal{R}_0^{o,\infty}(n))_{22}\big) + (\mathcal{R}_0^{o,\infty}(n))_{21}(\overset{o}{m}{}_2^{\infty})_{12} + (\mathcal{R}_1^{o,\infty}(n))_{21}(\overset{o}{m}{}_1^{\infty})_{12} \\
&\quad + (\mathcal{R}_1^{o,\infty}(n))_{22}(\overset{o}{m}{}_1^{\infty})_{22} + (\mathcal{R}_2^{o,\infty}(n))_{21}(\overset{o}{m}{}_0^{\infty})_{12} + (\mathcal{R}_2^{o,\infty}(n))_{22}(\overset{o}{m}{}_0^{\infty})_{22},
\end{aligned}$$

and

$$(\overset{o}{m}{}_0^{\infty})_{11} = \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0) + \boldsymbol{d}_o)\mathbb{E}^{-1}}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)} \left(\frac{\gamma_0^o + (\gamma_0^o)^{-1}}{2} \right) \frac{\theta_\infty^o(1,1,\boldsymbol{\Omega}^o)}{\theta_\infty^o(1,1,\mathbf{0})},$$

$$(\overset{o}{m}{}_0^{\infty})_{12} = \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0) + \boldsymbol{d}_o)\mathbb{E}^{-1}}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0) - \frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o + \boldsymbol{d}_o)} \left(\frac{\gamma_0^o - (\gamma_0^o)^{-1}}{2\mathrm{i}} \right) \frac{\theta_\infty^o(-1,1,\boldsymbol{\Omega}^o)}{\theta_\infty^o(-1,1,\mathbf{0})},$$

$$(\overset{o}{m}{}_0^{\infty})_{21} = -\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{2\mathrm{i}}\right)\frac{\theta_\infty^o(1,-1,\boldsymbol{\Omega}^o)}{\theta_\infty^o(1,-1,\mathbf{0})},$$

$$(\overset{o}{m}{}_0^{\infty})_{22} = \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{2}\right)\frac{\theta_\infty^o(-1,-1,\boldsymbol{\Omega}^o)}{\theta_\infty^o(-1,-1,\mathbf{0})},$$

$$(\overset{o}{m}{}_1^{\infty})_{11} = \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}^{-1}}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}\left(\left(\frac{\widetilde{\alpha}_\infty^o(1,1,\mathbf{0})\theta_\infty^o(1,1,\boldsymbol{\Omega}^o)-\widetilde{\alpha}_\infty^o(1,1,\boldsymbol{\Omega}^o)\theta_\infty^o(1,1,\mathbf{0})}{(\theta_\infty^o(1,1,\mathbf{0}))^2}\right)\right.$$
$$\left.\times\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{2}\right)-\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{8}\right)\left(\sum_{k=1}^{N+1}\left(a_k^o-b_{k-1}^o\right)\right)\frac{\theta_\infty^o(1,1,\boldsymbol{\Omega}^o)}{\theta_\infty^o(1,1,\mathbf{0})}\right),$$

$$(\overset{o}{m}{}_1^{\infty})_{12} = \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}^{-1}}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}\left(\left(\frac{\widetilde{\alpha}_\infty^o(-1,1,\mathbf{0})\theta_\infty^o(-1,1,\boldsymbol{\Omega}^o)-\widetilde{\alpha}_\infty^o(-1,1,\boldsymbol{\Omega}^o)\theta_\infty^o(-1,1,\mathbf{0})}{(\theta_\infty^o(-1,1,\mathbf{0}))^2}\right)\right.$$
$$\left.\times\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{2\mathrm{i}}\right)-\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{8\mathrm{i}}\right)\left(\sum_{k=1}^{N+1}\left(a_k^o-b_{k-1}^o\right)\right)\frac{\theta_\infty^o(-1,1,\boldsymbol{\Omega}^o)}{\theta_\infty^o(-1,1,\mathbf{0})}\right),$$

$$(\overset{o}{m}{}_1^{\infty})_{21} = -\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}\left(\left(\frac{\widetilde{\alpha}_\infty^o(1,-1,\mathbf{0})\theta_\infty^o(1,-1,\boldsymbol{\Omega}^o)-\widetilde{\alpha}_\infty^o(1,-1,\boldsymbol{\Omega}^o)\theta_\infty^o(1,-1,\mathbf{0})}{(\theta_\infty^o(1,-1,\mathbf{0}))^2}\right)\right.$$
$$\left.\times\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{2\mathrm{i}}\right)-\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{8\mathrm{i}}\right)\left(\sum_{k=1}^{N+1}\left(a_k^o-b_{k-1}^o\right)\right)\frac{\theta_\infty^o(1,-1,\boldsymbol{\Omega}^o)}{\theta_\infty^o(1,-1,\mathbf{0})}\right),$$

$$(\overset{o}{m}{}_1^{\infty})_{22} = \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}\left(\left(\frac{\widetilde{\alpha}_\infty^o(-1,-1,\mathbf{0})\theta_\infty^o(-1,-1,\boldsymbol{\Omega}^o)-\widetilde{\alpha}_\infty^o(-1,-1,\boldsymbol{\Omega}^o)\theta_\infty^o(-1,-1,\mathbf{0})}{(\theta_\infty^o(-1,-1,\mathbf{0}))^2}\right)\right.$$
$$\left.\times\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{2}\right)-\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{8}\right)\left(\sum_{k=1}^{N+1}\left(a_k^o-b_{k-1}^o\right)\right)\frac{\theta_\infty^o(-1,-1,\boldsymbol{\Omega}^o)}{\theta_\infty^o(-1,-1,\mathbf{0})}\right),$$

$$(\overset{o}{m}{}_2^{\infty})_{11} = \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}^{-1}}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}\left(\left(\theta_\infty^o(1,1,\boldsymbol{\Omega}^o)\left((\widetilde{\alpha}_\infty^o(1,1,\mathbf{0}))^2+\beta_\infty^o(1,1,\mathbf{0})\theta_\infty^o(1,1,\mathbf{0})\right)\right.\right.$$
$$\left.-\widetilde{\alpha}_\infty^o(1,1,\boldsymbol{\Omega}^o)\widetilde{\alpha}_\infty^o(1,1,\mathbf{0})\theta_\infty^o(1,1,\mathbf{0})-\beta_\infty^o(1,1,\boldsymbol{\Omega}^o)(\theta_\infty^o(1,1,\mathbf{0}))^2\right)\frac{1}{(\theta_\infty^o(1,1,\mathbf{0}))^3}$$
$$\times\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{2}\right)+\left(\frac{\widetilde{\alpha}_\infty^o(1,1,\mathbf{0})\theta_\infty^o(1,1,\boldsymbol{\Omega}^o)-\widetilde{\alpha}_\infty^o(1,1,\boldsymbol{\Omega}^o)\theta_\infty^o(1,1,\mathbf{0})}{(\theta_\infty^o(1,1,\mathbf{0}))^2}\right)$$
$$\times\left(\sum_{k=1}^{N+1}\left(b_{k-1}^o-a_k^o\right)\right)\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{8}\right)+\left(\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{16}\right)\sum_{k=1}^{N+1}\left((b_{k-1}^o)^2-(a_k^o)^2\right)\right.$$
$$\left.+\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{64}\right)\left(\sum_{k=1}^{N+1}\left(a_k^o-b_{k-1}^o\right)\right)^2\right)\frac{\theta_\infty^o(1,1,\boldsymbol{\Omega}^o)}{\theta_\infty^o(1,1,\mathbf{0})}\right),$$

$$(\overset{o}{m}{}_2^{\infty})_{12} = \frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}^{-1}}{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)}\left(\left(\theta_\infty^o(-1,1,\boldsymbol{\Omega}^o)\left((\widetilde{\alpha}_\infty^o(-1,1,\mathbf{0}))^2+\beta_\infty^o(-1,1,\mathbf{0})\theta_\infty^o(-1,1,\mathbf{0})\right)\right.\right.$$
$$\left.-\widetilde{\alpha}_\infty^o(-1,1,\boldsymbol{\Omega}^o)\widetilde{\alpha}_\infty^o(-1,1,\mathbf{0})\theta_\infty^o(-1,1,\mathbf{0})-\beta_\infty^o(-1,1,\boldsymbol{\Omega}^o)(\theta_\infty^o(-1,1,\mathbf{0}))^2\right)\frac{1}{(\theta_\infty^o(-1,1,\mathbf{0}))^3}$$
$$\times\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{2\mathrm{i}}\right)-\left(\frac{\widetilde{\alpha}_\infty^o(-1,1,\mathbf{0})\theta_\infty^o(-1,1,\boldsymbol{\Omega}^o)-\widetilde{\alpha}_\infty^o(-1,1,\boldsymbol{\Omega}^o)\theta_\infty^o(-1,1,\mathbf{0})}{(\theta_\infty^o(-1,1,\mathbf{0}))^2}\right)$$
$$\times\left(\sum_{k=1}^{N+1}\left(a_k^o-b_{k-1}^o\right)\right)\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{8\mathrm{i}}\right)-\left(\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{16\mathrm{i}}\right)\sum_{k=1}^{N+1}\left((a_k^o)^2-(b_{k-1}^o)^2\right)\right.$$
$$\left.-\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{64\mathrm{i}}\right)\left(\sum_{k=1}^{N+1}\left(a_k^o-b_{k-1}^o\right)\right)^2\right)\frac{\theta_\infty^o(-1,1,\boldsymbol{\Omega}^o)}{\theta_\infty^o(-1,1,\mathbf{0})}\right),$$

$$(\overset{o}{m}{}_2^{\infty})_{21} = -\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)\mathbb{E}}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o-\boldsymbol{d}_o)}\left(\left(\theta_\infty^o(1,-1,\boldsymbol{\Omega}^o)\left((\widetilde{\alpha}_\infty^o(1,-1,\mathbf{0}))^2+\beta_\infty^o(1,-1,\mathbf{0})\theta_\infty^o(1,-1,\mathbf{0})\right)\right.\right.$$

$$
\begin{aligned}
&-\widetilde{\alpha}^{o}_{\infty}(1,-1,\boldsymbol{\Omega}^{o})\widetilde{\alpha}^{o}_{\infty}(1,-1,\mathbf{0})\theta^{o}_{\infty}(1,-1,\mathbf{0})-\beta^{o}_{\infty}(1,-1,\boldsymbol{\Omega}^{o})(\theta^{o}_{\infty}(1,-1,\mathbf{0}))^{2}\Big)\tfrac{1}{(\theta^{o}_{\infty}(1,-1,\mathbf{0}))^{3}}\\
&\times\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{2\mathrm{i}}\right)+\left(\frac{\widetilde{\alpha}^{o}_{\infty}(1,-1,\mathbf{0})\theta^{o}_{\infty}(1,-1,\boldsymbol{\Omega}^{o})-\widetilde{\alpha}^{o}_{\infty}(1,-1,\boldsymbol{\Omega}^{o})\theta^{o}_{\infty}(1,-1,\mathbf{0})}{(\theta^{o}_{\infty}(1,-1,\mathbf{0}))^{2}}\right)\\
&\times\left(\sum_{k=1}^{N+1}\left(a_k^o-b_{k-1}^o\right)\right)\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{8\mathrm{i}}\right)+\left(\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{16\mathrm{i}}\right)\sum_{k=1}^{N+1}\left((a_k^o)^2-(b_{k-1}^o)^2\right)\right.\\
&\left.-\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{64\mathrm{i}}\right)\left(\sum_{k=1}^{N+1}\left(a_k^o-b_{k-1}^o\right)\right)^2\right)\frac{\theta^{o}_{\infty}(1,-1,\boldsymbol{\Omega}^{o})}{\theta^{o}_{\infty}(1,-1,\mathbf{0})}\Bigg),
\end{aligned}
$$

$$
\begin{aligned}
(\overset{o}{m}{}^{\infty}_{2})_{22}=\frac{\boldsymbol{\theta}^{o}(\boldsymbol{u}^{o}_{+}(0)+\boldsymbol{d}_{o})\mathbb{E}}{\boldsymbol{\theta}^{o}(-\boldsymbol{u}^{o}_{+}(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^{o}-\boldsymbol{d}_{o})}&\Big(\big(\theta^{o}_{\infty}(-1,-1,\boldsymbol{\Omega}^{o})\big((\widetilde{\alpha}^{o}_{\infty}(-1,-1,\mathbf{0}))^{2}+\beta^{o}_{\infty}(-1,-1,\mathbf{0})\theta^{o}_{\infty}(-1,-1,\mathbf{0})\big)\\
&-\widetilde{\alpha}^{o}_{\infty}(-1,-1,\boldsymbol{\Omega}^{o})\widetilde{\alpha}^{o}_{\infty}(-1,-1,\mathbf{0})\theta^{o}_{\infty}(-1,-1,\mathbf{0})-\beta^{o}_{\infty}(-1,-1,\boldsymbol{\Omega}^{o})(\theta^{o}_{\infty}(-1,-1,\mathbf{0}))^{2}\Big)\tfrac{1}{(\theta^{o}_{\infty}(-1,-1,\mathbf{0}))^{3}}\\
&\times\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{2}\right)+\left(\frac{\widetilde{\alpha}^{o}_{\infty}(-1,-1,\mathbf{0})\theta^{o}_{\infty}(-1,-1,\boldsymbol{\Omega}^{o})-\widetilde{\alpha}^{o}_{\infty}(-1,-1,\boldsymbol{\Omega}^{o})\theta^{o}_{\infty}(-1,-1,\mathbf{0})}{(\theta^{o}_{\infty}(-1,-1,\mathbf{0}))^{2}}\right)\\
&\times\left(\sum_{k=1}^{N+1}\left(b_{k-1}^o-a_k^o\right)\right)\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{8}\right)+\left(\left(\frac{\gamma_0^o-(\gamma_0^o)^{-1}}{16}\right)\sum_{k=1}^{N+1}\left((b_{k-1}^o)^2-(a_k^o)^2\right)\right.\\
&\left.+\left(\frac{\gamma_0^o+(\gamma_0^o)^{-1}}{64}\right)\left(\sum_{k=1}^{N+1}\left(a_k^o-b_{k-1}^o\right)\right)^2\right)\frac{\theta^{o}_{\infty}(-1,-1,\boldsymbol{\Omega}^{o})}{\theta^{o}_{\infty}(-1,-1,\mathbf{0})}\Bigg),
\end{aligned}
$$

where, for $\varepsilon_1, \varepsilon_2 = \pm 1$,

$$
\begin{aligned}
\theta^{o}_{\infty}(\varepsilon_1,\varepsilon_2,\mathfrak{Z}) &:= \boldsymbol{\theta}^{o}(\varepsilon_1\boldsymbol{u}^{o}_{+}(\infty)-\tfrac{1}{2\pi}(n+\tfrac{1}{2})\mathfrak{Z}+\varepsilon_2\boldsymbol{d}_{o}),\\
\widetilde{\alpha}^{o}_{\infty}(\varepsilon_1,\varepsilon_2,\mathfrak{Z}) &:= 2\pi\mathrm{i}\varepsilon_1\sum_{m\in\mathbb{Z}^N}(m,\widehat{\boldsymbol{\alpha}}^{o}_{\infty})\mathrm{e}^{2\pi\mathrm{i}(m,\varepsilon_1\boldsymbol{u}^{o}_{+}(\infty)-\frac{1}{2\pi}(n+\frac{1}{2})\mathfrak{Z}+\varepsilon_2\boldsymbol{d}_{o})+\pi\mathrm{i}(m,\tau^{o}m)},
\end{aligned}
$$

with $\boldsymbol{u}^{o}_{+}(\infty)=\int_{a^{o}_{N+1}}^{\infty^{+}}\boldsymbol{\omega}^{o}$, $\widehat{\boldsymbol{\alpha}}^{o}_{\infty}=(\widehat{\alpha}^{o}_{\infty,1},\widehat{\alpha}^{o}_{\infty,2},\ldots,\widehat{\alpha}^{o}_{\infty,N})$, *and* $\widehat{\alpha}^{o}_{\infty,j}:=c^{o}_{j1}$, $j=1,\ldots,N$,

$$
\beta^{o}_{\infty}(\varepsilon_1,\varepsilon_2,\mathfrak{Z}):=2\pi\sum_{m\in\mathbb{Z}^N}\left(\mathrm{i}\varepsilon_1(m,\widehat{\boldsymbol{\beta}}^{o}_{\infty})+\pi(m,\widehat{\boldsymbol{\alpha}}^{o}_{\infty})^2\right)\mathrm{e}^{2\pi\mathrm{i}(m,\varepsilon_1\boldsymbol{u}^{o}_{+}(\infty)-\frac{1}{2\pi}(n+\frac{1}{2})\mathfrak{Z}+\varepsilon_2\boldsymbol{d}_{o})+\pi\mathrm{i}(m,\tau^{o}m)},
$$

where $\widehat{\boldsymbol{\beta}}^{o}_{\infty}=(\widehat{\beta}^{o}_{\infty,1},\widehat{\beta}^{o}_{\infty,2},\ldots,\widehat{\beta}^{o}_{\infty,N})$, *with* $\widehat{\beta}^{o}_{\infty,j}:=\frac{1}{2}(c^{o}_{j2}+\frac{1}{2}c^{o}_{j1}\sum_{i=1}^{N+1}(b^{o}_{i-1}+a^{o}_{i}))$, $j=1,\ldots,N$, *where* c^{o}_{j1}, c^{o}_{j2}, $j=1,\ldots,N$, *are obtained from Equations* (O1) *and* (O2), *and* γ^{o}_{0} *is defined in Theorem* 3.1, *Equations* (3.53), *and*

$$
\begin{aligned}
\mathcal{R}^{o,\infty}_{0}(n)\underset{n\to\infty}{=}&\frac{1}{(n+\frac{1}{2})}\sum_{j=1}^{N+1}\Bigg(\frac{\left(\mathcal{B}^{o}(b^{o}_{j-1})\widehat{\alpha}^{o}_{0}(b^{o}_{j-1})-\mathcal{A}^{o}(b^{o}_{j-1})\left(\widehat{\alpha}^{o}_{1}(b^{o}_{j-1})+(b^{o}_{j-1})^{-1}\widehat{\alpha}^{o}_{0}(b^{o}_{j-1})\right)\right)}{b^{o}_{j-1}(\widehat{\alpha}^{o}_{0}(b^{o}_{j-1}))^{2}}\\
&+\frac{\left(\mathcal{B}^{o}(a^{o}_{j})\widehat{\alpha}^{o}_{0}(a^{o}_{j})-\mathcal{A}^{o}(a^{o}_{j})\left(\widehat{\alpha}^{o}_{1}(a^{o}_{j})+(a^{o}_{j})^{-1}\widehat{\alpha}^{o}_{0}(a^{o}_{j})\right)\right)}{a^{o}_{j}(\widehat{\alpha}^{o}_{0}(a^{o}_{j}))^{2}}\Bigg)+O\left(\frac{c(n)}{(n+\frac{1}{2})^{2}}\right),\\
\mathcal{R}^{o,\infty}_{1}(n)\underset{n\to\infty}{=}&\frac{1}{(n+\frac{1}{2})}\sum_{j=1}^{N+1}\Bigg(\frac{\left(\mathcal{B}^{o}(b^{o}_{j-1})\widehat{\alpha}^{o}_{0}(b^{o}_{j-1})-\mathcal{A}^{o}(b^{o}_{j-1})\widehat{\alpha}^{o}_{1}(b^{o}_{j-1})\right)}{(\widehat{\alpha}^{o}_{0}(b^{o}_{j-1}))^{2}}\\
&+\frac{\left(\mathcal{B}^{o}(a^{o}_{j})\widehat{\alpha}^{o}_{0}(a^{o}_{j})-\mathcal{A}^{o}(a^{o}_{j})\widehat{\alpha}^{o}_{1}(a^{o}_{j})\right)}{(\widehat{\alpha}^{o}_{0}(a^{o}_{j}))^{2}}\Bigg)+O\left(\frac{c(n)}{(n+\frac{1}{2})^{2}}\right),\\
\mathcal{R}^{o,\infty}_{2}(n)\underset{n\to\infty}{=}&\frac{1}{(n+\frac{1}{2})}\sum_{j=1}^{N+1}\Bigg(\frac{\left(\mathcal{B}^{o}(b^{o}_{j-1})\widehat{\alpha}^{o}_{0}(b^{o}_{j-1})b^{o}_{j-1}-\mathcal{A}^{o}(b^{o}_{j-1})\left(b^{o}_{j-1}\widehat{\alpha}^{o}_{1}(b^{o}_{j-1})-\widehat{\alpha}^{o}_{0}(b^{o}_{j-1})\right)\right)}{(\widehat{\alpha}^{o}_{0}(b^{o}_{j-1}))^{2}}\\
&+\frac{\left(\mathcal{B}^{o}(a^{o}_{j})\widehat{\alpha}^{o}_{0}(a^{o}_{j})a^{o}_{j}-\mathcal{A}^{o}(a^{o}_{j})\left(a^{o}_{j}\widehat{\alpha}^{o}_{1}(a^{o}_{j})-\widehat{\alpha}^{o}_{0}(a^{o}_{j})\right)\right)}{(\widehat{\alpha}^{o}_{0}(a^{o}_{j}))^{2}}\Bigg)+O\left(\frac{c(n)}{(n+\frac{1}{2})^{2}}\right),
\end{aligned}
$$

with all parameters defined in Theorem B.1.

Theorem B.3 ([22]**).** *Let all the conditions stated in Theorem* B.1 *be valid, and let all parameters be as defined therein. Let* $\overset{o}{\mathrm{Y}}\colon \mathbb{C}\setminus\mathbb{R}\to \mathrm{SL}_2(\mathbb{C})$ *be the unique solution of* **RHP2** *with integral representation* (1.30). *Let* $\boldsymbol{\pi}_{2n+1}(z)$ *be the odd degree monic orthogonal L-polynomial defined in Equations* (1.5) *with* $n\to\infty$ *asymptotics (in the entire complex plane) given by Theorem* 2.3.1 *of* [22]. *Then,*

$$\left(\xi_{-n-1}^{(2n+1)}\right)^2=\frac{1}{\|\boldsymbol{\pi}_{2n+1}(\cdot)\|^2_{\mathcal{L}}}=\frac{H_{2n+1}^{(-2n)}}{H_{2n+1}^{(-2n-2)}}\underset{n\to\infty}{=}\frac{\mathrm{e}^{-n\ell_o}}{\pi}\Xi^{\natural}\left(1+\frac{2}{n+\frac{1}{2}}\Xi^{\natural}(\mathfrak{Q}^{\natural})_{12}+O\left(\frac{c^{\natural}(n)}{(n+\frac{1}{2})^2}\right)\right),\tag{B.1}$$

where

$$\Xi^{\natural}:=2\mathbb{E}^2\left(\sum_{k=1}^{N+1}\left((b_{k-1}^o)^{-1}-(a_k^o)^{-1}\right)\right)^{-1}\frac{\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}{\boldsymbol{\theta}^o(-\boldsymbol{u}_+^o(0)-\frac{1}{2\pi}(n+\frac{1}{2})\boldsymbol{\Omega}^o+\boldsymbol{d}_o)\boldsymbol{\theta}^o(\boldsymbol{u}_+^o(0)+\boldsymbol{d}_o)}\quad(>0),$$

$\mathfrak{Q}^{\natural}$ *is defined in Theorem* 3.2, *Equations* (3.107), $(\mathfrak{Q}^{\natural})_{12}$ *denotes the* (1 2)*-element of* $\mathfrak{Q}^{\natural}$, $c^{\natural}(n)=_{n\to\infty}O(1)$, *and all relevant parameters are defined in Theorem* B.1. *Asymptotics for* $\xi_{-n-1}^{(2n+1)}$ *are obtained by taking the positive square root of both sides of Equation* (B.1).

Bibliography

[1] P. Deift and X. Zhou, "A Steepest Descent Method for Oscillatory Riemann-Hilbert Problems. Asymptotics for the MKdV Equation", Ann. of Math., Vol. 137, No. 2, pp. 295–368, 1993.

[2] P. Deift and X. Zhou, "Asymptotics for the Painlevé II Equation", Comm. Pure Appl. Math., Vol. 48, No. 3, pp. 277–337, 1995.

[3] P. Deift, S. Venakides, and X. Zhou, "New Results in Small Dispersion KdV by an Extension of the Steepest Descent Method for Riemann-Hilbert Problems", Int. Math. Res. Not., No. 6, pp. 285–299, 1997.

[4] W. B. Jones, W. J. Thron, and H. Waadeland, "A Strong Stieltjes Moment Problem", Trans. Amer. Math. Soc., Vol. 261, No. 2, pp. 503–528, 1980.

[5] W. B. Jones and W. J. Thron, "Survey of Continued Fractions Methods of Solving Moment Problems and Related Topics", pp. 4–37, in W. B. Jones, W. J. Thron, and H. Waadeland, eds., *Analytic Theory of Continued Fractions*, Lecture Notes in Mathematics, Vol. 932, Springer-Verlag, Berlin, 1982.

[6] W. B. Jones and W. J. Thron, "Orthogonal Laurent Polynomials and the Strong Hamburger Moment Problem", J. Math. Anal. Appl., Vol. 98, No. 2, pp. 528–554, 1984.

[7] O. Njåstad and W. J. Thron, "Unique Solvability of the Strong Hamburger Moment Problem", J. Austral. Math. Soc. Ser. A, Vol. 40, No. 1, pp. 5–19, 1986.

[8] O. Njåstad, "Solutions of the Strong Stieltjes Moment Problem", Methods Appl. Anal., Vol. 2, No. 3, pp. 320–347, 1995.

[9] O. Njåstad, "Extremal Solutions of the Strong Stieltjes Moment Problem", J. Comput. Appl. Math., Vol. 65, No. 1–3, pp. 309–318, 1995.

[10] O. Njåstad, "Solutions of the Strong Hamburger Moment Problem", J. Math. Anal. Appl., Vol. 197, No. 1, pp. 227–248, 1996.

[11] W. B. Jones and O. Njåstad, "Orthogonal Laurent Polynomials and Strong Moment Theory: A Survey", J. Comput. Appl. Math., Vol. 105, No. 1–2, pp. 51–91, 1999.

[12] T. J. Stieltjes, "Recherches sur les fractions continues", Ann. Fac. Sci. Univ. Toulouse, Vol. 8, J1–J122, 1894, Ann. Fac. Sci. Univ. Toulouse, Vol. 9, A1–A47, 1895 (in French).

[13] H. Hamburger, "Über eine Erweiterung des Stieltjesschen Momentproblems", Parts I, II, III, Math. Annalen, Vol. 81, pp. 235–319, 1920, Vol. 82, pp. 120–164, 1921, Vol. 82, pp. 168–187, 1921 (in German).

[14] G. Szegö, *Orthogonal Polynomials*, 4th edn., American Mathematical Society Colloquium Publications, Vol. 23, AMS, Providence, 1974.

[15] J. A. Shohat and J. D. Tamarkin, *The Problem of Moments*, American Mathematical Society Surveys, Vol. II, AMS, New York, 1943.

[16] E. Hendriksen and H. van Rossum, "Orthogonal Laurent Polynomials", Nederl. Akad. Wetensch. Indag. Math., Vol. 48, No. 1, pp. 17–36, 1986.

[17] L. Cochran and S. C. Cooper, "Orthogonal Laurent Polynomials on the Real Line", pp. 47–100, in S. C. Cooper and W. J. Thron, eds., *Continued Fractions and Orthogonal Functions: Theory and Applications*, Lecture Notes in Pure and Applied Mathematics, Vol. 154, Marcel Dekker, Inc., New York, 1994.

[18] M. E. H. Ismail and D. R. Masson, "Generalized Orthogonality and Continued Fractions", J. Approx. Theory, Vol. 83, No. 1, pp. 1–40, 1995.

[19] A. Zhedanov, "The "Classical" Laurent Biorthogonal Polynomials", J. Comput. Appl. Math., Vol. 98, No. 1, pp. 121–147, 1998.

[20] A. Zhedanov, "Biorthogonal Rational Functions and the Generalized Eigenvalue Problem", J. Approx. Theory, Vol. 101, No. 2, pp. 303–329, 1999.

[21] K. T.-R. McLaughlin, A. H. Vartanian, and X. Zhou, "Asymptotics of Laurent Polynomials of Even Degree Orthogonal with Respect to Varying Exponential Weights", Preprint, 2006.

[22] K. T.-R. McLaughlin, A. H. Vartanian, and X. Zhou, "Asymptotics of Laurent Polynomials of Odd Degree Orthogonal with Respect to Varying Exponential Weights", Preprint, 2006.

[23] A. Bultheel, P. Gonzalez-Vera, E. Hendriksen, and O. Njåstad, *Orthogonal Rational Functions*, Cambridge Monographs on Applied and Computational Mathematics, Vol. 5, Cambridge University Press, Cambridge, 1999.

[24] M. M. Djrbashian, "A Survey on the Theory of Orthogonal Systems and Some Open Problems", pp. 135–146, in P. Nevai, ed., *Orthogonal Polynomials. Theory and Practice*, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., Vol. 294, Kluwer Acad. Publ., Dordrecht, 1990.

[25] N. M. Ercolani and K. T.-R. McLaughlin, "Asymptotics of the Partition Function for Random Matrices via Riemann-Hilbert Techniques and Applications to Graphical Enumeration", Int. Math. Res. Not., No. 14, pp. 755–820, 2003.

[26] E. Hendriksen and C. Nijhuis, "Laurent-Jacobi Matrices and the Strong Hamburger Moment Problem", Acta Appl. Math., Vol. 61, No. 1–3, pp. 119–132, 2000.

[27] M. J. Cantero, L. Moral, and L. Velázquez, "Five-Diagonal Matrices and Zeros of Orthogonal Polynomials on the Unit Circle", Linear Algebra Appl., Vol. 362, pp. 29–56, 2003.

[28] B. Simon, "Orthogonal Polynomials on the Unit Circle: New Results", Int. Math. Res. Not., No. 53, pp. 2837–2880, 2004.

[29] A. Bultheel, P. González-Vera, E. Hendriksen, and O. Njåstad, "Quadrature and Orthogonal Rational Functions", J. Comput. Appl. Math., Vol. 127, No. 1–2, pp. 67–91, 2001.

[30] E. Hendriksen, "The Strong Hamburger Moment Problem and Self-Adjoint Operators in Hilbert Space", J. Comput. Appl. Math., Vol. 19, No. 1, pp. 79–88, 1987.

[31] P. González-Vera and O. Njåstad, "Convergence of Two-Point Padé Approximants to Series of Stieltjes", J. Comput. Appl. Math., Vol. 32, No. 1–2, pp. 97–105, 1990.

[32] J. Coussement, A. B. J. Kuijlaars, and W. Van Assche, "Direct and Inverse Spectral Transform for the Relativistic Toda Lattice and the Connection with Laurent Orthogonal Polynomials", Inverse Problems, Vol. 18, No. 3, pp. 923–942, 2002.

[33] J. Coussement and W. Van Assche, "A Continuum Limit of the Relativistic Toda Lattice: Asymptotic Theory of Discrete Laurent Orthogonal Polynomials with Varying Recurrence Coefficients", J. Phys. A: Math. Gen., Vol. 38, No. 15, pp. 3337–3366, 2005.

[34] J. Coussement and W. Van Assche, "An Extension of the Toda Lattice: A Direct and Inverse Spectral Transform Connected with Orthogonal Rational Functions", Inverse Problems, Vol. 20, No. 1, pp. 297–318, 2004.

[35] O. Njåstad, "A Modified Schur Algorithm and an Extended Hamburger Moment Problem", Trans. Amer. Math. Soc., Vol. 327, No. 1, pp. 283–311, 1991.

[36] E. B. Saff and V. Totik, *Logarithmic Potentials with External Fields*, Grundlehren der mathematischen Wissenschaften 316, Springer-Verlag, Berlin, 1997.

[37] P. Deift, T. Kriecherbauer, and K. T.-R. McLaughlin, "New Results on the Equilibrium Measure for Logarithmic Potentials in the Presence of an External Field", J. Approx. Theory, Vol. 95, No. 3, pp. 388–475, 1998.

[38] A. B. J. Kuijlaars and K. T.-R. McLaughlin, "Generic Behavior of the Density of States in Random Matrix Theory and Equilibrium Problems in the Presence of Real Analytic External Fields", Comm. Pure Appl. Math., Vol. 53, No. 6, pp. 736–785, 2000.

[39] K. Johansson, "On Fluctuations of Eigenvalues of Random Hermitian Matrices", Duke Math. J., Vol. 91, No. 1, pp. 151–204, 1998.

[40] P. Deift, T. Kriecherbauer, K. T.-R. McLaughlin, S. Venakides, and X. Zhou, "Uniform Asymptotics for Polynomials Orthogonal with Respect to Varying Exponential Weights and Applications to Universality Questions in Random Matrix Theory", Comm. Pure Appl. Math., Vol. 52, No. 11, pp. 1335–1425, 1999.

[41] P. Deift, T. Kriecherbauer, K. T.-R. McLaughlin, S. Venakides, and X. Zhou, "Strong Asymptotics of Orthogonal Polynomials with Respect to Exponential Weights", Comm. Pure Appl. Math., Vol. 52, No. 12, pp. 1491–1552, 1999.

[42] T. Kriecherbauer and K. T.-R. McLaughlin, "Strong Asymptotics of Polynomials Orthogonal with Respect to Freud Weights", Int. Math. Res. Not., No. 6, pp. 299–333, 1999.

[43] A. B. J. Kuijlaars, K. T.-R. McLaughlin, W. Van Assche, and M. Vanlessen, "The Riemann-Hilbert Approach to Strong Asymptotics for Orthogonal Polynomials on $[-1, 1]$", Adv. Math., Vol. 188, No. 2, pp. 337–398, 2004.

[44] M. Vanlessen, "Strong Asymptotics of the Recurrence Coefficients of Orthogonal Polynomials Associated to the Generalized Jacobi Weight", J. Approx. Theory, Vol. 125, No. 2, pp. 199–229, 2003.

[45] J. Baik, T. Kriecherbauer, K. T.-R. McLaughlin, and P. D. Miller, "Uniform Asymptotics for Polynomials Orthogonal with Respect to a General Class of Discrete Weights and Universality Results for Associated Ensembles", `arXiv:math.CA/0310278 v1`.

[46] P. Deift, *Orthogonal Polynomials and Random Matrices: A Riemann-Hilbert Approach*, Courant Lecture Notes in Mathematics, Vol. 3, Courant Institute of Mathematical Sciences, New York, 1999.

[47] A. B. J. Kuijlaars, H. Stahl, W. Van Assche, and F. Wielonsky, "Type II Hermite-Padé Approximation to the Exponential Function", `arXiv:math.CA/0510278 v1`.

[48] A. S. Fokas, A. R. Its, and A. V. Kitaev, "Discrete Painlevé Equations and Their Appearance in Quantum Gravity", Comm. Math. Phys., Vol. 142, No. 2, pp. 313–344, 1991.

[49] A. S. Fokas, A. R. Its, and A. V. Kitaev, "The Isomonodromy Approach to Matrix Models in $2D$ Quantum Gravity", Comm. Math. Phys., Vol. 147, No. 2, pp. 395–430, 1992.

[50] X. Zhou, "Direct and Inverse Scattering Transforms with Arbitrary Spectral Singularities", Comm. Pure Appl. Math., Vol. 42, No. 7, pp. 895–938, 1989.

[51] X. Zhou, "Inverse Scattering Transform for Systems with Rational Spectral Dependence", J. Differential Equations, Vol. 115, No. 2, pp. 277–303, 1995.

[52] K. T.-R. McLaughlin and P. D. Miller, "The $\overline{\partial}$ Steepest Descent Method and the Asymptotic Behavior of Polynomials Orthogonal on the Unit Circle with Fixed and Exponentially Varying Nonanalytic Weights", `arXiv:math.CA/0406484 v1`.

[53] P. A. Deift, A. R. Its, and X. Zhou, "A Riemann-Hilbert Approach to Asymptotic Problems Arising in the Theory of Random Matrix Models, and also in the Theory of Integrable Statistical Mechanics", Ann. of Math., Vol. 146, No. 1, pp. 149–235, 1997.

[54] J. Baik, P. Deift, and K. Johansson, "On the Distribution of the Length of the Longest Increasing Subsequence of Random Permutations", J. Amer. Math. Soc., Vol. 12, No. 4, pp. 1119–1178, 1999.

[55] S. Kamvissis, K. D. T.-R. McLaughlin, and P. D. Miller, *Semiclassical Soliton Ensembles for the Focusing Nonlinear Schrödinger Equation*, Annals of Mathematics Studies, Vol. 154, Princeton University Press, New Jersey, 2003.

[56] P. D. Miller, "Asymptotics of Semiclassical Soliton Ensembles: Rigorous Justification of the WKB Approximation", Int. Math. Res. Not., No. 8, pp. 383–454, 2002.

[57] A. I. Aptekarev, "Sharp Constants for Rational Approximations of Analytic Functions", Sb. Math., Vol. 193, No. 1, pp. 1–72, 2002.

[58] P. Deift and X. Zhou, "Perturbation Theory for Infinite-Dimensional Integrable Systems on the Line. A Case Study.", Acta Math., Vol. 188, No. 2, pp. 163–262, 2002 (Extended version: http://www.ml.kva.se/publications/acta/webarticles/deift).

[59] P. Deift and X. Zhou, "A Priori L^p-Estimates for Solutions of Riemann-Hilbert Problems", Int. Math. Res. Not., No. 40, pp. 2121–2154, 2002.

[60] P. Deift and X. Zhou, "Long-Time Asymptotics for Solutions of the NLS Equation with Initial Data in a Weighted Sobolev Space", Comm. Pure Appl. Math., Vol. 56, No. 8, pp. 1029–1077, 2003.

[61] A. B. J. Kuijlaars and M. Vanlessen, "Universality for Eigenvalue Correlations at the Origin of the Spectrum", Comm. Math. Phys., Vol. 243, No. 1, pp. 163–191, 2003.

[62] A. B. J. Kuijlaars, W. Van Assche, and F. Wielonsky, "Quadratic Hermite-Padé Approximation to the Exponential Function: a Riemann-Hilbert Approach", Constr. Approx., Vol. 21, No. 3, pp. 351–412, 2005.

[63] P. Deift and D. Gioev, 'Universality in Random Matrix Theory for Orthogonal and Symplectic Ensembles", arXiv:math-ph/0411075 v3.

[64] P. Deift and D. Gioev, 'Universality at the Edge of the Spectrum for Unitary, Orthogonal and Symplectic Ensembles of Random Matrices", arXiv:math-ph/0507023 v2.

[65] G. Lyng and P. D. Miller, "The N-Soliton of the Focusing Nonlinear Schrödinger Equation for N Large", arXiv:nlin.SI/0508007 v1.

[66] T. Claeys, A. B. J. Kuijlaars, and M. Vanlessen, "Multi-Critical Unitary Random Matrix Ensembles and the General Painlevé II Equation", arXiv:math-ph/0508062 v1.

[67] M. Vanlessen, "Strong Asymptotics of Laguerre-Type Orthogonal Polynomials and Applications in Random Matrix Theory", arXiv:math.CA/0504604 v2.

[68] G. Springer, *Introduction to Riemann Surfaces*, 2nd edn., Chelsea, New York, 1981.

[69] H. M. Farkas and I. Kra, *Riemann Surfaces*, 2nd edn., Graduate Texts in Mathematics, Vol. 71, Springer-Verlag, New York, 1992.