

```

> with(DifferentialGeometry) : with(Tensor) : with(Tools) : with(JetCalculus) :
  with(LinearAlgebra) : with(PDETools) :
Preferences("ShowFramePrompt", false) :
>

```

dKP hierarchy 4D

```

> DGsetup([x, y, t, z], [s, v], Js, 3) :
> Ssyst := {s2 = 1/2 s1^2 + v1, s3 = 1/3 s1^3 + v1 s1 + v2, s4 = 1/4 s1^4 + v1 s1^2 + v2 s1 + v3 + 1/2 v1^2} :
> Vsyst := map(Q -> coeffs(Q, {s1, s1,1}), expand(subs({s1,2 = TotalDiff(subs(Ssyst, s2),
  1), s1,3 = TotalDiff(subs(Ssyst, s3), 1), s1,4 = TotalDiff(subs(Ssyst, s4), 1)},
  {TotalDiff(subs(Ssyst, s3), 2) - TotalDiff(subs(Ssyst, s2), 3), TotalDiff(subs(Ssyst,
  s4), 2) - TotalDiff(subs(Ssyst, s2), 4), TotalDiff(subs(Ssyst, s4), 3)
  - TotalDiff(subs(Ssyst, s3), 4)}))) ; Vs2 := solve(Vsyst, {v1,2, v1,3, v1,4})

```

$$Vsyst := \{v_1 v_{1,1} - v_{1,3} + v_{2,2}, v_1 v_{1,2} + v_2 v_{1,1} - v_{1,4} + v_{2,3}, -v_1^2 v_{1,1} + v_2 v_{1,2} - v_{2,4} + v_{3,3}\}$$

$$Vs2 := \left\{ v_{1,2} = \frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2}, v_{1,3} = v_1 v_{1,1} + v_{2,2}, v_{1,4} = \frac{v_1^3 v_{1,1} + v_2^2 v_{1,1} + v_1 v_{2,4} - v_1 v_{3,3} + v_2 v_{2,3}}{v_2} \right\} \quad (1.1)$$

```

> Vs2a := solve(Vsyst, {v1,1, v1,2, v1,3}); Vs2b := solve(Vsyst, {v2,2, v2,3, v2,4}); Vs2c
  := solve(Vsyst, {v1,1, v2,2, v3,3})

```

$$Vs2a := \left\{ v_{1,1} = -\frac{v_1 v_{2,4} - v_1 v_{3,3} - v_2 v_{1,4} + v_2 v_{2,3}}{v_1^3 + v_2^2}, v_{1,2} = \frac{v_1^2 v_{1,4} - v_1^2 v_{2,3} + v_2 v_{2,4} - v_2 v_{3,3}}{v_1^3 + v_2^2}, v_{1,3} = \frac{v_1^3 v_{2,2} - v_1^2 v_{2,4} + v_1^2 v_{3,3} + v_1 v_2 v_{1,4} - v_1 v_2 v_{2,3} + v_2^2 v_{2,2}}{v_1^3 + v_2^2} \right\}$$

$$Vs2b := \{v_{2,2} = -v_1 v_{1,1} + v_{1,3}, v_{2,3} = -v_1 v_{1,2} - v_2 v_{1,1} + v_{1,4}, v_{2,4} = -v_1^2 v_{1,1} + v_2 v_{1,2} + v_{3,3}\}$$

$$Vs2c := \left\{ v_{1,1} = -\frac{v_1 v_{1,2} - v_{1,4} + v_{2,3}}{v_2}, v_{2,2} = \frac{v_1^2 v_{1,2} - v_1 v_{1,4} + v_1 v_{2,3} + v_2 v_{1,3}}{v_2}, v_{3,3} = -\frac{v_1^3 v_{1,2} - v_1^2 v_{1,4} + v_1^2 v_{2,3} + v_2^2 v_{1,2} - v_2 v_{2,4}}{v_2} \right\} \quad (1.2)$$

```

> Eq1 := Vsyst[1] : Eq2 := Vsyst[2] : Eq3 := Vsyst[3] :
> Vss3 := Vsyst :for i to 4 do for j to nops(Vsyst) do Vss3 := {op(Vss3), TotalDiff(Vsyst[j],
i)} :od:od:
nops(eliminate(Vss3, {v1,2, v1,3, v1,4, v1,1,2, v1,1,3, v1,1,4, v1,2,2, v1,2,3, v1,2,4, v1,3,3, v1,3,4,
v1,4,4}[2]))
1
(1.3)

```

```

> SymbEO2a := add(add(diff(Vsyst[1], vi,j) · pi · pj, j=i..4), i=1..4);
SymbEO2b := add(add(diff(Vsyst[2], vi,j) · pi · pj, j=i..4), i=1..4);
SymbEO2c := add(add(diff(Vsyst[3], vi,j) · pi · pj, j=i..4), i=1..4);
SymbEO2a := p1^2 v1 - p1 p3 + p2^2
SymbEO2b := p1^2 v2 + p1 p2 v1 - p1 p4 + p2 p3
SymbEO2c := -p1^2 v1^2 + p1 p2 v2 - p2 p4 + p3^2
(1.4)

```

```

> simplify(subs({p1=1, p2=q, p3=q^2+v1, p4=q^3+2 v1 q+v2}, {SymbEO2a, SymbEO2b,
SymbEO2c}))
{0}
(1.5)

```

```

> tmp := solve({coeffs(expand((a1·p1+a2·p2+a3·p3+a4·p4)·SymbEO2a+(b1·p1+b2
·p2+b3·p3+b4·p4)·SymbEO2b+(c1·p1+c2·p2+c3·p3+c4·p4)·SymbEO2c),
{p1,p2,p3,p4})}, {a1,a2,a3,a4,b1,b2,b3,b4,c1,c2,c3,c4})
tmp := {a1=c1 v1 - c2 v2, a2=0, a3=c1, a4=c2, b1=c2 v1, b2=-c1, b3=-c2, b4=0, c1
=c1, c2=c2, c3=0, c4=0}
(1.6)

```

```

> simplify([(v1 p1 + p3) · SymbEO2a - p2 · SymbEO2b + p1 · SymbEO2c, (-v2 p1 + p4)
· SymbEO2a + (v1 p1 - p3) · SymbEO2b + p2 · SymbEO2c])
[0, 0]
(1.7)

```

```

> simplify(subs(VsS2, v1 · TotalDiff(Eq1, 1) + TotalDiff(Eq1, 3) - TotalDiff(Eq2, 2)
+ TotalDiff(Eq3, 1))) ;
simplify(subs(VsS2, -v2 · TotalDiff(Eq1, 1) + TotalDiff(Eq1, 4) + v1 · TotalDiff(Eq2, 1)
- TotalDiff(Eq2, 3) + TotalDiff(Eq3, 2)))
0
0
(1.8)

```

```

> VsS3 := solve(Vss3, {v1,2, v1,3, v1,4, v1,1,2, v1,1,3, v1,1,4, v1,2,2, v1,2,3, v1,2,4, v1,3,3, v1,3,4,
v1,4,4, v2,2,2}) :
[nops({v1,2, v1,3, v1,4, v1,1,2, v1,1,3, v1,1,4, v1,2,2, v1,2,3, v1,2,4, v1,3,3, v1,3,4, v1,4,4,
v2,2,2}), nops(VsS3)]
[13, 13]
(1.9)

```

```

> Vss4 := Vss3 :for i to 4 do for k to i do for j to nops(Vsyst) do Vss4 := {op(Vss4),
TotalDiff(Vsyst[j], [i, k])} :od:od:od:
tmp := eliminate(Vss4, {v1,2, v1,3, v1,4, v1,1,2, v1,1,3, v1,1,4, v1,2,2, v1,2,3, v1,2,4, v1,3,3,
v1,3,4, v1,4,4, v2,2,2, v1,1,1,2, v1,1,1,3, v1,1,1,4, v1,1,2,2, v1,1,2,3, v1,1,2,4, v1,1,3,3,
v1,1,3,4, v1,1,4,4, v1,2,2,2, v1,2,2,3, v1,2,2,4, v1,2,3,3, v1,2,3,4, v1,2,4,4, v1,3,3,3, v1,3,3,4,

```

$$v_{1,3,4,4}, v_{1,4,4,4}, v_{2,2,2,2}, v_{2,2,2,3}, v_{2,2,2,4}\} : [nops(tmp[2]), nops(tmp[1])]]$$

[0, 35]

(1.10)

> VsS4 := tmp[1] :

This shows that dimensions of the symbol spaces (g[0],g[1],g[2],g[3],g[4],...) are equal to (1,4,7,10,13,...),

i.e. the Hilbert polynomial is given by the formula $\dim(g[k])=1+3k$, whence the general solution depends on 3 functions of 1 variable.

This is also what is expected from the characteristic variety (rational normal curve in our case) that has order 3 and dimension 1.

>

> DGsetup([x, y, t, z], [q, v], Jq, 3) :

> Qsyst := { $q_2 = q_{[1]} q_1 + v_{1,1}$, $q_3 = q_{[1]}^2 q_1 + v_{1,1} q_1 + v_{1,1} q_{[1]} + v_{1,2}$, $q_4 = q_{[1]}^3 q_1 + 2 v_{1,1} q_{[1]} q_1 + v_{1,1} q_{[1]}^2 + v_{2,1} q_1 + v_{1,2} q_{[1]} + v_{1,3} + v_{1,1} v_{1,1}$ } :

> simplify(subs(VsS3, expand(subs(Qsyst, subs({ $q_{1,2} = \text{TotalDiff}(\text{subs}(Qsyst, q_2), 1)$, $q_{1,3} = \text{TotalDiff}(\text{subs}(Qsyst, q_3), 1)$, $q_{1,4} = \text{TotalDiff}(\text{subs}(Qsyst, q_4), 1)$ }),
{ $\text{TotalDiff}(\text{subs}(Qsyst, q_3), 2) - \text{TotalDiff}(\text{subs}(Qsyst, q_2), 3)$,
 $\text{TotalDiff}(\text{subs}(Qsyst, q_4), 2) - \text{TotalDiff}(\text{subs}(Qsyst, q_2), 4)$, $\text{TotalDiff}(\text{subs}(Qsyst, q_4), 3) - \text{TotalDiff}(\text{subs}(Qsyst, q_3), 4)$ })))))

{0}

(1.11)

> DGsetup([x, y, t, z, q], [v], Jqv, 3) :

> Yf := f → TotalDiff(f, 2) - q · TotalDiff(f, 1) + $v_{1,1} \cdot \text{diff}(f, q)$:

Tf := f → TotalDiff(f, 3) - ($q^2 + v_{1,1}$) · TotalDiff(f, 1) + ($q \cdot v_{1,1} + v_{1,2}$) · diff(f, q) :

Zf := f → TotalDiff(f, 4) - ($q^3 + 2 v_{1,1} q + v_{2,1}$) · TotalDiff(f, 1) + ($q^2 v_{1,1} + q v_{1,2} + v_{1,3} + v_{1,1} v_{1,1}$) · diff(f, q) :

> simplify(subs(VsS3, {simplify(Yf(Tf(x)) - Tf(Yf(x))), simplify(Yf(Tf(q)) - Tf(Yf(q))), simplify(Yf(Zf(x)) - Zf(Yf(x))), simplify(Yf(Zf(q)) - Zf(Yf(q))), simplify(Tf(Zf(x)) - Zf(Tf(x))), simplify(Tf(Zf(q)) - Zf(Tf(q)))}))

{0}

(1.12)

> VSQtmp := solve(map(Q → coeffs(Q, {q}), {simplify(Yf(Tf(x)) - Tf(Yf(x))), simplify(Yf(Tf(q)) - Tf(Yf(q))), simplify(Yf(Zf(x)) - Zf(Yf(x))), simplify(Yf(Zf(q)) - Zf(Yf(q))), simplify(Tf(Zf(x)) - Zf(Tf(x))), simplify(Tf(Zf(q)) - Zf(Tf(q)))}), { $v_{1,2}, v_{1,3}, v_{1,4}, v_{1,1,2}, v_{1,1,3}, v_{1,1,4}$ })

$$VSQtmp := \begin{cases} v_{1,2} = v_{1,2}, v_{1,3} = v_{1,1} v_{1,1} + v_{2,2}, v_{1,4} = v_{1,1} v_{1,2} + v_{2,1} v_{1,1} + v_{2,3}, v_{1,1,2} \end{cases} \quad (1.13)$$

$$= \frac{v_1^2 v_{1,1,1} + 2 v_1 v_{1,1}^2 - v_{1,2}^2 + v_{1,2,4} - v_{1,3,3}}{v_2}, v_{1,1,3} = v_1 v_{1,1,1} + v_{1,1}^2 + v_{1,2,2} v_{1,1,4}$$

$$= \frac{1}{v_2} (v_1^3 v_{1,1,1} + 2 v_1^2 v_{1,1}^2 - v_1 v_{1,2}^2 + v_2^2 v_{1,1,1} + 2 v_2 v_{1,1} v_{1,2} + v_1 v_{1,2,4})$$

$$-v_1 v_{1,3,3} + v_2 v_{1,2,3} \Big\}$$

$$\begin{aligned} &> \text{simplify}([\text{subs}(VSQtmp, v_{1,3}) - \text{subs}(VsS2, v_{1,3}), \text{subs}(VSQtmp, v_{1,4}) - \text{subs}(VsS2, v_{1,4})]) \\ &\quad \left[0, -\frac{v_1 (v_1^2 v_{1,1} - v_2 v_{1,2} + v_{2,4} - v_{3,3})}{v_2} \right] \end{aligned} \quad (1.14)$$

$$\begin{aligned} &> \text{simplify}(\text{subs}(VSQtmp, \text{TotalDiff}(\text{subs}(VSQtmp, v_{1,3}), 1) - \text{subs}(VSQtmp, v_{1,1,3}))); \\ &\quad \text{simplify}(\text{subs}(VSQtmp, \text{TotalDiff}(\text{subs}(VSQtmp, v_{1,4}), 1) - \text{subs}(VSQtmp, v_{1,1,4}))) \\ &\quad \quad \quad 0 \\ &\quad \quad \quad 0 \end{aligned} \quad (1.15)$$

I. We first look for a connection (with torsion) preserving the field of GL2-curves and having α -surfaces totally geodesic + trace of torsion zero.

$$\begin{aligned} &> DGsetup([x, y, t, z], M) : \\ &> vM := [D_x, D_y, D_t, D_z] : dM := [dx, dy, dt, dz] : \\ &> C := \text{Connection}(\text{add}(\text{add}(\text{add}(((a \parallel i) \parallel j) \parallel k) \cdot vM[k] \& t dM[i] \& t dM[j], k = 1..4), j = 1..4), i = 1..4)) : \\ &> \text{DirectionalCovariantDerivative}(D_x, dt, C); \text{DirectionalCovariantDerivative}(D_x, D_t, C) \\ &\quad - (a_{113} dx) - (a_{213} dy) - (a_{313} dt) - (a_{413} dz) \\ &\quad \quad \quad a_{311} D_x + a_{312} D_y + a_{313} D_t + a_{314} D_z \end{aligned} \quad (1.16)$$

$$\begin{aligned} &> Y := \text{evalDG}(D_y - q_{[1]} \cdot D_x) : \\ &\quad T := \text{evalDG}(D_t - (q_{[1]}^2 + v_1) \cdot D_x) : \\ &\quad Z := \text{evalDG}(D_z - (q_{[1]}^3 + 2 v_1 q_{[1]} + v_2) \cdot D_x) : \\ &\quad \omega := \text{evalDG}(dx + q_{[1]} \cdot dy + (q_{[1]}^2 + v_1) \cdot dt + (q_{[1]}^3 + 2 v_1 q_{[1]} + v_2) \cdot dz) : \\ &\quad [Hook(Y, \omega), Hook(T, \omega), Hook(Z, \omega)] \\ &\quad \quad \quad [0, 0, 0] \end{aligned} \quad (1.17)$$

$$\begin{aligned} &> YY := \text{FromJet}(Y, \{v(x, y, t, z), q(x, y, t, z)\}) : TT := \text{FromJet}(T, \{v(x, y, t, z), q(x, y, t, z)\}) : \\ &\quad ZZ := \text{FromJet}(Z, \{v(x, y, t, z), q(x, y, t, z)\}) : \\ &\quad \omega\omega := \text{FromJet}(\omega, \{v(x, y, t, z), q(x, y, t, z)\}) : \\ &> DD := \{YY, ZZ, TT\} : \text{ChrCoe} := \{seq(seq(seq(((a \parallel i) \parallel j) \parallel k), i = 1..4), j = 1..4), k = 1..4)\} : \\ &\quad TGE := \{ \} : \text{for } i \text{ to } 3 \text{ do for } j \text{ to } 3 \text{ do } TGE := \{op(TGE), coeffs(\text{expand}(\text{subs}(VsS2, \text{subs}(Qsyst, \text{ToJet}(Hook(\text{convert}(\text{DirectionalCovariantDerivative}(DD[i], DD[j], C), \\ &\quad \text{'DGvector'}), \omega\omega), \{v(x, y, t, z), q(x, y, t, z)\}, \text{notation} = \text{jetnumbers}))\}), \{q_{[1]}, q_1\}))\} : \\ &\quad \text{od:od:} \\ &\quad [nops(\text{ChrCoe}), nops(TGE)] \\ &\quad \quad \quad [64, 59] \end{aligned} \quad (1.18)$$

$$\begin{aligned} &> Sconn := \text{solve}(TGE, \text{ChrCoe}) : \\ &> \text{FreePar} := \{ \} : \text{for } i \text{ to } nops(Sconn) \text{ do if } lhs(Sconn[i]) = rhs(Sconn[i]) \text{ then } \text{FreePar} \\ &\quad := \{op(\text{FreePar}), lhs(Sconn[i])\} \text{ fi:od: } \text{FreePar} \\ &\quad \quad \quad \{a_{141}, a_{144}, a_{244}, a_{344}, a_{411}, a_{413}, a_{414}, a_{423}, a_{424}, a_{434}\} \end{aligned} \quad (1.19)$$

$$\begin{aligned} &> \text{ChrCoe1} := \{a_{111}, a_{112}, a_{113}, a_{114}, a_{121}, a_{122}, a_{123}, a_{124}, a_{131}, a_{132}, a_{133}, a_{134}, \\ &\quad a_{142}, a_{143}, a_{211}, a_{212}, a_{213}, a_{214}, a_{221}, a_{222}, a_{223}, a_{224}, a_{231}, a_{232}, a_{233}, a_{234}, \end{aligned}$$

```

a241, a242, a243, a311, a312, a313, a314, a321, a322, a323, a324, a331, a332, a333,
a334, a341, a342, a343, a412, a421, a422, a431, a432, a433, a441, a442, a443, a444} :
> Sconn := solve(TGE, ChrCoe1) :
> FreePar := {a141, a144, a244, a344, a411, a413, a414, a423, a424, a434} :
> CC := DGsimplify(subs(Sconn, C)) :
> Quad1 := evalDG(add(add(diff(Eq1, vi,j)·vM[i] &s vM[j], j=i..4), i=1..4)) :
Quad2 := evalDG(add(add(diff(Eq2, vi,j)·vM[i] &s vM[j], j=i..4), i=1..4)) :
Quad3 := evalDG(add(add(diff(Eq3, vi,j)·vM[i] &s vM[j], j=i..4), i=1..4)) :
Quad1v := FromJet(Quad1, {v(x, y, t, z), q(x, y, t, z)}) : Quad2v := FromJet(Quad2, {v(x,
y, t, z), q(x, y, t, z)}) : Quad3v := FromJet(Quad3, {v(x, y, t, z), q(x, y, t, z)}) : Quadv
:= [Quad1v, Quad2v, Quad3v] :
> {seq(ContractIndices(Quad||i, ω &t ω, [[1, 1], [2, 2]]), i=1..3)}
{0} (1.20)

```

Let's check that ω uniquely determines the quadrics Quad||i.

```

> Qua := evalDG(add(add(xi[i, j]·vM[i] &s vM[j], j=i..4), i=1..4)) :
DGsimplify(evalDG(subs({xi[2, 2]=1, xi[2, 3]=0, xi[3, 3]=0},
subs(solve({coeffs(ContractIndices(Qua, ω &t ω, [[1, 1], [2, 2]]), {q[ ]}),
{seq(seq(xi[i, j], j=i..4), i=1..4)}), Qua)) - Quad1));
DGsimplify(evalDG(subs({xi[2, 2]=0, xi[2, 3]=1, xi[3, 3]=0},
subs(solve({coeffs(ContractIndices(Qua, ω &t ω, [[1, 1], [2, 2]]), {q[ ]}),
{seq(seq(xi[i, j], j=i..4), i=1..4)}), Qua)) - Quad2));
DGsimplify(evalDG(subs({xi[2, 2]=0, xi[2, 3]=0, xi[3, 3]=1},
subs(solve({coeffs(ContractIndices(Qua, ω &t ω, [[1, 1], [2, 2]]), {q[ ]}),
{seq(seq(xi[i, j], j=i..4), i=1..4)}), Qua)) - Quad3))
(0 Dx) Dx
(0 Dx) Dx
(0 Dx) Dx (1.21)

```

Explanation: the form ω has coefficients with q^i of degrees from 0 to 3, and the same for $\omega \otimes \omega$ with degrees from 0 to 6 - these are 7 quadrics on TM.

The annihilator in $S^2 TM$ of this 7D subspace of $S^2 T^*M$ has rank 3, consisting of <Quad1, Quad2, Quad3>.

```

> RNE := {} :for i to 4 do for j to 3 do RNE := {op(RNE),
coeffs(expand(numer(simplify(subs(VsS2, subs(Qsyst,
ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadv[j], CC), ω &t ω,
[[1, 1], [2, 2]]), {v(x, y, t, z)}, notation=jetnumbers))))), {q[ ]})} :od:od:
nops(RNE)
36 (1.22)

```

```

> sRNE := solve(RNE, FreePar)
sRNE := {a141=a141, a144=0, a244=0, a344=-v1,1, a411= $\frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2}$ , a413 (1.23)

```

$$= 0, a_{414} = a_{414}, a_{423} = 3 v_{1,1}, a_{424} = a_{424}, a_{434} = a_{434} \}$$

> sRNE := solve(RNE, {a144, a244, a344, a411, a413, a423})

$$sRNE := \left\{ a_{144} = 0, a_{244} = 0, a_{344} = -v_{1,1}, a_{411} = \frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2}, a_{413} = 0, a_{423} = 3 v_{1,1} \right\} \quad (1.24)$$

> CT := DGsimplify(evalDG(subs({a414 = α_1 , a424 = α_2 , a434 = $\alpha_3 + 3 v_{1,1}$, a441 = α_4 }, subs(sRNE, CC)))) :

$$\begin{aligned} & \text{simplify(ToJet(DGinfo(evalDG(convert(DirectionalCovariantDerivative(D_x, FromJet(\omega, \{v(x, y, t, z)\}), CT), 'DGform') \&w FromJet(\omega, \{v(x, y, t, z)\}), "CoefficientSet"), \\ & \{v(x, y, t, z)\}, notation = jetnumbers))} \\ & \left\{ 0, \frac{v_1^2 v_{1,1} - v_2 v_{1,2} + v_{2,4} - v_{3,3}}{v_2}, \frac{q_{[1]} (v_1^2 v_{1,1} - v_2 v_{1,2} + v_{2,4} - v_{3,3})}{v_2}, \right. \\ & \left. \frac{(q_{[1]}^2 + v_1) (v_1^2 v_{1,1} - v_2 v_{1,2} + v_{2,4} - v_{3,3})}{v_2} \right\} \end{aligned} \quad (1.25)$$

$$\begin{aligned} & \text{simplify(map(xx} \rightarrow \text{subs(VsS2, xx),} \\ & \text{seq(ToJet(DGinfo(evalDG(convert(DirectionalCovariantDerivative(vM[i],} \\ & \text{FromJet(\omega, \{v(x, y, t, z)\}), CT), 'DGform') \&w FromJet(\omega, \{v(x, y, t, z)\}),} \\ & \text{"CoefficientSet"), \{v(x, y, t, z)\}, notation = jetnumbers), i = 1..4))} \\ & \{0\} \end{aligned} \quad (1.26)$$

> # TorsionTensor(CT)

$$\begin{aligned} & \left\{ \text{seq(evalDG(CovariantDerivative(vM[j], CT) - subs(\{\alpha_1 = 0, \alpha_2 = 0, \alpha_3 = 0, \alpha_4 = 0\},} \right. \\ & \text{CovariantDerivative(vM[j], CT)) - vM[j] \&t (add(\alpha_i \cdot dM[i], i = 1..4))}, j = 1..4) \} \\ & \{ (0 D_x) dx \} \end{aligned} \quad (1.27)$$

This shows that the connection preserving the GL2-structure and having the Lax pair totally geodesic is determined up to arbitrary 1-form: $\nabla \rightarrow \nabla + Id \cdot \alpha$

$$\begin{aligned} & \text{ContractIndices(TorsionTensor(CT), [[1, 2]])} \\ & - (3 \alpha_1 dx) - (3 \alpha_2 dy) - ((3 \alpha_3 + 2 v_{1,1}) dt) \\ & - \left(\frac{3 (-v_1^2 v_{1,1} + \alpha_4 v_2 - v_{2,4} + v_{3,3})}{v_2} dz \right) \end{aligned} \quad (1.28)$$

> sCTa := solve(DGinfo(%, "CoefficientSet"), { $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ }); CTa
:= DGsimplify(evalDG(subs(sCTa, CT))) :

$$sCTa := \left\{ \alpha_1 = 0, \alpha_2 = 0, \alpha_3 = -\frac{2}{3} v_{1,1}, \alpha_4 = \frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2} \right\} \quad (1.29)$$

> # CTa

```

> RicCTA := RicciTensor(FromJet(CTa, {v(x, y, t, z)})) : RicCTa := ToJet(RicCTA, {v(x, y,
t, z)}), notation = jetnumbers) :
> RicCTaSkew := SymmetrizeIndices(RicCTa, [1, 2], "SkewSymmetric") :
RicCTaSymm := SymmetrizeIndices(RicCTa, [1, 2], "Symmetric") :

```

► Exact Solutions with symmetric Ricci of the canonical GL2-connection

```

[>

```

► Exact Solutions with skew symmetric Ricci of the canonical GL2-connection

```

[>

```

```

> ω0 := ω : ω0b := DGsimplify(evalDG(subs(q[1]=0, ω)));
ω1a := evalDG(1/ε · (subs(q[1]=q[1]+ε, ω) - ω)) : ω1 := evalDG(subs(ε=0, ω1a)) :
ω1b := DGsimplify(evalDG(subs(q[1]=0, ω1)));
ω2a := evalDG(1/ε · (subs(q[1]=q[1]+ε, ω1) - ω1)) : ω2 := evalDG(subs(ε=0,
ω2a)) : ω2b := DGsimplify(evalDG(1/2 · subs(q[1]=0, ω2)));
ω3a := evalDG(1/ε · (subs(q[1]=q[1]+ε, ω2) - ω2)) : ω3 := evalDG(subs(ε=0,
ω3a)) : ω3b := DGsimplify(evalDG(1/6 · subs(q[1]=0, ω3)))
ω0b := dx + v1 dt + v2 dz
ω1b := dy + 2 v1 dz
ω2b := dt
ω3b := dz
(1.30)

```

```

> evalDG(ω - ω0b - ω1b · q[1] - ω2b · q[1]^2 - ω3b · q[1]^3)
0 dx
(1.31)

```

```

> Quaω := evalDG([ω0b &s ω0b, ω0b &s ω1b, 2 ω0b &s ω2b + ω1b &s ω1b, ω0b &s ω3b
+ ω1b &s ω2b, 2 ω1b &s ω3b + ω2b &s ω2b, ω2b &s ω3b, ω3b &s ω3b]) : tmpΩ
:= { } :for i to nops(Quaω) do for j to 3 do tmpΩ := {op(tmpΩ),
ContractIndices(Quad||j, Quaω[i], [[1, 1], [2, 2]])} :od:od: tmpΩ
{0}
(1.32)

```

```

> eliminate(simplify(subs(VsS3, DGinfo(evalDG(RicCTaSymm - add(Λ[i] · Quaω[i], i = 1
..7))), "CoefficientSet")), {seq(Λ[i], i = 1..7)})[2]
{v1,1,1, 2 v1^7 v1,1^2 - v1^6 v1,1 v2,2 + v1^5 v2^2 v1,1,1 + 4 v1^4 v2^2 v1,1^2 + 2 v1^5 v1,1 v2,4 - 2 v1^5 v1,1 v3,3 -
v1^4 v2 v1,1 v2,3 + 3 v1^3 v2^2 v1,1 v2,2 + v1^2 v2^4 v1,1,1 + 2 v1 v2^4 v1,1^2 + v1^4 v2 v2,2,4 - v1^4 v2 v2,3,3 -
(1.33)

```

$$\begin{aligned}
& v_1^4 v_{2,2} v_{2,4} + v_1^4 v_{2,2} v_{3,3} - v_1^2 v_2^2 v_{1,1} v_{2,4} + 2 v_1 v_2^3 v_{1,1} v_{2,3} - v_2^4 v_{1,1} v_{2,2} + v_1^2 v_2^2 v_{2,3,4} - \\
& v_1^2 v_2^2 v_{3,3,3} - v_1^2 v_2 v_{2,3} v_{2,4} + v_1^2 v_2 v_{2,3} v_{3,3} - v_1 v_2^3 v_{2,2,4} + v_1 v_2^3 v_{2,3,3} + v_1 v_2^2 v_{2,2} v_{2,4} \\
& - v_1 v_2^2 v_{2,2} v_{3,3} - v_2^4 v_{2,2,3} + v_2^3 v_{2,4,4} - v_2^3 v_{3,3,4} - 2 v_2^2 v_{2,4}^2 + 3 v_2^2 v_{2,4} v_{3,3} - v_2^2 v_{3,3}^2 \\
& v_1^7 v_2 v_{1,1,1} + 2 v_1^6 v_2 v_{1,1}^2 - v_1^6 v_{1,1} v_{2,3} + 3 v_1^5 v_2 v_{1,1} v_{2,2} + v_1^4 v_2^3 v_{1,1,1} + 4 v_1^3 v_2^3 v_{1,1}^2 - \\
& v_1^4 v_2 v_{1,1} v_{2,4} + 2 v_1^3 v_2^2 v_{1,1} v_{2,3} - 2 v_1^2 v_2^3 v_{1,1} v_{2,2} - 2 v_1 v_2^5 v_{1,1,1} + 2 v_2^5 v_{1,1}^2 + \\
& v_1^4 v_2 v_{2,3,4} - v_1^4 v_2 v_{3,3,3} - v_1^4 v_{2,3} v_{2,4} + v_1^4 v_{2,3} v_{3,3} - v_1^3 v_2^2 v_{2,2,4} + v_1^3 v_2^2 v_{2,3,3} + \\
& v_1^3 v_2 v_{2,2} v_{2,4} - v_1^3 v_2 v_{2,2} v_{3,3} - v_1^2 v_2^3 v_{2,2,3} + 2 v_1 v_2^3 v_{1,1} v_{2,4} - 2 v_1 v_2^3 v_{1,1} v_{3,3} + v_1^2 \\
& v_2^2 v_{2,4,4} - v_1^2 v_2^2 v_{3,3,4} - 2 v_1^2 v_2 v_{2,4}^2 + 3 v_1^2 v_2 v_{2,4} v_{3,3} - v_1^2 v_2 v_{3,3}^2 + v_2^4 v_{2,2,4} - v_2^4 v_{2,3,3} \\
& - v_2^3 v_{2,2} v_{2,4} + v_2^3 v_{2,2} v_{3,3} \}
\end{aligned}$$

Thus the condition that $\text{sym}(\text{Ric}(\nabla))$ belongs to the span of quadrics defining the GL2 structure is non-vacous. It is another differential constraint, giving exact solutions.

But in fact, from what is computed below it follows that this is the same condition as $\text{sym}(\text{Ric})=0$.

> # ToJet(RicciTensor(FromJet(CTa, {v(x, y, t, z)})), {v(x, y, t, z)}, notation = jetnumbers)

> Vt0 := Annihilator([ω0, ω1, ω2])[1]; Vt1 := Annihilator([ω0, ω1])[2]; Vt2
:= Annihilator([ω0])[3]

$$Vt0 := - \left((q[]^3 - 3 v_1 q[] + v_2) D_x \right) + (3 q[]^2 - 2 v_1) D_y - (3 q[] D_t) + D_z$$

$$Vt1 := (q[]^2 - v_1) D_x - (2 q[] D_y) + D_t$$

$$Vt2 := -(q[] D_x) + D_y$$

(1.34)

> Vt01 := ContractIndices(TorsionTensor(CT), evalDG(Vt0 & Vt1), [[2, 1], [3, 2]]):
KrCond := DGinfo(evalDG(convert(Vt01, DGvector) - Hook(convert(Vt01, DGvector), dz)
· Vt0), "CoefficientSet"):

solve(map(Q → coeffs(expand(numer(Q)), {q[]}), KrCond), {α₁, α₂, α₃, α₄})

$$\left\{ \alpha_1 = 0, \alpha_2 = 0, \alpha_3 = -\frac{2}{3} v_{1,1}, \alpha_4 = \frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2} \right\}$$

(1.35)

> sCTa

$$\left\{ \alpha_1 = 0, \alpha_2 = 0, \alpha_3 = -\frac{2}{3} v_{1,1}, \alpha_4 = \frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2} \right\}$$

(1.36)

Thus there exist GL2-connections, with respect to which the null Lax surfaces are totally geodesic. They depend on 4 arbitrary functions.

The parameters can be fixed uniquely by the condition $\text{Trace}(\text{Torsion})=0$. This condition is equivalent to that one of Krynsky.

> simplify(subs(VsS3, ContractIndices(RicCTaSymm, evalDG(Vt0 & Vt0), [[1, 1], [2, 2]])))
0

(1.37)

> simplify(subs(VsS3, ContractIndices(RicCTaSymm, evalDG(Vt0 & Vt1), [[1, 1], [2, 2]])))
0

(1.38)

> simplify(subs(VsS3, ContractIndices(RicCTaSymm, evalDG(2 Vt0 & s Vt2 + 3 Vt1 & Vt1),
[[1, 1], [2, 2]])))

0

(1.39)

$$\begin{aligned} &> \text{seq}(\text{ContractIndices}(\text{Qua}\omega[i], \text{evalDG}(\text{Vt0 \&t Vt0}), [[1, 1], [2, 2]]), i = 1..7) \\ &\quad q_{[]}^6, -3 q_{[]}^5, 15 q_{[]}^4, -10 q_{[]}^3, 15 q_{[]}^2, -3 q_{[]}^1, 1 \end{aligned} \quad (1.40)$$

$$\begin{aligned} &> \text{evalDG}(\text{Vt0 \&t Vt0}) \\ &\left((q_{[]}^3 - 3 v_1 q_{[]} + v_2)^2 D_x \right) D_x - \left((q_{[]}^3 - 3 v_1 q_{[]} + v_2) (3 q_{[]}^2 \right. \\ &\quad - 2 v_1) D_x \right) D_y + \left(3 (q_{[]}^3 - 3 v_1 q_{[]} + v_2) q_{[]} D_x \right) D_t - \left((q_{[]}^3 \right. \\ &\quad - 3 v_1 q_{[]} + v_2) D_x \right) D_z - \left((q_{[]}^3 - 3 v_1 q_{[]} + v_2) (3 q_{[]}^2 - 2 v_1) D_y \right) D_x \\ &\quad + \left((3 q_{[]}^2 - 2 v_1)^2 D_y \right) D_y - \left((3 (3 q_{[]}^2 - 2 v_1) q_{[]} D_y) D_t \right) + \left((3 q_{[]}^2 \right. \\ &\quad - 2 v_1) D_y \right) D_z + \left(3 (q_{[]}^3 - 3 v_1 q_{[]} + v_2) q_{[]} D_t \right) D_x - \left((3 (3 q_{[]}^2 \right. \\ &\quad - 2 v_1) q_{[]} D_t) D_y \right) + (9 q_{[]}^2 D_t) D_t - (3 q_{[]} D_t) D_z - \left((q_{[]}^3 \right. \\ &\quad - 3 v_1 q_{[]} + v_2) D_z \right) D_x + \left((3 q_{[]}^2 - 2 v_1) D_z \right) D_y - \left((3 q_{[]} D_z) D_t \right) \\ &\quad + D_z D_z \end{aligned} \quad (1.41)$$

$$\begin{aligned} &> \text{Vt0} \\ &\quad - \left((q_{[]}^3 - 3 v_1 q_{[]} + v_2) D_x \right) + \left(3 q_{[]}^2 - 2 v_1 \right) D_y - (3 q_{[]} D_t) + D_z \end{aligned} \quad (1.42)$$

$$\begin{aligned} &> \text{QuT} := \text{evalDG}(\text{add}(\text{add}(\text{xi}[i, j] \cdot dM[i] \&s dM[j], j = i..4), i = 1..4)) : \\ &\text{DGsimplify}(\text{evalDG}(\text{subs}(\{\text{xi}[2, 2] = \Lambda_1, \text{xi}[2, 3] = \Lambda_2, \text{xi}[3, 3] = \Lambda_3\}, \\ &\quad \text{subs}(\text{solve}(\{\text{coeffs}(\text{ContractIndices}(\text{QuT}, \text{Vt0 \&t Vt0}), [[1, 1], [2, 2]]), \{q_{[]} \}), \\ &\quad \{\text{seq}(\text{seq}(\text{xi}[i, j], j = i..4), i = 1..4)\}, \text{QuT}))), \\ &\quad - \left(\left(\frac{3 \Lambda_1}{2} dx \right) dt \right) - \left(\left(\frac{9 \Lambda_2}{2} dx \right) dz \right) + (\Lambda_1 dy) dy + \left(\frac{\Lambda_2}{2} dy \right) dt - \left(\left(\left(\frac{5 \Lambda_1 v_1}{2} \right. \right. \right. \\ &\quad \left. \left. + \frac{3 \Lambda_3}{2} \right) dy \right) dz \right) - \left(\left(\left(\frac{3 \Lambda_1}{2} dt \right) dx \right) + \left(\frac{\Lambda_2}{2} dt \right) dy + (\Lambda_3 dt) dt - \left(\left(\left(\frac{7 \Lambda_2 v_1}{2} \right. \right. \right. \right. \\ &\quad \left. \left. + \frac{3 \Lambda_1 v_2}{2} \right) dt \right) dz \right) - \left(\left(\frac{9 \Lambda_2}{2} dz \right) dx \right) - \left(\left(\left(\frac{5 \Lambda_1 v_1}{2} + \frac{3 \Lambda_3}{2} \right) dz \right) dy \right) \\ &\quad - \left(\left(\left(\frac{7 \Lambda_2 v_1}{2} + \frac{3 \Lambda_1 v_2}{2} \right) dz \right) dt \right) - \left((14 \Lambda_1 v_1^2 + 9 \Lambda_2 v_2 + 6 \Lambda_3 v_1) dz \right) dz \end{aligned} \quad (1.43)$$

$$\begin{aligned} &> \text{eliminate}(\text{DGinfo}(\text{DGsimplify}(\text{evalDG}(\text{subs}(\text{VsS3}, \text{subs}(\{\text{xi}[2, 2] = \Lambda_1, \text{xi}[2, 3] = \Lambda_2, \\ &\quad \text{xi}[3, 3] = \Lambda_3\}, \text{subs}(\text{solve}(\{\text{coeffs}(\text{ContractIndices}(\text{QuT}, \text{Vt0 \&t Vt0}), [[1, 1], [2, 2]]), \\ &\quad \{q_{[]} \}), \{\text{seq}(\text{seq}(\text{xi}[i, j], j = i..4), i = 1..4)\}, \text{QuT})) - \text{RicCTaSymm})), \\ &\quad \text{"CoefficientSet"}), \{\Lambda_1, \Lambda_2, \Lambda_3\}) \end{aligned}$$

$$\left[\left\{ \Lambda_1 = v_{1,1,1}, \Lambda_2 = -\frac{1}{v_2 (v_1^6 - v_2^4)} (2 v_1^7 v_{1,1}^2 - v_1^6 v_{1,1} v_{2,2} + v_1^5 v_2^2 v_{1,1,1} + 4 v_1^4 v_2^2 v_{1,1}^2 + 2 \right. \right. \quad (1.44)$$

$$\left. v_1^5 v_{1,1} v_{2,4} - 2 v_1^5 v_{1,1} v_{3,3} - v_1^4 v_2 v_{1,1} v_{2,3} + 3 v_1^3 v_2^2 v_{1,1} v_{2,2} + v_1^2 v_2^4 v_{1,1,1} + 2 v_1 v_2^4 v_{1,1}^2 \right.$$

$$\begin{aligned}
& + v_1^4 v_2 v_{2,2,4} - v_1^4 v_2 v_{2,3,3} - v_1^4 v_{2,2} v_{2,4} + v_1^4 v_{2,2} v_{3,3} - v_1^2 v_2^2 v_{1,1} v_{2,4} + 2 v_1 v_2^3 v_{1,1} v_{2,3} \\
& - v_2^4 v_{1,1} v_{2,2} + v_1^2 v_2^2 v_{2,3,4} - v_1^2 v_2^2 v_{3,3,3} - v_1^2 v_2 v_{2,3} v_{2,4} + v_1^2 v_2 v_{2,3} v_{3,3} - v_1 v_2^3 v_{2,2,4} \\
& + v_1 v_2^3 v_{2,3,3} + v_1 v_2^2 v_{2,2} v_{2,4} - v_1 v_2^2 v_{2,2} v_{3,3} - v_2^4 v_{2,2,3} + v_2^3 v_{2,4,4} - v_2^3 v_{3,3,4} - 2 v_2^2 \\
& v_{2,4}^2 + 3 v_2^2 v_{2,4} v_{3,3} - v_2^2 v_{3,3}^2), \Lambda_3 = -\frac{1}{v_2 (v_1^6 - v_2^4)} (4 v_1^7 v_2 v_{1,1,1} + 2 v_1^6 v_2 v_{1,1}^2 - \\
& v_1^6 v_{1,1} v_{2,3} + 3 v_1^5 v_2 v_{1,1} v_{2,2} + v_1^4 v_2^3 v_{1,1,1} + 4 v_1^3 v_2^3 v_{1,1}^2 - v_1^4 v_2 v_{1,1} v_{2,4} + 2 v_1^3 \\
& v_2^2 v_{1,1} v_{2,3} - 2 v_1^2 v_2^3 v_{1,1} v_{2,2} - 3 v_1 v_2^5 v_{1,1,1} + 2 v_2^5 v_{1,1}^2 + v_1^4 v_2 v_{2,3,4} - v_1^4 v_2 v_{3,3,3} - \\
& v_1^4 v_{2,3} v_{2,4} + v_1^4 v_{2,3} v_{3,3} - v_1^3 v_2^2 v_{2,2,4} + v_1^3 v_2^2 v_{2,3,3} + v_1^3 v_2 v_{2,2} v_{2,4} - v_1^3 v_2 v_{2,2} v_{3,3} - \\
& v_1^2 v_2^3 v_{2,2,3} + 2 v_1 v_2^3 v_{1,1} v_{2,4} - 2 v_1 v_2^3 v_{1,1} v_{3,3} + v_1^2 v_2^2 v_{2,4,4} - v_1^2 v_2^2 v_{3,3,4} - 2 v_1^2 v_2 v_{2,2}^2 \\
& + 3 v_1^2 v_2 v_{2,4} v_{3,3} - v_1^2 v_2 v_{3,3}^2 + v_2^4 v_{2,2,4} - v_2^4 v_{2,3,3} - v_2^3 v_{2,2} v_{2,4} + v_2^3 v_{2,2} v_{3,3}) \}, \{ \}]
\end{aligned}$$

This computation has shown that symmetrized Ricci tensor of the canonical GL2-connection is a section of the bundle of quadrics annihilating the dual (tangent) to the canonical (cotangent) rational normal curve.

>

Below we show that the GL2+totally geodesic conditions are actually equivalent to the hierarchy system of 3 equations in 4D

$$\begin{aligned}
> \text{RNE} := \{ \} : \text{for } i \text{ to } 4 \text{ do for } j \text{ to } 3 \text{ do } \text{RNE} := \{ \text{op}(\text{RNE}), \\
& \text{coeffs}(\text{expand}(\text{numer}(\text{simplify}(\text{subs}(\text{VsS2}, \text{subs}(\text{Qsyst}, \\
& \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(vM[i], \text{Quadv}[j], \text{CT}), \omega \& t \omega, \\
& [[1, 1], [2, 2]]), \{v(x, y, t, z)\}, \text{notation} = \text{jetnumbers}))))) \}, \{q_{[]}\} \} : \text{od:od: RNE} \\
& \{0\} \tag{1.45}
\end{aligned}$$

$$\begin{aligned}
> \text{RNE} := \{ \} : \text{for } i \text{ to } 4 \text{ do for } j \text{ to } 3 \text{ do } \text{RNE} := \{ \text{op}(\text{RNE}), \\
& \text{coeffs}(\text{expand}(\text{numer}(\text{simplify}(\text{subs}(\text{Qsyst}, \\
& \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(vM[i], \text{Quadv}[j], \text{CT}), \omega \& t \omega, \\
& [[1, 1], [2, 2]]), \{v(x, y, t, z)\}, \text{notation} = \text{jetnumbers}))))) \}, \{q_{[]}\} \} : \text{od:od: RNE} \\
& \{0, -v_1 v_{1,1} + v_{1,3} - v_{2,2}, 2 v_1^2 v_{1,1} - 2 v_1 v_{1,3} + 2 v_1 v_{2,2}, -v_1^2 v_{1,1} + v_2 v_{1,2} - v_{2,4} + v_{3,3}, 2 v_1^3 v_{1,1} - 2 v_1 v_2 v_{1,2} + 2 v_1 v_{2,4} - 2 v_1 v_{3,3}, -v_1^3 v_{1,1} - v_2^2 v_{1,1} - v_1 v_{2,4} + v_1 v_{3,3} + v_2 v_{1,4} \\
& - v_2 v_{2,3}, 2 v_1^4 v_{1,1} + 2 v_1 v_2^2 v_{1,1} + 2 v_1^2 v_{2,4} - 2 v_1^2 v_{3,3} - 2 v_1 v_2 v_{1,4} + 2 v_1 v_2 v_{2,3} \} \tag{1.46}
\end{aligned}$$

$$\begin{aligned}
> \text{solve}(\%, \{v_{1,2}, v_{1,3}, v_{1,4}\}) \\
& \left\{ v_{1,2} = \frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2}, v_{1,3} = v_1 v_{1,1} + v_{2,2} v_{1,4} \right. \tag{1.47}
\end{aligned}$$

$$= \frac{v_1^3 v_{1,1} + v_2^2 v_{1,1} + v_1 v_{2,4} - v_1 v_{3,3} + v_2 v_{2,3}}{v_2} \}$$

Finally, let us write the formulas for the Christofel symbols of the canonical connection.

> vM

$$[D_x, D_y, D_t, D_z] \quad (1.48)$$

> VsS2

$$\left\{ v_{1,2} = \frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2}, v_{1,3} = v_1 v_{1,1} + v_{2,2} v_{1,4} \right. \\ \left. = \frac{v_1^3 v_{1,1} + v_2^2 v_{1,1} + v_1 v_{2,4} - v_1 v_{3,3} + v_2 v_{2,3}}{v_2} \right\} \quad (1.49)$$

> $\Gamma_{113} := v_{1,1} :$

$$\Gamma_{114} := v_{1,2} : \Gamma_{214} := 2 v_{1,1} :$$

$$\Gamma_{122} := v_{1,1} :$$

$$\Gamma_{123} := v_{1,2} : \Gamma_{223} := 2 v_{1,1} :$$

$$\Gamma_{124} := 2 v_{2,2} - v_{1,3} : \Gamma_{224} := 2 v_{1,2} : \Gamma_{324} := 3 v_{1,1} :$$

$$\Gamma_{131} := -\frac{2}{3} v_{1,1} :$$

$$\Gamma_{132} := v_{1,2} : \Gamma_{232} := \frac{1}{3} v_{1,1} :$$

$$\Gamma_{133} := 2 v_{2,2} - v_{1,3} : \Gamma_{233} := 2 v_{1,2} : \Gamma_{333} := \frac{4}{3} v_{1,1} :$$

$$\Gamma_{134} := 2 v_{2,3} - v_{1,4} - 2 v_2 v_{1,1} : \Gamma_{234} := 4 v_{2,2} - 2 v_{1,3} : \Gamma_{334} := 3 v_{1,2} : \Gamma_{434} \\ := \frac{7}{3} v_{1,1} :$$

$$\Gamma_{141} := v_{1,2} : \Gamma_{241} := -3 v_{1,1} :$$

$$\Gamma_{142} := 4 v_{1,3} - 3 v_{2,2} : \Gamma_{242} := 2 v_{1,2} : \Gamma_{342} := -2 v_{1,1} :$$

$$\Gamma_{143} := 2 v_{2,3} - v_{1,4} + 3 v_2 v_{1,1} : \Gamma_{243} := 3 v_{1,3} - v_{2,2} : \Gamma_{343} := 3 v_{1,2} : \Gamma_{443} := -v_{1,1} :$$

$$\Gamma_{144} := 3 v_{3,3} - v_1 v_{1,3} - 2 v_{2,4} : \Gamma_{244} := 4 v_{2,3} - 2 v_{1,4} + v_2 v_{1,1} : \Gamma_{344} := 2 v_{1,3} \\ + v_{2,2} : \Gamma_{444} := 4 v_{1,2} :$$

> DirectionalCovariantDerivative(vM[1], vM[1], CTa);

DirectionalCovariantDerivative(vM[1], vM[2], CTa);

evalDG(convert(DirectionalCovariantDerivative(vM[1], vM[3], CTa), DGvector) - Γ_{113} · vM[1]);

simplify(subs(VsS2, evalDG(convert(DirectionalCovariantDerivative(vM[1], vM[4], CTa), DGvector) - (Γ_{114} · vM[1] + Γ_{214} · vM[2]))));

DirectionalCovariantDerivative(vM[2], vM[1], CTa);

simplify(subs(VsS2, evalDG(convert(DirectionalCovariantDerivative(vM[2], vM[2], CTa), DGvector) - Γ_{122} · vM[1])));

simplify(subs(VsS2, evalDG(convert(DirectionalCovariantDerivative(vM[2], vM[3], CTa), DGvector) - (Γ_{123} · vM[1] + Γ_{223} · vM[2]))));

```

DGsimplify(subs(VsS2, evalDG(convert(DirectionalCovariantDerivative(vM[2], vM[4],
    CTa), DGvector) - (I124·vM[1] + I224·vM[2] + I324·vM[3]))));
evalDG(convert(DirectionalCovariantDerivative(vM[3], vM[1], CTa), DGvector) - I131
    ·vM[1]);
simplify(subs(VsS2, evalDG(convert(DirectionalCovariantDerivative(vM[3], vM[2],
    CTa), DGvector) - (I132·vM[1] + I232·vM[2]))));
DGsimplify(subs(VsS2, evalDG(convert(DirectionalCovariantDerivative(vM[3], vM[3],
    CTa), DGvector) - (I133·vM[1] + I233·vM[2] + I333·vM[3]))));
DGsimplify(subs(VsS2, evalDG(convert(DirectionalCovariantDerivative(vM[3], vM[4],
    CTa), DGvector) - (I134·vM[1] + I234·vM[2] + I334·vM[3] + I434·vM[4]))));
;
DGsimplify(subs(VsS2, evalDG(convert(DirectionalCovariantDerivative(vM[4], vM[1],
    CTa), DGvector) - (I141·vM[1] + I241·vM[2]))));
DGsimplify(subs(VsS2, evalDG(convert(DirectionalCovariantDerivative(vM[4], vM[2],
    CTa), DGvector) - (I142·vM[1] + I242·vM[2] + I342·vM[3]))));
DGsimplify(subs(VsS2, evalDG(convert(DirectionalCovariantDerivative(vM[4], vM[3],
    CTa), DGvector) - (I143·vM[1] + I243·vM[2] + I343·vM[3] + I443·vM[4]))));
;
DGsimplify(subs(VsS2, evalDG(convert(DirectionalCovariantDerivative(vM[4], vM[4],
    CTa), DGvector) - (I144·vM[1] + I244·vM[2] + I344·vM[3] + I444·vM[4]))));
;

```

0 D_x

0 D_x

0 D_x

0 D_x

0 D_x

0 D_x

0 D_x

0 D_x

0 D_x

0 D_x

0 D_x

0 D_x

0 D_x

0 D_x

0 D_x

0 D_x

(1.50)

II. Let us check that there exists a unique symmetric connection preserving GL2-structure

```

> ChrSeq := {seq(seq(seq(((a||i)||j)||k), k=1..4), j=i..4), i=1..4)}: nops(ChrSeq);
CsC := Connection(subs({seq(seq(seq(((a||i)||j)||k)=(((a||j)||i)||k), k=1..4), j=1
    ..(i-1)), i=1..4)}, add(add(add(((a||i)||j)||k)·vM[k]&t dM[i]&t dM[j], k=1
    ..4), j=1..4), i=1..4)): TorsionTensor(CsC)

```

$$((0 D_x) dx) dx$$

(1.51)

> RNE := { } :for i to 4 do for j to 3 do RNE := {op(RNE),
coeffs(expand(numer(simplify(subs(VsS2, subs(Qsyst,
ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadv[j], CsC), ω
&t ω, [[1, 1], [2, 2]]), {v(x, y, t, z)}, notation = jetnumbers))))), {q[_]}) :od:od:

> Sss1 := solve(RNE, ChrSeq); nops(Sss1)

$$\begin{aligned} Sss1 := & \left\{ a111=0, a112=0, a113=0, a114=0, a121=0, a122=0, a123=0, a124=0, \right. \\ & a131=v_{1,1}, a132=0, a133=0, a134=0, a141=\frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2}, a142=2 v_{1,1}, \\ & a143=0, a144=0, a221=\frac{4}{9} v_{1,1}, a222=0, a223=0, a224=0, a231 \\ & =\frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2}, a232=\frac{8}{9} v_{1,1}, a233=0, a234=0, a241=-\frac{4}{9} v_1 v_{1,1} + v_{2,2}, \\ & a242=\frac{2(v_1^2 v_{1,1} + v_{2,4} - v_{3,3})}{v_2}, a243=\frac{4}{3} v_{1,1}, a244=0, a331=\frac{11}{9} v_1 v_{1,1} + v_{2,2}, \\ & a332=\frac{2(v_1^2 v_{1,1} + v_{2,4} - v_{3,3})}{v_2}, a333=\frac{7}{9} v_{1,1}, a334=0, a341= \\ & -\frac{1}{3} \frac{3 v_1^3 v_{1,1} - v_2^2 v_{1,1} + 3 v_1 v_{2,4} - 3 v_1 v_{3,3} - 3 v_2 v_{2,3}}{v_2}, a342=\frac{22}{9} v_1 v_{1,1} + 2 v_{2,2}, \\ & a343=\frac{3(v_1^2 v_{1,1} + v_{2,4} - v_{3,3})}{v_2}, a344=\frac{2}{3} v_{1,1}, a441=-\frac{59}{9} v_1^2 v_{1,1} - v_1 v_{2,2} - 2 v_{2,4} \\ & + 3 v_{3,3}, a442=-\frac{2(v_1^3 v_{1,1} - 2 v_2^2 v_{1,1} + v_1 v_{2,4} - v_1 v_{3,3} - v_2 v_{2,3})}{v_2}, a443 \\ & =\frac{16}{3} v_1 v_{1,1} + 3 v_{2,2}, a444=\frac{4(v_1^2 v_{1,1} + v_{2,4} - v_{3,3})}{v_2} \left. \right\} \end{aligned}$$

40

(1.52)

> CsS := DGsimplify(evalDG(subs(Sss1, CsC)))

$$\begin{aligned} CsS := & - \left(\left(\left(\frac{3 v_1^3 v_{1,1} - v_2^2 v_{1,1} + 3 v_1 v_{2,4} - 3 v_1 v_{3,3} - 3 v_2 v_{2,3}}{3 v_2} D_x \right) dt \right) dz \right) \\ & + \left(\left(\left(\frac{11 v_1 v_{1,1}}{9} + v_{2,2} \right) D_x \right) dt \right) dt - \left(\left(\left(\left(\frac{4 v_1 v_{1,1}}{9} - v_{2,2} \right) D_x \right) dz \right) dy \right) \\ & + \left(\left(\left(\frac{22 v_1 v_{1,1}}{9} + 2 v_{2,2} \right) D_y \right) dt \right) dz + \left(\left(\left(\frac{3(v_1^2 v_{1,1} + v_{2,4} - v_{3,3})}{v_2} D_t \right) dz \right) dt \right) \end{aligned}$$

(1.53)

$$\begin{aligned}
& + \left(\left(\frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2} D_x \right) dz \right) dx + \left(\left(\frac{4 v_{1,1}}{3} D_t \right) dy \right) dz \\
& - \left(\left(\left(\frac{3 v_1^3 v_{1,1} - v_2^2 v_{1,1} + 3 v_1 v_{2,4} - 3 v_1 v_{3,3} - 3 v_2 v_{2,3}}{3 v_2} D_x \right) dz \right) dt \right) \\
& + \left(\left(\frac{2 v_{1,1}}{3} D_z \right) dz \right) dt + \left(\left(\frac{4 v_{1,1}}{9} D_x \right) dy \right) dy + \left(\left(\frac{7 v_{1,1}}{9} D_t \right) dt \right) dt \\
& + \left(\left(\frac{2 (v_1^2 v_{1,1} + v_{2,4} - v_{3,3})}{v_2} D_y \right) dy \right) dz + \left(\left(\frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2} D_x \right) dy \right) dt \\
& - \left(\left(\left(\frac{2 (v_1^3 v_{1,1} - 2 v_2^2 v_{1,1} + v_1 v_{2,4} - v_1 v_{3,3} - v_2 v_{2,3})}{v_2} D_y \right) dz \right) dz \right) \\
& + ((2 v_{1,1} D_y) dx) dz - \left(\left(\left(\frac{59}{9} v_1^2 v_{1,1} + v_1 v_{2,2} + 2 v_{2,4} - 3 v_{3,3} \right) D_x \right) dz \right) dz \\
& + \left(\left(\frac{4 (v_1^2 v_{1,1} + v_{2,4} - v_{3,3})}{v_2} D_z \right) dz \right) dz + \left(\left(\frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2} D_x \right) dx \right) dz \\
& + ((v_{1,1} D_x) dt) dx + ((v_{1,1} D_x) dx) dt + \left(\left(\left(\frac{22 v_1 v_{1,1}}{9} + 2 v_{2,2} \right) D_y \right) dz \right) dt \\
& + \left(\left(\frac{v_1^2 v_{1,1} + v_{2,4} - v_{3,3}}{v_2} D_x \right) dt \right) dy + \left(\left(\frac{8 v_{1,1}}{9} D_y \right) dy \right) dt \\
& + \left(\left(\frac{8 v_{1,1}}{9} D_y \right) dt \right) dy + \left(\left(\frac{2 (v_1^2 v_{1,1} + v_{2,4} - v_{3,3})}{v_2} D_y \right) dt \right) dt \\
& + \left(\left(\frac{4 v_{1,1}}{3} D_t \right) dz \right) dy + \left(\left(\frac{2 (v_1^2 v_{1,1} + v_{2,4} - v_{3,3})}{v_2} D_y \right) dz \right) dy \\
& + \left(\left(\frac{3 (v_1^2 v_{1,1} + v_{2,4} - v_{3,3})}{v_2} D_t \right) dt \right) dz + ((2 v_{1,1} D_y) dz) dx \\
& - \left(\left(\left(\left(\frac{4 v_1 v_{1,1}}{9} - v_{2,2} \right) D_x \right) dy \right) dz \right) + \left(\left(\frac{2 v_{1,1}}{3} D_z \right) dt \right) dz + \left(\left(\left(\frac{16 v_1 v_{1,1}}{3} \right. \right. \right. \\
& \left. \left. \left. + 3 v_{2,2} \right) D_t \right) dz \right) dz
\end{aligned}$$

> seq(seq(simplify(subs(VsS2,
 ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadv[j], CsS), ω
 &t ω, [[1, 1], [2, 2]]), {v(x, y, t, z)}, notation=jetnumbers))), i=1..4), j=1..3)
 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0

(1.54)

> # simplify(subs(VsS2, subs(Qsyst, ToJet(evalDG(DirectionalCovariantDerivative(vM[1],
 ωω, CsS) &w ωω), {v(x, y, t, z), q(x, y, t, z)}, notation=jetnumbers))))

> TorsionTensor(CsS)

$$((0 D_x) dx) dx$$

(1.55)

$$\begin{aligned}
& \text{DirectionalCovariantDerivative}(vM[1], vM[1], CsS); \\
& \text{DirectionalCovariantDerivative}(vM[1], vM[2], CsS); \\
& \text{evalDG}(\text{convert}(\text{DirectionalCovariantDerivative}(vM[1], vM[3], CsS), 'DGvector') - v[1, 1] \\
& \quad \cdot vM[1]); \\
& \text{simplify}(\text{subs}(VsS2, \text{evalDG}(\text{convert}(\text{DirectionalCovariantDerivative}(vM[1], vM[4], CsS), \\
& \quad 'DGvector') - (v[1, 2] \cdot vM[1] + 2 v[1, 1] \cdot vM[2])))) \\
& \quad 0 D_x \\
& \quad 0 D_x \\
& \quad 0 D_x \\
& \quad 0 D_x
\end{aligned} \tag{1.56}$$

$$\begin{aligned}
& \text{evalDG}\left(\text{convert}(\text{DirectionalCovariantDerivative}(vM[2], vM[2], CsS), 'DGvector')\right. \\
& \quad \left. - \frac{4}{9} v[1, 1] \cdot vM[1]\right); \\
& \text{simplify}\left(\text{subs}\left(VsS2, \text{evalDG}\left(\text{convert}(\text{DirectionalCovariantDerivative}(vM[2], vM[3], CsS),\right.\right.\right. \\
& \quad \left.\left.\left.'DGvector') - \left(v[1, 2] \cdot vM[1] + \frac{8}{9} v[1, 1] \cdot vM[2]\right)\right)\right)\right); \\
& \text{DGsimplify}\left(\text{simplify}\left(\text{subs}\left(VsS2, \text{evalDG}\left(\text{convert}(\text{DirectionalCovariantDerivative}(vM[2],\right.\right.\right.\right. \\
& \quad \left.\left.\left.vM[4], CsS), 'DGvector') - \left(\frac{1}{9} (13 v[2, 2] - 4 v[1, 3]) \cdot vM[1] + 2 v[1, 2] \cdot vM[2]\right.\right.\right.\right. \\
& \quad \left.\left.\left.+ \frac{4}{3} v[1, 1] \cdot vM[3]\right)\right)\right)\right) \\
& \quad 0 D_x \\
& \quad 0 D_x \\
& \quad 0 D_x
\end{aligned} \tag{1.57}$$

$$\begin{aligned}
& \text{DGsimplify}\left(\text{simplify}\left(\text{subs}\left(VsS2, \text{evalDG}\left(\text{convert}(\text{DirectionalCovariantDerivative}(vM[3],\right.\right.\right.\right. \\
& \quad \left.\left.\left.vM[3], CsS), 'DGvector') - \left(\frac{1}{9} (11 v[1, 3] - 2 v[2, 2]) \cdot vM[1] + 2 v[1, 2] \cdot vM[2]\right.\right.\right.\right. \\
& \quad \left.\left.\left.+ \frac{7}{9} v[1, 1] \cdot vM[3]\right)\right)\right)\right); \\
& \text{DGsimplify}\left(\text{simplify}\left(\text{subs}\left(VsS2, \text{evalDG}\left(\text{convert}(\text{DirectionalCovariantDerivative}(vM[3],\right.\right.\right.\right. \\
& \quad \left.\left.\left.vM[4], CsS), 'DGvector') - \left(\left(2 v[2, 3] - v[1, 4] + \frac{4}{3} v[2] \cdot v[1, 1]\right) \cdot vM[1]\right.\right.\right.\right. \\
& \quad \left.\left.\left.+ \frac{2}{9} (11 v[1, 3] - 2 v[2, 2]) \cdot vM[2] + 3 v[1, 2] \cdot vM[3] + \frac{2}{3} v[1, 1] \cdot vM[4]\right)\right)\right)\right) \\
& \quad 0 D_x \\
& \quad 0 D_x
\end{aligned} \tag{1.58}$$

$$\begin{aligned}
& \text{DGsimplify}\left(\text{simplify}\left(\text{subs}\left(VsS2, \text{evalDG}\left(\text{convert}(\text{DirectionalCovariantDerivative}(vM[4],\right.\right.\right.\right. \\
& \quad \left.\left.\left.vM[4], CsS), 'DGvector') - \left(\left(v[3, 3] + \frac{1}{9} v[1] (32 v[2, 2] - 41 v[1, 3]) - 2 v[2] \cdot v[1,\right.\right.\right.\right. \\
& \quad \left.\left.\left.2\right)\right) \cdot vM[1] + (4 v[2, 3] - 2 v[1, 4] + 6 v[2] v[1, 1]) \cdot vM[2] + \frac{1}{3} (16 v[1, 3] \right.
\end{aligned}$$

==>

```

> CsC := Connection(subs( {seq(seq(seq( (( (a || i) || j) || k) = (( (a || j) || i) || k), k = 1 .. 4), j = 1
.. (i - 1)), i = 1 .. 4)}, add(add(add( (( (a || i) || j) || k) · vM[k] &t dM[i] &t dM[j], k = 1
.. 4), j = 1 .. 4), i = 1 .. 4)):

```

$$TGE := \{ \} : \text{for } i \text{ to } 3 \text{ do for } j \text{ to } 3 \text{ do } TGE := \{ op(TGE), coeffs(expand(subs(VsS2, \\ subs(Qsyst, ToJet(Hook(convert(DirectionalCovariantDerivative(DD[i], DD[j], CsC), \\ 'DGvector'), \omega\omega), \{ v(x, y, t, z), q(x, y, t, z) \}, notation = jetnumbers))), \{ q_{\lceil}, q_{\lceil} \}) \} :$$

(1.60)

(1.61)

(1.62)

(1.63)

(1.64)

$$-v_{1,2}$$

$$\begin{array}{c} -\frac{1}{2} v_{1,1} \\ 0 \\ 0 \end{array}$$

(1.65)

> simplify(subs(VsS2b, subs(Sss20, a221))); subs(Sss20, a222); subs(Sss20, a223);
subs(Sss20, a224)

$$\begin{array}{c} v_{1,1} \\ 0 \\ 0 \\ 0 \end{array}$$

(1.66)

> simplify(subs(VsS2b, subs(Sss20, a231))); subs(Sss20, a232); subs(Sss20, a233);
subs(Sss20, a234)

$$\begin{array}{c} v_{1,2} \\ \frac{1}{2} v_{1,1} \\ 0 \\ 0 \end{array}$$

(1.67)

> simplify(subs(VsS2, subs(Sss20, a241))); subs(Sss20, a242); subs(Sss20, a243);
subs(Sss20, a244)

$$\begin{array}{c} \frac{3}{2} v_1 v_{1,1} + v_{2,2} \\ 0 \\ \frac{1}{2} v_{1,1} \\ 0 \end{array}$$

(1.68)

> simplify(subs(VsS2, subs(Sss20, a331))); simplify(subs(VsS2b, subs(Sss20, a332)));
subs(Sss20, a333); subs(Sss20, a334)

$$\begin{array}{c} -v_1 v_{1,1} + v_{2,2} \\ 2 v_{1,2} \\ 0 \\ 0 \end{array}$$

(1.69)

> simplify(subs(VsS2b, subs(Sss20, a341))); simplify(subs(VsS2, subs(Sss20, a342)));
simplify(subs(VsS2b, subs(Sss20, a343))); subs(Sss20, a344)

$$\begin{array}{c} -2 v_1 v_{1,2} - \frac{3}{2} v_2 v_{1,1} + v_{1,4} \\ \frac{1}{2} v_1 v_{1,1} + 2 v_{2,2} \\ v_{1,2} \\ 0 \end{array}$$

(1.70)

> simplify(subs(VsS2, subs(Sss20, a441))); simplify(subs(VsS2b, subs(Sss20, a442)));
simplify(subs(VsS2b, subs(Sss20, a443))); subs(Sss20, a444)

$$-v_1^2 v_{1,1} - v_1 v_{2,2} - 2 v_{2,4} + 3 v_{3,3}$$

$$\begin{aligned} & -4 v_1 v_{1,2} - 3 v_2 v_{1,1} + 2 v_{1,4} \\ & -v_1 v_{1,1} + 3 v_{1,3} \\ & 0 \end{aligned}$$

(1.71)

$$\text{> simplify}\left(\text{subs}\left(\text{VsS2}, \left(-v_1^2 v_{1,1} - v_1 v_{2,2} - 2 v_{2,4} + 3 v_{3,3}\right) - \left(2 v_{3,3} - v_2 v_{1,2} - v_{2,4} - v_1 v_{2,2}\right)\right)\right)$$

$$0$$

(1.72)

Veronese web hierarchy 4D

$$\text{> DGsetup}([x1, x2, x3, x4], [u], Ju, 3) :$$

$$\text{> X2f} := f \rightarrow \text{TotalDiff}(f, 2) - \frac{(q - a_1) u_2}{(q - a_2) u_1} \cdot \text{TotalDiff}(f, 1) :$$

$$\text{X3f} := f \rightarrow \text{TotalDiff}(f, 3) - \frac{(q - a_1) u_3}{(q - a_3) u_1} \cdot \text{TotalDiff}(f, 1) :$$

$$\text{X4f} := f \rightarrow \text{TotalDiff}(f, 4) - \frac{(q - a_1) u_4}{(q - a_4) u_1} \cdot \text{TotalDiff}(f, 1) :$$

$$\text{> numer(X2f(X3f(x1))) - X3f(X2f(x1)); numer(X2f(X4f(x1))) - X4f(X2f(x1));}$$

$$\begin{aligned} & \text{numer(X3f(X4f(x1))) - X4f(X3f(x1))} \\ & - (q - a_1) u_1 (a_1 u_2 u_{1,3} - a_1 u_3 u_{1,2} - a_2 u_1 u_{2,3} + a_2 u_3 u_{1,2} + a_3 u_1 u_{2,3} - a_3 u_2 u_{1,3}) \\ & - (q - a_1) u_1 (a_1 u_2 u_{1,4} - a_1 u_4 u_{1,2} - a_2 u_1 u_{2,4} + a_2 u_4 u_{1,2} + a_4 u_1 u_{2,4} - a_4 u_2 u_{1,4}) \\ & - (q - a_1) u_1 (a_1 u_3 u_{1,4} - a_1 u_4 u_{1,3} - a_3 u_1 u_{3,4} + a_3 u_4 u_{1,3} + a_4 u_1 u_{3,4} - a_4 u_3 u_{1,4}) \end{aligned}$$

(2.1)

$$\text{> Eq1} := \text{collect}(a_1 u_2 u_{1,3} - a_1 u_3 u_{1,2} - a_2 u_1 u_{2,3} + a_2 u_3 u_{1,2} + a_3 u_1 u_{2,3} - a_3 u_2 u_{1,3},$$

$$\{u_{1,2}, u_{1,3}, u_{1,4}, u_{2,3}, u_{2,4}, u_{3,4}\};$$

$$\text{Eq2} := \text{collect}(a_1 u_2 u_{1,4} - a_1 u_4 u_{1,2} - a_2 u_1 u_{2,4} + a_2 u_4 u_{1,2} + a_4 u_1 u_{2,4} - a_4 u_2 u_{1,4},$$

$$\{u_{1,2}, u_{1,3}, u_{1,4}, u_{2,3}, u_{2,4}, u_{3,4}\};$$

$$\text{Eq3} := \text{collect}(a_1 u_3 u_{1,4} - a_1 u_4 u_{1,3} - a_3 u_1 u_{3,4} + a_3 u_4 u_{1,3} + a_4 u_1 u_{3,4} - a_4 u_3 u_{1,4},$$

$$\{u_{1,2}, u_{1,3}, u_{1,4}, u_{2,3}, u_{2,4}, u_{3,4}\})$$

$$\text{Eq1} := (-a_1 u_3 + a_2 u_3) u_{1,2} + (a_1 u_2 - a_3 u_2) u_{1,3} + (-a_2 u_1 + a_3 u_1) u_{2,3}$$

$$\text{Eq2} := (-a_1 u_4 + a_2 u_4) u_{1,2} + (a_1 u_2 - a_4 u_2) u_{1,4} + (-a_2 u_1 + a_4 u_1) u_{2,4}$$

$$\text{Eq3} := (-a_1 u_4 + a_3 u_4) u_{1,3} + (a_1 u_3 - a_4 u_3) u_{1,4} + (-a_3 u_1 + a_4 u_1) u_{3,4}$$

(2.2)

$$\text{> collect}\left(\text{simplify}\left(\frac{u_4 \cdot \text{Eq1} - u_3 \cdot \text{Eq2} + u_2 \cdot \text{Eq3}}{u_1}\right), \{u_{1,2}, u_{1,3}, u_{1,4}, u_{2,3}, u_{2,4}, u_{3,4}\}\right)$$

$$(-a_2 u_4 + a_3 u_4) u_{2,3} + (a_2 u_3 - a_4 u_3) u_{2,4} + (-a_3 u_2 + a_4 u_2) u_{3,4}$$

(2.3)

$$\text{> VsS2} := \text{solve}(\{\text{Eq1}, \text{Eq2}, \text{Eq3}\}, \{u_{2,3}, u_{2,4}, u_{3,4}\})$$

$$\text{VsS2} := \left\{ u_{2,3} = \frac{a_1 u_2 u_{1,3} - a_1 u_3 u_{1,2} + a_2 u_3 u_{1,2} - a_3 u_2 u_{1,3}}{u_1 (a_2 - a_3)}, u_{2,4} \right.$$

(2.4)

$$= \frac{a_1 u_2 u_{1,4} - a_1 u_4 u_{1,2} + a_2 u_4 u_{1,2} - a_4 u_2 u_{1,4}}{u_1 (a_2 - a_4)}, u_{3,4}$$

$$= \frac{a_1 u_3 u_{1,4} - a_1 u_4 u_{1,3} + a_3 u_4 u_{1,3} - a_4 u_3 u_{1,4}}{u_1 (a_3 - a_4)} \Bigg\}$$

> VsS2a := solve({Eq1, Eq2, Eq3}, {u_{1,3}, u_{1,4}, u_{3,4}}); VsS2b := solve({Eq1, Eq2, Eq3}, {u_{1,2}, u_{1,4}, u_{2,4}}); VsS2c := solve({Eq1, Eq2, Eq3}, {u_{1,2}, u_{1,3}, u_{2,3}})

$$\text{VsS2a} := \left\{ u_{1,3} = \frac{a_1 u_3 u_{1,2} + a_2 u_1 u_{2,3} - a_2 u_3 u_{1,2} - a_3 u_1 u_{2,3}}{u_2 (a_1 - a_3)}, u_{1,4} \right.$$

$$= \frac{a_1 u_4 u_{1,2} + a_2 u_1 u_{2,4} - a_2 u_4 u_{1,2} - a_4 u_1 u_{2,4}}{u_2 (a_1 - a_4)}, u_{3,4}$$

$$= \frac{a_2 u_3 u_{2,4} - a_2 u_4 u_{2,3} + a_3 u_4 u_{2,3} - a_4 u_3 u_{2,4}}{u_2 (a_3 - a_4)} \Bigg\}$$

$$\text{VsS2b} := \left\{ u_{1,2} = \frac{a_1 u_2 u_{1,3} - a_2 u_1 u_{2,3} + a_3 u_1 u_{2,3} - a_3 u_2 u_{1,3}}{u_3 (a_1 - a_2)}, u_{1,4} \right.$$

$$= \frac{a_1 u_4 u_{1,3} + a_3 u_1 u_{3,4} - a_3 u_4 u_{1,3} - a_4 u_1 u_{3,4}}{u_3 (a_1 - a_4)}, u_{2,4}$$

$$= \frac{a_2 u_4 u_{2,3} + a_3 u_2 u_{3,4} - a_3 u_4 u_{2,3} - a_4 u_2 u_{3,4}}{u_3 (a_2 - a_4)} \Bigg\}$$

$$\text{VsS2c} := \left\{ u_{1,2} = \frac{a_1 u_2 u_{1,4} - a_2 u_1 u_{2,4} + a_4 u_1 u_{2,4} - a_4 u_2 u_{1,4}}{(a_1 - a_2) u_4}, u_{1,3} \right. \quad (2.5)$$

$$= \frac{a_1 u_3 u_{1,4} - a_3 u_1 u_{3,4} + a_4 u_1 u_{3,4} - a_4 u_3 u_{1,4}}{u_4 (a_1 - a_3)}, u_{2,3}$$

$$= \frac{a_2 u_3 u_{2,4} - a_3 u_2 u_{3,4} + a_4 u_2 u_{3,4} - a_4 u_3 u_{2,4}}{u_4 (a_2 - a_3)} \Bigg\}$$

> simplify(subs(VsS2, u₄·TotalDiff(Eq1, 1) - u₁·TotalDiff(Eq1, 4) - u₃·TotalDiff(Eq2, 1) + u₁·TotalDiff(Eq2, 3) + u₂·TotalDiff(Eq3, 1) - u₁·TotalDiff(Eq3, 2)));
simplify(subs(VsS2, (a₁ - a₄)·TotalDiff(Eq1, 4) + (a₃ - a₁)·TotalDiff(Eq2, 3) + (a₁ - a₂)·TotalDiff(Eq3, 2)))

0
0

(2.6)

> Vss3 := {Eq1, Eq2, Eq3} :for i to 4 do for j to 3 do Vss3 := {op(Vss3), TotalDiff(Eq|| j, i) } :od:od:
tmp := eliminate(Vss3, {u_{2,3}, u_{2,4}, u_{3,4}, u_{1,2,3}, u_{1,2,4}, u_{1,3,4}, u_{2,2,3}, u_{2,2,4}, u_{2,3,4}, u_{2,3,3}, u_{3,3,4}, u_{3,4,4}, u_{4,4,4}}) : [nops(tmp[1]), nops(tmp[2])];

[13, 1]

(2.7)

> VsS3 := solve(Vss3, {u_{2,3}, u_{2,4}, u_{3,4}, u_{1,2,3}, u_{1,2,4}, u_{1,3,4}, u_{2,2,3}, u_{2,2,4}, u_{2,3,3}, u_{2,3,4}, u_{2,4,4}, u_{3,3,4}, u_{3,4,4}, u_{4,4,4}}) :

> nops(%)

14

(2.8)

> # ... #

> SymbE1 := add(add(diff(Eq1, u_{i,j})·p_i·p_j, j=i..4), i=1..4);

SymbE2 := add(add(diff(Eq2, u_{i,j})·p_i·p_j, j=i..4), i=1..4);

SymbE3 := add(add(diff(Eq3, u_{i,j})·p_i·p_j, j=i..4), i=1..4);

SymbE1 := (-a₁ u₃ + a₂ u₃) p₁ p₂ + (a₁ u₂ - a₃ u₂) p₁ p₃ + (-a₂ u₁ + a₃ u₁) p₂ p₃

SymbE2 := (-a₁ u₄ + a₂ u₄) p₁ p₂ + (a₁ u₂ - a₄ u₂) p₁ p₄ + (-a₂ u₁ + a₄ u₁) p₂ p₄

SymbE3 := (-a₁ u₄ + a₃ u₄) p₁ p₃ + (a₁ u₃ - a₄ u₃) p₁ p₄ + (-a₃ u₁ + a₄ u₁) p₃ p₄ (2.9)

> collect(expand($\frac{p_4 \cdot \text{SymbE1} - p_3 \cdot \text{SymbE2} + p_2 \cdot \text{SymbE3}}{p_1}$), {p₁, p₂, p₃, p₄})

((-a₂ u₄ + a₃ u₄) p₃ + (a₂ u₃ - a₄ u₃) p₄) p₂ + (-a₃ u₂ + a₄ u₂) p₄ p₃ (2.10)

> collect(expand($\frac{u_4 \cdot \text{SymbE1} - u_3 \cdot \text{SymbE2} + u_2 \cdot \text{SymbE3}}{u_1}$), {p₁, p₂, p₃, p₄})

((-a₂ u₄ + a₃ u₄) p₃ + (a₂ u₃ - a₄ u₃) p₄) p₂ + (-a₃ u₂ + a₄ u₂) p₄ p₃ (2.11)

> tmp := solve({coeffs(expand((a₁·p₁ + a₂·p₂ + a₃·p₃ + a₄·p₄)·SymbE1 + (b₁·p₁ + b₂·p₂ + b₃·p₃ + b₄·p₄)·SymbE2 + (c₁·p₁ + c₂·p₂ + c₃·p₃ + c₄·p₄)·SymbE3), {p₁, p₂, p₃, p₄}), {a₁, a₂, a₃, a₄, b₁, b₂, b₃, b₄, c₁, c₂, c₃, c₄})

tmp := {a₁ = $\frac{c_1 u_4}{u_2}$, a₂ = 0, a₃ = 0, a₄ = $\frac{c_1 a_2 u_1 - c_1 a_4 u_1 + c_2 a_1 u_2 - c_2 a_4 u_2}{u_2 (a_1 - a_2)}$, b₁ = (2.12)

- $\frac{c_1 u_3}{u_2}$, b₂ = 0, b₃ = - $\frac{c_1 a_2 u_1 - c_1 a_3 u_1 + c_2 a_1 u_2 - c_2 a_3 u_2}{u_2 (a_1 - a_2)}$, b₄ = 0, c₁ = c₁, c₂

= c₂, c₃ = 0, c₄ = 0}

> simplify((u₄ p₁ - u₁ p₄)·SymbE1 - (u₃ p₁ - u₁ p₃)·SymbE2 + (u₂ p₁ - u₁ p₂)·SymbE3)

0

(2.13)

> simplify((a₁ - a₄) p₄·SymbE1 + (a₃ - a₁) p₃·SymbE2 + (a₁ - a₂) p₂·SymbE3)

0

(2.14)

This shows that the Veronese system is compatible and the Hilbert polynomial is dim(g[k])=1+3k, whence the general solution depends on 3 functions of 1 variable.

This is also what is expected from the characteristic variety (rational normal curve in our case) that has order 3 and dimension 1.

>

> DGsetup([x1, x2, x3, x4], M) :

> vM := [D_x1, D_x2, D_x3, D_x4] : dM := [dx1, dx2, dx3, dx4] :

> C := Connection(add(add(add(((a||i)||j)||k)·vM[k] &t dM[i] &t dM[j], k=1..4), j=1..4), i=1..4)):

> X2 := evalDG($D_{x2} - \frac{(q - a_1) u_2}{(q - a_2) u_1} \cdot D_{x1}$):

$$\begin{aligned}
X3 &:= \text{evalDG} \left(D_{x3} - \frac{(q - a_1) u_3}{(q - a_3) u_1} \cdot D_{x1} \right) : \\
X4 &:= \text{evalDG} \left(D_{x4} - \frac{(q - a_1) u_4}{(q - a_4) u_1} \cdot D_{x1} \right) : \\
\omega &:= \text{evalDG} \left(\frac{u_1}{q - a_1} \cdot dx1 + \frac{u_2}{q - a_2} \cdot dx2 + \frac{u_3}{q - a_3} \cdot dx3 + \frac{u_4}{q - a_4} \cdot dx4 \right) : \\
&[\text{Hook}(X2, \omega), \text{Hook}(X3, \omega), \text{Hook}(X4, \omega)] \\
&[0, 0, 0] \tag{2.15}
\end{aligned}$$

$$\begin{aligned}
> XX2 &:= \text{FromJet}(X2, \{u(x1, x2, x3, x4)\}) : XX3 := \text{FromJet}(X3, \{u(x1, x2, x3, x4)\}) : XX4 \\
&:= \text{FromJet}(X4, \{u(x1, x2, x3, x4)\}) : \\
\omega\omega &:= \text{FromJet}(\omega, \{u(x1, x2, x3, x4)\}) : \\
> DD &:= \{XX2, XX3, XX4\} : \text{ChrCoe} := \{ \text{seq}(\text{seq}(\text{seq}(((a \parallel i) \parallel j) \parallel k), i = 1 \dots 4), j = 1 \dots 4), k \\
&= 1 \dots 4) \} : \\
TGE &:= \{ \} : \text{for } i \text{ to } 3 \text{ do for } j \text{ to } 3 \text{ do } TGE := \{ \text{op}(TGE), \text{coeffs}(\text{expand}(\text{numer}(\text{subs}(\text{VsS2}, \\
&\text{ToJet}(\text{Hook}(\text{convert}(\text{DirectionalCovariantDerivative}(DD[i], DD[j], C), 'DGvector'), \\
&\omega\omega), \{u(x1, x2, x3, x4)\}, \text{notation} = \text{jetnumbers})) \}, \{q\}) \} : \text{od:od:} \\
&[\text{nops}(\text{ChrCoe}), \text{nops}(TGE)] \\
&[64, 54] \tag{2.16}
\end{aligned}$$

$$\begin{aligned}
> Ck &:= \text{Connection} \left(\text{evalDG} \left(\text{add} \left(\text{add} \left(\left(\frac{u_{i,j}}{u_i} + \alpha_j \right) \cdot vM[i] \&t dM[i] \&t dM[j], j = i \dots 4 \right), i \right. \right. \right. \\
&= 1 \dots 4 \left. \right) + \text{add} \left(\text{add} \left(\left(\frac{u_{j,i}}{u_i} + \alpha_j \right) \cdot vM[i] \&t dM[i] \&t dM[j], j = 1 \dots (i - 1) \right), i = 1 \right. \\
&\left. \left. \left. \dots 4 \right) \right) \right) : CK := \text{FromJet}(Ck, \{u(x1, x2, x3, x4)\}) : \\
> \text{ToJet}(\text{DirectionalCovariantDerivative}(vM[1], vM[4], CK), \{u(x1, x2, x3, x4)\}, \text{notation} \\
&= \text{jetnumbers}) \\
&\frac{\alpha_1 u_4 + u_{1,4}}{u_4} D_{x4} \tag{2.17}
\end{aligned}$$

$$\begin{aligned}
> \{ \text{seq}(\text{evalDG}(\text{convert}(\text{DirectionalCovariantDerivative}(vM[i], \omega\omega, CK), 'DGform') \\
&\&w \omega\omega), i = 1 \dots 4) \} \\
&\{(0 \, dx1) \wedge dx2\} \tag{2.18}
\end{aligned}$$

$$\begin{aligned}
> Sconn &:= \text{solve}(TGE, \text{ChrCoe}) : \\
> \text{FreePar} &:= \{ \} : \text{for } i \text{ to } \text{nops}(Sconn) \text{ do if } \text{lhs}(Sconn[i]) = \text{rhs}(Sconn[i]) \text{ then } \text{FreePar} \\
&:= \{ \text{op}(\text{FreePar}), \text{lhs}(Sconn[i]) \} \text{ fi:od: } \text{FreePar} \\
&\{a133, a144, a244, a333, a414, a422, a424, a433, a434, a444\} \tag{2.19}
\end{aligned}$$

$$\begin{aligned}
> \text{FreePar} &:= \{ \} : \text{for } i \text{ to } \text{nops}(Sconn) \text{ do if } \text{lhs}(Sconn[i]) = \text{rhs}(Sconn[i]) \text{ then } \text{FreePar} \\
&:= \{ \text{op}(\text{FreePar}), \text{lhs}(Sconn[i]) \} \text{ fi:od: } \text{FreePar} \\
&\{a144, a232, a244, a313, a333, a343, a344, a411, a424, a444\} \tag{2.20}
\end{aligned}$$

$$\begin{aligned}
> \text{ChrCoe1} &:= \{a111, a112, a113, a114, a121, a122, a123, a124, a131, a132, a134, a141, \\
&a142, a143, a211, a212, a213, a214, a221, a222, a223, a224, a231, a232, a233, a234, \\
&a241, a242, a243, a311, a312, a313, a314, a321, a322, a323, a324, a331, a332, a334, \\
&a341, a342, a343, a344, a411, a412, a413, a421, a423, a431, a432, a441, a442, a443\} :
\end{aligned}$$

```

> Sconn := solve(TGE, ChrCoe1) :
> FreePar := {a133, a144, a244, a333, a414, a422, a424, a433, a434, a444} :
> CC := DGsimplify(subs(Sconn, C)) :
>
> Quad1 := evalDG(add(add(diff(Eq1, ui,j) · vM[i] &s vM[j], j = 1..4), i = 1..4)) :
Quad2 := evalDG(add(add(diff(Eq2, ui,j) · vM[i] &s vM[j], j = 1..4), i = 1..4)) :
Quad3 := evalDG(add(add(diff(Eq3, ui,j) · vM[i] &s vM[j], j = 1..4), i = 1..4)) :
Quad1u := FromJet(Quad1, {u(x1, x2, x3, x4)}) : Quad2u := FromJet(Quad2, {u(x1, x2,
x3, x4)}) : Quad3u := FromJet(Quad3, {u(x1, x2, x3, x4)}) :
Quadu := [Quad1u, Quad2u, Quad3u] :
> {seq(ContractIndices(Quad||i, ω &t ω, [[1, 1], [2, 2]]), i = 1..3)}

```

{0} (2.21)

```

> RNE := { } :for i to 4 do for j to 3 do RNE := {op(RNE),
coeffs(expand(numer(simplify(subs(VsS2,
ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CC), ωω
&t ωω, [[1, 1], [2, 2]]), {u(x1, x2, x3, x4)}, notation = jetnumbers)) )), {q}))} :od:
od: nops(RNE)

```

33 (2.22)

```

> sRNE := solve(RNE, FreePar)

```

$$sRNE := \left\{ a133 = 0, a144 = 0, a244 = 0, a333 = \frac{1}{u_1 u_3 u_4 (a_3 - a_4)} \left(a434 a_3 u_1 u_3 u_4 \right. \right. \\ \left. \left. - a434 a_4 u_1 u_3 u_4 - a_1 u_3^2 u_{1,4} + a_1 u_3 u_4 u_{1,3} + a_3 u_1 u_4 u_{3,3} - a_3 u_3 u_4 u_{1,3} \right. \right. \\ \left. \left. - a_4 u_1 u_4 u_{3,3} + a_4 u_3^2 u_{1,4} \right), a414 = a414, a422 = 0, a424 = a424, a433 = 0, a434 \right. \\ \left. = a434, a444 = a444 \right\} \quad (2.23)$$

```

> sRNE := solve(RNE, {a133, a144, a244, a333, a422, a433})

```

$$sRNE := \left\{ a133 = 0, a144 = 0, a244 = 0, a333 = \frac{1}{u_1 u_3 u_4 (a_3 - a_4)} \left(a434 a_3 u_1 u_3 u_4 \right. \right. \\ \left. \left. - a434 a_4 u_1 u_3 u_4 - a_1 u_3^2 u_{1,4} + a_1 u_3 u_4 u_{1,3} + a_3 u_1 u_4 u_{3,3} - a_3 u_3 u_4 u_{1,3} \right. \right. \\ \left. \left. - a_4 u_1 u_4 u_{3,3} + a_4 u_3^2 u_{1,4} \right), a422 = 0, a433 = 0 \right\} \quad (2.24)$$

```

> CT := DGsimplify(evalDG(subs(
{a414 = u1,4/u4 + α1, a424 = u2,4/u4 + α2, a434 = u3,4/u4
+ α3, a444 = u4,4/u4 + α4}, subs(sRNE, CC)))) :

```

```

> {seq(seq(simplify(subs(VsS2, evalDG(DirectionalCovariantDerivative(vM[i], vM[j], CT)
- DirectionalCovariantDerivative(vM[i], vM[j], Ck))))), i = 1..4), j = 1..4)}
{0, 0 D_x1, 0 D_x2, 0 D_x3} (2.25)

```

```

> DirectionalCovariantDerivative(D_x1, D_x4, CT); DirectionalCovariantDerivative(D_x4,
D_x4, CT)

```

$$\frac{\alpha_1 u_4 + u_{1,4}}{u_4} D_{x4}$$

$$\frac{\alpha_4 u_4 + u_{4,4}}{u_4} D_{x4} \quad (2.26)$$

> # TorsionTensor(CT)

> sCTa := solve(DGinfo(ContractIndices(TorsionTensor(CT), [[1, 2]]), "CoefficientSet"),
 $\{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$: CTa := DGsimplify(evalDG(subs(sCTa, CT))) :

> simplify($\left(\text{subs}\left(\text{VsS2}, \text{subs}\left(sCTa, \left\{ 3\alpha_1 + \frac{u_{1,2}}{u_2} + \frac{u_{1,3}}{u_3} + \frac{u_{1,4}}{u_4}, 3\alpha_2 + \frac{u_{1,2}}{u_1} + \frac{u_{2,3}}{u_3} + \frac{u_{2,4}}{u_4}, 3\alpha_3 + \frac{u_{1,3}}{u_1} + \frac{u_{2,3}}{u_2} + \frac{u_{3,4}}{u_4}, 3\alpha_4 + \frac{u_{1,4}}{u_1} + \frac{u_{2,4}}{u_2} + \frac{u_{3,4}}{u_3} \right\} \right) \right) \right)$)
 $\{0\}$ (2.27)

> DirectionalCovariantDerivative(D_x1, D_x4, CTa); DirectionalCovariantDerivative(D_x1,
D_x1, CTa)

$$\frac{2u_{1,4}u_3u_2 - u_{1,3}u_4u_2 - u_{1,2}u_3u_4}{3u_2u_3u_4} D_{x4} - \left(\frac{u_1u_2u_3u_{1,4} + u_1u_2u_4u_{1,3} + u_1u_3u_4u_{1,2} - 3u_{1,1}u_2u_3u_4}{3u_3u_1u_4u_2} D_{x1} \right) \quad (2.28)$$

I.

Thus there exist GL2-connections, with respect to which the null Lax surfaces are totally geodesic. They depend on 4 arbitrary functions.

The parameters can be fixed uniquely by the condition Trace(Torsion)=0.

> RNE := { } :for i to 4 do for j to 3 do RNE := {op(RNE),
coeffs(expand(numer(simplify(subs(VsS2,
ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CT), ω &t ω,
[[1, 1], [2, 2]]), {u(x1, x2, x3, x4)}, notation = jetnumbers))))) , {q}) } :od:od: RNE
 $\{0\}$ (2.29)

> RNE := { } :for i to 4 do for j to 3 do RNE := {op(RNE),
coeffs(expand(numer(simplify(ToJet(ContractIndices(DirectionalCovariantDerivative(
vM[i], Quadu[j], CT), ω &t ω, [[1, 1], [2, 2]]), {u(x1, x2, x3, x4)}, notation
= jetnumbers))))) , {q}) } :od:od: nops(RNE)
13 (2.30)

> solve(RNE, {u_{2,3}, u_{2,4}, u_{3,4}})
 $\left\{ u_{2,3} = \frac{a_1 u_2 u_{1,3} - a_1 u_3 u_{1,2} + a_2 u_3 u_{1,2} - a_3 u_2 u_{1,3}}{u_1 (a_2 - a_3)}, u_{2,4} \right.$
 $= \frac{a_1 u_2 u_{1,4} - a_1 u_4 u_{1,2} + a_2 u_4 u_{1,2} - a_4 u_2 u_{1,4}}{u_1 (a_2 - a_4)}, u_{3,4}$
 $= \left. \frac{a_1 u_3 u_{1,4} - a_1 u_4 u_{1,3} + a_3 u_4 u_{1,3} - a_4 u_3 u_{1,4}}{u_1 (a_3 - a_4)} \right\}$ (2.31)

> VsS2

$$\left\{ \begin{aligned} u_{2,3} &= \frac{a_1 u_2 u_{1,3} - a_1 u_3 u_{1,2} + a_2 u_3 u_{1,2} - a_3 u_2 u_{1,3}}{u_1 (a_2 - a_3)}, u_{2,4} \\ &= \frac{a_1 u_2 u_{1,4} - a_1 u_4 u_{1,2} + a_2 u_4 u_{1,2} - a_4 u_2 u_{1,4}}{u_1 (a_2 - a_4)}, u_{3,4} \\ &= \frac{a_1 u_3 u_{1,4} - a_1 u_4 u_{1,3} + a_3 u_4 u_{1,3} - a_4 u_3 u_{1,4}}{u_1 (a_3 - a_4)} \end{aligned} \right\} \quad (2.32)$$

```
> # ContractIndices(TorsionTensor(CT), [[1, 2]])
> sCTa := solve(DGinfo(ContractIndices(TorsionTensor(CT), [[1, 2]]), "CoefficientSet"),
  {α1, α2, α3, α4}) : CTa := DGsimplify(evalDG(subs(sCTa, CT))) :
```

```
>
```

```
> α1 = subs(sCTa, α1);
α2 = simplify(subs(solve(VsS2, {u1,3, u1,4, u3,4}), subs(sCTa, α2)));
α3 = simplify(subs(solve(VsS2, {u1,2, u1,4, u2,4}), subs(sCTa, α3)));
α4 = simplify(subs(solve(VsS2, {u1,2, u1,3, u2,3}), subs(sCTa, α4)));
```

$$\alpha_1 = -\frac{1}{3} \frac{u_2 u_3 u_{1,4} + u_2 u_4 u_{1,3} + u_3 u_4 u_{1,2}}{u_2 u_3 u_4}$$

$$\alpha_2 = -\frac{1}{3} \frac{u_1 u_3 u_{2,4} + u_1 u_4 u_{2,3} + u_3 u_4 u_{1,2}}{u_1 u_3 u_4}$$

$$\alpha_3 = -\frac{1}{3} \frac{u_1 u_2 u_{3,4} + u_1 u_4 u_{2,3} + u_2 u_4 u_{1,3}}{u_1 u_4 u_2}$$

$$\alpha_4 = -\frac{1}{3} \frac{u_1 u_2 u_{3,4} + u_1 u_3 u_{2,4} + u_2 u_3 u_{1,4}}{u_1 u_2 u_3} \quad (2.33)$$

```
> sCTa2 := {α1 = -1/3 * (u2 u3 u1,4 + u2 u4 u1,3 + u3 u4 u1,2) / (u2 u3 u4), α2 =
  -1/3 * (u1 u3 u2,4 + u1 u4 u2,3 + u3 u4 u1,2) / (u1 u3 u4), α3 = -1/3 * (u1 u2 u3,4 + u1 u4 u2,3 + u2 u4 u1,3) / (u1 u4 u2),
  α4 = -1/3 * (u1 u2 u3,4 + u1 u3 u2,4 + u2 u3 u1,4) / (u3 u1 u2) } : CTa2
:= DGsimplify(evalDG(subs(sCTa2, CT))) :
```

```
> # CTa2
```

```
> [ContractIndices(TorsionTensor(CTa), [[1, 2]]), DGsimplify(simplify(subs(VsS2,
  ContractIndices(TorsionTensor(CTa2), [[1, 2]]))) ) ]
[0 dx1, 0 dx1] \quad (2.34)
```

```
>
```

```
> RicCTA := RicciTensor(FromJet(CTa, {u(x1, x2, x3, x4)})) : RicCTa := ToJet(RicCTA,
  {u(x1, x2, x3, x4)}, notation = jetnumbers) :
> RicCTaSkew := SymmetrizeIndices(RicCTa, [1, 2], "SkewSymmetric") :
RicCTaSymm := SymmetrizeIndices(RicCTa, [1, 2], "Symmetric") :
```

$$\{ \text{seq}(\text{ContractIndices}(\text{Quad} \parallel i, \text{evalDG}(\omega \& t \omega), [[1, 1], [2, 2]]), i = 1 \dots 3) \} \\ \{0\} \quad (2.35)$$

$$\begin{aligned} \omega 0 &:= \text{evalDG}(-\omega) : \omega 0b := \text{DGsimplify}(\text{evalDG}(\text{subs}(q=0, \omega 0))) ; \\ \omega 1a &:= \text{evalDG}\left(\frac{1}{\varepsilon} \cdot (\text{subs}(q=q+\varepsilon, \omega 0) - \omega 0)\right) : \omega 1 := \text{evalDG}(\text{subs}(\varepsilon=0, \omega 1a)) : \\ \omega 1b &:= \text{DGsimplify}(\text{evalDG}(\text{subs}(q=0, \omega 1))) ; \\ \omega 2a &:= \text{evalDG}\left(\frac{1}{\varepsilon} \cdot (\text{subs}(q=q+\varepsilon, \omega 1) - \omega 1)\right) : \omega 2 := \text{evalDG}(\text{subs}(\varepsilon=0, \omega 2a)) : \\ \omega 2b &:= \text{DGsimplify}\left(\text{evalDG}\left(\frac{1}{2} \cdot \text{subs}(q=0, \omega 2)\right)\right) ; \\ \omega 3a &:= \text{evalDG}\left(\frac{1}{\varepsilon} \cdot (\text{subs}(q=q+\varepsilon, \omega 2) - \omega 2)\right) : \omega 3 := \text{evalDG}(\text{subs}(\varepsilon=0, \omega 3a)) : \\ \omega 3b &:= \text{DGsimplify}\left(\text{evalDG}\left(\frac{1}{6} \cdot \text{subs}(q=0, \omega 3)\right)\right) \end{aligned}$$

$$\begin{aligned} \omega 0b &:= \frac{u_1}{a_1} dx1 + \frac{u_2}{a_2} dx2 + \frac{u_3}{a_3} dx3 + \frac{u_4}{a_4} dx4 \\ \omega 1b &:= \frac{u_1}{a_1^2} dx1 + \frac{u_2}{a_2^2} dx2 + \frac{u_3}{a_3^2} dx3 + \frac{u_4}{a_4^2} dx4 \\ \omega 2b &:= \frac{u_1}{a_1^3} dx1 + \frac{u_2}{a_2^3} dx2 + \frac{u_3}{a_3^3} dx3 + \frac{u_4}{a_4^3} dx4 \\ \omega 3b &:= \frac{u_1}{a_1^4} dx1 + \frac{u_2}{a_2^4} dx2 + \frac{u_3}{a_3^4} dx3 + \frac{u_4}{a_4^4} dx4 \end{aligned} \quad (2.36)$$

$$\begin{aligned} \# \text{Qua} \omega &:= \text{evalDG}([\omega 0b \& s \omega 0b, \omega 0b \& s \omega 1b, 2 \omega 0b \& s \omega 2b + \omega 1b \& s \omega 1b, \omega 0b \& s \omega 3b \\ &+ \omega 1b \& s \omega 2b, 2 \omega 1b \& s \omega 3b + \omega 2b \& s \omega 2b, \omega 2b \& s \omega 3b, \omega 3b \& s \omega 3b]) : \text{tmp} \Omega \\ &:= \{ \} : \text{for } i \text{ to nops}(\text{Qua} \omega) \text{ do for } j \text{ to 3 do tmp} \Omega := \{ \text{op}(\text{tmp} \Omega), \\ &\text{ContractIndices}(\text{Quad} \parallel j, \text{Qua} \omega[i], [[1, 1], [2, 2]]) \} : \text{od} : \text{od} : \text{tmp} \Omega \end{aligned}$$

$$\begin{aligned} \omega \omega &:= \text{evalDG}((q - a_1)(q - a_2)(q - a_3)(q - a_4)\omega) \\ \omega \omega &:= (q - a_2)(q - a_3)(q - a_4)u_1 dx1 + (q - a_1)(q - a_3)(q - a_4)u_2 dx2 + (q - a_1)(q - a_2)(q - a_4)u_3 dx3 + (q - a_1)(q - a_2)(q - a_3)u_4 dx4 \end{aligned} \quad (2.37)$$

$$\begin{aligned} \omega 0 &:= \text{evalDG}(-\omega \omega) : \omega 0b := \text{DGsimplify}(\text{evalDG}(\text{subs}(q=0, \omega 0))) ; \\ \omega 1a &:= \text{evalDG}\left(\frac{1}{\varepsilon} \cdot (\text{subs}(q=q+\varepsilon, \omega 0) - \omega 0)\right) : \omega 1 := \text{evalDG}(\text{subs}(\varepsilon=0, \omega 1a)) : \\ \omega 1b &:= \text{DGsimplify}(\text{evalDG}(\text{subs}(q=0, \omega 1))) ; \\ \omega 2a &:= \text{evalDG}\left(\frac{1}{\varepsilon} \cdot (\text{subs}(q=q+\varepsilon, \omega 1) - \omega 1)\right) : \omega 2 := \text{evalDG}(\text{subs}(\varepsilon=0, \omega 2a)) : \\ \omega 2b &:= \text{DGsimplify}\left(\text{evalDG}\left(\frac{1}{2} \cdot \text{subs}(q=0, \omega 2)\right)\right) ; \\ \omega 3a &:= \text{evalDG}\left(\frac{1}{\varepsilon} \cdot (\text{subs}(q=q+\varepsilon, \omega 2) - \omega 2)\right) : \omega 3 := \text{evalDG}(\text{subs}(\varepsilon=0, \omega 3a)) : \end{aligned}$$

$$\omega 3b := DG\text{simplify}\left(\text{evalDG}\left(\frac{1}{6} \cdot \text{subs}(q=0, \omega 3)\right)\right)$$

$$\omega 0b := a_2 a_3 a_4 u_1 dx1 + a_1 a_3 a_4 u_2 dx2 + a_1 a_2 a_4 u_3 dx3 + a_1 a_2 a_3 u_4 dx4$$

$$\omega 1b := -(u_1 (a_2 a_3 + a_2 a_4 + a_3 a_4) dx1) - (u_2 (a_3 a_1 + a_4 a_1 + a_3 a_4) dx2) - (u_3 (a_2 a_1 + a_4 a_1 + a_2 a_4) dx3) - (u_4 (a_2 a_1 + a_3 a_1 + a_2 a_3) dx4)$$

$$\omega 2b := u_1 (a_2 + a_3 + a_4) dx1 + u_2 (a_1 + a_3 + a_4) dx2 + u_3 (a_1 + a_2 + a_4) dx3 + u_4 (a_1 + a_2 + a_3) dx4$$

$$\omega 3b := -(u_1 dx1) - (u_2 dx2) - (u_3 dx3) - (u_4 dx4) \quad (2.38)$$

$$\begin{aligned} &> \text{Qua}\omega := \text{evalDG}([\omega 0b \&s \omega 0b, \omega 0b \&s \omega 1b, 2 \omega 0b \&s \omega 2b + \omega 1b \&s \omega 1b, \omega 0b \&s \omega 3b \\ &\quad + \omega 1b \&s \omega 2b, 2 \omega 1b \&s \omega 3b + \omega 2b \&s \omega 2b, \omega 2b \&s \omega 3b, \omega 3b \&s \omega 3b]) : \text{tmp}\Omega \\ &\quad := \{ \} : \text{for } i \text{ to nops(Qua}\omega) \text{ do for } j \text{ to 3 do tmp}\Omega := \{ \text{op}(\text{tmp}\Omega), \\ &\quad \text{ContractIndices(Quad}||j, \text{Qua}\omega[i], [[1, 1], [2, 2]]) \} : \text{od:od: tmp}\Omega \\ &\quad \{ 0 \} \end{aligned} \quad (2.39)$$

$$\begin{aligned} &> \text{eliminate}(\text{simplify}(\text{subs}(\text{VsS3}, \text{DGinfo}(\text{evalDG}(\text{RicCTaSymm} - \text{add}(\Lambda[i] \cdot \text{Qua}\omega[i], i = 1 \\ &\quad \dots 7))), \text{"CoefficientSet"}))), \{ \text{seq}(\Lambda[i], i = 1 \dots 7) \}) [2] \\ &\quad \{ \} \end{aligned} \quad (2.40)$$

Thus the condition that $\text{sym}(\text{Ric}(\nabla))$ belongs to the span of quadrics defining the GL2 structure is non-vacous. It is another differential constraint, giving exact solutions.

But in fact, from what is computed below it follows that this is the same condition as $\text{sym}(\text{Ric})=0$.

$$\begin{aligned} &> \# \text{ToJet}(\text{RicciTensor}(\text{FromJet}(\text{CTa}, \{u(x1, x2, x3, x4)\})), \{u(x1, x2, x3, x4)\}, \text{notation} \\ &\quad = \text{jetnumbers}) \end{aligned}$$

$$\begin{aligned} &> \text{Vt0} := \text{factor}(\text{Annihilator}([\omega 0, \omega 1, \omega 2])[1]); \text{Vt1} := \text{factor}(\text{Annihilator}([\omega 0, \omega 1])[2]); \\ &\quad \text{Vt2} := \text{factor}(\text{Annihilator}([\omega 0])[3]) \end{aligned}$$

$$\begin{aligned} \text{Vt0} := & - \left(\frac{(q - a_1)^3 (a_3 - a_4) (a_2 - a_4) u_4}{(q - a_4)^3 u_1 (a_1 - a_3) (a_1 - a_2)} D_{x1} \right) \\ & + \frac{u_4 (q - a_2)^3 (a_3 - a_4) (a_1 - a_4)}{u_2 (a_1 - a_2) (q - a_4)^3 (a_2 - a_3)} D_{x2} \\ & - \left(\frac{(a_2 - a_4) (a_1 - a_4) (q - a_3)^3 u_4}{(q - a_4)^3 (a_2 - a_3) (a_1 - a_3) u_3} D_{x3} \right) + D_{x4} \end{aligned}$$

$$\text{Vt1} := \frac{u_4 (q - a_1)^2 (a_2 - a_4)}{(a_1 - a_2) (q - a_4)^2 u_1} D_{x1} - \left(\frac{u_4 (a_1 - a_4) (q - a_2)^2}{u_2 (a_1 - a_2) (q - a_4)^2} D_{x2} \right) + D_{x4}$$

$$\text{Vt2} := - \left(\frac{(q - a_1) u_2}{(q - a_2) u_1} D_{x1} \right) + D_{x2} \quad (2.41)$$

$$\begin{aligned} &> \text{Vt01} := \text{ContractIndices}(\text{TorsionTensor}(\text{CT}), \text{evalDG}(\text{Vt0} \&t \text{Vt1}), [[2, 1], [3, 2]]) : \\ &\quad \text{KrCond} := \text{DGinfo}(\text{evalDG}(\text{convert}(\text{Vt01}, \text{DGvector}) - \text{Hook}(\text{convert}(\text{Vt01}, \text{DGvector}), dx4) \\ &\quad \cdot \text{Vt0}), \text{"CoefficientSet"}) : \\ &\quad \text{sCTa3} := \text{map}(T \rightarrow \text{factor}(T), \text{solve}(\text{map}(Q \rightarrow \text{coeffs}(\text{expand}(\text{numer}(Q)), \{q\}), \text{subs}(\text{VsS3}, \end{aligned}$$

$$\text{KrCond})) , \{ \alpha_1, \alpha_2, \alpha_3, \alpha_4 \}) : \\ \rightarrow \text{simplify}(\{ \text{seq}(\text{subs}(\text{VsS2}, \text{subs}(\text{sCTa}, \alpha[i])) - \text{subs}(\text{sCTa2}, \alpha[i])), i = 1..4), \\ \text{seq}(\text{subs}(\text{VsS2}, \text{subs}(\text{sCTa}, \alpha[i])) - \text{subs}(\text{sCTa3}, \alpha[i])), i = 1..4) \}) \\ \{0\} \quad (2.42)$$

$$\rightarrow \text{sCTa2} \\ \left\{ \alpha_1 = -\frac{1}{3} \frac{u_2 u_3 u_{1,4} + u_2 u_4 u_{1,3} + u_3 u_4 u_{1,2}}{u_2 u_3 u_4}, \alpha_2 = \right. \\ \left. -\frac{1}{3} \frac{u_1 u_3 u_{2,4} + u_1 u_4 u_{2,3} + u_3 u_4 u_{1,2}}{u_1 u_3 u_4}, \alpha_3 = \right. \\ \left. -\frac{1}{3} \frac{u_1 u_2 u_{3,4} + u_1 u_4 u_{2,3} + u_2 u_4 u_{1,3}}{u_1 u_4 u_2}, \alpha_4 = \right. \\ \left. -\frac{1}{3} \frac{u_1 u_2 u_{3,4} + u_1 u_3 u_{2,4} + u_2 u_3 u_{1,4}}{u_1 u_2 u_3} \right\} \quad (2.43)$$

$$\rightarrow \text{evalDG}((q - a_4)^3 \cdot \text{Vt0}) \\ - \left(\frac{(q - a_1)^3 (a_3 - a_4) (a_2 - a_4) u_4}{u_1 (a_1 - a_3) (a_1 - a_2)} D_{x1} \right) + \frac{u_4 (q - a_2)^3 (a_3 - a_4) (a_1 - a_4)}{u_2 (a_1 - a_2) (a_2 - a_3)} D_{x2} \quad (2.44) \\ - \left(\frac{(a_2 - a_4) (a_1 - a_4) (q - a_3)^3 u_4}{(a_2 - a_3) (a_1 - a_3) u_3} D_{x3} \right) + (q - a_4)^3 D_{x4}$$

$\rightarrow$ Thus there exist GL2-connections, with respect to which the null Lax surfaces are totally geodesic. They depend on 4 arbitrary functions. The parameters can be fixed uniquely by the condition $\text{Trace}(\text{Torsion})=0$. This condition is equivalent to that one of Krynsky.

$$\rightarrow \text{simplify}(\text{subs}(\text{VsS3}, \text{ContractIndices}(\text{RicCTaSymm}, \text{evalDG}(\text{Vt0} \& \text{Vt0}), [[1, 1], [2, 2]]))) \\ 0 \quad (2.45)$$

$$\rightarrow \text{simplify}(\text{subs}(\text{VsS3}, \text{ContractIndices}(\text{RicCTaSymm}, \text{evalDG}(\text{Vt0} \& \text{Vt1}), [[1, 1], [2, 2]]))) \\ 0 \quad (2.46)$$

$$\rightarrow \text{simplify}(\text{subs}(\text{VsS3}, \text{ContractIndices}(\text{RicCTaSymm}, \text{evalDG}(2 \text{ Vt0} \& \text{s Vt2} + 3 \text{ Vt1} \& \text{Vt1}), \\ [[1, 1], [2, 2]]))) \\ 0 \quad (2.47)$$

$$\rightarrow \text{QuT} := \text{evalDG}(\text{add}(\text{add}(\xi[i, j] \cdot dM[i] \& s dM[j], j = i..4), i = 1..4)) : \\ \text{SuT} := \text{solve}(\{ \text{coeffs}(\text{expand}(\text{numer}(\text{ContractIndices}(\text{QuT}, \text{Vt0} \& \text{Vt0}), [[1, 1], [2, 2]]))) , \\ \{q\} \}) , \{ \xi_{1,1}, \xi_{1,2}, \xi_{1,3}, \xi_{1,4}, \xi_{2,2}, \xi_{3,3}, \xi_{4,4} \}) : \\ \text{TuT} := \text{DGsimplify}(\text{evalDG}(\text{subs}(\{ \xi_{2,3} = \Lambda_1, \xi_{2,4} = \Lambda_2, \xi_{3,4} = \Lambda_3 \}, \text{subs}(\text{SuT}, \text{QuT})))) :$$

$\rightarrow$ # RicCTaSymm

$$\rightarrow \text{eliminate}(\text{DGinfo}(\text{DGsimplify}(\text{evalDG}(\text{subs}(\text{VsS3}, \text{TuT} - \text{RicCTaSymm}}))), \\ \text{"CoefficientSet"}), \{ \Lambda_1, \Lambda_2, \Lambda_3 \} \\ [\{ \Lambda_1 = 0, \Lambda_2 = 0, \Lambda_3 = 0 \}, \{ \}] \quad (2.48)$$

This computation has shown that symmetrized Ricci tensor of the canonical GL2-connection is a section of the bundle of quadrics annihilating the dual (tangent) to the canonical (cotangent) rational normal curve.

>

Finally, let us compute the Christoffel symbols of the canonical connection.

Due to full S[4] symmetry in the indices (1,2,3,4) it is enough to compute only $\nabla_j \partial_j = \Gamma_{jj}^i \partial_i$ and

$\nabla_j \partial_k = \Gamma_{jk}^i \partial_i$ for $j \neq k$. So we compute $\nabla_1 \partial_1$ and $\nabla_1 \partial_2$; the full result follows.

$$\begin{aligned} > \left\{ u_{2,3} = \frac{(a_1 - a_3) u_2 u_{1,3} - (a_1 - a_2) u_3 u_{1,2}}{u_1 (a_2 - a_3)}, u_{2,4} \right. \\ &= \frac{(a_1 - a_4) u_2 u_{1,4} - (a_1 - a_2) u_4 u_{1,2}}{u_1 (a_2 - a_4)}, u_{3,4} \\ &= \left. \frac{(a_1 - a_4) u_3 u_{1,4} - (a_1 - a_3) u_4 u_{1,3}}{u_1 (a_3 - a_4)} \right\} : \end{aligned}$$

$$\begin{aligned} > \Gamma_{11} := \frac{u_{1,1}}{u_1} - \frac{1}{3} \cdot \left(\frac{u_{1,2}}{u_2} + \frac{u_{1,3}}{u_3} + \frac{u_{1,4}}{u_4} \right) : \Gamma_{212} := \frac{1}{3} \cdot \left(2 \frac{u_{1,2}}{u_2} - \frac{u_{1,3}}{u_3} \right. \\ &\quad \left. - \frac{u_{1,4}}{u_4} \right) : \end{aligned}$$

```
> DGSimplify(evalDG(convert(DirectionalCovariantDerivative(D_x1, D_x1, CTa), DGvector)
- Γ111·D_x1));
DGSimplify(evalDG(convert(DirectionalCovariantDerivative(D_x1, D_x2, CTa), DGvector)
- Γ212·D_x2));
```

0 D_x1

0 D_x1

(2.49)

>

>

II.

Let us check that there exists a unique symmetric connection preserving GL2-structure

```
> ChrSeq := {seq(seq(seq(((a||i)||j)||k), k=1..4), j=i..4), i=1..4)}:
CsC := Connection(subs({seq(seq(seq(((a||i)||j)||k)=(((a||j)||i)||k), k=1..4), j=1
..(i-1)), i=1..4)}, add(add(add(((a||i)||j)||k)·vM[k]&t dM[i]&t dM[j], k=1
..4), j=1..4), i=1..4)): TorsionTensor(CsC)
```

(2.50)

```
> RNE := {} :for i to 4 do for j to 3 do RNE := {op(RNE),
coeffs(expand(numer(simplify(subs(VsS2,
ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CsC), ω
&t ω, [[1, 1], [2, 2]]), {u(x1, x2, x3, x4)}, notation=jetnumbers)))}, {q})} :od:od:
```

```
> Sss1 := solve(RNE, ChrSeq) : nops(Sss1)
```

40

(2.51)

```
> CsS := DGSimplify(evalDG(subs(Sss1, CsC))) :
```

```
> seq(seq(simplify(subs(VsS2,
ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CsS), ω
```

& t ω, [[1, 1], [2, 2]], {u(x1, x2, x3, x4)}, notation = jetnumbers)) , i = 1 ..4), j = 1 ..3)

0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0

(2.52)

>
>
>
>

for i to 4 do for j to 4 do a || i || j := a_i - a_j :od:od:

> simplify $\left(\text{subs}(Sss1, a111) - \left(\frac{u_{1,1}}{u_1} - \frac{1}{9} \cdot \left(\frac{(a13 \cdot a24 + a14 \cdot a23)^2}{a13 \cdot a14 \cdot a23 \cdot a24} \frac{u_{1,2}}{u_2} \right. \right. \right.$
 $\left. \left. + \frac{(a12 \cdot a34 + a14 \cdot a32)^2}{a12 \cdot a14 \cdot a32 \cdot a34} \frac{u_{1,3}}{u_3} + \frac{(a12 \cdot a43 + a13 \cdot a42)^2}{a12 \cdot a13 \cdot a42 \cdot a43} \frac{u_{1,4}}{u_4} \right) \right) \Bigg);$
simplify $\left(\text{subs}(Sss1, a112) - \frac{1}{9} \frac{a23 \cdot a24}{a12} \frac{u_1}{u_2} \cdot \left(\frac{1}{a43} \frac{u_{1,3}}{u_3} + \frac{1}{a34} \frac{u_{1,4}}{u_4} \right) \right);$
simplify $\left(\text{subs}(Sss1, a113) - \frac{1}{9} \frac{a32 \cdot a34}{a13} \frac{u_1}{u_3} \cdot \left(\frac{1}{a42} \frac{u_{1,2}}{u_2} + \frac{1}{a24} \frac{u_{1,4}}{u_4} \right) \right);$
simplify $\left(\text{subs}(Sss1, a114) - \frac{1}{9} \frac{a42 \cdot a43}{a14} \frac{u_1}{u_4} \cdot \left(\frac{1}{a32} \frac{u_{1,2}}{u_2} + \frac{1}{a23} \frac{u_{1,3}}{u_3} \right) \right);$

0
0
0
0

(2.53)

> simplify $\left(\text{subs} \left(VsS2, \text{subs}(Sss1, a222) - \left(\frac{u_{2,2}}{u_2} - \frac{1}{9} \cdot \left(\frac{(a23 \cdot a14 + a24 \cdot a13)^2}{a13 \cdot a14 \cdot a23 \cdot a24} \frac{u_{1,2}}{u_1} \right. \right. \right. \right.$
 $\left. \left. + \frac{(a21 \cdot a34 + a24 \cdot a31)^2}{a21 \cdot a24 \cdot a31 \cdot a34} \frac{u_{2,3}}{u_3} + \frac{(a21 \cdot a43 + a23 \cdot a41)^2}{a21 \cdot a23 \cdot a41 \cdot a43} \frac{u_{2,4}}{u_4} \right) \right) \Bigg);$
simplify $\left(\text{subs} \left(VsS2, \text{subs}(Sss1, a221) - \frac{1}{9} \frac{a13 \cdot a14}{a21} \frac{u_2}{u_1} \cdot \left(\frac{1}{a43} \frac{u_{2,3}}{u_3} + \frac{1}{a34} \frac{u_{2,4}}{u_4} \right) \right) \right);$
simplify $\left(\text{subs} \left(VsS2, \text{subs}(Sss1, a223) - \frac{1}{9} \frac{a31 \cdot a34}{a23} \frac{u_2}{u_3} \cdot \left(\frac{1}{a41} \frac{u_{1,2}}{u_1} + \frac{1}{a14} \frac{u_{2,4}}{u_4} \right) \right) \right);$
simplify $\left(\text{subs} \left(VsS2, \text{subs}(Sss1, a224) - \frac{1}{9} \frac{a41 \cdot a43}{a24} \frac{u_2}{u_4} \cdot \left(\frac{1}{a31} \frac{u_{1,2}}{u_1} + \frac{1}{a13} \frac{u_{2,3}}{u_3} \right) \right) \right);$

0
0
0
0

(2.54)

> simplify $\left(\text{subs} \left(VsS2, \text{subs}(Sss1, a222) - \left(\frac{u_{2,2}}{u_2} \right. \right. \right.$
 $\left. \left. + \frac{u_2}{9 u_1} \left(\frac{3(a12 \cdot a23 + a32 \cdot a24 + a42 \cdot a21)}{a23 \cdot a24} \frac{u_{1,2}}{u_2} - \frac{(a13 \cdot a24 + a12 \cdot a34)^2}{a23 \cdot a21 \cdot a24 \cdot a43} \frac{u_{1,3}}{u_3} \right) \right) \right);$

>

>

Finally let us compute the canonical projective totally geodesic connection

$\geq$

$$\begin{aligned} &> \text{Sss2} := \text{solve}(\text{TGE}, \text{ChrCoe}) : \text{nops}(\text{Sss2}) \\ &\quad \quad \quad 40 \end{aligned} \tag{2.59}$$

$$\begin{aligned} &> \text{FreePar} := \{ \} : \text{for } i \text{ to } \text{nops}(\text{Sss2}) \text{ do if } \text{lhs}(\text{Sss2}[i]) = \text{rhs}(\text{Sss2}[i]) \text{ then } \text{FreePar} \\ &\quad := \{ \text{op}(\text{FreePar}), \text{lhs}(\text{Sss2}[i]) \} \text{ fi:od: } \text{FreePar} \\ &\quad \quad \quad \{a144, a244, a344, a444\} \end{aligned} \tag{2.60}$$

$$\begin{aligned} &> \text{nops}(\text{Sss2} \setminus \{a144 = a144, a244 = a244, a344 = a344, a444 = a444\}) \\ &\quad \quad \quad 36 \end{aligned} \tag{2.61}$$

$$\begin{aligned} &> \text{pt} := \left\{ a144 = \frac{1}{2} \frac{u_{1,4}}{u_4}, a244 = \frac{1}{2} \frac{u_{2,4}}{u_4}, a344 = \frac{1}{2} \frac{u_{3,4}}{u_4}, a444 = \frac{u_{4,4}}{u_4} \right\} : \\ &> \text{Sss20} := \{ \text{op}(\text{subs}(\text{pt}, \text{Sss2} \setminus \{a144 = a144, a244 = a244, a344 = a344, a444 = a444\})), \\ &\quad \text{op}(\text{pt}) \} : \\ &> \text{subs}(\text{Sss2}, a111); \text{subs}(\text{Sss2}, a122); \\ &\quad \quad \quad \frac{2 a144 u_1 u_4 - u_1 u_{1,4} + u_4 u_{1,1}}{u_1 u_4} \\ &\quad \quad \quad \frac{1}{2} \frac{2 a144 u_2 u_4 - u_2 u_{1,4} + u_4 u_{1,2}}{u_2 u_4} \end{aligned} \tag{2.62}$$

$$\begin{aligned} &> \text{subs}(\text{Sss20}, a111); \text{subs}(\text{Sss20}, a112); \text{subs}(\text{Sss20}, a113); \text{subs}(\text{Sss20}, a114) \\ &\quad \quad \quad \frac{u_{1,1}}{u_1} \\ &\quad \quad \quad 0 \\ &\quad \quad \quad 0 \\ &\quad \quad \quad 0 \end{aligned} \tag{2.63}$$

$$\begin{aligned} &> \text{simplify}(\text{subs}(\text{VsS2a}, \text{subs}(\text{Sss20}, a121))); \text{subs}(\text{Sss20}, a122); \text{subs}(\text{Sss20}, a123); \\ &\quad \text{subs}(\text{Sss20}, a124) \\ &\quad \quad \quad \frac{1}{2} \frac{u_{1,2}}{u_1} \\ &\quad \quad \quad \frac{1}{2} \frac{u_{1,2}}{u_2} \\ &\quad \quad \quad 0 \\ &\quad \quad \quad 0 \end{aligned} \tag{2.64}$$

$$\begin{aligned} &> \text{simplify}(\text{subs}(\text{VsS2}, \text{subs}(\text{Sss20}, a131))); \text{subs}(\text{Sss20}, a132); \text{subs}(\text{Sss20}, a133); \\ &\quad \text{subs}(\text{Sss20}, a134) \\ &\quad \quad \quad \frac{1}{2} \frac{u_{1,3}}{u_1} \\ &\quad \quad \quad 0 \\ &\quad \quad \quad \frac{1}{2} \frac{u_{1,3}}{u_3} \\ &\quad \quad \quad 0 \end{aligned} \tag{2.65}$$

$$\begin{aligned} &> \text{simplify}(\text{subs}(\text{VsS2}, \text{subs}(\text{Sss20}, a141))); \text{subs}(\text{Sss20}, a142); \text{subs}(\text{Sss20}, a143); \\ &\quad \text{subs}(\text{Sss20}, a144) \end{aligned}$$

$$\begin{array}{c} \frac{1}{2} \frac{u_{1,4}}{u_1} \\ 0 \\ 0 \\ \frac{1}{2} \frac{u_{1,4}}{u_4} \end{array} \quad (2.66)$$

Adler-Shabat triples 4D

- > $DGsetup([x1, x2, x3, x4], [s, u], Js, 3) :$
- > $Ssyst := \{s_2 = \ln(u_2 - s_1), s_3 = \ln(u_3 - s_1), s_4 = \ln(u_4 - s_1)\} :$
- > $Vsyst := map(Q \rightarrow coeffs(Q, \{s_1, s_{1,1}\}), expand(numer(subs(\{s_{1,2} = TotalDiff(subs(Ssyst, s_2), 1), s_{1,3} = TotalDiff(subs(Ssyst, s_3), 1), s_{1,4} = TotalDiff(subs(Ssyst, s_4), 1)\}, \{TotalDiff(subs(Ssyst, s_3), 2) - TotalDiff(subs(Ssyst, s_2), 3), TotalDiff(subs(Ssyst, s_4), 2) - TotalDiff(subs(Ssyst, s_2), 4), TotalDiff(subs(Ssyst, s_4), 3) - TotalDiff(subs(Ssyst, s_3), 4)\}))) ; VsS2 := solve(Vsyst, \{u_{2,3}, u_{2,4}, u_{3,4}\})$
- $$Vsyst := \{u_2 u_{2,3} - u_3 u_{2,3} - u_{1,2} + u_{1,3}, u_2 u_{2,4} - u_4 u_{2,4} - u_{1,2} + u_{1,4}, u_3 u_{3,4} - u_4 u_{3,4} - u_{1,3} + u_{1,4}\}$$
- $$VsS2 := \left\{ u_{2,3} = \frac{u_{1,2} - u_{1,3}}{u_2 - u_3}, u_{2,4} = \frac{u_{1,2} - u_{1,4}}{u_2 - u_4}, u_{3,4} = \frac{u_{1,3} - u_{1,4}}{u_3 - u_4} \right\} \quad (3.1)$$
- > $Eq1 := Vsyst[1] : Eq2 := Vsyst[2] : Eq3 := Vsyst[3] :$
- > $eliminate(Vsyst, \{u_{1,2}, u_{1,3}, u_{1,4}\})$
- $$\left[\{u_{1,2} = u_2 u_{2,4} - u_4 u_{2,4} + u_{1,4} u_{1,3} = u_3 u_{3,4} - u_4 u_{3,4} + u_{1,4} u_{1,4} = u_{1,4}\}, \{u_2 u_{2,3} - u_2 u_{2,4} - u_3 u_{2,3} + u_3 u_{3,4} + u_4 u_{2,4} - u_4 u_{3,4}\} \right] \quad (3.2)$$
- > $simplify(subs(VsS2, (u_2 - u_3) u_{2,3} + (u_3 - u_4) u_{3,4} + (u_4 - u_2) u_{2,4}))$
- $$0 \quad (3.3)$$
- > $Vss3 := Vsyst : \text{for } i \text{ to } 4 \text{ do for } j \text{ to } nops(Vsyst) \text{ do } Vss3 := \{op(Vss3), TotalDiff(Vsyst[j], i)\} : \text{od} : \text{od} :$
- $$nops(eliminate(Vss3, \{u_{2,3}, u_{2,4}, u_{3,4}, u_{1,2,3}, u_{1,2,4}, u_{1,3,4}, u_{2,2,3}, u_{2,2,4}, u_{2,3,3}, u_{2,3,4}, u_{2,4,3}, u_{2,4,4}, u_{3,3,3}, u_{3,3,4}\})[2])$$
- $$0 \quad (3.4)$$
- > $VsS3 := solve(Vss3, \{u_{2,3}, u_{2,4}, u_{3,4}, u_{1,2,3}, u_{1,2,4}, u_{1,3,4}, u_{2,2,3}, u_{2,3,3}, u_{2,3,4}, u_{2,4,3}, u_{2,4,4}, u_{3,3,3}, u_{3,3,4}\}) : nops(VsS3)$
- $$13 \quad (3.5)$$
- > $SymbEO2a := add(add(diff(Vsyst[1], u_{i,j}) \cdot p_i \cdot p_j, j = i..4), i = 1..4) ;$
- $$SymbEO2b := add(add(diff(Vsyst[2], u_{i,j}) \cdot p_i \cdot p_j, j = i..4), i = 1..4) ;$$

$$\begin{aligned}
\text{SymbEO2c} &:= \text{add}(\text{add}(\text{diff}(\text{Vsyst}[3], u_{i,j}) \cdot p_i \cdot p_j, j=i..4), i=1..4); \\
\text{SymbEO2a} &:= -p_1 p_2 + p_1 p_3 + (u_2 - u_3) p_2 p_3 \\
\text{SymbEO2b} &:= -p_1 p_2 + p_1 p_4 + (u_2 - u_4) p_2 p_4 \\
\text{SymbEO2c} &:= -p_1 p_3 + p_1 p_4 + (u_3 - u_4) p_3 p_4
\end{aligned} \tag{3.6}$$

$$\begin{aligned}
> \text{simplify}\left(\text{subs}\left(\left\{p_1=1, p_2=\frac{1}{q-u_2}, p_3=\frac{1}{q-u_3}, p_4=\frac{1}{q-u_4}\right\}, \{\text{SymbEO2a}, \text{SymbEO2b}, \right. \right. \\
\left. \left. \text{SymbEO2c}\}\right)\right) \\
\{0\}
\end{aligned} \tag{3.7}$$

$$\begin{aligned}
> \text{tmp} &:= \text{solve}(\{\text{coeffs}(\text{expand}((a1 \cdot p_1 + a2 \cdot p_2 + a3 \cdot p_3 + a4 \cdot p_4) \cdot \text{SymbEO2a} + (b1 \cdot p_1 + b2 \\
&\cdot p_2 + b3 \cdot p_3 + b4 \cdot p_4) \cdot \text{SymbEO2b} + (c1 \cdot p_1 + c2 \cdot p_2 + c3 \cdot p_3 + c4 \cdot p_4) \cdot \text{SymbEO2c}), \\
&\{p_1, p_2, p_3, p_4\})\}, \{a1, a2, a3, a4, b1, b2, b3, b4, c1, c2, c3, c4\}) \\
\text{tmp} &:= \{a1=c1, a2=0, a3=0, a4=-c1 u_2 + c1 u_4 + c2, b1=-c1, b2=0, b3=c1 u_2 \\
&- c1 u_3 - c2, b4=0, c1=c1, c2=c2, c3=0, c4=0\}
\end{aligned} \tag{3.8}$$

$$\begin{aligned}
> \text{simplify}([(p_1 + (u_4 - u_2) \cdot p_4) \cdot \text{SymbEO2a} + (-p_1 + (u_2 - u_3) \cdot p_3) \cdot \text{SymbEO2b} + p_1 \\
\cdot \text{SymbEO2c}, p_4 \cdot \text{SymbEO2a} - p_3 \cdot \text{SymbEO2b} + p_2 \cdot \text{SymbEO2c}]) \\
[0, 0]
\end{aligned} \tag{3.9}$$

$$\begin{aligned}
> \text{simplify}(\text{subs}(\text{VsS2}, \text{TotalDiff}(\text{Eq1}, 1) + (u_4 - u_2) \cdot \text{TotalDiff}(\text{Eq1}, 4) - \text{TotalDiff}(\text{Eq2}, 1) \\
+ (u_2 - u_3) \cdot \text{TotalDiff}(\text{Eq2}, 3) + \text{TotalDiff}(\text{Eq3}, 1))) \\
\text{simplify}(\text{subs}(\text{VsS2}, \text{TotalDiff}(\text{Eq1}, 4) - \text{TotalDiff}(\text{Eq2}, 3) + \text{TotalDiff}(\text{Eq3}, 2))) \\
0 \\
0
\end{aligned} \tag{3.10}$$

$$\begin{aligned}
> \text{Vss4} &:= \text{Vss3} : \text{for } i \text{ to } 4 \text{ do for } k \text{ to } i \text{ do for } j \text{ to } \text{nops}(\text{Vsyst}) \text{ do } \text{Vss4} := \{\text{op}(\text{Vss4}), \\
&\text{TotalDiff}(\text{Vsyst}[j], [i, k])\} : \text{od} : \text{od} : \text{od} : \\
\text{tmp} &:= \text{eliminate}(\text{Vss4}, \{u_{2,3}, u_{2,4}, u_{3,4}, u_{1,2,3}, u_{1,2,4}, u_{1,3,4}, u_{2,2,3}, u_{2,3,3}, u_{2,3,4}, u_{2,2,4}, \\
&u_{2,4,4}, u_{3,3,4}, u_{3,4,4}, u_{1,1,2,3}, u_{1,1,2,4}, u_{1,1,3,4}, u_{1,2,2,3}, u_{1,2,2,4}, u_{1,2,3,4}, u_{1,2,4,4}, \\
&u_{1,2,3,3}, u_{1,2,3,4}, u_{1,3,3,4}, u_{1,3,4,4}, u_{2,2,2,3}, u_{2,2,2,4}, u_{2,2,3,4}, u_{2,2,3,3}, u_{2,3,3,4}, u_{2,2,4,4}, \\
&u_{2,3,4,4}, u_{2,3,3,3}, u_{3,3,3,4}, u_{3,3,4,4}, u_{2,4,4,4}, u_{3,4,4,4}\}) : [\text{nops}(\text{tmp}[2]), \text{nops}(\text{tmp}[1])] \\
[0, 35]
\end{aligned} \tag{3.11}$$

This shows that dimensions of the symbol spaces (g[0],g[1],g[2],g[3],g[4],...) are equal to (1,4,7,10,13,...),

i.e. the Hilbert polynomial is given by the formula $\dim(g[k])=1+3k$, whence the general solution depends on 3 functions of 1 variable.

This is also what is expected from the characteristic variety (rational normal curve in our case) that has order 3 and dimension 1.

>

$$> \text{DGsetup}([x1, x2, x3, x4], [q, u], \text{Jq}, 3) :$$

$$> \text{Qsyst} := \left\{ q_2 = \frac{u_{1,2} - q_1}{u_2 - q_{[1]}}, q_3 = \frac{u_{1,3} - q_1}{u_3 - q_{[1]}}, q_4 = \frac{u_{1,4} - q_1}{u_4 - q_{[1]}} \right\} :$$

$$> \text{simplify}(\text{subs}(\text{VsS3}, \text{expand}(\text{subs}(\text{Qsyst}, \text{subs}(\{q_{1,2} = \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_2), 1), q_{1,3}$$

$$\begin{aligned}
&= \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_3), 1), q_{1,4} = \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_4), 1) \}, \\
&\{ \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_3), 2) - \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_2), 3), \\
&\text{TotalDiff}(\text{subs}(\text{Qsyst}, q_4), 2) - \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_2), 4), \text{TotalDiff}(\text{subs}(\text{Qsyst}, \\
&q_4), 3) - \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_3), 4) \} \} \} \} \\
&\{0\}
\end{aligned} \tag{3.12}$$

> DGsetup([x1, x2, x3, x4, q], [u], Jqv, 3) :

$$> X2f := f \rightarrow \text{TotalDiff}(f, 2) + \frac{1}{u_2 - q} \cdot \text{TotalDiff}(f, 1) + \frac{u_{1,2}}{u_2 - q} \cdot \text{diff}(f, q) :$$

$$X3f := f \rightarrow \text{TotalDiff}(f, 3) + \frac{1}{u_3 - q} \cdot \text{TotalDiff}(f, 1) + \frac{u_{1,3}}{u_3 - q} \cdot \text{diff}(f, q) :$$

$$X4f := f \rightarrow \text{TotalDiff}(f, 4) + \frac{1}{u_4 - q} \cdot \text{TotalDiff}(f, 1) + \frac{u_{1,4}}{u_4 - q} \cdot \text{diff}(f, q) :$$

$$\begin{aligned}
> \text{simplify}(\text{subs}(\text{VsS3}, \{ \text{simplify}(X2f(X3f(x1)) - X3f(X2f(x1))), \text{simplify}(X2f(X3f(q)) \\
&- X3f(X2f(q))), \text{simplify}(X2f(X4f(x1)) - X4f(X2f(x1))), \text{simplify}(X2f(X4f(q)) \\
&- X4f(X2f(q))), \text{simplify}(X3f(X4f(x1)) - X4f(X3f(x1))), \text{simplify}(X3f(X4f(q)) \\
&- X4f(X3f(q))) \} \} \} \} \\
&\{0\}
\end{aligned} \tag{3.13}$$

>

1. We first look for a connection (with torsion) preserving the field of GL2-curves and having α -surfaces totally geodesic + trace of torsion zero.

> DGsetup([x1, x2, x3, x4], M) :

> vM := [D_x1, D_x2, D_x3, D_x4] : dM := [dx1, dx2, dx3, dx4] :

> C := Connection(add(add(add(((a||i)||j)||k)·vM[k]&t dM[i]&t dM[j], k=1..4), j=1..4), i=1..4)) :

$$> X2 := \text{evalDG}\left(D_{x2} + \frac{1}{u_2 - q_{[]}} \cdot D_{x1}\right) :$$

$$X3 := \text{evalDG}\left(D_{x3} + \frac{1}{u_3 - q_{[]}} \cdot D_{x1}\right) :$$

$$X4 := \text{evalDG}\left(D_{x4} + \frac{1}{u_4 - q_{[]}} \cdot D_{x1}\right) :$$

$$\omega := \text{evalDG}\left(dx1 - \frac{1}{u_2 - q_{[]}} \cdot dx2 - \frac{1}{u_3 - q_{[]}} \cdot dx3 - \frac{1}{u_4 - q_{[]}} \cdot dx4\right) :$$

$$[\text{Hook}(X2, \omega), \text{Hook}(X3, \omega), \text{Hook}(X4, \omega)]$$

$$[0, 0, 0]$$

(3.14)

> XX2 := FromJet(X2, {u(x1, x2, x3, x4), q(x1, x2, x3, x4)}) :

XX3 := FromJet(X3, {u(x1, x2, x3, x4), q(x1, x2, x3, x4)}) :

XX4 := FromJet(X4, {u(x1, x2, x3, x4), q(x1, x2, x3, x4)}) :

$\omega\omega$:= FromJet(ω , {u(x1, x2, x3, x4), q(x1, x2, x3, x4)}) :

> DD := {XX2, XX3, XX4} : ChrCoe := {seq(seq(seq(((a||i)||j)||k), i=1..4), j=1..4), k=1..4)} :

TGE := {} : for i to 3 do for j to 3 do TGE := {op(TGE), coeffs(expand(numer(subs(VsS2, subs(Qsyst, ToJet(Hook(convert(DirectionalCovariantDerivative(DD[i], DD[j], C),

'DGvector'), $\omega\omega$), $\{u(x1, x2, x3, x4), q(x1, x2, x3, x4)\}$, notation = jetnumbers)))),
 $\{q_{[]}, q_1\}$ } } :od:od:
 [nops(ChrCoe), nops(TGE)]

[64, 57] (3.15)

> Sconn := solve(TGE, ChrCoe) :
 > FreePar := { } :for i to nops(Sconn) do if lhs(Sconn[i]) = rhs(Sconn[i]) then FreePar
 := {op(FreePar), lhs(Sconn[i])} fi:od: FreePar
 {a144, a244, a323, a344, a411, a414, a424, a433, a434, a444}

(3.16)

> ChrCoe1 := {a111, a112, a113, a114, a121, a122, a123, a124, a131, a132, a133, a134,
 a141, a142, a143, a211, a212, a213, a214, a221, a222, a223, a224, a231, a232, a233,
 a234, a241, a242, a243, a311, a312, a313, a314, a321, a322, a324, a331, a332, a333,
 a334, a341, a342, a343, a412, a413, a414, a421, a422, a423, a431, a432, a441, a442,
 a443} :
 > Sconn := solve(TGE, ChrCoe1) :
 > FreePar := {a144, a244, a323, a344, a411, a414, a424, a433, a434, a444} :
 > CC := DGsimplify(subs(Sconn, C)) :
 > Quad1 := evalDG(add(add(diff(Eq1, u_{i,j}) · vM[i] &s vM[j], j = i..4), i = 1..4) :
 Quad2 := evalDG(add(add(diff(Eq2, u_{i,j}) · vM[i] &s vM[j], j = i..4), i = 1..4) :
 Quad3 := evalDG(add(add(diff(Eq3, u_{i,j}) · vM[i] &s vM[j], j = i..4), i = 1..4) :
 Quad1u := FromJet(Quad1, {u(x1, x2, x3, x4), q(x1, x2, x3, x4)}) : Quad2u
 := FromJet(Quad2, {u(x1, x2, x3, x4), q(x1, x2, x3, x4)}) : Quad3u
 := FromJet(Quad3, {u(x1, x2, x3, x4), q(x1, x2, x3, x4)}) : Quadu := [Quad1u,
 Quad2u, Quad3u] :
 > {seq(ContractIndices(Quad||i, ω &t ω , [[1, 1], [2, 2]]), i = 1..3) }
 {0}

(3.17)

> RNE := { } :for i to 4 do for j to 3 do RNE := {op(RNE),
 coeffs(expand(numer(simplify(subs(VsS2, subs(Qsyst,
 ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CC), ω
 &t ω , [[1, 1], [2, 2]]), {u(x1, x2, x3, x4), q(x1, x2, x3, x4)}, notation = jetnumbers))))
))), {q_[], q₁}) } } :od:od: nops(RNE)

9 (3.18)

> sRNE := solve(RNE, FreePar) :
 > RNE := { } :for i to 4 do for j to 3 do RNE := {op(RNE),
 coeffs(expand(numer(simplify(subs(VsS2, subs(Qsyst,
 ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CC), ω
 &t ω , [[1, 1], [2, 2]]), {u(x1, x2, x3, x4)}, notation = jetnumbers)))))), {q_[]}) } } :od:
 od: nops(RNE)

9 (3.19)

> sRNE := solve(RNE, FreePar) :
 > FreePar
 {a144, a244, a323, a344, a411, a414, a424, a433, a434, a444}

(3.20)

> sRNE := solve(RNE, {a144, a244, a323, a344, a411, a433}) :

```

> CT := DGsimplify( evalDG( subs( { a414 =  $\alpha_1$ , a424 =  $\alpha_2$ , a434 =  $\alpha_3$ , a444 =  $\alpha_4$  }, subs(sRNE,
    CC) ) ) ) :
> # simplify( ToJet( DGinfo( evalDG( convert( DirectionalCovariantDerivative( D_x1,
    FromJet( $\omega$ , {u(x1, x2, x3, x4)} ), CT), 'DGform') &w FromJet( $\omega$ , {u(x1, x2, x3, x4)} ) )
    , "CoefficientSet" ), {u(x1, x2, x3, x4)} , notation = jetnumbers) )
> RNE := { } :for i to 4 do for j to 3 do RNE := { op(RNE),
    coeffs( expand( numer( simplify( subs( VsS2, subs( Qsyst,
    ToJet( ContractIndices( DirectionalCovariantDerivative(vM[i], Quadu[j], CT),  $\omega$  &t  $\omega$ ,
    [[1, 1], [2, 2]] ), {u(x1, x2, x3, x4)} , notation = jetnumbers) ) ) ) ), {q[_]} ) } :od:od:
    simplify(RNE)
    {0}
    (3.21)

```

```

> # simplify( map( xx  $\rightarrow$  subs( VsS2, xx ),
    seq( ToJet( DGinfo( evalDG( convert( DirectionalCovariantDerivative( vM[i],
    FromJet( $\omega$ , {u(x1, x2, x3, x4)} ), CT), 'DGform') &w FromJet( $\omega$ , {u(x1, x2, x3, x4)} ) )
    , "CoefficientSet" ), {u(x1, x2, x3, x4)} , notation = jetnumbers), i = 1 ..4) ) )
>
> sCTa := solve( DGinfo( ContractIndices( TorsionTensor(CT), [[1, 2]]), "CoefficientSet"),
    { $\alpha_1$ ,  $\alpha_2$ ,  $\alpha_3$ ,  $\alpha_4$ } ) : CTa := DGsimplify( evalDG( subs(sCTa, CT) ) ) :
> ContractIndices( TorsionTensor(CTa), [[1, 2]])
    0 dx1
    (3.22)

```

```

> collect( numer( subs( sCTa,  $\alpha_1$  ) ), {u1,1, u1,2, u1,3, u1,4, u2,2, u3,3, u4,4} )
(4 u2 u3 - 4 u2 u4 + u32 - 6 u3 u4 + 5 u42) u1,2 + (-u22 - 4 u2 u3 + 6 u2 u4 + 4 u3 u4 - 5
u42) u1,3 + (u22 - 2 u2 u4 - u32 + 2 u3 u4) u1,4 + (-u22 u3 + u22 u4 + u2 u32 - u2 u42 - u32 u4
+ u3 u42) u2,2 + (-u22 u3 + u22 u4 + u2 u32 - u2 u42 - u32 u4 + u3 u42) u3,3 + (-u22 u3 + u22 u4
+ u2 u32 - u2 u42 - u32 u4 + u3 u42) u4,4
(3.23)

```

```

> VsS2
{ u2,3 =  $\frac{u_{1,2} - u_{1,3}}{u_2 - u_3}$ , u2,4 =  $\frac{u_{1,2} - u_{1,4}}{u_2 - u_4}$ , u3,4 =  $\frac{u_{1,3} - u_{1,4}}{u_3 - u_4}$  }
(3.24)

```

```

>
> factor( - (u3 - u4) ( (u3 - u2) + 5 (u2 - u4) ) u1,3 + u1,3 (u2 - u4) (u2 + 4 u3 - 5 u4)
    - (u2 - u3) (u2 - 2 u4 + u3) u1,3 )
    0
    (3.25)

```

```

> simplify( subs( VsS2, (u3 - u4) (u2 - u4) (u2 - u3) (u2,2 + u3,3 + u3,4 + u4,4 - 5 u2,3)
    + (u3 - u4) (u2 - u3)2 u2,3 + (u2 - u3) (u3 - u4)2 u3,4) )
u22 u3 u2,2 + u22 u3 u3,3 + u22 u3 u4,4 - u22 u4 u2,2 - u22 u4 u3,3 - u22 u4 u4,4 - u2 u32 u2,2 - u2
u32 u3,3 - u2 u32 u4,4 + u2 u42 u2,2 + u2 u42 u3,3 + u2 u42 u4,4 + u32 u4 u2,2 + u32 u4 u3,3 +
(3.26)

```

$$\begin{aligned}
& u_3^2 u_4 u_{4,4} - u_3 u_4^2 u_{2,2} - u_3 u_4^2 u_{3,3} - u_3 u_4^2 u_{4,4} + u_2^2 u_{1,3} - u_2^2 u_{1,4} - 4 u_2 u_3 u_{1,2} \\
& + 4 u_2 u_3 u_{1,3} + 4 u_2 u_4 u_{1,2} - 6 u_2 u_4 u_{1,3} + 2 u_2 u_4 u_{1,4} - u_3^2 u_{1,2} + u_3^2 u_{1,4} \\
& + 6 u_3 u_4 u_{1,2} - 4 u_3 u_4 u_{1,3} - 2 u_3 u_4 u_{1,4} - 5 u_4^2 u_{1,2} + 5 u_4^2 u_{1,3}
\end{aligned}$$

$$\begin{aligned}
& \text{factor}(\text{subs}(sCTa, \alpha_1)); \\
& - \frac{1}{3} \frac{1}{(u_3 - u_4)(u_2 - u_4)(u_2 - u_3)} (u_2^2 u_3 u_{2,2} + u_2^2 u_3 u_{3,3} + u_2^2 u_3 u_{4,4} - u_2^2 u_4 u_{2,2} - \quad (3.27) \\
& u_2^2 u_4 u_{3,3} - u_2^2 u_4 u_{4,4} - u_2 u_3^2 u_{2,2} - u_2 u_3^2 u_{3,3} - u_2 u_3^2 u_{4,4} + u_2 u_4^2 u_{2,2} + u_2 u_4^2 u_{3,3} \\
& + u_2 u_4^2 u_{4,4} + u_3^2 u_4 u_{2,2} + u_3^2 u_4 u_{3,3} + u_3^2 u_4 u_{4,4} - u_3 u_4^2 u_{2,2} - u_3 u_4^2 u_{3,3} - u_3 u_4^2 u_{4,4} \\
& + u_2^2 u_{1,3} - u_2^2 u_{1,4} - 4 u_2 u_3 u_{1,2} + 4 u_2 u_3 u_{1,3} + 4 u_2 u_4 u_{1,2} - 6 u_2 u_4 u_{1,3} \\
& + 2 u_2 u_4 u_{1,4} - u_3^2 u_{1,2} + u_3^2 u_{1,4} + 6 u_3 u_4 u_{1,2} - 4 u_3 u_4 u_{1,3} - 2 u_3 u_4 u_{1,4} - 5 u_4^2 u_{1,2} \\
& + 5 u_4^2 u_{1,3})
\end{aligned}$$

$$\begin{aligned}
& \text{> } sCTa2 := \left\{ \alpha_1 = -\frac{1}{3} (u_{2,2} + u_{3,3} + u_{3,4} + u_{4,4} - 5 u_{2,3}) - \frac{(u_2 - u_3)}{3(u_2 - u_4)} u_{2,3} \right. \\
& \quad \left. - \frac{(u_3 - u_4)}{3(u_2 - u_4)} u_{3,4}, \right. \\
& \alpha_2 = \frac{(2 u_{2,2} - 3 u_{2,4} + u_{4,4})}{3(u_2 - u_4)} - \frac{(u_{2,2} - u_{3,3})}{3(u_2 - u_3)} - \frac{(u_{2,3} - u_{2,4})}{3(u_3 - u_4)}, \\
& \alpha_3 = \frac{(2 u_{3,3} - 3 u_{3,4} + u_{4,4})}{3(u_3 - u_4)} - \frac{(u_{2,2} - u_{3,3})}{3(u_2 - u_3)} - \frac{(u_{2,3} - u_{3,4})}{3(u_2 - u_4)}, \\
& \alpha_4 = \left. \frac{2 u_{2,3} - u_{2,2} - 2 u_{3,4} + u_{4,4}}{3(u_2 - u_4)} - \frac{(u_{3,3} - u_{4,4})}{3(u_3 - u_4)} \right\}; \\
& \text{> } \text{simplify}(\{ \text{seq}(\text{subs}(VsS2, \text{subs}(sCTa, \alpha[i])) - \text{subs}(sCTa2, \alpha[i])), i = 1..4) \}) \\
& \quad \{0\} \quad (3.28)
\end{aligned}$$

$$\begin{aligned}
& \text{> } \text{factor}(\text{DirectionalCovariantDerivative}(D_{x1}, D_{x1}, CTa)) \\
& \quad (3.29)
\end{aligned}$$

Thus there exist GL2-connections, with respect to which the null Lax surfaces are totally geodesic. They depend on 4 arbitrary functions.
The parameters can be fixed uniquely by the condition $\text{Trace}(\text{Torsion})=0$.

$$\begin{aligned}
& \text{> } RNE := \{ \} : \text{for } i \text{ to } 4 \text{ do for } j \text{ to } 3 \text{ do } RNE := \{ \text{op}(RNE), \\
& \quad \text{coeffs}(\text{expand}(\text{numer}(\text{simplify}(\text{subs}(VsS2, \text{subs}(Qsyst, \\
& \quad \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(vM[i], \text{Quadu}[j], CT), \omega \& t \omega, \\
& \quad [[1, 1], [2, 2]]), \{u(x1, x2, x3, x4)\}, \text{notation} = \text{jetnumbers}))) \}, \{q_{[j]}\}) \} : \text{od:od:} \\
& \quad \text{simplify}(RNE) \\
& \quad \{0\} \quad (3.30)
\end{aligned}$$

$$\begin{aligned}
& \text{> } RNE := \{ \} : \text{for } i \text{ to } 4 \text{ do for } j \text{ to } 3 \text{ do } RNE := \{ \text{op}(RNE), \\
& \quad \text{coeffs}(\text{expand}(\text{numer}(\text{simplify}(\text{subs}(VsS2, \text{subs}(Qsyst,
\end{aligned}$$

$ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CTA), \omega$
 $\&t \omega, [[1, 1], [2, 2]], \{u(x1, x2, x3, x4)\}, notation = jetnumbers))))) , \{q_{[]}\} \} : od:$
 $od: simplify(RNE)$
 $\{0\}$ (3.31)

$> RNE := \{ \} : for i to 4 do for j to 3 do RNE := \{ op(RNE),$
 $coeffs(expand(numer(simplify(subs(Qsyst,$
 $ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CT), \omega \&t \omega,$
 $[[1, 1], [2, 2]]), \{u(x1, x2, x3, x4)\}, notation = jetnumbers))))) , \{q_{[]}\} \} : od: od:$
 $simplify(RNE)$

$$\begin{aligned}
 &\{0, -u_2 u_{2,3} + u_3 u_{2,3} + u_{1,2} - u_{1,3}, u_{2,3} u_2 - u_{2,3} u_3 - u_{1,2} + u_{1,3}, -u_2 u_{2,4} + u_4 u_{2,4} \\
 &+ u_{1,2} - u_{1,4}, u_{2,4} u_2 - u_{2,4} u_4 - u_{1,2} + u_{1,4}, -u_3 u_{3,4} + u_4 u_{3,4} + u_{1,3} - u_{1,4}, u_{3,4} u_3 \\
 &- u_{3,4} u_4 - u_{1,3} + u_{1,4}, u_2^2 u_{2,3} - u_2^2 u_{2,4} - u_2 u_3 u_{2,3} + u_2 u_3 u_{2,4} - u_2 u_4 u_{2,3} \\
 &+ u_2 u_4 u_{2,4} + u_3 u_4 u_{2,3} - u_3 u_4 u_{2,4} + u_2 u_{1,3} - u_2 u_{1,4} - u_3 u_{1,2} + u_3 u_{1,4} + u_4 u_{1,2} \\
 &- u_4 u_{1,3}, u_2 u_3 u_{2,3} - u_2 u_3 u_{3,4} - u_2 u_4 u_{2,3} + u_2 u_4 u_{3,4} - u_3^2 u_{2,3} + u_3^2 u_{3,4} \\
 &+ u_3 u_4 u_{2,3} - u_3 u_4 u_{3,4} + u_2 u_{1,3} - u_2 u_{1,4} - u_3 u_{1,2} + u_3 u_{1,4} + u_4 u_{1,2} - u_4 u_{1,3}, \\
 &u_2 u_3 u_{2,4} - u_2 u_3 u_{3,4} - u_2 u_4 u_{2,4} + u_2 u_4 u_{3,4} - u_3 u_4 u_{2,4} + u_3 u_4 u_{3,4} + u_4^2 u_{2,4} - \\
 &u_4^2 u_{3,4} + u_2 u_{1,3} - u_2 u_{1,4} - u_3 u_{1,2} + u_3 u_{1,4} + u_4 u_{1,2} - u_4 u_{1,3}\}
 \end{aligned} \quad (3.32)$$

$> solve(\%, \{u_{2,3}, u_{2,4}, u_{3,4}\})$

$$\left\{ u_{2,3} = \frac{u_{1,2} - u_{1,3}}{u_2 - u_3}, u_{2,4} = \frac{u_{1,2} - u_{1,4}}{u_2 - u_4}, u_{3,4} = \frac{u_{1,3} - u_{1,4}}{u_3 - u_4} \right\} \quad (3.33)$$

$> VsS2$

$$\left\{ u_{2,3} = \frac{u_{1,2} - u_{1,3}}{u_2 - u_3}, u_{2,4} = \frac{u_{1,2} - u_{1,4}}{u_2 - u_4}, u_{3,4} = \frac{u_{1,3} - u_{1,4}}{u_3 - u_4} \right\} \quad (3.34)$$

$> RicCTA := RicciTensor(FromJet(CTA, \{u(x1, x2, x3, x4)\})) : RicCTa := ToJet(RicCTA,$
 $\{u(x1, x2, x3, x4)\}, notation = jetnumbers) :$

$> RicCTaSkew := SymmetrizeIndices(RicCTa, [1, 2], "SkewSymmetric") :$
 $RicCTaSymm := SymmetrizeIndices(RicCTa, [1, 2], "Symmetric") :$

$>$

$> \omega\omega := evalDG((q_{[]} - u_2) \cdot (q_{[]} - u_3) \cdot (q_{[]} - u_4) \cdot \omega)$

$$\omega\omega := (-u_2 + q_{[]}) (-u_3 + q_{[]}) (-u_4 + q_{[]}) dx1 + (-u_3 + q_{[]}) (-u_4 + q_{[]}) dx2 + (-u_2 + q_{[]}) (-u_4 + q_{[]}) dx3 + (-u_2 + q_{[]}) (-u_3 + q_{[]}) dx4 \quad (3.35)$$

$> \omega0 := -\omega\omega : \omega0b := DGsimplify(evalDG(subs(q_{[]} = 0, \omega0))) ;$
 $\omega1a := evalDG\left(\frac{1}{\epsilon} \cdot (subs(q_{[]} = q_{[]} + \epsilon, \omega0) - \omega0)\right) : \omega1 := evalDG(subs(\epsilon = 0,$
 $\omega1a)) : \omega1b := DGsimplify(evalDG(subs(q_{[]} = 0, \omega1))) ;$

$$\begin{aligned}
\omega 2a &:= evalDG\left(\frac{1}{\varepsilon} \cdot (subs(q_{[1]} = q_{[1]} + \varepsilon, \omega 1) - \omega 1)\right) : \omega 2 := evalDG(subs(\varepsilon = 0, \\
&\omega 2a)) : \omega 2b := DGsimplify\left(evalDG\left(\frac{1}{2} \cdot subs(q_{[1]} = 0, \omega 2)\right)\right); \\
\omega 3a &:= evalDG\left(\frac{1}{\varepsilon} \cdot (subs(q_{[1]} = q_{[1]} + \varepsilon, \omega 2) - \omega 2)\right) : \omega 3 := evalDG(subs(\varepsilon = 0, \\
&\omega 3a)) : \omega 3b := DGsimplify\left(evalDG\left(\frac{1}{6} \cdot subs(q_{[1]} = 0, \omega 3)\right)\right) \\
\omega 0b &:= u_4 u_3 u_2 dx1 - (u_3 u_4 dx2) - (u_2 u_4 dx3) - (u_2 u_3 dx4) \\
\omega 1b &:= -((u_2 u_3 + u_2 u_4 + u_3 u_4) dx1) + (u_3 + u_4) dx2 + (u_2 + u_4) dx3 + (u_2 + u_3) dx4 \\
\omega 2b &:= (u_2 + u_3 + u_4) dx1 - dx2 - dx3 - dx4 \\
\omega 3b &:= -dx1
\end{aligned} \tag{3.36}$$

$$\begin{aligned}
> evalDG(\omega 0 - \omega 0b - \omega 1b \cdot q_{[1]} - \omega 2b \cdot q_{[1]}^2 - \omega 3b \cdot q_{[1]}^3) \\
0 dx1
\end{aligned} \tag{3.37}$$

$$\begin{aligned}
> Qua\omega &:= evalDG([\omega 0b \&s \omega 0b, \omega 0b \&s \omega 1b, 2 \omega 0b \&s \omega 2b + \omega 1b \&s \omega 1b, \omega 0b \&s \omega 3b \\
&+ \omega 1b \&s \omega 2b, 2 \omega 1b \&s \omega 3b + \omega 2b \&s \omega 2b, \omega 2b \&s \omega 3b, \omega 3b \&s \omega 3b]) : tmp\Omega \\
&:= \{ \} : \text{for } i \text{ to } nops(Qua\omega) \text{ do for } j \text{ to } 3 \text{ do } tmp\Omega := \{op(tmp\Omega), \\
&ContractIndices(Quad||j, Qua\omega[i], [[1, 1], [2, 2]])\} : \text{od:od: } tmp\Omega \\
\{0\}
\end{aligned} \tag{3.38}$$

$$\begin{aligned}
> \# eliminate(simplify(subs(VsS3, DGinfo(evalDG(RicCTaSymm - add(\Lambda[i] \cdot Qua\omega[i], i = 1 \\
..7))), "CoefficientSet"))), \{seq(\Lambda[i], i = 1 ..7)\}) [2]
\end{aligned}$$

Thus the condition that $\text{sym}(\text{Ric}(\nabla))$ belongs to the span of quadrics defining the GL2 structure is non-vacous. It is another differential constraint, giving exact solutions.

But in fact, from what is computed below it follows that this is the same condition as $\text{sym}(\text{Ric})=0$.

$$\begin{aligned}
> Vt0 &:= Annihilator([\omega 0, \omega 1, \omega 2]) [1]; Vt1 := Annihilator([\omega 0, \omega 1]) [2]; Vt2 \\
&:= Annihilator([\omega 0]) [3] \\
Vt0 &:= \\
Vt1 &:= -\left(\frac{u_2 - u_3}{(-u_3 + q_{[1]})^2} D_{x1}\right) - \left(\frac{(-u_2 + q_{[1]})^2}{(-u_3 + q_{[1]})^2} D_{x2}\right) + D_{x3} \\
Vt2 &:= -\left(\frac{1}{-u_2 + q_{[1]}} D_{x1}\right) + D_{x2}
\end{aligned} \tag{3.39}$$

$$\begin{aligned}
> Vt01 &:= ContractIndices(TorsionTensor(CT), evalDG(Vt0 \&t Vt1), [[2, 1], [3, 2]]) : \\
KrCond &:= DGinfo(evalDG(convert(Vt01, DGvector) - Hook(convert(Vt01, DGvector), dx4) \\
&\cdot Vt0), "CoefficientSet") : \\
sCTa3 &:= solve(map(Q \rightarrow coeffs(expand(numer(Q)), \{q_{[1]}\}), KrCond), \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}) : \\
> simplify(\{seq(subs(VsS2, subs(sCTa, \alpha[i])) - subs(sCTa2, \alpha[i])), i = 1 ..4), \\
&seq(subs(VsS2, subs(sCTa, \alpha[i])) - subs(sCTa3, \alpha[i])), i = 1 ..4\}) \\
\{0\}
\end{aligned} \tag{3.40}$$

Thus there exist GL2-connections, with respect to which the null Lax surfaces are totally geodesic. They depend on 4 arbitrary functions.

The parameters can be fixed uniquely by the condition $\text{Trace}(\text{Torsion})=0$. This condition is

equivalent to that one of Krynsky.

$$\text{> simplify(subs(VsS3, ContractIndices(RicCTaSymm, evalDG(Vt0 \&t Vt0), [[1, 1], [2, 2]])))}$$

0

(3.41)

$$\text{> simplify(subs(VsS3, ContractIndices(RicCTaSymm, evalDG(Vt0 \&t Vt1), [[1, 1], [2, 2]])))}$$

0

(3.42)

$$\begin{aligned} \text{> QuT} &:= \text{evalDG}(\text{add}(\text{add}(\text{xi}[i, j] \cdot dM[j], j = i..4), i = 1..4)) : \\ \text{SuT} &:= \text{solve}(\{\text{coeffs}(\text{expand}(\text{numer}(\text{ContractIndices}(\text{QuT}, \text{Vt0} \&t \text{Vt0}, [[1, 1], [2, 2]]))), \\ &\quad \{q[\]\}\}, \{\xi_{1,1}, \xi_{1,2}, \xi_{1,3}, \xi_{1,4}, \xi_{2,2}, \xi_{2,3}, \xi_{2,4}\}) : \\ \text{TuT} &:= \text{DGsimplify}(\text{evalDG}(\text{subs}(\{\xi_{3,3} = \Lambda_1, \xi_{3,4} = \Lambda_2, \xi_{4,4} = \Lambda_3\}, \text{subs}(\text{SuT}, \text{QuT})))) : \end{aligned}$$

> # RicCTaSymm

$$\begin{aligned} \text{> eliminate}(\text{DGinfo}(\text{DGsimplify}(\text{evalDG}(\text{subs}(\text{VsS3}, \text{TuT} - \text{RicCTaSymm}}))), \\ \text{"CoefficientSet"}, \{\Lambda_1, \Lambda_2, \Lambda_3\})[2] \end{aligned}$$

{ }

(3.43)

This computation has shown that symmetrized Ricci tensor of the canonical GL2-connection is a section of the bundle of quadrics annihilating the dual (tangent) to the canonical (cotangent) rational normal curve.

>

Let us compute the formulae for the Christoffel coefficients of the canonical connection.

$$\text{> VsS2a} := \text{solve}\left(\left\{u_{2,3} = \frac{u_{1,2} - u_{1,3}}{u_2 - u_3}, u_{2,4} = \frac{u_{1,2} - u_{1,4}}{u_2 - u_4}, u_{3,4} = \frac{u_{1,3} - u_{1,4}}{u_3 - u_4}\right\}, \{u_{1,3}, u_{1,4}, u_{3,4}\}\right);$$

$$\text{VsS2b} := \text{solve}\left(\left\{u_{2,3} = \frac{u_{1,2} - u_{1,3}}{u_2 - u_3}, u_{2,4} = \frac{u_{1,2} - u_{1,4}}{u_2 - u_4}, u_{3,4} = \frac{u_{1,3} - u_{1,4}}{u_3 - u_4}\right\}, \{u_{1,2}, u_{2,4}, u_{1,4}\}\right);$$

$$\text{VsS2c} := \text{solve}\left(\left\{u_{2,3} = \frac{u_{1,2} - u_{1,3}}{u_2 - u_3}, u_{2,4} = \frac{u_{1,2} - u_{1,4}}{u_2 - u_4}, u_{3,4} = \frac{u_{1,3} - u_{1,4}}{u_3 - u_4}\right\}, \{u_{1,2}, u_{1,3}, u_{2,3}\}\right);$$

$$\begin{aligned} \text{VsS2a} &:= \left\{u_{1,3} = -u_2 u_{2,3} + u_3 u_{2,3} + u_{1,2} u_{1,4} = -u_2 u_{2,4} + u_4 u_{2,4} + u_{1,2} u_{3,4} = \right. \\ &\quad \left. - \frac{u_2 u_{2,3} - u_2 u_{2,4} - u_3 u_{2,3} + u_4 u_{2,4}}{u_3 - u_4}\right\} \end{aligned}$$

$$\begin{aligned} \text{VsS2b} &:= \left\{u_{1,2} = u_2 u_{2,3} - u_3 u_{2,3} + u_{1,3} u_{1,4} = -u_3 u_{3,4} + u_4 u_{3,4} + u_{1,3} u_{2,4} \right. \\ &\quad \left. = \frac{u_2 u_{2,3} - u_3 u_{2,3} + u_3 u_{3,4} - u_4 u_{3,4}}{u_2 - u_4}\right\} \end{aligned}$$

$$VsS2c := \left\{ \begin{aligned} u_{1,2} &= u_2 u_{2,4} - u_4 u_{2,4} + u_{1,4}, u_{1,3} = u_3 u_{3,4} - u_4 u_{3,4} + u_{1,4}, u_{2,3} \\ &= \frac{u_2 u_{2,4} - u_3 u_{3,4} - u_4 u_{2,4} + u_4 u_{3,4}}{u_2 - u_3} \end{aligned} \right\} \quad (3.44)$$

$$> R_a := u_{2,2} + u_{3,3} + u_{4,4} :$$

$$R_b := u_{2,3} + u_{2,4} + u_{3,4} :$$

$$R_c := \frac{u_2}{(u_2 - u_3)(u_2 - u_4)} \cdot u_{3,4} + \frac{u_3}{(u_3 - u_2)(u_3 - u_4)} \cdot u_{2,4} + \frac{u_4}{(u_4 - u_2)(u_4 - u_3)} \cdot u_{2,3} :$$

$$\varPi_{11} := \frac{1}{3} \cdot (2 R_b - R_a) :$$

$$\varPi_{12} := R_c : \varPi_{13} := R_c : \varPi_{14} := R_c :$$

$$\begin{aligned} \varPi_{21} := & -\frac{1}{3} \cdot \left(\frac{(u_{2,2} - u_{3,3})}{(u_2 - u_3)} + \frac{(u_{2,2} - u_{4,4})}{(u_2 - u_4)} + 5 R_c \right) : \varPi_{31} := -\frac{1}{3} \cdot \left(\frac{(u_{3,3} - u_{2,2})}{(u_3 - u_2)} \right. \\ & + \frac{(u_{3,3} - u_{4,4})}{(u_3 - u_4)} + 5 R_c \Big) : \varPi_{41} := -\frac{1}{3} \cdot \left(\frac{(u_{4,4} - u_{2,2})}{(u_4 - u_2)} + \frac{(u_{4,4} - u_{3,3})}{(u_4 - u_3)} \right. \\ & \left. + 5 R_c \right) : \end{aligned}$$

$$\varGamma_{12} := 2 u_{3,4} - \frac{1}{3} \cdot (R_a + R_b) : \varGamma_{313} := 2 u_{2,4} - \frac{1}{3} \cdot (R_a + R_b) : \varGamma_{414} := 2 u_{2,3} - \frac{1}{3} \cdot (R_a + R_b) :$$

$$\begin{aligned} \varGamma_{221} := & -u_{2,2} + \frac{(u_2 - u_4)}{(u_3 - u_4)} \cdot u_{2,3} + \frac{(u_2 - u_3)}{(u_4 - u_3)} \cdot u_{2,4} : \varGamma_{331} := -u_{3,3} + \frac{(u_3 - u_4)}{(u_2 - u_4)} \cdot u_{2,3} \\ & + \frac{(u_3 - u_2)}{(u_4 - u_2)} \cdot u_{3,4} : \varGamma_{441} := -u_{4,4} + \frac{(u_4 - u_3)}{(u_2 - u_3)} \cdot u_{2,4} + \frac{(u_4 - u_2)}{(u_3 - u_2)} \cdot u_{3,4} : \end{aligned}$$

$$\begin{aligned} \varGamma_{222} := & -\frac{1}{3} \cdot \left(\frac{(u_{2,2} - u_{3,3})}{(u_2 - u_3)} + \frac{(u_{2,2} - u_{4,4})}{(u_2 - u_4)} + 2 R_c \right) : \varGamma_{333} := -\frac{1}{3} \cdot \left(\frac{(u_{3,3} - u_{2,2})}{(u_3 - u_2)} \right. \\ & + \frac{(u_{3,3} - u_{4,4})}{(u_3 - u_4)} + 2 R_c \Big) : \varGamma_{444} := -\frac{1}{3} \cdot \left(\frac{(u_{4,4} - u_{2,2})}{(u_4 - u_2)} + \frac{(u_{4,4} - u_{3,3})}{(u_4 - u_3)} \right. \\ & \left. + 2 R_c \right) : \end{aligned}$$

$$\begin{aligned} \varGamma_{223} := & \frac{(R_b - u_{2,2} - 2 u_{3,4})}{(u_2 - u_3)} : \varGamma_{323} := \frac{(2 u_{2,2} + u_{3,3} - 3 u_{2,3} - u_{2,4} + u_{3,4})}{3 (u_2 - u_3)} \\ & - \frac{(u_{2,2} - u_{4,4})}{3 (u_2 - u_4)} : \end{aligned}$$

$$\varGamma_{232} := \frac{(2 u_{3,3} + u_{2,2} - 3 u_{2,3} - u_{3,4} + u_{2,4})}{3 (u_3 - u_2)} - \frac{(u_{3,3} - u_{4,4})}{3 (u_3 - u_4)} : \varGamma_{332}$$

$$\begin{aligned}
&:= \frac{(R_b - u_{3,3} - 2 u_{2,4})}{(u_3 - u_2)} : \\
\Gamma_{224} &:= \frac{(R_b - u_{2,2} - 2 u_{3,4})}{(u_2 - u_4)} : \Gamma_{424} := \frac{(2 u_{2,2} + u_{4,4} - 3 u_{2,4} - u_{2,3} + u_{3,4})}{3(u_2 - u_4)} \\
&- \frac{(u_{2,2} - u_{3,3})}{3(u_2 - u_3)} : \\
\Gamma_{242} &:= \frac{(2 u_{4,4} + u_{2,2} - 3 u_{2,4} - u_{3,4} + u_{2,3})}{3(u_4 - u_2)} - \frac{(u_{4,4} - u_{3,3})}{3(u_4 - u_3)} : \Gamma_{442} \\
&:= \frac{(R_b - u_{4,4} - 2 u_{2,3})}{(u_4 - u_2)} : \\
\Gamma_{334} &:= \frac{(R_b - u_{3,3} - 2 u_{2,4})}{(u_3 - u_4)} : \Gamma_{434} := \frac{(2 u_{3,3} + u_{4,4} - 3 u_{3,4} - u_{2,3} + u_{2,4})}{3(u_3 - u_4)} \\
&- \frac{(u_{3,3} - u_{2,2})}{3(u_3 - u_2)} : \\
\Gamma_{343} &:= \frac{(2 u_{4,4} + u_{3,3} - 3 u_{3,4} - u_{2,4} + u_{2,3})}{3(u_4 - u_3)} - \frac{(u_{4,4} - u_{2,2})}{3(u_4 - u_2)} : \Gamma_{443} \\
&:= \frac{(R_b - u_{4,4} - 2 u_{2,3})}{(u_4 - u_3)} :
\end{aligned}$$

> simplify(evalDG(subs(VsS2, convert(DirectionalCovariantDerivative(D_x1, D_x1, CTa),
 DGvector) - $\Gamma_{111} \cdot D_{x1}$)));
 simplify(evalDG(subs(VsS2, convert(DirectionalCovariantDerivative(D_x1, D_x2, CTa),
 DGvector) - $\Gamma_{112} \cdot D_{x1}$ - $\Gamma_{212} \cdot D_{x2}$)));
 simplify(evalDG(subs(VsS2, convert(DirectionalCovariantDerivative(D_x1, D_x3, CTa),
 DGvector) - $\Gamma_{113} \cdot D_{x1}$ - $\Gamma_{313} \cdot D_{x3}$)));
 simplify(evalDG(subs(VsS2, convert(DirectionalCovariantDerivative(D_x1, D_x4, CTa),
 DGvector) - $\Gamma_{114} \cdot D_{x1}$ - $\Gamma_{414} \cdot D_{x4}$)));
 simplify(evalDG(subs(VsS2, convert(DirectionalCovariantDerivative(D_x2, D_x1, CTa),
 DGvector) - $\Gamma_{121} \cdot D_{x1}$ - $\Gamma_{221} \cdot D_{x2}$)));
 simplify(evalDG(subs(VsS2, convert(DirectionalCovariantDerivative(D_x3, D_x1, CTa),
 DGvector) - $\Gamma_{131} \cdot D_{x1}$ - $\Gamma_{331} \cdot D_{x3}$)));
 simplify(evalDG(subs(VsS2, convert(DirectionalCovariantDerivative(D_x4, D_x1, CTa),
 DGvector) - $\Gamma_{141} \cdot D_{x1}$ - $\Gamma_{441} \cdot D_{x4}$)));
 simplify(evalDG(subs(VsS2, convert(DirectionalCovariantDerivative(D_x2, D_x2, CTa),
 DGvector) - $\Gamma_{222} \cdot D_{x2}$)));
 simplify(evalDG(subs(VsS2, convert(DirectionalCovariantDerivative(D_x3, D_x3, CTa),
 DGvector) - $\Gamma_{333} \cdot D_{x3}$)));
 simplify(evalDG(subs(VsS2, convert(DirectionalCovariantDerivative(D_x4, D_x4, CTa),
 DGvector) - $\Gamma_{444} \cdot D_{x4}$)));
 simplify(evalDG(subs(VsS2, convert(DirectionalCovariantDerivative(D_x2, D_x3, CTa),
 DGvector) - $\Gamma_{223} \cdot D_{x2}$ - $\Gamma_{323} \cdot D_{x3}$)));

$$\begin{aligned}
& \text{seq}(\text{seq}(\text{simplify}(\text{subs}(\text{VsS2}, \text{subs}(\text{Qsyst}, \\
& \quad \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(\text{vM}[i], \text{Quadu}[j], \text{CsS}), \omega \\
& \quad \& \omega, [[1, 1], [2, 2]]), \{u(x1, x2, x3, x4)\}, \text{notation} = \text{jetnumbers}))\)), i = 1..4), j = 1 \\
& \quad \dots) \\
& \quad 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0
\end{aligned} \tag{3.48}$$

$$\begin{aligned}
& \text{VsS2} \\
& \left\{ u_{2,3} = \frac{u_{1,2} - u_{1,3}}{u_2 - u_3}, u_{2,4} = \frac{u_{1,2} - u_{1,4}}{u_2 - u_4}, u_{3,4} = \frac{u_{1,3} - u_{1,4}}{u_3 - u_4} \right\}
\end{aligned} \tag{3.49}$$

$$\begin{aligned}
& R_a := u_{2,2} + u_{3,3} + u_{4,4} : \\
& R_b := u_{2,3} + u_{2,4} + u_{3,4} : \\
& R_c := \frac{u_2}{(u_2 - u_3)(u_2 - u_4)} \cdot u_{3,4} + \frac{u_3}{(u_3 - u_2)(u_3 - u_4)} \cdot u_{2,4} + \frac{u_4}{(u_4 - u_2)(u_4 - u_3)} \\
& \quad \cdot u_{2,3} :
\end{aligned}$$

$$\begin{aligned}
& \gamma_{23} := \frac{1}{u_2 - u_3} : \gamma_{32} := \frac{1}{u_3 - u_2} : \gamma_{24} := \frac{1}{u_2 - u_4} : \gamma_{42} := \frac{1}{u_4 - u_2} : \gamma_{34} \\
& \quad := \frac{1}{u_3 - u_4} : \gamma_{43} := \frac{1}{u_4 - u_3} :
\end{aligned}$$

$$\begin{aligned}
& R_d := (\gamma_{23} + \gamma_{24}) \cdot u_{1,2} + (\gamma_{32} + \gamma_{34}) \cdot u_{1,3} + (\gamma_{42} + \gamma_{43}) \cdot u_{1,4} : \\
& R_e := ((\gamma_{23} + \gamma_{24}) \cdot u_{2,2} + (\gamma_{32} + \gamma_{34}) \cdot u_{3,3} + (\gamma_{42} + \gamma_{43}) \cdot u_{4,4}) : \\
& R_f := (\gamma_{23} \gamma_{24} u_{2,2} + \gamma_{32} \gamma_{34} u_{3,3} + \gamma_{42} \gamma_{43} u_{4,4}) :
\end{aligned}$$

$$\begin{aligned}
& \Pi_{11} := \left(\frac{2}{3} \cdot R_d - \frac{2}{9} \cdot R_a - \frac{1}{9} \cdot \left(\left(\frac{\gamma_{24}}{\gamma_{23}} + \frac{\gamma_{23}}{\gamma_{24}} \right) \cdot u_{2,2} + \left(\frac{\gamma_{32}}{\gamma_{34}} + \frac{\gamma_{34}}{\gamma_{32}} \right) \cdot u_{3,3} \right. \right. \\
& \quad \left. \left. + \left(\frac{\gamma_{42}}{\gamma_{43}} + \frac{\gamma_{43}}{\gamma_{42}} \right) \cdot u_{4,4} \right) \right) : \\
& \Gamma_{211} := \frac{1}{9} (u_2 - u_3)(u_2 - u_4) (\gamma_{34} \cdot (u_{3,3} - 4 u_{2,3}) + \gamma_{43} \cdot (u_{4,4} - 4 u_{2,4})) : \\
& \Gamma_{311} := \frac{1}{9} (u_3 - u_2)(u_3 - u_4) (\gamma_{24} \cdot (u_{2,2} - 4 u_{2,3}) + \gamma_{42} \cdot (u_{4,4} - 4 u_{3,4})) : \\
& \Gamma_{411} := \frac{1}{9} (u_4 - u_2)(u_4 - u_3) (\gamma_{23} \cdot (u_{2,2} - 4 u_{2,4}) + \gamma_{32} \cdot (u_{3,3} - 4 u_{3,4})) : \\
& \Pi_{12} := -\frac{1}{9} \cdot R_e + \frac{1}{3} \cdot (\gamma_{34} \cdot u_{2,3} + \gamma_{43} \cdot u_{2,4}) : \Pi_{13} := -\frac{1}{9} \cdot R_e + \frac{1}{3} \cdot (\gamma_{24} \cdot u_{2,3} + \gamma_{42} \\
& \quad \cdot u_{3,4}) : \Pi_{14} := -\frac{1}{9} \cdot R_e + \frac{1}{3} \cdot (\gamma_{23} \cdot u_{2,4} + \gamma_{32} \cdot u_{3,4}) : \\
& \Gamma_{212} := \left(-\frac{4}{9} \cdot u_{2,2} - \frac{2}{9} \cdot R_a + \frac{1}{9} \frac{\gamma_{34}}{\gamma_{32}} \cdot (u_{3,3} - u_{2,3}) + \frac{1}{9} \frac{\gamma_{43}}{\gamma_{42}} \cdot (u_{4,4} - u_{2,4}) + \frac{5}{9} \right. \\
& \quad \left. \cdot (u_{2,3} + u_{2,4}) \right) : \\
& \Gamma_{313} := \left(-\frac{4}{9} \cdot u_{3,3} - \frac{2}{9} \cdot R_a + \frac{1}{9} \frac{\gamma_{24}}{\gamma_{23}} \cdot (u_{2,2} - u_{2,3}) + \frac{1}{9} \frac{\gamma_{42}}{\gamma_{43}} \cdot (u_{4,4} - u_{3,4}) + \frac{5}{9} \right.
\end{aligned}$$

$$\begin{aligned}
& \cdot (u_{2,3} + u_{3,4}) \Big) : \\
\Gamma_{414} &:= \left(-\frac{4}{9} \cdot u_{4,4} - \frac{2}{9} \cdot R_a + \frac{1}{9} \frac{\gamma_{23}}{\gamma_{24}} \cdot (u_{2,2} - u_{2,4}) + \frac{1}{9} \frac{\gamma_{32}}{\gamma_{34}} \cdot (u_{3,3} - u_{3,4}) + \frac{5}{9} \right. \\
& \quad \left. \cdot (u_{2,4} + u_{3,4}) \right) : \\
\Gamma_{312} &:= \frac{1}{9} \cdot \frac{\gamma_{42}}{\gamma_{43}} \cdot (u_{2,2} - 4 u_{2,3} + 4 u_{3,4} - u_{4,4}) : \Gamma_{412} := \frac{1}{9} \cdot \frac{\gamma_{32}}{\gamma_{34}} \cdot (u_{2,2} - 4 u_{2,4} \\
& \quad + 4 u_{3,4} - u_{3,3}) : \\
\Gamma_{213} &:= \frac{1}{9} \cdot \frac{\gamma_{43}}{\gamma_{42}} \cdot (u_{3,3} - 4 u_{2,3} + 4 u_{2,4} - u_{4,4}) : \Gamma_{413} := \frac{1}{9} \cdot \frac{\gamma_{23}}{\gamma_{24}} \cdot (u_{3,3} - 4 u_{3,4} \\
& \quad + 4 u_{2,4} - u_{2,2}) : \\
\Gamma_{214} &:= \frac{1}{9} \cdot \frac{\gamma_{34}}{\gamma_{32}} \cdot (u_{4,4} - 4 u_{2,4} + 4 u_{2,3} - u_{3,3}) : \Gamma_{314} := \frac{1}{9} \cdot \frac{\gamma_{24}}{\gamma_{23}} \cdot (u_{4,4} - 4 u_{3,4} \\
& \quad + 4 u_{2,3} - u_{2,2}) : \\
\Gamma_{122} &:= -\frac{1}{9} \cdot R_f : \Gamma_{133} := -\frac{1}{9} \cdot R_f : \Gamma_{144} := -\frac{1}{9} \cdot R_f : \\
\Gamma_{222} &:= -\frac{1}{3} \cdot R_e + \frac{1}{9} \cdot \left(\frac{\gamma_{23}}{\gamma_{24}} + 2 + \frac{\gamma_{24}}{\gamma_{23}} \right) \cdot (\gamma_{34} \cdot u_{3,3} + \gamma_{43} \cdot u_{4,4}) + \frac{2}{9} \cdot \left(\gamma_{34} - \gamma_{23} \right. \\
& \quad \left. + \gamma_{24} - \frac{\gamma_{23} \cdot \gamma_{24}}{\gamma_{34}} \right) \cdot (u_{2,3} - u_{2,4}) : \\
\Gamma_{333} &:= -\frac{1}{3} \cdot R_e + \frac{1}{9} \cdot \left(\frac{\gamma_{32}}{\gamma_{34}} + 2 + \frac{\gamma_{34}}{\gamma_{32}} \right) \cdot (\gamma_{24} \cdot u_{2,2} + \gamma_{42} \cdot u_{4,4}) + \frac{2}{9} \cdot \left(\gamma_{24} - \gamma_{32} \right. \\
& \quad \left. + \gamma_{34} - \frac{\gamma_{32} \cdot \gamma_{34}}{\gamma_{24}} \right) \cdot (u_{2,3} - u_{3,4}) : \\
\Gamma_{444} &:= -\frac{1}{3} \cdot R_e + \frac{1}{9} \cdot \left(\frac{\gamma_{42}}{\gamma_{43}} + 2 + \frac{\gamma_{43}}{\gamma_{42}} \right) \cdot (\gamma_{23} \cdot u_{2,2} + \gamma_{32} \cdot u_{3,3}) + \frac{2}{9} \cdot \left(\gamma_{23} - \gamma_{42} \right. \\
& \quad \left. + \gamma_{43} - \frac{\gamma_{42} \cdot \gamma_{43}}{\gamma_{23}} \right) \cdot (u_{2,4} - u_{3,4}) : \\
\Gamma_{322} &:= -\frac{\gamma_{23}}{9} \cdot \frac{\gamma_{42}}{\gamma_{43}} \cdot (u_{2,2} - 4 u_{2,3} + 4 u_{3,4} - u_{4,4}) : \Gamma_{422} := -\frac{\gamma_{24}}{9} \cdot \frac{\gamma_{32}}{\gamma_{34}} \cdot (u_{2,2} \\
& \quad - 4 u_{2,4} + 4 u_{3,4} - u_{3,3}) : \\
\Gamma_{233} &:= -\frac{\gamma_{32}}{9} \cdot \frac{\gamma_{43}}{\gamma_{42}} \cdot (u_{3,3} - 4 u_{2,3} + 4 u_{2,4} - u_{4,4}) : \Gamma_{433} := -\frac{\gamma_{34}}{9} \cdot \frac{\gamma_{23}}{\gamma_{24}} \cdot (u_{3,3} \\
& \quad - 4 u_{3,4} + 4 u_{2,4} - u_{2,2}) : \\
\Gamma_{244} &:= -\frac{\gamma_{42}}{9} \cdot \frac{\gamma_{34}}{\gamma_{32}} \cdot (u_{4,4} - 4 u_{2,4} + 4 u_{2,3} - u_{3,3}) : \Gamma_{344} := -\frac{\gamma_{23}}{9} \cdot \frac{\gamma_{42}}{\gamma_{43}} \cdot (u_{2,2} \\
& \quad - 4 u_{2,3} + 4 u_{3,4} - u_{4,4}) : \\
\Gamma_{123} &:= -\frac{1}{9} \cdot R_f : \Gamma_{124} := -\frac{1}{9} \cdot R_f : \Gamma_{134} := -\frac{1}{9} \cdot R_f :
\end{aligned}$$

$$\begin{aligned}\Gamma_{223} &:= -\frac{2}{3} \cdot \gamma_{23} \cdot R_a - \frac{1}{9} \cdot (\gamma_{34} - 4 \gamma_{23}) \cdot (u_{3,3} + u_{2,4}) + \frac{1}{9} \cdot (\gamma_{34} + 5 \gamma_{23}) \cdot (u_{4,4} \\ &\quad + u_{2,3}) : \\ \Gamma_{323} &:= -\frac{2}{3} \cdot \gamma_{32} \cdot R_a - \frac{1}{9} \cdot (\gamma_{24} - 4 \gamma_{32}) \cdot (u_{2,2} + u_{3,4}) + \frac{1}{9} \cdot (\gamma_{24} + 5 \gamma_{32}) \cdot (u_{4,4} \\ &\quad + u_{2,3}) : \\ \Gamma_{423} &:= \frac{\gamma_{23}}{9} \cdot (u_{2,2} - 4 u_{2,4} + 4 u_{3,4} - u_{3,3}) : \end{aligned}$$

```
> simplify(evalDG(subs(VsS2, convert(DirectionalCovariantDerivative(D_x1, D_x1, CsS),
    DGvector) - \Gamma_{111} \cdot D_x1 - \Gamma_{211} \cdot D_x2 - \Gamma_{311} \cdot D_x3 - \Gamma_{411} \cdot D_x4))) ;
simplify(evalDG(subs(VsS2a, convert(DirectionalCovariantDerivative(D_x1, D_x2, CsS),
    DGvector) - \Gamma_{112} \cdot D_x1 - \Gamma_{212} \cdot D_x2 - \Gamma_{312} \cdot D_x3 - \Gamma_{412} \cdot D_x4))) ;
simplify(evalDG(subs(VsS2a, convert(DirectionalCovariantDerivative(D_x1, D_x3, CsS),
    DGvector) - \Gamma_{113} \cdot D_x1 - \Gamma_{213} \cdot D_x2 - \Gamma_{313} \cdot D_x3 - \Gamma_{413} \cdot D_x4))) ;
simplify(evalDG(subs(VsS2a, convert(DirectionalCovariantDerivative(D_x1, D_x4, CsS),
    DGvector) - \Gamma_{114} \cdot D_x1 - \Gamma_{214} \cdot D_x2 - \Gamma_{314} \cdot D_x3 - \Gamma_{414} \cdot D_x4))) ;
simplify(evalDG(subs(VsS2a, convert(DirectionalCovariantDerivative(D_x2, D_x2, CsS),
    DGvector) - \Gamma_{122} \cdot D_x1 - \Gamma_{222} \cdot D_x2 - \Gamma_{322} \cdot D_x3 - \Gamma_{422} \cdot D_x4))) ;
simplify(evalDG(subs(VsS2a, convert(DirectionalCovariantDerivative(D_x2, D_x3, CsS),
    DGvector) - \Gamma_{123} \cdot D_x1 - \Gamma_{223} \cdot D_x2 - \Gamma_{323} \cdot D_x3 - \Gamma_{423} \cdot D_x4))) ;
0 D_x1
0 D_x1
0 D_x1
0 D_x1
0 D_x1
0 D_x1
0 D_x1
```

(3.50)

>

III. Finally let us compute the canonical projective totally geodesic connection

```
> CsC := Connection(subs({seq(seq(seq(((a||i)||j)||k)=(((a||j)||i)||k), k=1..4), j=1
    ..(i-1)), i=1..4}), add(add(add(((a||i)||j)||k) \cdot vM[k] &t dM[i] &t dM[j], k=1
    ..4), j=1..4), i=1..4))) ;
> DD := {XX2, XX3, XX4} : ChrCoe := {seq(seq(seq(((a||i)||j)||k), k=1..4), j=i..4), i
    =1..4)} :
TGE := { } : for i to 3 do for j to 3 do TGE := {op(TGE), coeffs(expand(numer(subs(VsS2,
    subs(Qsyst, ToJet(Hook(convert(DirectionalCovariantDerivative(DD[i], DD[j], CsC),
    'DGvector'), \omega\omega), {u(x1, x2, x3, x4), q(x1, x2, x3, x4)}, notation=jetnumbers))))),
    {q[, q1})} : od: od: [nops(ChrCoe), nops(TGE)]
```

[40, 39]

(3.51)

```
> Sss2 := solve(TGE, ChrCoe \ {a144, a244, a344, a444}) : nops(Sss2)
```

36

(3.52)

$$> pt := \left\{ a144 = -\frac{1}{2} u_{4,4}, a244 = \frac{1}{2} \frac{u_{2,2} - 2 u_{2,4} + u_{4,4}}{u_2 - u_4}, a344 = \frac{1}{2} \frac{u_{3,3} - 2 u_{3,4} + u_{4,4}}{u_3 - u_4}, \right. \\ \left. a444 = 0 \right\} :$$

$$> Sss20 := \{op(subs(pt, Sss2)), op(pt)\} : CsC0 := subs(Sss20, CsC) :$$

$$> subs(Sss20, a111); subs(Sss20, a112); subs(Sss20, a113); subs(Sss20, a114)$$

$$0$$

$$0$$

$$0$$

$$0$$

(3.53)

$$> simplify(subs(VsS2, subs(Sss20, a121))); subs(Sss20, a122); subs(Sss20, a123); \\ subs(Sss20, a124)$$

$$0$$

$$-\frac{1}{2} u_{2,2}$$

$$0$$

$$0$$

(3.54)

$$> factor(subs(VsS2, subs(Sss20, a131))); subs(Sss20, a132); subs(Sss20, a133); subs(Sss20, \\ a134)$$

$$0$$

$$0$$

$$-\frac{1}{2} u_{3,3}$$

$$0$$

(3.55)

$$> simplify(subs(VsS2c, subs(Sss20, a141))); subs(Sss20, a142); subs(Sss20, a143); \\ subs(Sss20, a144)$$

$$0$$

$$0$$

$$0$$

$$-\frac{1}{2} u_{4,4}$$

(3.56)

$$> simplify(subs(VsS2b, subs(Sss20, a221))); simplify(subs(VsS2, subs(Sss20, a222))); \\ subs(Sss20, a223); subs(Sss20, a224)$$

$$0$$

$$0$$

$$0$$

$$0$$

(3.57)

$$> factor(subs(VsS2b, subs(Sss20, a231))); factor(subs(VsS2b, subs(Sss20, a232))); \\ simplify(subs(VsS2a, subs(Sss20, a233))); subs(Sss20, a234)$$

$$0$$

$$-\frac{1}{2} \frac{u_{2,2} - 2 u_{2,3} + u_{3,3}}{u_2 - u_3}$$

$$\frac{1}{2} \frac{u_{2,2} - 2u_{2,3} + u_{3,3}}{u_2 - u_3} \quad (3.58)$$

> factor(subs(VsS2, subs(Sss20, a241))); factor(subs(VsS2a, subs(Sss20, a242)));
subs(Sss20, a243); subs(Sss20, a244)

$$0$$

$$-\frac{1}{2} \frac{u_{2,2} - 2u_{2,4} + u_{4,4}}{u_2 - u_4}$$

$$0$$

$$\frac{1}{2} \frac{u_{2,2} - 2u_{2,4} + u_{4,4}}{u_2 - u_4} \quad (3.59)$$

> simplify(subs(VsS2, subs(Sss20, a331))); simplify(subs(VsS2b, subs(Sss20, a332)));
simplify(subs(VsS2b, subs(Sss20, a333))); subs(Sss20, a334)

$$0$$

$$0$$

$$0$$

$$0 \quad (3.60)$$

> simplify(subs(VsS2b, subs(Sss20, a341))); simplify(subs(VsS2, subs(Sss20, a342)));
simplify(subs(VsS2c, subs(Sss20, a343))); subs(Sss20, a344)

$$0$$

$$0$$

$$-\frac{1}{2} \frac{u_{3,3} - 2u_{3,4} + u_{4,4}}{u_3 - u_4}$$

$$\frac{1}{2} \frac{u_{3,3} - 2u_{3,4} + u_{4,4}}{u_3 - u_4} \quad (3.61)$$

> simplify(subs(VsS2, subs(Sss20, a441))); simplify(subs(VsS2b, subs(Sss20, a442)));
simplify(subs(VsS2b, subs(Sss20, a443))); subs(Sss20, a444)

$$0$$

$$0$$

$$0$$

$$0 \quad (3.62)$$

> DGsimplify(evalDG(subs(VsS2a, convert(DirectionalCovariantDerivative(D_x2, D_x3,
CsC0), DGvector))))

$$-\left(\frac{u_{2,2} - 2u_{2,3} + u_{3,3}}{2(u_2 - u_3)} D_{x2} \right) + \frac{u_{2,2} - 2u_{2,3} + u_{3,3}}{2(u_2 - u_3)} D_{x3} \quad (3.63)$$

> DGsimplify(evalDG(subs(VsS2a, convert(DirectionalCovariantDerivative(D_x3, D_x2,
CsC0), DGvector))))

$$-\left(\frac{u_{2,2} - 2u_{2,3} + u_{3,3}}{2(u_2 - u_3)} D_{x2} \right) + \frac{u_{2,2} - 2u_{2,3} + u_{3,3}}{2(u_2 - u_3)} D_{x3} \quad (3.64)$$

>

L>

MASh universal hierarchy 4D

$$\begin{aligned}
 & \text{DGsetup}([x, y, t, z], [u], J, 3) : \\
 & \text{Eq1} := u_{1,3} - u_{2,2} + u_2 u_{1,1} - u_1 u_{1,2} : \\
 & \text{Eq2} := u_{1,4} - u_{2,3} + u_3 u_{1,1} - u_1 u_{1,3} : \\
 & \text{Eq3} := u_{2,4} - u_{3,3} + u_3 u_{1,2} - u_2 u_{1,3} : \\
 & S := \text{solve}(\{\text{Eq1}, \text{Eq2}, \text{Eq3}\}, \{u_{1,2}, u_{1,3}, u_{1,4}\}) \\
 & S := \left\{ u_{1,2} = \frac{u_2^2 u_{1,1} - u_2 u_{2,2} + u_{2,4} - u_{3,3}}{u_1 u_2 - u_3}, u_{1,3} \right. \\
 & \quad = \frac{u_2 u_3 u_{1,1} + u_1 u_{2,4} - u_1 u_{3,3} - u_3 u_{2,2}}{u_1 u_2 - u_3}, u_{1,4} \\
 & \quad \left. = \frac{u_1^2 u_{2,4} - u_1^2 u_{3,3} + u_1 u_2 u_{2,3} - u_1 u_3 u_{2,2} + u_3^2 u_{1,1} - u_3 u_{2,3}}{u_1 u_2 - u_3} \right\} \tag{4.1}
 \end{aligned}$$

$$\begin{aligned}
 & \text{Vss3} := \{\text{Eq1}, \text{Eq2}, \text{Eq3}\} \text{ for } i \text{ to } 4 \text{ do } \text{Vss3} := \{\text{op}(\text{Vss3}), \text{TotalDiff}(\text{Eq1}, i), \\
 & \quad \text{TotalDiff}(\text{Eq2}, i), \text{TotalDiff}(\text{Eq3}, i)\} \text{ od:} \\
 & \text{nops}(\text{eliminate}(\text{Vss3}, \{u_{1,2}, u_{1,3}, u_{1,4}, u_{1,1}, u_{1,2}, u_{1,3}, u_{1,4}, u_{1,2}, u_{1,3}, u_{1,4}, u_{1,3}, \\
 & \quad u_{1,3}, u_{1,4}, u_{2,3}, u_{2,4}\})[2]) \\
 & \quad \quad \quad 0 \tag{4.2}
 \end{aligned}$$

$$\begin{aligned}
 & \text{S3} := \text{factor}(\text{solve}(\text{Vss3}, \{u_{1,2}, u_{1,3}, u_{1,4}, u_{1,1}, u_{1,2}, u_{1,3}, u_{1,4}, u_{1,2}, u_{1,3}, u_{1,4}, u_{1,3}, \\
 & \quad u_{1,3}, u_{1,4}, u_{2,3}, u_{2,4}\})) : \text{nops}(\text{S3}) \\
 & \quad \quad \quad 13 \tag{4.3}
 \end{aligned}$$

$$\begin{aligned}
 & \text{Symb1} := \text{add}(\text{add}(\text{diff}(\text{Eq1}, u_{i,j}) \cdot p_i \cdot p_j, j = 1..4), i = 1..4); \\
 & \text{Symb2} := \text{add}(\text{add}(\text{diff}(\text{Eq2}, u_{i,j}) \cdot p_i \cdot p_j, j = 1..4), i = 1..4); \\
 & \text{Symb3} := \text{add}(\text{add}(\text{diff}(\text{Eq3}, u_{i,j}) \cdot p_i \cdot p_j, j = 1..4), i = 1..4); \\
 & \quad \text{Symb1} := p_1^2 u_2 - p_1 p_2 u_1 + p_1 p_3 - p_2^2 \\
 & \quad \text{Symb2} := p_1^2 u_3 - p_1 p_3 u_1 + p_1 p_4 - p_2 p_3 \\
 & \quad \text{Symb3} := p_1 p_2 u_3 - p_1 p_3 u_2 + p_2 p_4 - p_3^2 \tag{4.4}
 \end{aligned}$$

$$\begin{aligned}
 & \text{simplify}(\text{subs}(\{p_1 = 1, p_2 = q - u_1, p_3 = q^2 - u_1 q - u_2, p_4 = q^3 - u_1 q^2 - u_2 q - u_3\}, \\
 & \quad \{\text{Symb1}, \text{Symb2}, \text{Symb3}\})) \\
 & \quad \quad \quad \{0\} \tag{4.5}
 \end{aligned}$$

$$\begin{aligned}
 & \text{tmp} := \text{solve}(\{\text{coeffs}(\text{expand}((a1 \cdot p_1 + a2 \cdot p_2 + a3 \cdot p_3 + a4 \cdot p_4) \cdot \text{Symb1} + (b1 \cdot p_1 + b2 \cdot p_2 \\
 & \quad + b3 \cdot p_3 + b4 \cdot p_4) \cdot \text{Symb2} + (c1 \cdot p_1 + c2 \cdot p_2 + c3 \cdot p_3 + c4 \cdot p_4) \cdot \text{Symb3}), \{p_1, p_2, p_3, p_4\}) \\
 & \quad \{a1, a2, a3, a4, b1, b2, b3, b4, c1, c2, c3, c4\}) \\
 & \text{tmp} := \{a1 = c2 u_3, a2 = 0, a3 = -c2 u_1 + c1, a4 = c2, b1 = -c2 u_2, b2 = c2 u_1 - c1, b3 = \\
 & \quad -c2, b4 = 0, c1 = c1, c2 = c2, c3 = 0, c4 = 0\} \tag{4.6}
 \end{aligned}$$

$$\begin{aligned} & \text{simplify}([p_3 \cdot \text{Symb1} - p_2 \cdot \text{Symb2} + p_1 \cdot \text{Symb3}, (p_4 + u_3 p_1) \cdot \text{Symb1} - (p_3 + u_2 p_1) \cdot \text{Symb2} \\ & \quad + (p_2 + u_1 p_1) \cdot \text{Symb3}]) \\ & \quad [0, 0] \end{aligned} \quad (4.7)$$

compatibility conditions:

$$\begin{aligned} & \text{simplify}(\text{TotalDiff}(\text{Eq1}, 3) - \text{TotalDiff}(\text{Eq2}, 2) + \text{TotalDiff}(\text{Eq3}, 1)); \\ & \text{simplify}(\text{subs}(S, (\text{TotalDiff}(\text{Eq1}, 4) + u_3 \text{TotalDiff}(\text{Eq1}, 1)) - (\text{TotalDiff}(\text{Eq2}, 3) \\ & \quad + u_2 \text{TotalDiff}(\text{Eq2}, 1)) + (\text{TotalDiff}(\text{Eq3}, 2) + u_1 \text{TotalDiff}(\text{Eq3}, 1)))) \\ & \quad 0 \\ & \quad 0 \end{aligned} \quad (4.8)$$

This shows that dimensions of the symbol spaces $(g[0], g[1], g[2], g[3], g[4], \dots)$ are equal to $(1, 4, 7, 10, 13, \dots)$,

i.e. the Hilbert polynomial is given by the formula $\dim(g[k]) = 1 + 3k$, whence the general solution depends on 3 functions of 1 variable.

This also is confirmed by the characteristic variety = rational normal curve that has degree 3 and dimension 1.

$$\begin{aligned} & \text{DGsetup}([x, y, t, z], [q, u], Jq, 3) : \\ & \text{Qsyst} := \{q_2 = (q_{[1]} - u_1) q_1, q_3 = (q_{[1]}^2 - u_1 q_{[1]} - u_2) q_1, q_4 = (q_{[1]}^3 - u_1 q_{[1]}^2 - u_2 q_{[1]} \\ & \quad - u_3) q_1\} : \\ & \text{simplify}(\text{subs}(S3, \text{expand}(\text{subs}(\text{Qsyst}, \text{subs}(\{q_{1,2} = \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_2), 1), q_{1,3} \\ & \quad = \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_3), 1), q_{1,4} = \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_4), 1)\}, \\ & \quad \{\text{TotalDiff}(\text{subs}(\text{Qsyst}, q_3), 2) - \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_2), 3), \\ & \quad \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_4), 2) - \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_2), 4), \text{TotalDiff}(\text{subs}(\text{Qsyst}, \\ & \quad q_4), 3) - \text{TotalDiff}(\text{subs}(\text{Qsyst}, q_3), 4)\}))) \\ & \quad \{0\} \end{aligned} \quad (4.9)$$

$$\begin{aligned} & \text{DGsetup}([x, y, t, z, q], [u], Jqu, 3) : \\ & X2f := f \rightarrow \text{TotalDiff}(f, 2) - (q - u_1) \cdot \text{TotalDiff}(f, 1) : \\ & X3f := f \rightarrow \text{TotalDiff}(f, 3) - (q^2 - u_1 q - u_2) \cdot \text{TotalDiff}(f, 1) : \\ & X4f := f \rightarrow \text{TotalDiff}(f, 4) - (q^3 - u_1 q^2 - u_2 q - u_3) \cdot \text{TotalDiff}(f, 1) : \\ & \text{simplify}(\text{subs}(S3, \{\text{simplify}(X2f(X3f(x1)) - X3f(X2f(x1))), \text{simplify}(X2f(X3f(q)) \\ & \quad - X3f(X2f(q))), \text{simplify}(X2f(X4f(x1)) - X4f(X2f(x1))), \text{simplify}(X2f(X4f(q)) \\ & \quad - X4f(X2f(q))), \text{simplify}(X3f(X4f(x1)) - X4f(X3f(x1))), \text{simplify}(X3f(X4f(q)) \\ & \quad - X4f(X3f(q)))\})) \\ & \quad \{0\} \end{aligned} \quad (4.10)$$

I. We first look for a connection (with torsion) preserving the field of GL2-curves and having α -surfaces totally geodesic + trace of torsion zero.

$$\begin{aligned} & \text{DGsetup}([x, y, t, z], M) : \\ & vM := [D_x, D_y, D_t, D_z] : dM := [dx, dy, dt, dz] : \\ & C := \text{Connection}(\text{add}(\text{add}(\text{add}(((a \parallel i) \parallel j) \parallel k) \cdot vM[k] \& t dM[i] \& t dM[j], k = 1 .. 4), j = 1 \\ & \quad .. 4), i = 1 .. 4)) : \\ & X2 := \text{evalDG}(vM[2] - (q_{[1]} - u_1) \cdot vM[1]) : \end{aligned}$$

$$\begin{aligned}
X3 &:= \text{evalDG}(vM[3] - (q_{[1]}^2 - u_1 q_{[1]} - u_2) \cdot vM[1]) : \\
X4 &:= \text{evalDG}(vM[4] - (q_{[1]}^3 - u_1 q_{[1]}^2 - u_2 q_{[1]} - u_3) \cdot vM[1]) : \\
\omega &:= \text{evalDG}(dM[1] + (q_{[1]} - u_1) \cdot dM[2] + (q_{[1]}^2 - u_1 q_{[1]} - u_2) \cdot dM[3] + (q_{[1]}^3 - u_1 q_{[1]}^2 \\
&\quad - u_2 q_{[1]} - u_3) \cdot dM[4]) : \\
&[Hook(X2, \omega), Hook(X3, \omega), Hook(X4, \omega)] \\
&\quad [0, 0, 0]
\end{aligned} \tag{4.11}$$

$$\begin{aligned}
> XX2 &:= \text{FromJet}(X2, \{u(x, y, t, z), q(x, y, t, z)\}) : \\
XX3 &:= \text{FromJet}(X3, \{u(x, y, t, z), q(x, y, t, z)\}) : \\
XX4 &:= \text{FromJet}(X4, \{u(x, y, t, z), q(x, y, t, z)\}) : \\
\omega\omega &:= \text{FromJet}(\omega, \{u(x, y, t, z), q(x, y, t, z)\}) : \\
> DD &:= \{XX2, XX3, XX4\} : \text{ChrCoe} := \{seq(seq(seq(((a \parallel i) \parallel j) \parallel k), i = 1..4), j = 1..4), k \\
&\quad = 1..4)\} : \\
TGE &:= \{ \} : \text{for } i \text{ to } 3 \text{ do for } j \text{ to } 3 \text{ do } TGE := \{op(TGE), coeffs(expand(numer(subs(S3, \\
&\quad subs(Qsyst, ToJet(Hook(convert(DirectionalCovariantDerivative(DD[i], DD[j], C), \\
&\quad 'DGvector'), \omega\omega), \{u(x, y, t, z), q(x, y, t, z)\}, notation = jetnumbers)))), \{q_{[1]}, q_1\})\} : \\
&\text{od:od:} \\
&[nops(ChrCoe), nops(TGE)] \\
&\quad [64, 59]
\end{aligned} \tag{4.12}$$

$$\begin{aligned}
> Sconn &:= \text{solve}(TGE, ChrCoe) : \\
> FreePar &:= \{ \} : \text{for } i \text{ to } nops(Sconn) \text{ do if } lhs(Sconn[i]) = rhs(Sconn[i]) \text{ then } FreePar \\
&\quad := \{op(FreePar), lhs(Sconn[i])\} \text{ fi:od: } FreePar \\
&\quad \{a144, a244, a344, a413, a414, a423, a424, a433, a434, a444\}
\end{aligned} \tag{4.13}$$

$$\begin{aligned}
> ChrCoe1 &:= \{a111, a112, a113, a114, a121, a122, a123, a124, a131, a132, a133, a134, \\
&\quad a141, a142, a143, a211, a212, a213, a214, a221, a222, a223, a224, a231, a232, a233, \\
&\quad a234, a241, a242, a243, a311, a312, a313, a314, a321, a322, a323, a324, a331, a332, \\
&\quad a333, a334, a341, a342, a343, a411, a412, a421, a422, a431, a432, a441, a442, a443\} : \\
&\quad nops(\%) \\
&\quad 54
\end{aligned} \tag{4.14}$$

$$\begin{aligned}
> Sconn &:= \text{solve}(TGE, ChrCoe1) : \\
> FreePar &:= \{a144, a244, a344, a413, a414, a423, a424, a433, a434, a444\} : \\
> CC &:= \text{DGsimplify}(subs(Sconn, C)) : \\
> Quad1 &:= \text{evalDG}(add(add(diff(Eq1, u_{i,j}) \cdot vM[i] \&s vM[j], j = i..4), i = 1..4)) : \\
Quad2 &:= \text{evalDG}(add(add(diff(Eq2, u_{i,j}) \cdot vM[i] \&s vM[j], j = i..4), i = 1..4)) : \\
Quad3 &:= \text{evalDG}(add(add(diff(Eq3, u_{i,j}) \cdot vM[i] \&s vM[j], j = i..4), i = 1..4)) : \\
Quad1u &:= \text{FromJet}(Quad1, \{u(x, y, t, z), q(x, y, t, z)\}) : Quad2u := \text{FromJet}(Quad2, \{u(x, \\
&\quad y, t, z), q(x, y, t, z)\}) : Quad3u := \text{FromJet}(Quad3, \{u(x, y, t, z), q(x, y, t, z)\}) : Quadu \\
&\quad := [Quad1u, Quad2u, Quad3u] : \\
> \{seq(ContractIndices(Quad \parallel i, \omega \&t \omega, [[1, 1], [2, 2]]), i = 1..3)\} \\
&\quad \{0\}
\end{aligned} \tag{4.15}$$

$$\begin{aligned}
> RNE &:= \{ \} : \text{for } i \text{ to } 4 \text{ do for } j \text{ to } 3 \text{ do } RNE := \{op(RNE), \\
&\quad coeffs(expand(numer(simplify(subs(S3, subs(Qsyst, \\
&\quad ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CC), \omega
\end{aligned}$$

&t ω, [[1, 1], [2, 2]], {u(x, y, t, z), q(x, y, t, z)}, notation = jetnumbers))))), {q_[],
q₁} } :od:od: nops(RNE)

43

(4.16)

> sRNE := solve(RNE, FreePar) :

> RNE := { } :for i to 4 do for j to 3 do RNE := { op(RNE),
coeffs(expand(numer(simplify(subs(S3, subs(Qsyst,
ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CC), ω
&t ω, [[1, 1], [2, 2]], {u(x, y, t, z)}, notation = jetnumbers)))))), {q_[]} } :od:od:
nops(RNE)

43

(4.17)

> sRNE := solve(RNE, {a144, a244, a344, a413, a423, a433}) :

> CT := DGsimplify(evalDG(subs({ a414 = α₁, a424 = α₂, a434 = α₃, a444 = α₄ }, subs(sRNE,
CC)))) :

> RNE := { } :for i to 4 do for j to 3 do RNE := { op(RNE),
coeffs(expand(numer(simplify(subs(S3, subs(Qsyst,
ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CT), ω &t ω,
[[1, 1], [2, 2]], {u(x, y, t, z)}, notation = jetnumbers)))))), {q_[]} } :od:od:
simplify(RNE)

{0}

(4.18)

> sCTa := solve(DGinfo(ContractIndices(TorsionTensor(CT), [[1, 2]]), "CoefficientSet"),
{α₁, α₂, α₃, α₄}) : CTa := DGsimplify(evalDG(subs(sCTa, CT))) :

> ContractIndices(TorsionTensor(CTa), [[1, 2]])

0 dx

(4.19)

> sCTa

$$\left\{ \alpha_1 = 0, \alpha_2 = -\frac{1}{3} u_{1,1}, \alpha_3 = \right.$$

(4.20)

$$-\frac{1}{3} \frac{u_1^2 u_{2,1} u_{1,1} - u_1 u_3 u_{1,1} + 2 u_2^2 u_{1,1} - 2 u_2 u_{2,2} + 2 u_{2,4} - 2 u_{3,3}}{u_1 u_2 - u_3}, \alpha_4 =$$

$$-\frac{1}{3} \frac{1}{u_1 u_2 - u_3} (u_1^3 u_{2,1} u_{1,1} - u_1^2 u_3 u_{1,1} + 3 u_1 u_2^2 u_{1,1} - u_1 u_2 u_{2,2} + u_2 u_3 u_{1,1}$$

$$+ 4 u_1 u_{2,4} - 4 u_1 u_{3,3} - 3 u_3 u_{2,2}) \left. \right\}$$

> CTa :

> simplify(subs(solve(S[1..3], {u_{1,1}, u_{2,3}, u_{3,3}}), subs(sCTa, α₃)));
simplify(subs(solve(S[1..3], {u_{1,1}, u_{2,3}, u_{3,3}}), subs(sCTa, α₄)))
$$-\frac{1}{3} \frac{u_1^2 u_{1,2} - u_1 u_{1,3} + u_1 u_{2,2} + 2 u_2 u_{1,2}}{u_2}$$

$$-\frac{1}{3} \frac{u_1^3 u_{1,2} - u_1^2 u_{1,3} + u_1^2 u_{2,2} + 3 u_1 u_2 u_{1,2} + u_2 u_{1,3} + 2 u_2 u_{2,2}}{u_2} \quad (4.21)$$

$$> sCTa2 := \left\{ \alpha_1 = 0, \alpha_2 = -\frac{1}{3} u_{1,1}, \alpha_3 = -\frac{1}{3} \frac{u_1^2 u_{1,2} - u_1 u_{1,3} + u_1 u_{2,2} + 2 u_2 u_{1,2}}{u_2}, \alpha_4 = \right. \\ \left. -\frac{1}{3} \frac{u_1^3 u_{1,2} - u_1^2 u_{1,3} + u_1^2 u_{2,2} + 3 u_1 u_2 u_{1,2} + u_2 u_{1,3} + 2 u_2 u_{2,2}}{u_2} \right\}.$$

$$> \text{simplify}(\{ \text{seq}(\text{subs}(S, \text{subs}(sCTa, \alpha[i])) - \text{subs}(sCTa2, \alpha[i])), i = 1..4) \}) \\ \{0\} \quad (4.22)$$

$$> \text{factor}(\text{DirectionalCovariantDerivative}(D_x, D_y, CTa)) \\ - (u_{1,1} D_x) \quad (4.23)$$

Thus there exist GL2-connections, with respect to which the null Lax surfaces are totally geodesic. They depend on 4 arbitrary functions.

The parameters can be fixed uniquely by the condition $\text{Trace}(\text{Torsion})=0$.

$$> RNE := \{ \} : \text{for } i \text{ to } 4 \text{ do for } j \text{ to } 3 \text{ do } RNE := \{ \text{op}(RNE), \\ \text{coeffs}(\text{expand}(\text{numer}(\text{simplify}(\text{subs}(S, \text{subs}(Qsyst, \\ \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(vM[i], \text{Quadu}[j], CTa), \omega \\ \& \omega, [[1, 1], [2, 2]]), \{u(x, y, t, z)\}, \text{notation} = \text{jetnumbers}))))) , \{q_{[]}\}) \} : \text{od:od:} \\ \text{simplify}(RNE) \\ \{0\} \quad (4.24)$$

$$> RNE := \{ \} : \text{for } i \text{ to } 4 \text{ do for } j \text{ to } 3 \text{ do } RNE := \{ \text{op}(RNE), \\ \text{coeffs}(\text{expand}(\text{numer}(\text{simplify}(\text{subs}(Qsyst, \\ \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(vM[i], \text{Quadu}[j], CT), \omega \& \omega, \\ [[1, 1], [2, 2]]), \{u(x, y, t, z)\}, \text{notation} = \text{jetnumbers}))))) , \{q_{[]}\}) \} : \text{od:od:} \\ \text{simplify}(RNE) :$$

$$> \text{solve}(\%, \{u_{1,2}, u_{1,3}, u_{1,4}\}) \\ \left\{ u_{1,2} = \frac{u_2^2 u_{1,1} - u_2 u_{2,2} + u_{2,4} - u_{3,3}}{u_1 u_2 - u_3}, u_{1,3} = \frac{u_2 u_3 u_{1,1} + u_1 u_{2,4} - u_1 u_{3,3} - u_3 u_{2,2}}{u_1 u_2 - u_3}, u_{1,4} \right. \\ \left. = \frac{u_1^2 u_{2,4} - u_1^2 u_{3,3} + u_1 u_2 u_{2,3} - u_1 u_3 u_{2,2} + u_3^2 u_{1,1} - u_3 u_{2,3}}{u_1 u_2 - u_3} \right\} \quad (4.25)$$

$$> S \\ \left\{ u_{1,2} = \frac{u_2^2 u_{1,1} - u_2 u_{2,2} + u_{2,4} - u_{3,3}}{u_1 u_2 - u_3}, u_{1,3} = \frac{u_2 u_3 u_{1,1} + u_1 u_{2,4} - u_1 u_{3,3} - u_3 u_{2,2}}{u_1 u_2 - u_3}, u_{1,4} \right. \\ \left. = \frac{u_1^2 u_{2,4} - u_1^2 u_{3,3} + u_1 u_2 u_{2,3} - u_1 u_3 u_{2,2} + u_3^2 u_{1,1} - u_3 u_{2,3}}{u_1 u_2 - u_3} \right\} \quad (4.26)$$

$$> \text{RicCTA} := \text{RicciTensor}(\text{FromJet}(CTa, \{u(x, y, t, z)\})) : \text{RicCTa} := \text{ToJet}(\text{RicCTA}, \{u(x, y, \\ t, z)\}, \text{notation} = \text{jetnumbers}) :$$

$$> \text{RicCTaSkew} := \text{SymmetrizeIndices}(\text{RicCTa}, [1, 2], \text{"SkewSymmetric"}) :$$

```

RicCTaSymm := SymmetrizeIndices(RicCTa, [1, 2], "Symmetric") :
>
>  $\omega\omega := evalDG(\omega)$ 
 $\omega\omega := dx + (q[ ] - u_1) dy + (q[ ]^2 - q[ ] u_1 - u_2) dt + (q[ ]^3 - q[ ]^2 u_1 - q[ ] u_2 - u_3) dz$  (4.27)

```

```

>  $\omega0 := -\omega\omega$ ;  $\omega0b := DGsimplify(evalDG(subs(q[ ] = 0, \omega0)))$ ;
 $\omega1a := evalDG\left(\frac{1}{\epsilon} \cdot (subs(q[ ] = q[ ] + \epsilon, \omega0) - \omega0)\right)$ ;  $\omega1 := evalDG(subs(\epsilon = 0, \omega1a))$ ;
 $\omega1b := DGsimplify(evalDG(subs(q[ ] = 0, \omega1)))$ ;
 $\omega2a := evalDG\left(\frac{1}{\epsilon} \cdot (subs(q[ ] = q[ ] + \epsilon, \omega1) - \omega1)\right)$ ;  $\omega2 := evalDG(subs(\epsilon = 0, \omega2a))$ ;
 $\omega2b := DGsimplify\left(evalDG\left(\frac{1}{2} \cdot subs(q[ ] = 0, \omega2)\right)\right)$ ;
 $\omega3a := evalDG\left(\frac{1}{\epsilon} \cdot (subs(q[ ] = q[ ] + \epsilon, \omega2) - \omega2)\right)$ ;  $\omega3 := evalDG(subs(\epsilon = 0, \omega3a))$ ;
 $\omega3b := DGsimplify\left(evalDG\left(\frac{1}{6} \cdot subs(q[ ] = 0, \omega3)\right)\right)$ 
 $\omega0b := -dx + u_1 dy + u_2 dt + u_3 dz$ 
 $\omega1b := -dy + u_1 dt + u_2 dz$ 
 $\omega2b := -dt + u_1 dz$ 
 $\omega3b := -dz$  (4.28)

```

```

>  $evalDG(\omega0 - \omega0b - \omega1b \cdot q[ ] - \omega2b \cdot q[ ]^2 - \omega3b \cdot q[ ]^3)$ 
 $0 dx$  (4.29)

```

```

>  $Qua\omega := evalDG([ \omega0b \&s \omega0b, \omega0b \&s \omega1b, 2 \omega0b \&s \omega2b + \omega1b \&s \omega1b, \omega0b \&s \omega3b$ 
 $+ \omega1b \&s \omega2b, 2 \omega1b \&s \omega3b + \omega2b \&s \omega2b, \omega2b \&s \omega3b, \omega3b \&s \omega3b ])$ ;  $tmp\Omega$ 
 $:= \{ \} : \text{for } i \text{ to } nops(Qua\omega) \text{ do for } j \text{ to } 3 \text{ do } tmp\Omega := \{op(tmp\Omega),$ 
 $ContractIndices(Quad||j, Qua\omega[i], [[1, 1], [2, 2]])\} : \text{od:od: } tmp\Omega$ 
 $\{0\}$  (4.30)

```

```

> # eliminate(simplify(subs(S3, DGinfo(evalDG(RicCTaSymm - add( $\Lambda[i] \cdot Qua\omega[i]$ ,  $i = 1$ 
 $..7$ ))), "CoefficientSet"))), {seq( $\Lambda[i]$ ,  $i = 1 .. 7$ )})[2]

```

Thus the condition that $\text{sym}(\text{Ric}(\nabla))$ belongs to the span of quadrics defining the GL2 structure is non-vacous. It is another differential constraint, giving exact solutions.

But in fact, from what is computed below it follows that this is the same condition as $\text{sym}(\text{Ric})=0$.

```

>  $Vt0 := Annihilator([ \omega0, \omega1, \omega2 ])[1]$ ;  $Vt1 := Annihilator([ \omega0, \omega1 ])[2]$ ;  $Vt2$ 
 $:= Annihilator([ \omega0 ])[3]$ 
 $Vt0 := -((q[ ]^3 - 3 q[ ]^2 u_1 + 3 q[ ] u_1^2 - u_1^3 + 3 q[ ] u_2 - 2 u_1 u_2 - u_3) D_x) + (3 q[ ]^2$ 
 $- 3 q[ ] u_1 + u_1^2 + u_2) D_y - ((3 q[ ] - u_1) D_t) + D_z$ 
 $Vt1 := (q[ ]^2 - 2 q[ ] u_1 + u_1^2 + u_2) D_x - ((2 q[ ] - u_1) D_y) + D_t$ 
 $Vt2 := -((q[ ] - u_1) D_x) + D_y$  (4.31)

```

$$\begin{aligned}
& \text{Vt01} := \text{ContractIndices}(\text{TorsionTensor}(CT), \text{evalDG}(\text{Vt0} \& \text{Vt1}), [[2, 1], [3, 2]]) : \\
& \text{KrCond} := \text{DGinfo}(\text{evalDG}(\text{convert}(\text{Vt01}, \text{DGvector}) - \text{Hook}(\text{convert}(\text{Vt01}, \text{DGvector}), \text{dz} \\
& \quad \cdot \text{Vt0}), \text{"CoefficientSet"})) : \\
& \text{sCTa3} := \text{solve}(\text{map}(Q \rightarrow \text{coeffs}(\text{expand}(\text{numer}(Q)), \{q_{[]}\}), \text{KrCond}), \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}) : \\
& \text{simplify}(\{ \text{seq}(\text{subs}(S, \text{subs}(sCTa, \alpha[i]) - \text{subs}(sCTa2, \alpha[i])), i = 1..4), \text{seq}(\text{subs}(S, \\
& \quad \text{subs}(sCTa, \alpha[i]) - \text{subs}(sCTa3, \alpha[i])), i = 1..4) \} \\
& \quad \{0\} \tag{4.32}
\end{aligned}$$

Thus there exist GL2-connections, with respect to which the null Lax surfaces are totally geodesic. They depend on 4 parameters that can be fixed uniquely by the condition $\text{Trace}(\text{Torsion})=0$. This latter condition is equivalent to that of Krynski.

$$\text{simplify}(\text{subs}(S3, \text{ContractIndices}(\text{RicCTaSymm}, \text{evalDG}(\text{Vt0} \& \text{Vt0}), [[1, 1], [2, 2]]))) \tag{4.33}$$

$$\text{simplify}(\text{subs}(S3, \text{ContractIndices}(\text{RicCTaSymm}, \text{evalDG}(\text{Vt0} \& \text{Vt1}), [[1, 1], [2, 2]]))) \tag{4.34}$$

$$\begin{aligned}
& \text{QuT} := \text{evalDG}(\text{add}(\text{add}(\text{xi}[i, j] \cdot \text{dM}[i] \& \text{s dM}[j], j = i..4), i = 1..4)) : \\
& \text{SuT} := \text{solve}(\{ \text{coeffs}(\text{expand}(\text{numer}(\text{ContractIndices}(\text{QuT}, \text{Vt0} \& \text{Vt0}), [[1, 1], [2, 2]]))), \\
& \quad \{q_{[]}\} \}, \{ \xi_{1, 1}, \xi_{1, 2}, \xi_{1, 3}, \xi_{1, 4}, \xi_{2, 2}, \xi_{2, 3}, \xi_{2, 4} \}) : \\
& \text{TuT} := \text{DGsimplify}(\text{evalDG}(\text{subs}(\{ \xi_{3, 3} = \Lambda_1, \xi_{3, 4} = \Lambda_2, \xi_{4, 4} = \Lambda_3 \}, \text{subs}(\text{SuT}, \text{QuT})))) : \\
& \text{eliminate}(\text{DGinfo}(\text{DGsimplify}(\text{evalDG}(\text{subs}(S3, \text{TuT} - \text{RicCTaSymm}))), \\
& \quad \text{"CoefficientSet"}), \{ \Lambda_1, \Lambda_2, \Lambda_3 \})[2] \\
& \quad \{ \} \tag{4.35}
\end{aligned}$$

This computation has shown that symmetrized Ricci tensor of the canonical GL2-connection is a section of the bundle of quadrics annihilating the dual (tangent) to the canonical (cotangent) rational normal curve.

Let us compute the formulae for the Christoffel coefficients of the canonical connection.

$$\begin{aligned}
& S \\
& \left\{ u_{1, 2} = \frac{u_1^2 u_{1, 1} - u_2 u_{2, 2} + u_{2, 4} - u_{3, 3}}{u_1 u_2 - u_3}, u_{1, 3} = \frac{u_2 u_3 u_{1, 1} + u_1 u_{2, 4} - u_1 u_{3, 3} - u_3 u_{2, 2}}{u_1 u_2 - u_3}, u_{1, 4} \right. \tag{4.36} \\
& \quad \left. = \frac{u_1^2 u_{2, 4} - u_1^2 u_{3, 3} + u_1 u_2 u_{2, 3} - u_1 u_3 u_{2, 2} + u_3^2 u_{1, 1} - u_3 u_{2, 3}}{u_1 u_2 - u_3} \right\}
\end{aligned}$$

$$\begin{aligned}
& S2a := \text{solve}(\{ \text{Eq1}, \text{Eq2}, \text{Eq3} \}, \{ u_{1, 1}, u_{2, 2}, u_{3, 3} \}); \\
& S2b := \text{solve}(\{ \text{Eq1}, \text{Eq2}, \text{Eq3} \}, \{ u_{2, 2}, u_{2, 3}, u_{2, 4} \}); \\
& S2c := \text{solve}(\{ \text{Eq1}, \text{Eq2}, \text{Eq3} \}, \{ u_{1, 1}, u_{1, 2}, u_{1, 3} \}) : \\
& S2a := \left\{ u_{1, 1} = \frac{u_1 u_{1, 3} - u_{1, 4} + u_{2, 3}}{u_3}, u_{2, 2} \right. \\
& \quad \left. = \frac{u_1 u_2 u_{1, 3} - u_1 u_3 u_{1, 2} - u_2 u_{1, 4} + u_2 u_{2, 3} + u_3 u_{1, 3}}{u_3}, u_{3, 3} = -u_2 u_{1, 3} + u_3 u_{1, 2} + u_{2, 4} \right\} \\
& S2b := \{ u_{2, 2} = -u_1 u_{1, 2} + u_2 u_{1, 1} + u_{1, 3}, u_{2, 3} = -u_1 u_{1, 3} + u_3 u_{1, 1} + u_{1, 4}, u_{2, 4} = u_2 u_{1, 3} \\
& \quad - u_3 u_{1, 2} + u_{3, 3} \} \tag{4.37}
\end{aligned}$$

$$\begin{aligned}
& \triangleright \Gamma_{112} := -u_{1,1} : \\
& \Gamma_{113} := -(u_1 u_{1,1} + u_{1,2}) : \Gamma_{213} := -u_{1,1} : \\
& \Gamma_{114} := -(u_1^2 u_{1,1} + u_1 u_{1,2} + u_2 u_{1,1} + u_{1,3}) : \Gamma_{214} := -(u_1 u_{1,1} + u_{1,2}) : \Gamma_{314} := \\
& \quad -u_{1,1} : \\
& \Gamma_{121} := -\frac{1}{3} u_{1,1} : \\
& \Gamma_{122} := -u_{1,2} : \Gamma_{222} := -\frac{1}{3} u_{1,1} : \\
& \Gamma_{123} := -(u_2 u_{1,1} + u_{1,3}) : \Gamma_{223} := -u_{1,2} : \Gamma_{323} := -\frac{1}{3} u_{1,1} : \\
& \Gamma_{124} := -(u_1 u_2 u_{1,1} + u_2 u_{1,2} + u_3 u_{1,1} + u_{1,4}) : \Gamma_{224} := -(u_2 u_{1,1} + u_{1,3}) : \Gamma_{324} := \\
& \quad -u_{1,2} : \Gamma_{424} := -\frac{1}{3} u_{1,1} : \\
& \Gamma_{131} := -\frac{1}{3} (u_1 u_{1,1} + 2 u_{1,2}) : \Gamma_{132} := -u_{1,3} : \\
& \Gamma_{232} := -\frac{1}{3} (u_1 u_{1,1} + 2 u_{1,2}) : \Gamma_{233} := -u_{1,3} : \\
& \Gamma_{133} := -(u_3 u_{1,1} + u_{1,4}) : \Gamma_{333} := -\frac{1}{3} (u_1 u_{1,1} + 2 u_{1,2}) : \Gamma_{334} := -u_{1,3} : \\
& \Gamma_{134} := -(u_1 u_3 u_{1,1} + u_1 u_{1,4} + u_2 u_{1,3} + u_{3,3}) : \Gamma_{234} := -(u_3 u_{1,1} + u_{1,4}) : \Gamma_{434} := \\
& \quad -\frac{1}{3} (u_1 u_{1,1} + 2 u_{1,2}) : \\
& \Gamma_{141} := -\frac{1}{3} (u_1^2 u_{1,1} + u_1 u_{1,2} + 2 u_2 u_{1,1} + 3 u_{1,3}) : \\
& \Gamma_{142} := -u_{1,4} : \Gamma_{242} := -\frac{1}{3} (u_1^2 u_{1,1} + u_1 u_{1,2} + 2 u_2 u_{1,1} + 3 u_{1,3}) : \\
& \Gamma_{143} := -u_1 u_{1,4} - u_{2,4} : \Gamma_{243} := -u_{1,4} : \Gamma_{343} := -\frac{1}{3} (u_1^2 u_{1,1} + u_1 u_{1,2} + 2 u_2 u_{1,1} \\
& \quad + 3 u_{1,3}) : \\
& \Gamma_{144} := -(u_1^2 u_{1,4} + u_1 u_{2,4} + u_2 u_{1,4} + u_{3,4}) : \Gamma_{244} := -u_1 u_{1,4} - u_{2,4} : \Gamma_{344} := -u_{1,4} : \\
& \Gamma_{444} := -\frac{1}{3} (u_1^2 u_{1,1} + u_1 u_{1,2} + 2 u_2 u_{1,1} + 3 u_{1,3}) :
\end{aligned}$$

$$\begin{aligned}
& \triangleright \{ \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[1], vM[1], CTa), \\
& \quad \text{DGvector}))) , \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[1], vM[2], CTa), \\
& \quad \text{DGvector}) - \Gamma_{112} \cdot vM[1])) , \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[1], vM[3], CTa), \\
& \quad \text{DGvector}) - \Gamma_{113} \cdot vM[1] - \Gamma_{213} \cdot vM[2])) , \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[1], vM[4], CTa), \\
& \quad \text{DGvector}) - \Gamma_{114} \cdot vM[1] - \Gamma_{214} \cdot vM[2] - \Gamma_{314} \cdot vM[3])) , \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[2], vM[1], CTa), \\
& \quad \text{DGvector}) - \Gamma_{121} \cdot vM[1])) , \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[2], vM[2], CTa), \\
& \quad \text{DGvector}) - \Gamma_{122} \cdot vM[1] - \Gamma_{222} \cdot vM[2])) ,
\end{aligned}$$

$$\begin{aligned}
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[2], vM[3], CTa), \\
& \quad \text{DGvector}) - \Gamma_{123} \cdot vM[1] - \Gamma_{223} \cdot vM[2] - \Gamma_{323} \cdot vM[3]))), \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[2], vM[4], CTa), \\
& \quad \text{DGvector}) - \Gamma_{124} \cdot vM[1] - \Gamma_{224} \cdot vM[2] - \Gamma_{324} \cdot vM[3] - \Gamma_{424} \cdot vM[4]))), \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[3], vM[1], CTa), \\
& \quad \text{DGvector}) - \Gamma_{131} \cdot vM[1]))), \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[3], vM[2], CTa), \\
& \quad \text{DGvector}) - \Gamma_{132} \cdot vM[1] - \Gamma_{232} \cdot vM[2]))), \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[3], vM[3], CTa), \\
& \quad \text{DGvector}) - \Gamma_{133} \cdot vM[1] - \Gamma_{233} \cdot vM[2] - \Gamma_{333} \cdot vM[3]))), \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[3], vM[4], CTa), \\
& \quad \text{DGvector}) - \Gamma_{134} \cdot vM[1] - \Gamma_{234} \cdot vM[2] - \Gamma_{334} \cdot vM[3] - \Gamma_{434} \cdot vM[4]))), \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[4], vM[1], CTa), \\
& \quad \text{DGvector}) - \Gamma_{141} \cdot vM[1]))), \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[4], vM[2], CTa), \\
& \quad \text{DGvector}) - \Gamma_{142} \cdot vM[1] - \Gamma_{242} \cdot vM[2]))), \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[4], vM[3], CTa), \\
& \quad \text{DGvector}) - \Gamma_{143} \cdot vM[1] - \Gamma_{243} \cdot vM[2] - \Gamma_{343} \cdot vM[3]))), \\
& \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(vM[4], vM[4], CTa), \\
& \quad \text{DGvector}) - \Gamma_{144} \cdot vM[1] - \Gamma_{244} \cdot vM[2] - \Gamma_{344} \cdot vM[3] - \Gamma_{444} \cdot vM[4])))) \} \\
& \quad \{0 \ D_x\} \tag{4.38}
\end{aligned}$$

Thus we found all coefficients $\Gamma[ijk]=\Gamma_{jk}^i$ (current convention: $\nabla_j \partial_k = \Gamma_{jk}^i \partial_i$).

II. Let us check that there exists a unique symmetric connection preserving GL2-structure.

$$\begin{aligned}
& \text{ChrSeq} := \{\text{seq}(\text{seq}(\text{seq}(((a \parallel i) \parallel j) \parallel k), k = 1 \dots 4), j = i \dots 4), i = 1 \dots 4)\}; \\
& \text{CsC} := \text{Connection}(\text{subs}(\{\text{seq}(\text{seq}(\text{seq}(((a \parallel i) \parallel j) \parallel k) = ((a \parallel j) \parallel i) \parallel k), k = 1 \dots 4), j = 1 \\
& \quad \dots (i - 1)), i = 1 \dots 4\}, \text{add}(\text{add}(\text{add}(((a \parallel i) \parallel j) \parallel k) \cdot vM[k] \& \text{dM}[i] \& \text{dM}[j], k = 1 \\
& \quad \dots 4), j = 1 \dots 4), i = 1 \dots 4)): \text{TorsionTensor}(\text{CsC}) \\
& \quad ((0 \ D_x) \ dx) \ dx \tag{4.39}
\end{aligned}$$

$$\begin{aligned}
& \text{RNE} := \{ \} : \text{for } i \text{ to } 4 \text{ do for } j \text{ to } 3 \text{ do RNE} := \{ \text{op}(\text{RNE}), \\
& \quad \text{coeffs}(\text{expand}(\text{numer}(\text{simplify}(\text{subs}(S, \text{subs}(\text{Qsyst}, \\
& \quad \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(vM[i], \text{Quadu}[j], \text{CsC}), \omega \\
& \quad \& \omega, [[1, 1], [2, 2]]), \{u(x, y, t, z)\}, \text{notation} = \text{jetnumbers}))))), \{q_{[]}\} \} : \text{od:od:} \\
& \text{Sss1} := \text{solve}(\text{RNE}, \text{ChrSeq}) : \text{nops}(\text{Sss1}) \tag{4.40}
\end{aligned}$$

$$\begin{aligned}
& \text{CsS} := \text{DGsimplify}(\text{evalDG}(\text{subs}(\text{Sss1}, \text{CsC}))) : \\
& \text{seq}(\text{seq}(\text{simplify}(\text{subs}(S, \text{subs}(\text{Qsyst}, \\
& \quad \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(vM[i], \text{Quadu}[j], \text{CsS}), \omega \\
& \quad \& \omega, [[1, 1], [2, 2]]), \{u(x, y, t, z)\}, \text{notation} = \text{jetnumbers}))))), i = 1 \dots 4), j = 1 \dots 3) \\
& \quad 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0 \tag{4.41}
\end{aligned}$$

> S

$$\left\{ u_{1,2} = \frac{u_2^2 u_{1,1} - u_2 u_{2,2} + u_{2,4} - u_{3,3}}{u_1 u_2 - u_3}, u_{1,3} = \frac{u_2 u_3 u_{1,1} + u_1 u_{2,4} - u_1 u_{3,3} - u_3 u_{2,2}}{u_1 u_2 - u_3}, u_{1,4} \right. \quad (4.42)$$

$$\left. = \frac{u_1^2 u_{2,4} - u_1^2 u_{3,3} + u_1 u_2 u_{2,3} - u_1 u_3 u_{2,2} + u_3^2 u_{1,1} - u_3 u_{2,3}}{u_1 u_2 - u_3} \right\}$$

$$> S2a := \text{solve}(\{Eq1, Eq2, Eq3\}, \{u_{1,1}, u_{2,2}, u_{3,3}\});$$

$$S2b := \text{solve}(\{Eq1, Eq2, Eq3\}, \{u_{2,2}, u_{2,3}, u_{2,4}\});$$

$$S2c := \text{solve}(\{Eq1, Eq2, Eq3\}, \{u_{1,1}, u_{1,2}, u_{1,3}\});$$

$$S2a := \left\{ u_{1,1} = \frac{u_1 u_{1,3} - u_{1,4} + u_{2,3}}{u_3}, u_{2,2} \right.$$

$$\left. = \frac{u_1 u_2 u_{1,3} - u_1 u_3 u_{1,2} - u_2 u_{1,4} + u_2 u_{2,3} + u_3 u_{1,3}}{u_3}, u_{3,3} = -u_2 u_{1,3} + u_3 u_{1,2} + u_{2,4} \right\}$$

$$S2b := \{u_{2,2} = -u_1 u_{1,2} + u_2 u_{1,1} + u_{1,3}, u_{2,3} = -u_1 u_{1,3} + u_3 u_{1,1} + u_{1,4}, u_{2,4} = u_2 u_{1,3} - u_3 u_{1,2} + u_{3,3}\} \quad (4.43)$$

$$> \Pi_{12} := -\frac{2}{3} u_{1,1} :$$

$$\Pi_{13} := -\frac{2}{3} u_1 u_{1,1} - u_{1,2} : \Gamma_{213} := -\frac{1}{3} u_{1,1} :$$

$$\Pi_{14} := -\frac{2}{3} u_1^2 u_{1,1} - u_1 u_{1,2} - \frac{1}{3} u_2 u_{1,1} - u_{1,3} : \Gamma_{214} := -\frac{2}{3} u_1 u_{1,1} - u_{1,2} :$$

$$\Pi_{22} := -\frac{8}{9} u_{1,2} : \Gamma_{222} := -\frac{4}{9} u_{1,1} : \Pi_{23} := \frac{1}{9} u_1 u_{1,2} - \frac{5}{9} u_2 u_{1,1} - u_{1,3} : \Gamma_{223} :=$$

$$-\frac{2}{9} u_1 u_{1,1} - \frac{7}{9} u_{1,2} : \Gamma_{323} := -\frac{2}{9} u_{1,1} :$$

$$\Pi_{24} := \frac{1}{9} u_1^2 u_{1,2} - \frac{7}{9} u_1 u_2 u_{1,1} - \frac{7}{9} u_2 u_{1,2} - \frac{1}{3} u_3 u_{1,1} - u_{1,4} : \Gamma_{224} := \frac{1}{9} u_1 u_{1,2}$$

$$- \frac{5}{9} u_2 u_{1,1} - u_{1,3} : \Gamma_{324} := -\frac{4}{9} u_1 u_{1,1} - \frac{2}{3} u_{1,2} :$$

$$\Pi_{33} := \frac{1}{9} u_1^2 u_{1,2} + \frac{4}{9} u_2 u_{1,2} - \frac{10}{9} u_3 u_{1,1} - u_{1,4} : \Gamma_{233} := -\frac{1}{9} u_1^2 u_{1,1} - \frac{4}{9} u_2 u_{1,1}$$

$$- u_{1,3} : \Gamma_{333} := -\frac{2}{9} u_1 u_{1,1} - \frac{5}{9} u_{1,2} : \Gamma_{433} := -\frac{1}{9} u_{1,1} :$$

$$\Pi_{34} := \frac{1}{9} u_1^3 u_{1,2} - \frac{1}{9} u_1^2 u_2 u_{1,1} + \frac{4}{9} u_1 u_2 u_{1,2} - \frac{5}{9} u_1 u_3 u_{1,1} - \frac{4}{9} u_2^2 u_{1,1} - u_1 u_{1,4}$$

$$- \frac{1}{3} u_3 u_{1,2} - u_{2,4} :$$

$$\Gamma_{234} := \frac{1}{9} u_1^2 u_{1,2} + \frac{4}{9} u_2 u_{1,2} - \frac{2}{3} u_3 u_{1,1} - u_{1,4} : \Gamma_{334} := -\frac{2}{9} u_1^2 u_{1,1} - \frac{1}{9} u_1 u_{1,2}$$

$$- \frac{7}{9} u_2 u_{1,1} - u_{1,3} : \Gamma_{434} := -\frac{2}{9} u_1 u_{1,1} - \frac{1}{3} u_{1,2} :$$

$$\Pi_{44} := \frac{1}{9} u_1^4 u_{1,2} - \frac{1}{9} u_1^3 u_2 u_{1,1} + \frac{5}{9} u_1^2 u_2 u_{1,2} - \frac{1}{9} u_1^2 u_3 u_{1,1} - \frac{4}{9} u_1 u_2^2 u_{1,1} -$$

$$u_1^2 u_{1,4} - u_1 u_{2,4} + \frac{2}{3} u_1 u_3 u_{1,2} + \frac{4}{9} u_2^2 u_{1,2} - \frac{4}{3} u_2 u_{1,1} u_3 - u_2 u_{1,4} - u_{3,4} :$$

$$\begin{aligned}
\Gamma_{244} &:= \frac{1}{9} u_1^3 u_{1,2} - \frac{1}{9} u_1^2 u_2 u_{1,1} + \frac{4}{9} u_1 u_2 u_{1,2} + \frac{1}{3} u_1 u_3 u_{1,1} - \frac{4}{9} u_2^2 u_{1,1} - u_1 u_{1,4} \\
&\quad + u_3 u_{1,2} - u_{2,4} : \\
\Gamma_{344} &:= -\frac{1}{9} u_1 u_{2,2} - \frac{8}{9} u_1 u_{1,3} + \frac{1}{3} u_2 u_{1,2} - u_{2,3} : \Gamma_{444} := -\frac{4}{9} u_1^2 u_{1,1} - \frac{4}{3} u_1 u_{1,2} \\
&\quad - u_{2,2} :
\end{aligned}$$

$$\begin{aligned}
> \{ \text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(\text{vM}[1], \text{vM}[1], \text{CsS}), \\
&\quad \text{DGvector}))) , \\
&\text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(\text{vM}[1], \text{vM}[2], \text{CsS}), \\
&\quad \text{DGvector}) - \Gamma_{112} \cdot \text{vM}[1])) , \\
&\text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(\text{vM}[1], \text{vM}[3], \text{CsS}), \\
&\quad \text{DGvector}) - \Gamma_{113} \cdot \text{vM}[1] - \Gamma_{213} \cdot \text{vM}[2])) , \\
&\text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(\text{vM}[1], \text{vM}[4], \text{CsS}), \\
&\quad \text{DGvector}) - \Gamma_{114} \cdot \text{vM}[1] - \Gamma_{214} \cdot \text{vM}[2])) , \\
&\text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(\text{vM}[2], \text{vM}[2], \text{CsS}), \\
&\quad \text{DGvector}) - \Gamma_{122} \cdot \text{vM}[1] - \Gamma_{222} \cdot \text{vM}[2])) , \\
&\text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(\text{vM}[2], \text{vM}[3], \text{CsS}), \\
&\quad \text{DGvector}) - \Gamma_{123} \cdot \text{vM}[1] - \Gamma_{223} \cdot \text{vM}[2] - \Gamma_{323} \cdot \text{vM}[3])) , \\
&\text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(\text{vM}[2], \text{vM}[4], \text{CsS}), \\
&\quad \text{DGvector}) - \Gamma_{124} \cdot \text{vM}[1] - \Gamma_{224} \cdot \text{vM}[2] - \Gamma_{324} \cdot \text{vM}[3])) , \\
&\text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(\text{vM}[3], \text{vM}[3], \text{CsS}), \\
&\quad \text{DGvector}) - \Gamma_{133} \cdot \text{vM}[1] - \Gamma_{233} \cdot \text{vM}[2] - \Gamma_{333} \cdot \text{vM}[3] - \Gamma_{433} \cdot \text{vM}[4])) , \\
&\text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(\text{vM}[3], \text{vM}[4], \text{CsS}), \\
&\quad \text{DGvector}) - \Gamma_{134} \cdot \text{vM}[1] - \Gamma_{234} \cdot \text{vM}[2] - \Gamma_{334} \cdot \text{vM}[3] - \Gamma_{434} \cdot \text{vM}[4])) , \\
&\text{simplify}(\text{evalDG}(\text{subs}(S, \text{convert}(\text{DirectionalCovariantDerivative}(\text{vM}[4], \text{vM}[4], \text{CsS}), \\
&\quad \text{DGvector}) - \Gamma_{144} \cdot \text{vM}[1] - \Gamma_{244} \cdot \text{vM}[2] - \Gamma_{344} \cdot \text{vM}[3] - \Gamma_{444} \cdot \text{vM}[4]))) \} \\
&\quad \{0 \ D_x\} \tag{4.44}
\end{aligned}$$

>

III. Finally let us compute the canonical projective totally geodesic connection

$$\begin{aligned}
> \text{CsC} &:= \text{Connection}(\text{subs}(\{ \text{seq}(\text{seq}(\text{seq}(((a \parallel i) \parallel j) \parallel k) = (((a \parallel j) \parallel i) \parallel k), k = 1..4), j = 1 \\
&\quad ..(i-1)), i = 1..4) \}, \text{add}(\text{add}(\text{add}(((a \parallel i) \parallel j) \parallel k) \cdot \text{vM}[k] \&t \text{dM}[i] \&t \text{dM}[j], k = 1 \\
&\quad ..4), j = 1..4), i = 1..4)) : \\
> \text{ChrCoe} &:= \{ \text{seq}(\text{seq}(\text{seq}(((a \parallel i) \parallel j) \parallel k), k = 1..4), j = i..4), i = 1..4 \} : \\
&\text{TGE} := \{ \} : \text{for } i \text{ to } 3 \text{ do for } j \text{ to } 3 \text{ do } \text{TGE} := \{ \text{op}(\text{TGE}), \text{coeffs}(\text{expand}(\text{subs}(S, \\
&\quad \text{subs}(\text{Qsyst}, \text{ToJet}(\text{Hook}(\text{convert}(\text{DirectionalCovariantDerivative}(\text{DD}[i], \text{DD}[j], \text{CsC}), \\
&\quad \text{'DGvector'}), \omega\omega), \{u(x, y, t, z), q(x, y, t, z)\}, \text{notation} = \text{jetnumbers})) , \{q_{[1]}, q_1\}) \} : \\
&\quad \text{od:od:} \\
&\quad [\text{nops}(\text{ChrCoe}), \text{nops}(\text{TGE})] \tag{4.45}
\end{aligned}$$

[40, 45]

$$\begin{aligned}
> \text{Sss2} &:= \text{solve}(\text{TGE}, \text{ChrCoe} \setminus \{a_{144}, a_{244}, a_{344}, a_{444}\}) : \text{nops}(\text{Sss2}) \\
&\quad 36 \tag{4.46}
\end{aligned}$$

$$\begin{aligned}
> \text{Sss20} &:= \text{simplify} \left(\left\{ \text{op} \left(\text{subs} \left(\left\{ a_{144} = 0, a_{244} = 0, a_{344} = \frac{1}{2} u_{1,2}, a_{444} = u_{1,3} \right\}, \text{subs}(S2b, \right. \right. \right.
\end{aligned}$$

$$\begin{aligned}
\Gamma_{141} &:= -\frac{1}{2} u_1^2 u_{1,1} - \frac{1}{2} u_1 u_{1,2} - \frac{1}{2} u_2 u_{1,1}; \Gamma_{142} := -\frac{1}{2} u_1 u_{1,1} - \frac{1}{2} u_{1,2}; \Gamma_{143} := \\
&\quad -\frac{1}{2} u_{1,1}; \\
\Gamma_{221} &:= -u_{1,2}; \\
\Gamma_{231} &:= -\frac{1}{2} u_2 u_{1,1} - u_{1,3}; \\
\Gamma_{241} &:= -\frac{1}{2} u_1 u_2 u_{1,1} - \frac{1}{2} u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} - u_{1,4}; \Gamma_{242} := -\frac{1}{2} u_2 u_{1,1}; \Gamma_{243} := \\
&\quad -\frac{1}{2} u_{1,2}; \\
\Gamma_{331} &:= -u_3 u_{1,1} - u_{1,4}; \Gamma_{332} := -u_{1,3}; \Gamma_{333} := u_{1,2}; \\
\Gamma_{341} &:= -\frac{1}{2} u_1 u_3 u_{1,1} - u_1 u_{1,4} - u_{2,4} - \frac{1}{2} u_3 u_{1,2}; \Gamma_{342} := -\frac{1}{2} u_3 u_{1,1} - u_{1,4}; \Gamma_{344} \\
&:= \frac{1}{2} u_{1,2}; \\
\Gamma_{441} &:= -u_1^2 u_{1,4} - u_1 u_{2,4} - u_2 u_{1,4} - u_{3,4}; \Gamma_{442} := -u_1 u_{1,4} - u_{2,4}; \Gamma_{443} := -u_{1,4}; \Gamma_{444} \\
&:= u_{1,3};
\end{aligned}$$

> CsC0 := subs(Sss20, CsC) :

> TGE := { } :for i to 3 do for j to 3 do TGE := {op(TGE), coeffs(expand(subs(S, subs(Qsyst, ToJet(Hook(convert(DirectionalCovariantDerivative(DD[i], DD[j], FromJet(CsC0, {u(x, y, t, z), q(x, y, t, z)})), 'DGvector'), ωω), {u(x, y, t, z), q(x, y, t, z)}, notation = jetnumbers))), {q_[j], q_[i]))} :od:od: simplify(subs(S, TGE))
{0} (4.50)

> CovariantDerivative(vM[1], CsC0)

$$\begin{aligned}
& - \left(\left(\frac{u_{1,1}}{2} D_{-x} \right) dy \right) - \left(\left(\frac{u_1 u_{1,1}}{2} D_{-x} \right) dt \right) - \left(\left(\left(\frac{1}{2} u_1^2 u_{1,1} + \frac{1}{2} u_1 u_{1,2} \right. \right. \right. \\
& \quad \left. \left. + \frac{1}{2} u_2 u_{1,1} \right) D_{-x} \right) dz \right) - \left(\left(\frac{u_{1,1}}{2} D_{-y} \right) dt \right) - \left(\left(\left(\frac{u_1 u_{1,1}}{2} + \frac{u_{1,2}}{2} \right) D_{-y} \right) dz \right) \\
& \quad - \left(\left(\frac{u_{1,1}}{2} D_{-t} \right) dz \right)
\end{aligned} \tag{4.51}$$

> DirectionalCovariantDerivative(vM[1], vM[1], CsC0);
DirectionalCovariantDerivative(vM[1], vM[2], CsC0);
DirectionalCovariantDerivative(vM[1], vM[3], CsC0);
DirectionalCovariantDerivative(vM[1], vM[4], CsC0);
DirectionalCovariantDerivative(vM[2], vM[2], CsC0);
DirectionalCovariantDerivative(vM[2], vM[3], CsC0);
DirectionalCovariantDerivative(vM[2], vM[4], CsC0);
DirectionalCovariantDerivative(vM[3], vM[3], CsC0);
DirectionalCovariantDerivative(vM[3], vM[4], CsC0);
DirectionalCovariantDerivative(vM[4], vM[4], CsC0);
0 D_{-x}

$$\begin{aligned}
& - \left(\frac{u_{1,1}}{2} D_{-x} \right) \\
& - \left(\frac{u_1 u_{1,1}}{2} D_{-x} \right) - \left(\frac{u_{1,1}}{2} D_{-y} \right) \\
& - \left(\left(\frac{1}{2} u_1^2 u_{1,1} + \frac{1}{2} u_1 u_{1,2} + \frac{1}{2} u_2 u_{1,1} \right) D_{-x} \right) - \left(\left(\frac{u_1 u_{1,1}}{2} + \frac{u_{1,2}}{2} \right) D_{-y} \right) \\
& - \left(\frac{u_{1,1}}{2} D_{-t} \right) \\
& - (u_{1,2} D_{-x}) \\
& - \left(\left(\frac{u_2 u_{1,1}}{2} + u_{1,3} \right) D_{-x} \right) \\
& - \left(\left(\frac{1}{2} u_1 u_2 u_{1,1} + \frac{1}{2} u_2 u_{1,2} + \frac{1}{2} u_3 u_{1,1} + u_{1,4} \right) D_{-x} \right) - \left(\frac{u_2 u_{1,1}}{2} D_{-y} \right) \\
& - \left(\frac{u_{1,2}}{2} D_{-t} \right) \\
& - ((u_3 u_{1,1} + u_{1,4}) D_{-x}) - (u_{1,3} D_{-y}) + u_{1,2} D_{-t} \\
& - \left(\left(\frac{1}{2} u_1 u_3 u_{1,1} + u_1 u_{1,4} + u_2 u_{1,3} - \frac{1}{2} u_3 u_{1,2} + u_{3,3} \right) D_{-x} \right) - \left(\left(\frac{u_3 u_{1,1}}{2} \right. \right. \\
& \quad \left. \left. + u_{1,4} \right) D_{-y} \right) + \frac{u_{1,2}}{2} D_{-z} \\
& - \left((u_1^2 u_{1,4} + u_1 u_2 u_{1,3} - u_1 u_3 u_{1,2} + u_1 u_{3,3} + u_2 u_{1,4} + u_{3,4}) D_{-x} \right) - ((u_1 u_{1,4} + u_2 u_{1,3} \\
& \quad - u_3 u_{1,2} + u_{3,3}) D_{-y}) - (u_{1,4} D_{-t}) + u_{1,3} D_{-z} \quad (4.52)
\end{aligned}$$

dKP hierarchy 5D

> DGsetup([x,y,t,z,s],[u],J,4):

> Eq1 := u_{1,3} - u_{2,2} - u₁ u_{1,1}:

Eq2 := u_{1,4} - u_{2,3} - u₁ u_{1,2} - u₂ u_{1,1}:

Eq3 := u_{2,4} - u_{3,3} + u₁² u_{1,1} - u₂ u_{1,2}:

Eq4 := u_{1,5} - u_{2,4} - u₁ u_{2,2} - u₂ u_{1,2} - (u₃ + u₁²) u_{1,1}:

Eq5 := u_{2,5} - u_{3,4} + u₁ u₂ u_{1,1} + (u₁² - u₃) u_{1,2} - u₂ u_{2,2}:

Eq6 := u_{3,5} - u_{4,4} + (u₂² - u₁³ - u₁ u₃) u_{1,1} + 2 u₁ u₂ u_{1,2} - u₃ u_{2,2}:

EQ := {Eq1, Eq2, Eq3, Eq4, Eq5, Eq6}:

> S2 := solve(EQ, {u_{1,3}, u_{1,4}, u_{1,5}, u_{2,4}, u_{2,5}, u_{3,5}}):

S2 := {u_{1,3} = u₁ u_{1,1} + u_{2,2}, u_{1,4} = u₁ u_{1,2} + u₂ u_{1,1} + u_{2,3}, u_{1,5} = u₁ u_{2,2} + 2 u₂ u_{1,2}

(5.1)

$$+ u_3 u_{1,1} + u_{3,3} u_{2,4} = -u_1^2 u_{1,1} + u_2 u_{1,2} + u_{3,3} u_{2,5} = -u_1^2 u_{1,2} - u_1 u_2 u_{1,1} + u_2 u_{2,2} \\ + u_3 u_{1,2} + u_{3,4} u_{3,5} = u_1^3 u_{1,1} - 2 u_1 u_2 u_{1,2} + u_1 u_3 u_{1,1} - u_2^2 u_{1,1} + u_3 u_{2,2} + u_{4,4} \}$$

> S2a := solve(EQ, {u_{2,2}, u_{2,3}, u_{2,4}, u_{3,3}, u_{3,4}, u_{4,4}})

$$S2a := \{u_{2,2} = -u_1 u_{1,1} + u_{1,3}, u_{2,3} = -u_1 u_{1,2} - u_2 u_{1,1} + u_{1,4}, u_{2,4} = -u_1 u_{1,3} - u_2 u_{1,2} \quad (5.2)$$

$$- u_3 u_{1,1} + u_{1,5}, u_{3,3} = u_1^2 u_{1,1} - u_1 u_{1,3} - 2 u_2 u_{1,2} - u_3 u_{1,1} + u_{1,5}, u_{3,4} = u_1^2 u_{1,2} \\ + 2 u_1 u_2 u_{1,1} - u_2 u_{1,3} - u_3 u_{1,2} + u_{2,5}, u_{4,4} = -u_1^3 u_{1,1} + 2 u_1 u_2 u_{1,2} + u_2^2 u_{1,1} \\ - u_3 u_{1,3} + u_{3,5} \}$$

> Symb1 := add(add(diff(Eq1, u_{i,j}) · p_i · p_j, j = 1..5), i = 1..5);

Symb2 := add(add(diff(Eq2, u_{i,j}) · p_i · p_j, j = 1..5), i = 1..5);

Symb3 := add(add(diff(Eq3, u_{i,j}) · p_i · p_j, j = 1..5), i = 1..5);

Symb4 := add(add(diff(Eq4, u_{i,j}) · p_i · p_j, j = 1..5), i = 1..5);

Symb5 := add(add(diff(Eq5, u_{i,j}) · p_i · p_j, j = 1..5), i = 1..5);

Symb6 := add(add(diff(Eq6, u_{i,j}) · p_i · p_j, j = 1..5), i = 1..5);

$$Symb1 := -p_1^2 u_1 + p_1 p_3 - p_2^2$$

$$Symb2 := -p_1^2 u_2 - p_1 p_2 u_1 + p_1 p_4 - p_2 p_3$$

$$Symb3 := p_1^2 u_1^2 - p_1 p_2 u_2 + p_2 p_4 - p_3^2$$

$$Symb4 := (-u_1^2 - u_3) p_1^2 - u_2 p_1 p_2 + p_1 p_5 - u_1 p_2^2 - p_2 p_4$$

$$Symb5 := u_1 u_2 p_1^2 + (u_1^2 - u_3) p_1 p_2 - u_2 p_2^2 + p_2 p_5 - p_3 p_4$$

$$Symb6 := (-u_1^3 - u_1 u_3 + u_2^2) p_1^2 + 2 u_1 u_2 p_1 p_2 - u_3 p_2^2 + p_3 p_5 - p_4^2 \quad (5.3)$$

> simplify(subs({p₁ = 1, p₂ = q, p₃ = q² + u₁, p₄ = q³ + 2 u₁ q + u₂, p₅ = q⁴ + 3 u₁ q² + 2 u₂ q + u₃ + u₁²}, {Symb1, Symb2, Symb3, Symb4, Symb5, Symb6}))

$$\{0\} \quad (5.4)$$

> tmp := solve({coeffs(expand((a1 · p₁ + a2 · p₂ + a3 · p₃ + a4 · p₄ + a5 · p₅) · Symb1 + (b1 · p₁ + b2 · p₂ + b3 · p₃ + b4 · p₄ + b5 · p₅) · Symb2 + (c1 · p₁ + c2 · p₂ + c3 · p₃ + c4 · p₄ + c5 · p₅) · Symb3 + (d1 · p₁ + d2 · p₂ + d3 · p₃ + d4 · p₄ + d5 · p₅) · Symb4 + (e1 · p₁ + e2 · p₂ + e3 · p₃ + e4 · p₄ + e5 · p₅) · Symb5 + (f1 · p₁ + f2 · p₂ + f3 · p₃ + f4 · p₄ + f5 · p₅) · Symb6), {p₁, p₂, p₃, p₄, p₅}), {a1, a2, a3, a4, a5, b1, b2, b3, b4, b5, c1, c2, c3, c4, c5, d1, d2, d3, d4, d5, e1, e2, e3, e4, e5, f1, f2, f3, f4, f5})

$$tmp := \{a1 = f3 u_1^3 - e2 u_1^2 - e3 u_1 u_2 - f1 u_1^2 + f3 u_1 u_3 - f3 u_2^2 + c1 u_1 - c2 u_2 - e2 u_3 \quad (5.5)$$

$$- f1 u_3, a2 = -e3 u_1^2 - f2 u_1^2 - f3 u_1 u_2 + e1 u_1 - e2 u_2 - f2 u_3, a3 = c1, a4 = c2 + e1, a5$$

$$= e2, b1 = -f2 u_1^2 + f3 u_1 u_2 + c2 u_1 + e1 u_1 - e3 u_3 + f1 u_2 - f2 u_3, b2 = e2 u_1 - e3 u_2$$

$$+ f1 u_1 - f3 u_3 - c1, b3 = -c2, b4 = f1, b5 = e3 + f2, c1 = c1, c2 = c2, c3 = 0, c4 = -e3,$$

$$c5 = f3, d1 = -f3 u_1^2 + e2 u_1 + e3 u_2 + f2 u_2, d2 = e3 u_1 + f2 u_1 + f3 u_2 - e1, d3 = -f1$$

$-e_2, d_4 = -e_3 - f_2, d_5 = 0, e_1 = e_1, e_2 = e_2, e_3 = e_3, e_4 = -f_3, e_5 = 0, f_1 = f_1, f_2 = f_2, f_3 = f_3, f_4 = 0, f_5 = 0\}$

so 8 compatibility conditions parametrized by $(c_1, c_2, e_1, e_2, e_3, f_1, f_2, f_3)$ - we check them at one step

```
> simplify(subs(S2, subs(tmp, add((a||i)·TotalDiff(Eq1, i), i = 1..5) + add((b||i)
·TotalDiff(Eq2, i), i = 1..5) + add((c||i)·TotalDiff(Eq3, i), i = 1..5) + add((d||i)
·TotalDiff(Eq4, i), i = 1..5) + add((e||i)·TotalDiff(Eq5, i), i = 1..5) + add((f||i)
·TotalDiff(Eq6, i), i = 1..5))))
0 (5.6)
```

```
>
> Vss3 := EQ:for i to 5 do for j to nops(EQ) do Vss3 := {op(Vss3), TotalDiff(EQ[j], i)}:od:
od:
var3 := {u1,3, u1,4, u1,5, u2,4, u2,5, u3,5, u1,1,2, u1,1,3, u1,1,4, u1,1,5, u1,2,3, u1,2,4, u1,2,5,
u1,3,3, u1,3,4, u1,3,5, u1,4,4, u1,4,5, u2,2,4, u2,3,3, u2,3,4, u2,3,5, u2,4,4, u2,5,5, u3,3,3,
u3,3,5, u3,4,4, u3,4,5}:
Vvv3 := eliminate(Vss3, var3):[nops(Vss3), nops(var3), nops(Vvv3[1]), nops(Vvv3[2])]
[36, 28, 28, 0] (5.7)
```

```
> S3 := Vvv3[1]:
> tmp := {}:for i to nops(S3) do if lhs(S3[i]) = rhs(S3[i]) then tmp := {op(tmp),
lhs(S3[i])}:fi:od: tmp
{} (5.8)
```

```
> 35 - (28 - 6)
13 (5.9)
```

This shows that dimensions of the symbol spaces $(g[0], g[1], g[2], g[3], g[4], \dots)$ are equal to $(1, 5, 9, 13, 17, \dots)$,

i.e. the Hilbert polynomial is given by the formula $\dim(g[k]) = 1 + 4k$, whence the general solution depends on 4 functions of 1 variable.

This is also what is expected from the characteristic variety (rational normal curve in our case) that has order 4 and dimension 1.

```
>
> DGsetup([x, y, t, z, s], [u, q], Jq, 3):
> Qsyst := {q2 = q[ ] q1 + u1,1, q3 = (q[ ]2 + u1) q1 + u1,1 q[ ] + u1,2, q4 = (q[ ]3 + 2 u1 q[ ]
+ u2) q1 + u1,1 q[ ]2 + u1,2 q[ ] + u1,3 + u1 u1,1, q5 = (q[ ]4 + 3 u1 q[ ]2 + 2 u2 q[ ] + u3 +
u12) q1 + u1,1 q[ ]3 + u1,2 q[ ]2 + (u1,3 + 2 u1 u1,1) q[ ] + u1,4 + u2 u1,1 + u1 u1,2}:
> simplify(subs(S3, expand(subs(Qsyst, subs({q1,2 = TotalDiff(subs(Qsyst, q2), 1), q1,3
= TotalDiff(subs(Qsyst, q3), 1), q1,4 = TotalDiff(subs(Qsyst, q4), 1), q1,5
= TotalDiff(subs(Qsyst, q5), 1)}, {TotalDiff(subs(Qsyst, q3), 2)
- TotalDiff(subs(Qsyst, q2), 3), TotalDiff(subs(Qsyst, q4), 2)
- TotalDiff(subs(Qsyst, q2), 4), TotalDiff(subs(Qsyst, q5), 2)
- TotalDiff(subs(Qsyst, q2), 5), TotalDiff(subs(Qsyst, q4), 3)
- TotalDiff(subs(Qsyst, q3), 4), TotalDiff(subs(Qsyst, q5), 3)
- TotalDiff(subs(Qsyst, q3), 5), TotalDiff(subs(Qsyst, q5), 4)
- TotalDiff(subs(Qsyst, q4), 5)}))))
{0} (5.10)
```

```

> DGsetup([x, y, t, z, s, q], [u], Jqu, 3) :
> Yf := f → TotalDiff(f, 2) - q · TotalDiff(f, 1) + u1,1 · diff(f, q) :
Tf := f → TotalDiff(f, 3) - (q2 + u1) · TotalDiff(f, 1) + (q · u1,1 + u1,2) · diff(f, q) :
Zf := f → TotalDiff(f, 4) - (q3 + 2 u1 q + u2) · TotalDiff(f, 1) + (q2 u1,1 + q u1,2 + u1,3
+ u1 u1,1) · diff(f, q) :
Sf := f → TotalDiff(f, 5) - (q4 + 3 u1 q2 + 2 u2 q + u3 + u12) · TotalDiff(f, 1) + (q3 u1,1 +
q2 u1,2 + q (u1,3 + 2 u1 u1,1) + u1,4 + u2 u1,1 + u1 u1,2) · diff(f, q) :
> simplify(subs(S3, {simplify(Yf(Tf(x)) - Tf(Yf(x))), simplify(Yf(Tf(q)) - Tf(Yf(q))),
simplify(Yf(Zf(x)) - Zf(Yf(x))), simplify(Yf(Zf(q)) - Zf(Yf(q))),
simplify(Yf(Sf(x)) - Sf(Yf(x))), simplify(Yf(Sf(q)) - Sf(Yf(q))),
simplify(Tf(Zf(x)) - Zf(Tf(x))), simplify(Tf(Zf(q)) - Zf(Tf(q))),
simplify(Tf(Sf(x)) - Sf(Tf(x))), simplify(Tf(Sf(q)) - Sf(Tf(q))),
simplify(Zf(Sf(x)) - Sf(Zf(x))), simplify(Zf(Sf(q)) - Sf(Zf(q))) })),
{0})
(5.11)

```

I. We first look for a connection (with torsion) preserving the field of GL2-curves and having α -surfaces totally geodesic + trace of torsion zero.

```

> DGsetup([x, y, t, z, s], M) :
> vM := [D_x, D_y, D_t, D_z, D_s] : dM := [dx, dy, dt, dz, ds] : var := x, y, t, z, s :
> C := Connection(add(add(add(((a || i) || j) || k) · vM[k] &t dM[i] &t dM[j], k = 1 .. 5), j = 1
.. 5), i = 1 .. 5)) :
> DirectionalCovariantDerivative(D_x, dt, C); DirectionalCovariantDerivative(D_x, D_t, C)
- (a113 dx) - (a213 dy) - (a313 dt) - (a413 dz) - (a513 ds)
a311 D_x + a312 D_y + a313 D_t + a314 D_z + a315 D_s
(5.12)
> Y := evalDG(D_y - q[ ] · D_x) :
T := evalDG(D_t - (q[ ]2 + u1) · D_x) :
Z := evalDG(D_z - (q[ ]3 + 2 u1 q[ ] + u2) · D_x) :
S := evalDG(D_s - (q[ ]4 + 3 u1 q[ ]2 + 2 u2 q[ ] + u3 + u12) · D_x) :
ω := evalDG(dx + q[ ] · dy + (q[ ]2 + u1) · dt + (q[ ]3 + 2 u1 q[ ] + u2) · dz + (q[ ]4 + 3 u1 q[ ]2
+ 2 u2 q[ ] + u3 + u12) · ds) :
[Hook(Y, ω), Hook(T, ω), Hook(Z, ω), Hook(S, ω)]
[0, 0, 0, 0]
(5.13)

```

```

> YY := FromJet(Y, {u(x, y, t, z, s), q(x, y, t, z, s)}) : TT := FromJet(T, {u(x, y, t, z, s), q(x, y,
t, z, s)}) : ZZ := FromJet(Z, {u(x, y, t, z, s), q(x, y, t, z, s)}) : SS := FromJet(S, {u(x, y,
t, z, s), q(x, y, t, z, s)}) : ωω := FromJet(ω, {u(x, y, t, z, s), q(x, y, t, z, s)}) :
> DD := {YY, ZZ, TT, SS} : ChrCoe := {seq(seq(seq(((a || i) || j) || k), i = 1 .. 5), j = 1 .. 5), k
= 1 .. 5)} :
TGE := { } : for i to 4 do for j to 4 do TGE := {op(TGE), coeffs(expand(subs(S2, subs
(Qsyst, ToJet(Hook(convert(DirectionalCovariantDerivative(DD[i], DD[j], C),
'DGvector'), ωω), {u(x, y, t, z, s), q(x, y, t, z, s)}, notation = jetnumbers)) ), {q[ ], q1})
} : od : od :

```

$$[nops(ChrCoe), nops(TGE)] \quad [125, 132] \quad (5.14)$$

$$\begin{aligned} &> Sconn := solve(TGE, ChrCoe) : \\ &> FreePar := \{ \} : \text{for } i \text{ to } nops(Sconn) \text{ do if } lhs(Sconn[i]) = rhs(Sconn[i]) \text{ then } FreePar \\ &\quad := \{op(FreePar), lhs(Sconn[i])\} \text{ fi:od: } FreePar \\ &\quad \{a155, a255, a355, a455, a514, a515, a524, a525, a534, a535, a544, a545, a555\} \end{aligned} \quad (5.15)$$

$$\begin{aligned} &> ChrCoe1 := ChrCoe \setminus FreePar \\ ChrCoe1 &:= \{a111, a112, a113, a114, a115, a121, a122, a123, a124, a125, a131, a132, \\ &\quad a133, a134, a135, a141, a142, a143, a144, a145, a151, a152, a153, a154, a211, a212, \\ &\quad a213, a214, a215, a221, a222, a223, a224, a225, a231, a232, a233, a234, a235, a241, \\ &\quad a242, a243, a244, a245, a251, a252, a253, a254, a311, a312, a313, a314, a315, a321, \\ &\quad a322, a323, a324, a325, a331, a332, a333, a334, a335, a341, a342, a343, a344, a345, \\ &\quad a351, a352, a353, a354, a411, a412, a413, a414, a415, a421, a422, a423, a424, a425, \\ &\quad a431, a432, a433, a434, a435, a441, a442, a443, a444, a445, a451, a452, a453, a454, \\ &\quad a511, a512, a513, a521, a522, a523, a531, a532, a533, a541, a542, a543, a551, a552, \\ &\quad a553, a554\} \end{aligned} \quad (5.16)$$

$$\begin{aligned} &> [nops(ChrCoe), nops(FreePar), nops(ChrCoe1)] \\ &\quad [125, 13, 112] \end{aligned} \quad (5.17)$$

$$\begin{aligned} &> FreePar := \{a155, a255, a355, a455, a514, a515, a524, a525, a534, a535, a544, a545, \\ &\quad a555\} : \\ &> ChrCoe1 := \{a111, a112, a113, a114, a115, a121, a122, a123, a124, a125, a131, a132, \\ &\quad a133, a134, a135, a141, a142, a143, a144, a145, a151, a152, a153, a154, a211, a212, \\ &\quad a213, a214, a215, a221, a222, a223, a224, a225, a231, a232, a233, a234, a235, a241, \\ &\quad a242, a243, a244, a245, a251, a252, a253, a254, a311, a312, a313, a314, a315, a321, \\ &\quad a322, a323, a324, a325, a331, a332, a333, a334, a335, a341, a342, a343, a344, a345, \\ &\quad a351, a352, a353, a354, a411, a412, a413, a414, a415, a421, a422, a423, a424, a425, \\ &\quad a431, a432, a433, a434, a435, a441, a442, a443, a444, a445, a451, a452, a453, a454, \\ &\quad a511, a512, a513, a521, a522, a523, a531, a532, a533, a541, a542, a543, a551, a552, \\ &\quad a553, a554\} : \end{aligned}$$

$$\begin{aligned} &> Sconn := solve(TGE, ChrCoe1) : \\ &> CC := DGsimplify(subs(Sconn, C)) : \\ &> \text{for } k \text{ to } 6 \text{ do } Quad||k := evalDG(add(add(diff(Eq||k, u_{i,j}) \cdot vM[i] \&s vM[j], j=i..5), i=1 \\ &\quad ..5)) : \text{od:} \\ &\quad \text{for } k \text{ to } 6 \text{ do } Quad||k := FromJet(Quad||k, \{u(x, y, t, z, s), q(x, y, t, z, s)\}, notation \\ &\quad = jetnumbers) : \text{od:} \\ &\quad Quad|| := [seq(Quad||k, k=1..6)] : \\ &> \{seq(ContractIndices(Quad||i, \omega \&t \omega, [[1, 1], [2, 2]]), i=1..6)\} \\ &\quad \{0\} \end{aligned} \quad (5.18)$$

$$\begin{aligned} &> ContractIndices(TorsionTensor(CC), [[1, 2]]) \\ &\quad (-4 a515 + 4 a155) dx + (-4 a525 + 4 a255 + 2 a514) dy + (-4 a535 + 4 a355 \\ &\quad + 4 a524 - 2 u_{1,1}) dt + (-2 a514 u_1 + 4 a455 + 6 a534 - 4 a545 - 6 u_{1,2}) dz \\ &\quad - ((4 a514 u_2 + 4 a524 u_1 + 6 u_1 u_{1,1} - 8 a544 + 4 a555 + 12 u_{2,2}) ds) \end{aligned} \quad (5.19)$$

Let's check that ω uniquely determines the quadrics $Quad||i$.

$$> Qua := evalDG(add(add(xi[i, j] \cdot vM[i] \&s vM[j], j=i..5), i=1..5)) :$$

$$\begin{aligned}
& \text{sxi} := \text{solve}(\{ \text{coeffs}(\text{ContractIndices}(\text{Qua}, \omega \& t \omega, [[1, 1], [2, 2]]), \{q_{[]}\}) \}, \\
& \quad \{ \text{seq}(\text{seq}(\text{xi}[i, j], j = i..5), i = 1..5) \}) \\
& \text{sxi} := \{ \xi_{1,1} = u_1^3 \xi_{4,4} - u_1^2 \xi_{3,3} - 2 u_1 u_2 \xi_{3,4} - u_2^2 \xi_{4,4} + u_1 \xi_{2,2} + u_2 \xi_{2,3} + u_3 \xi_{2,4} + u_3 \xi_{3,3}, \quad (5.20) \\
& \quad \xi_{1,2} = -u_1^2 \xi_{3,4} - 2 u_1 u_2 \xi_{4,4} + u_1 \xi_{2,3} + u_2 \xi_{2,4} + 2 u_2 \xi_{3,3} + u_3 \xi_{3,4}, \xi_{1,3} = u_1 \xi_{2,4} \\
& \quad + u_1 \xi_{3,3} + u_2 \xi_{3,4} + u_3 \xi_{4,4} - \xi_{2,2}, \xi_{1,4} = -\xi_{2,3}, \xi_{1,5} = -\xi_{2,4} - \xi_{3,3}, \xi_{2,2} = \xi_{2,2}, \xi_{2,3} \\
& \quad = \xi_{2,3}, \xi_{2,4} = \xi_{2,4}, \xi_{2,5} = -\xi_{3,4}, \xi_{3,3} = \xi_{3,3}, \xi_{3,4} = \xi_{3,4}, \xi_{3,5} = -\xi_{4,4}, \xi_{4,4} = \xi_{4,4}, \xi_{4,5} = 0, \\
& \quad \xi_{5,5} = 0 \}
\end{aligned}$$

$$\begin{aligned}
& \text{DGsimplify}(\text{evalDG}(\text{subs}(\{ \xi_{2,2} = -1, \xi_{2,3} = 0, \xi_{2,4} = 0, \xi_{3,3} = 0, \xi_{3,4} = 0, \xi_{4,4} = 0 \}, \text{subs}(\text{sxi}, \\
& \quad \text{Qua})) - \text{Quad1})), \\
& \text{DGsimplify}(\text{evalDG}(\text{subs}(\{ \xi_{2,2} = 0, \xi_{2,3} = -1, \xi_{2,4} = 0, \xi_{3,3} = 0, \xi_{3,4} = 0, \xi_{4,4} = 0 \}, \text{subs}(\text{sxi}, \\
& \quad \text{Qua})) - \text{Quad2})), \\
& \text{DGsimplify}(\text{evalDG}(\text{subs}(\{ \xi_{2,2} = 0, \xi_{2,3} = 0, \xi_{2,4} = 1, \xi_{3,3} = -1, \xi_{3,4} = 0, \xi_{4,4} = 0 \}, \text{subs}(\text{sxi}, \\
& \quad \text{Qua})) - \text{Quad3})), \\
& \text{DGsimplify}(\text{evalDG}(\text{subs}(\{ \xi_{2,2} = -u_1, \xi_{2,3} = 0, \xi_{2,4} = -1, \xi_{3,3} = 0, \xi_{3,4} = 0, \xi_{4,4} = 0 \}, \\
& \quad \text{subs}(\text{sxi}, \text{Qua})) - \text{Quad4})), \\
& \text{DGsimplify}(\text{evalDG}(\text{subs}(\{ \xi_{2,2} = -u_2, \xi_{2,3} = 0, \xi_{2,4} = 0, \xi_{3,3} = 0, \xi_{3,4} = -1, \xi_{4,4} = 0 \}, \\
& \quad \text{subs}(\text{sxi}, \text{Qua})) - \text{Quad5})), \\
& \text{DGsimplify}(\text{evalDG}(\text{subs}(\{ \xi_{2,2} = -u_3, \xi_{2,3} = 0, \xi_{2,4} = 0, \xi_{3,3} = 0, \xi_{3,4} = 0, \xi_{4,4} = -1 \}, \\
& \quad \text{subs}(\text{sxi}, \text{Qua})) - \text{Quad6})) \\
& (0 D_x) D_x, (0 D_x) D_x, (0 D_x) D_x, (0 D_x) D_x, (0 D_x) D_x, (0 D_x) D_x \quad (5.21)
\end{aligned}$$

Explanation: the form ω has coefficients with q^i of degrees from 0 to 4, and the same for $\omega \otimes \omega$ with degrees from 0 to 8 - these are 9 quadrics on TM.
The annihilator in $S^2 TM$ of this 9D subspace of $S^2 T^*M$ has rank 6, consisting of $\langle \text{Quad1}, \dots, \text{Quad6} \rangle$.

$$\begin{aligned}
& \text{RNE} := \{ \} : \text{for } i \text{ to } 5 \text{ do for } j \text{ to } 6 \text{ do RNE} := \{ \text{op}(\text{RNE}), \\
& \quad \text{coeffs}(\text{expand}(\text{numer}(\text{simplify}(\text{subs}(\text{S2}, \text{subs}(\text{Qsyst}, \\
& \quad \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(\text{vM}[i], \text{Quadu}[j], \text{CC}), \omega \\
& \quad \& t \omega, [[1, 1], [2, 2]]), \{u(x, y, t, z, s)\}, \text{notation} = \text{jetnumbers})) \} \} : \text{od:od:} \\
& \quad \text{nops}(\text{RNE}) \\
& \quad \quad \quad 118 \quad (5.22)
\end{aligned}$$

$$\begin{aligned}
& \text{RNE}[1..20] \\
& \{ a155, a255, a355, -2 a255 - a514, -a514 - a255, -2 a355 - a524 + u_{1,1}, a355 + a524 \quad (5.23) \\
& \quad -2 u_{1,1}, -2 a455 - a534 + 2 u_{1,2}, -a524 + u_{1,1} - a355, 3 a155 u_1 - u_{1,1}, 4 a155 u_1 \\
& \quad - u_{1,1}, 5 a155 u_1 - u_{1,1}, 6 a155 u_1 - u_{1,1}, -2 a255 u_1 - a514 u_1, a155 u_1 - a355 \\
& \quad - u_{1,1}, 2 a155 u_1 - a355 - u_{1,1}, 3 a155 u_1 - a355 - u_{1,1}, 4 a155 u_1 - a355 - u_{1,1},
\end{aligned}$$

$$\{a155 u_1^2 - a355 u_1 - u_1 u_{1,1}, a155 u_2 - 4 a255 u_1 - 2 a514 u_1\}$$

> FreePar

$$\{a155, a255, a355, a455, a514, a515, a524, a525, a534, a535, a544, a545, a555\} \quad (5.24)$$

> tmp := eliminate(RNE, FreePar)

$$tmp := [\{a155 = 0, a255 = 0, a355 = 0, a455 = -u_{1,2}, a514 = 0, a515 = a515, a524 = u_{1,1}, \quad (5.25)$$

$$\begin{aligned} & a525 = a525, a534 = 4 u_{1,2}, a535 = a535, a544 = 4 u_{2,2}, a545 = a545, a555 = a555\}, \\ & \{u_{1,1}, u_1 u_{1,1}, u_2 u_{1,1}, (u_1^2 + u_3) u_{1,1}, u_1 u_2 u_{1,1}, u_1 (u_1^2 + u_3) u_{1,1}, u_2 (u_1^2 + u_3) u_{1,1}, \\ & -8 u_1 u_{1,1}, -7 u_1 u_{1,1}, -6 u_1 u_{1,1}, -5 u_1 u_{1,1}, -3 u_1 u_{1,1}, -u_1 u_{1,1}, -5 u_1^2 u_{1,1}, -u_1^2 u_{1,1}, \\ & -5 u_1 u_2 u_{1,1}, -u_1 u_{1,1} (6 u_1^2 + u_3), -5 u_1^2 u_2 u_{1,1}, -u_2 u_{1,1} (6 u_1^2 + u_3), -16 u_1^2 u_{1,1} \\ & -u_3 u_{1,1}, -15 u_1^2 u_{1,1} - u_3 u_{1,1}, -12 u_1^2 u_{1,1} - u_3 u_{1,1}, -9 u_1^2 u_{1,1} - u_3 u_{1,1}, -6 u_1^2 u_{1,1} \\ & -u_3 u_{1,1}, -5 u_1^3 u_{1,1} - 5 u_1 u_3 u_{1,1}, -6 u_1^4 u_{1,1} - 7 u_1^2 u_3 u_{1,1} - u_3^2 u_{1,1}, -21 u_1^3 u_{1,1} \\ & -6 u_1 u_3 u_{1,1} + 4 u_2^2 u_{1,1}, -6 u_1^3 u_{1,1} - u_1 u_3 u_{1,1} + u_2^2 u_{1,1}, -u_1^3 u_{1,1} - u_1 u_3 u_{1,1} + 2 \\ & u_2^2 u_{1,1}\} \end{aligned}$$

$$\begin{aligned} & \text{DGsimplify}(\text{subs}(tmp[1], \text{ContractIndices}(\text{TorsionTensor}(CC), [[1, 2]]))) \\ & - (4 a515 dx) - (4 a525 dy) - ((4 a535 - 2 u_{1,1}) dt) - ((-14 u_{1,2} + 4 a545) dz) \\ & - ((10 u_1 u_{1,1} + 4 a555 - 20 u_{2,2}) ds) \end{aligned} \quad (5.26)$$

> eliminate({op(RNE), op(DGinfo(ContractIndices(TorsionTensor(CC), [[1, 2]]), "CoefficientSet"))}, FreePar)

$$\left[\left\{ a155 = 0, a255 = 0, a355 = 0, a455 = -u_{1,2}, a514 = 0, a515 = 0, a524 = u_{1,1}, a525 = 0, \quad (5.27) \right. \right.$$

$$\begin{aligned} & a534 = 4 u_{1,2}, a535 = \frac{1}{2} u_{1,1}, a544 = 4 u_{2,2}, a545 = \frac{7}{2} u_{1,2}, a555 = -\frac{5}{2} u_1 u_{1,1} \\ & + 5 u_{2,2} \}, \{u_{1,1}, u_1 u_{1,1}, u_2 u_{1,1}, u_{1,1} (6 u_1^2 + u_3), u_{1,1} (9 u_1^2 + u_3), u_{1,1} (12 u_1^2 + u_3), \\ & u_{1,1} (15 u_1^2 + u_3), u_{1,1} (16 u_1^2 + u_3), u_{1,1} (6 u_1^4 + 7 u_1^2 u_3 + u_3^2), u_{1,1} (u_1^3 + u_1 u_3 - 2 \\ & u_2^2), u_{1,1} (6 u_1^3 + u_1 u_3 - u_2^2), u_{1,1} (21 u_1^3 + 6 u_1 u_3 - 4 u_2^2), (u_1^2 + u_3) u_{1,1}, u_1 u_2 u_{1,1}, \\ & u_1 u_{1,1} (6 u_1^2 + u_3), u_1 (u_1^2 + u_3) u_{1,1}, u_2 u_{1,1} (6 u_1^2 + u_3), u_2 (u_1^2 + u_3) u_{1,1} \} \end{aligned}$$

$$\begin{aligned} & \text{subs}(\{u_{1,1} = 0\}, \%[2]) \\ & \{0\} \end{aligned} \quad (5.28)$$

Thus GL2-connection having α -surfaces totally geodesic does not exist in 5D (for dKP) ...

Let us start looking for a connection by dropping the totally geodesic condition, but concentrating on GL2-property and Ricci tensor.

> RNE := { } :for i to 5 do for j to 6 do RNE := {op(RNE),
coeffs(expand(numer(simplify(subs(S2, subs(Qsyst,
ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], C), ω & t ω ,

$[[1, 1], [2, 2]]), \{u(x, y, t, z, s)\}, notation = jetnumbers))))) , \{q_{[]}\} : od: od: nops(RNE)$

230

(5.29)

> RNE[1..20]
 $\{a_{115}, a_{125}, a_{135}, a_{145}, a_{155}, a_{215}, a_{225}, a_{235}, a_{245}, a_{255}, a_{315}, a_{325}, a_{335}, a_{345},$
 $a_{355}, a_{114} - 2 a_{215}, a_{114} - a_{215}, a_{124} - 2 a_{225}, -a_{225} + a_{124}, -2 a_{235} + a_{134}\}$ (5.30)

> Sconn := solve(RNE, ChrCoe) :
 > FreePar := { } :for i to nops(Sconn) do if lhs(Sconn[i]) = rhs(Sconn[i]) then FreePar
 := {op(FreePar), lhs(Sconn[i])} fi:od: FreePar, nops(%)
 $\{a_{413}, a_{414}, a_{415}, a_{423}, a_{424}, a_{425}, a_{433}, a_{434}, a_{435}, a_{443}, a_{444}, a_{445}, a_{453}, a_{454},$
 $a_{455}, a_{515}, a_{525}, a_{535}, a_{545}, a_{555}\}, 20$ (5.31)

> ChrCoe1 := ChrCoe\FreePar
 ChrCoe1 := $\{a_{111}, a_{112}, a_{113}, a_{114}, a_{115}, a_{121}, a_{122}, a_{123}, a_{124}, a_{125}, a_{131}, a_{132},$
 $a_{133}, a_{134}, a_{135}, a_{141}, a_{142}, a_{143}, a_{144}, a_{145}, a_{151}, a_{152}, a_{153}, a_{154}, a_{155}, a_{211},$
 $a_{212}, a_{213}, a_{214}, a_{215}, a_{221}, a_{222}, a_{223}, a_{224}, a_{225}, a_{231}, a_{232}, a_{233}, a_{234}, a_{235},$
 $a_{241}, a_{242}, a_{243}, a_{244}, a_{245}, a_{251}, a_{252}, a_{253}, a_{254}, a_{255}, a_{311}, a_{312}, a_{313}, a_{314},$
 $a_{315}, a_{321}, a_{322}, a_{323}, a_{324}, a_{325}, a_{331}, a_{332}, a_{333}, a_{334}, a_{335}, a_{341}, a_{342}, a_{343},$
 $a_{344}, a_{345}, a_{351}, a_{352}, a_{353}, a_{354}, a_{355}, a_{411}, a_{412}, a_{421}, a_{422}, a_{431}, a_{432}, a_{441},$
 $a_{442}, a_{451}, a_{452}, a_{511}, a_{512}, a_{513}, a_{514}, a_{521}, a_{522}, a_{523}, a_{524}, a_{531}, a_{532}, a_{533},$
 $a_{534}, a_{541}, a_{542}, a_{543}, a_{544}, a_{551}, a_{552}, a_{553}, a_{554}\}$ (5.32)

> [nops(ChrCoe), nops(FreePar), nops(ChrCoe1)]
 [125, 20, 105] (5.33)

> FreePar := $\{a_{413}, a_{414}, a_{415}, a_{423}, a_{424}, a_{425}, a_{433}, a_{434}, a_{435}, a_{443}, a_{444}, a_{445},$
 $a_{453}, a_{454}, a_{455}, a_{515}, a_{525}, a_{535}, a_{545}, a_{555}\} :$
 > ChrCoe1 := $\{a_{111}, a_{112}, a_{113}, a_{114}, a_{115}, a_{121}, a_{122}, a_{123}, a_{124}, a_{125}, a_{131}, a_{132},$
 $a_{133}, a_{134}, a_{135}, a_{141}, a_{142}, a_{143}, a_{144}, a_{145}, a_{151}, a_{152}, a_{153}, a_{154}, a_{155}, a_{211},$
 $a_{212}, a_{213}, a_{214}, a_{215}, a_{221}, a_{222}, a_{223}, a_{224}, a_{225}, a_{231}, a_{232}, a_{233}, a_{234}, a_{235},$
 $a_{241}, a_{242}, a_{243}, a_{244}, a_{245}, a_{251}, a_{252}, a_{253}, a_{254}, a_{255}, a_{311}, a_{312}, a_{313}, a_{314},$
 $a_{315}, a_{321}, a_{322}, a_{323}, a_{324}, a_{325}, a_{331}, a_{332}, a_{333}, a_{334}, a_{335}, a_{341}, a_{342}, a_{343},$
 $a_{344}, a_{345}, a_{351}, a_{352}, a_{353}, a_{354}, a_{355}, a_{411}, a_{412}, a_{421}, a_{422}, a_{431}, a_{432}, a_{441},$
 $a_{442}, a_{451}, a_{452}, a_{511}, a_{512}, a_{513}, a_{514}, a_{521}, a_{522}, a_{523}, a_{524}, a_{531}, a_{532}, a_{533},$
 $a_{534}, a_{541}, a_{542}, a_{543}, a_{544}, a_{551}, a_{552}, a_{553}, a_{554}\} :$

> Sconn := solve(RNE, ChrCoe1) :

> CC := DGsimplify(subs(Sconn, C)) :

> TorC := TorsionTensor(CC) :

REN := {seq(seq(coeffs(expand(numer(ToJet(Hook(convert(ContractIndices(DD[i]
 &t DD[j], TorC, [[1, 2], [2, 3]]), 'DGvector'), ω), {u(x, y, t, z, s), q(x, y, t, z, s)},
 notation = jetnumbers)) , {q_[]}, j = i + 1 .. 4), i = 1 .. 4) : nops(REN)

38

(5.34)

> tmp := eliminate(REN union DGinfo(ContractIndices(TorsionTensor(CC), [[1, 2]]),
 "CoefficientSet"), FreePar)

tmp := $\left\{ a_{413} = 0, a_{414} = 0, a_{415} = 0, a_{423} = \frac{9}{4} u_{1, 1}, a_{424} = 0, a_{425} = 0, a_{433} = 3 u_{1, 2}, \right.$ (5.35)
 $a_{434} = u_{1, 1}, a_{435} = 0, a_{443} = \frac{15}{4} u_{1, 1} u_{1, 1} + 3 u_{2, 2}, a_{444} = \frac{5}{2} u_{1, 2}, a_{445} = -\frac{1}{4} u_{1, 1},$

$$a453 = 3 u_1 u_{1,2} + \frac{9}{2} u_2 u_{1,1} + 3 u_{2,3}, a454 = 3 u_1 u_{1,1} + 4 u_{2,2}, a455 = -u_{1,2}, a515 = 0, a525 = 0, a535 = \frac{3}{2} u_{1,1}, a545 = \frac{7}{2} u_{1,2}, a555 = \frac{9}{2} u_1 u_{1,1} + 5 u_{2,2}, \{ \}$$

$$\begin{aligned} &> \text{Sconn2} := \text{tmp}[1] : \text{CC} := \text{DGsimplify}(\text{subs}(\text{Sconn2}, \text{CC})) : \\ &> \{ \text{seq}(\text{seq}(\text{coeffs}(\text{expand}(\text{numer}(\text{simplify}(\text{subs}(S2, \text{subs}(Qsyst, \\ &\quad \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(\text{vM}[i], \text{Quadu}[j], \text{CC}), \omega \\ &\quad \&t \omega, [[1, 1], [2, 2]]), \{u(x, y, t, z, s)\}, \text{notation} = \text{jetnumbers})))), \{q_{[i]}\}, i = 1 \\ &\quad \dots 5), j = 1 \dots 6) \} \text{ union } \text{DGinfo}(\text{ContractIndices}(\text{TorsionTensor}(\text{CC}), [[1, 2]]), \\ &\quad \text{"CoefficientSet"}) \end{aligned} \quad \{0\} \quad (5.36)$$

$$\begin{aligned} &> \text{coeffs}(\text{expand}(\text{subs}(S2, \text{subs}(Qsyst, \\ &\quad \text{ToJet}(\text{Hook}(\text{convert}(\text{DirectionalCovariantDerivative}(\text{DD}[3], \text{DD}[1], \text{CC}), \text{'DGvector'}), \omega\omega), \{u(x, y, t, z, s), q(x, y, t, z, s)\}, \text{notation} = \text{jetnumbers}))), \{q_{[1]}, q_1\}) \\ &\quad - \frac{1}{2} u_1 u_{1,1}, - \frac{3}{4} u_{1,1} \end{aligned} \quad (5.37)$$

$$\begin{aligned} &> \text{RicCC} := \text{RicciTensor}(\text{FromJet}(\text{CC}, \{u(x, y, t, z, s)\})) : \text{RicCc} := \text{DGsimplify}(\text{subs}(S3, \\ &\quad \text{ToJet}(\text{RicCC}, \{u(x, y, t, z, s)\}, \text{notation} = \text{jetnumbers}))) : \\ &> \text{RicCcSkew} := \text{SymmetrizeIndices}(\text{RicCc}, [1, 2], \text{"SkewSymmetric"}) : \\ &\quad \text{RicCcSymm} := \text{SymmetrizeIndices}(\text{RicCc}, [1, 2], \text{"Symmetric"}) : \\ &> \omega \\ &\quad dx + q_{[1]} dy + (q_{[1]}^2 + u_1) dt + (q_{[1]}^3 + 2 q_{[1]} u_1 + u_2) dz + (q_{[1]}^4 + 3 q_{[1]}^2 u_1 \\ &\quad + 2 q_{[1]} u_2 + u_1^2 + u_3) ds \end{aligned} \quad (5.38)$$

$$\begin{aligned} &> \omega 0 := \omega : \omega 0b := \text{DGsimplify}(\text{evalDG}(\text{subs}(q_{[1]} = 0, \omega))) : \\ &\quad \omega 1a := \text{evalDG}\left(\frac{1}{\varepsilon} \cdot (\text{subs}(q_{[1]} = q_{[1]} + \varepsilon, \omega) - \omega)\right) : \omega 1 := \text{evalDG}(\text{subs}(\varepsilon = 0, \omega 1a)) : \\ &\quad \omega 1b := \text{DGsimplify}(\text{evalDG}(\text{subs}(q_{[1]} = 0, \omega 1))) : \\ &\quad \omega 2a := \text{evalDG}\left(\frac{1}{\varepsilon} \cdot (\text{subs}(q_{[1]} = q_{[1]} + \varepsilon, \omega 1) - \omega 1)\right) : \omega 2 := \text{evalDG}(\text{subs}(\varepsilon = 0, \\ &\quad \omega 2a)) : \omega 2b := \text{DGsimplify}\left(\text{evalDG}\left(\frac{1}{2} \cdot \text{subs}(q_{[1]} = 0, \omega 2)\right)\right) : \\ &\quad \omega 3a := \text{evalDG}\left(\frac{1}{\varepsilon} \cdot (\text{subs}(q_{[1]} = q_{[1]} + \varepsilon, \omega 2) - \omega 2)\right) : \omega 3 := \text{evalDG}(\text{subs}(\varepsilon = 0, \\ &\quad \omega 3a)) : \omega 3b := \text{DGsimplify}\left(\text{evalDG}\left(\frac{1}{6} \cdot \text{subs}(q_{[1]} = 0, \omega 3)\right)\right) : \\ &\quad \omega 4a := \text{evalDG}\left(\frac{1}{\varepsilon} \cdot (\text{subs}(q_{[1]} = q_{[1]} + \varepsilon, \omega 3) - \omega 3)\right) : \omega 4 := \text{evalDG}(\text{subs}(\varepsilon = 0, \\ &\quad \omega 4a)) : \omega 4b := \text{DGsimplify}\left(\text{evalDG}\left(\frac{1}{24} \cdot \text{subs}(q_{[1]} = 0, \omega 4)\right)\right) : \\ &\quad \omega 0b := dx + u_1 dt + u_2 dz + (u_1^2 + u_3) ds \end{aligned}$$

$$\omega 1b := dy + 2 u_1 dz + 2 u_2 ds$$

$$\omega 2b := dt + 3 u_1 ds$$

$$\omega 3b := dz$$

$$\omega 4b := ds$$

(5.39)

$$> evalDG(\omega - \omega 0b - \omega 1b \cdot q_{[1]} - \omega 2b \cdot q_{[1]}^2 - \omega 3b \cdot q_{[1]}^3 - \omega 4b \cdot q_{[1]}^4) \\ 0 dx$$

(5.40)

$$> Qua\omega := evalDG([\omega 0b \&s \omega 0b, \omega 0b \&s \omega 1b, 2 \omega 0b \&s \omega 2b + \omega 1b \&s \omega 1b, \omega 0b \&s \omega 3b \\ + \omega 1b \&s \omega 2b, 2 \omega 0b \&s \omega 4b + 2 \omega 1b \&s \omega 3b + \omega 2b \&s \omega 2b, \omega 1b \&s \omega 4b + \omega 2b \\ \&s \omega 3b, 2 \omega 2b \&s \omega 4b + \omega 3b \&s \omega 3b, \omega 3b \&s \omega 4b, \omega 4b \&s \omega 4b]): tmp\Omega := \{ \} : \text{for } i \\ \text{to nops}(Qua\omega) \text{ do for } j \text{ to } 6 \text{ do } tmp\Omega := \{ op(tmp\Omega), ContractIndices(Quad || j, \\ Qua\omega[i], [[1, 1], [2, 2]]) \} : \text{od:od: } tmp\Omega \\ \{0\}$$

(5.41)

$$> eliminate(simplify(subs(S3, DGinfo(evalDG(RicCcSymm - add(\Lambda[i] \cdot Qua\omega[i], i = 1 \\ ..nops(Qua\omega))), "CoefficientSet"))), \{seq(\Lambda[i], i = 1 ..nops(Qua\omega))\})[2]$$

$$\{u_{1,1,1}, 6u_{1,1}^2 + 7u_{1,2,2}, 28u_1u_{1,1,1} + 3u_{1,1}^2 - 14u_{1,2,2}, 3u_1^5u_{1,1,1} + 12u_1^4u_{1,1}^2 + 6 \\ u_1^4u_{1,2,2} + 6u_1^3u_3u_{1,1,1} + 17u_1^3u_{1,1}u_{2,2} + 6u_1^3u_{1,2}^2 - u_1^2u_2^2u_{1,1,1} + 10u_1^2u_2u_{1,1}u_{1,2} \\ + 19u_1^2u_3u_{1,1}^2 - 4u_1u_2^2u_{1,1}^2 + 3u_1^3u_{2,2,3} + 6u_1^2u_3u_{1,2,2} + 7u_1^2u_{1,1}u_{3,3} + 2u_1^2u_{2,2}^2 \\ - 2u_1u_2^2u_{1,2,2} + 3u_1u_2^2u_{1,1,1} + 16u_1u_3u_{1,1}u_{2,2} + 6u_1u_3u_{1,2}^2 - u_2^2u_3u_{1,1,1} - 6 \\ u_2^2u_{1,1}u_{2,2} - 6u_2^2u_{1,2}^2 - 2u_2u_3u_{1,1}u_{1,2} + u_2^2u_{1,1}^2 + 2u_1^2u_{2,2,5} + 3u_1u_3u_{2,2,3} \\ + 3u_1u_{1,1}u_{4,4} + 2u_1u_{2,2}u_{3,3} - u_2^2u_{2,2,3} - 2u_2u_{1,1}u_{3,4} - 2u_2u_{1,2}u_{3,3} \\ + u_3u_{1,1}u_{3,3} + u_3u_{2,2}^2 - 2u_1u_{1,5,5} + 2u_1u_{2,4,5} + u_3u_{2,2,5} + u_{2,2}u_{4,4} - u_{3,5,5} \\ + u_{4,4,5}, 126u_1^5u_{1,1,1} + 258u_1^4u_{1,1}^2 + 63u_1^4u_{1,2,2} + 168u_1^3u_3u_{1,1,1} + 238u_1^3u_{1,1}u_{2,2} \\ + 126u_1^3u_{1,2}^2 - 42u_1^2u_2^2u_{1,1,1} + 140u_1^2u_2u_{1,1}u_{1,2} + 356u_1^2u_3u_{1,1}^2 - 86u_1u_2^2u_{1,1}^2 \\ + 63u_1^3u_{2,2,3} + 63u_1^2u_3u_{1,2,2} + 98u_1^2u_{1,1}u_{3,3} + 28u_1^2u_{2,2}^2 - 21u_1u_2^2u_{1,2,2} + 42u_1 \\ u_3^2u_{1,1,1} + 224u_1u_3u_{1,1}u_{2,2} + 126u_1u_3u_{1,2}^2 - 14u_2^2u_3u_{1,1,1} - 84u_2^2u_{1,1}u_{2,2} - 98 \\ u_2^2u_{1,2}^2 - 28u_2u_3u_{1,1}u_{1,2} + 14u_2^2u_{1,1}^2 + 28u_1^2u_{2,2,5} + 63u_1u_3u_{2,2,3} + 42u_1u_{1,1}u_{4,4} \\ + 28u_1u_{2,2}u_{3,3} - 21u_2^2u_{2,2,3} - 28u_2u_{1,1}u_{3,4} - 28u_2u_{1,2}u_{3,3} + 14u_3u_{1,1}u_{3,3} \\ + 14u_3u_{2,2}^2 - 28u_1u_{1,5,5} + 28u_1u_{2,4,5} + 14u_3u_{2,2,5} + 14u_{2,2}u_{4,4} - 14u_{3,5,5} \\ + 14u_{4,4,5}, 3u_1^6u_{1,1,1} + 12u_1^5u_{1,1}^2 + 6u_1^5u_{1,2,2} + 6u_1^4u_3u_{1,1,1} + 17u_1^4u_{1,1}u_{2,2} + 6 \\ u_1^4u_{1,2}^2 - 7u_1^3u_2^2u_{1,1,1} + 19u_1^3u_2u_{1,1}u_{1,2} + 19u_1^3u_3u_{1,1}^2 - 4u_1^2u_2^2u_{1,1}^2 + 3u_1^4u_{2,2,3} \\ + 6u_1^3u_2u_{2,2,2} + 6u_1^3u_3u_{1,2,2} + 7u_1^3u_{1,1}u_{3,3} + 2u_1^3u_{2,2}^2 - 2u_1^2u_2^2u_{1,2,2} + 3u_1^2 \\ u_3^2u_{1,1,1} + 16u_1^2u_3u_{1,1}u_{2,2} + 6u_1^2u_3u_{1,2}^2 - 7u_1u_2^2u_3u_{1,1,1} - 6u_1u_2^2u_{1,1}u_{2,2} - 6u_1 \\ u_2^2u_{1,2}^2 + 7u_1u_2u_3u_{1,1}u_{1,2} + u_1u_2^2u_{1,1}^2 + 2u_2^4u_{1,1,1} - 3u_2^3u_{1,1}u_{1,2} + 2u_1^3u_{2,2,5} + 3$$

(5.42)

$$\begin{aligned}
& u_1^2 u_3 u_{2,2,3} + 3 u_1^2 u_{1,1} u_{4,4} + 2 u_1^2 u_{2,2} u_{3,3} - u_1 u_2^2 u_{2,2,3} + 6 u_1 u_2 u_3 u_{2,2,2} \\
& - 2 u_1 u_2 u_{1,1} u_{3,4} - 2 u_1 u_2 u_{1,2} u_{3,3} + u_1 u_3 u_{1,1} u_{3,3} + u_1 u_3 u_{2,2}^2 - 2 u_2^3 u_{2,2,2} - 2 \\
& u_1^2 u_{1,5,5} + 2 u_1^2 u_{2,4,5} + u_1 u_3 u_{2,2,5} + u_1 u_{2,2} u_{4,4} - u_1 u_{3,5,5} + u_1 u_{4,4,5} \}
\end{aligned}$$

Thus the condition that $\text{sym}(\text{Ric}(\nabla))$ belongs to the span of quadrics defining the GL2 structure is non-vacous. It is another differential constraint, giving exact solutions.

$$\begin{aligned}
& \triangleright \text{Vt0} := \text{Annihilator}([\omega 0, \omega 1, \omega 2, \omega 3] [1]; \text{Vt1} := \text{Annihilator}([\omega 0, \omega 1, \omega 2] [2]; \text{Vt2} \\
& \quad := \text{Annihilator}([\omega 0, \omega 1] [3]; \text{Vt3} := \text{Annihilator}([\omega 0] [4]; \\
& \quad \text{Vt01} := \text{ContractIndices}(\text{TorsionTensor}(\text{CC}), \text{evalDG}(\text{Vt0} \& \text{Vt1}), [[2, 1], [3, 2]]) : \\
& \text{Vt0} := (q[]^4 - 6 q[]^2 u_1 + 4 q[] u_2 + 2 u_1^2 - u_3) D_x - ((4 q[]^3 - 8 q[] u_1 \\
& \quad + 2 u_2) D_y) + (6 q[]^2 - 3 u_1) D_t - (4 q[] D_z) + D_s \\
& \text{Vt1} := -((q[]^3 - 3 q[] u_1 + u_2) D_x) + (3 q[]^2 - 2 u_1) D_y - (3 q[] D_t) + D_z \\
& \text{Vt2} := (3 q[]^4 + 3 q[]^2 u_1 - u_1^2 - u_3) D_x - ((4 q[]^3 + 6 q[] u_1 + 2 u_2) D_y) + D_s \\
& \quad \text{Vt3} := -(q[] D_x) + D_y \tag{5.43}
\end{aligned}$$

Thus there exist GL2-connections, with respect to which the null Lax surfaces are totally geodesic. The 4 parameters can be fixed uniquely by the condition $\text{Trace}(\text{Torsion})=0$.

$$\begin{aligned}
& \triangleright \text{seq}(\text{ContractIndices}(\text{Qua}\omega[i], \text{evalDG}(\text{Vt0} \& \text{Vt0}), [[1, 1], [2, 2]]), i = 1 .. \text{nops}(\text{Qua}\omega)) \\
& \quad q_{[]}^8, -4 q_{[]}^7, 28 q_{[]}^6, -28 q_{[]}^5, 70 q_{[]}^4, -28 q_{[]}^3, 28 q_{[]}^2, -4 q_{[]}, 1 \tag{5.44}
\end{aligned}$$

$$\begin{aligned}
& \triangleright \text{QuT} := \text{evalDG}(\text{add}(\text{add}(\text{xi}[i, j] \cdot dM[i] \& s dM[j], j = i .. 5), i = 1 .. 5)) : \\
& \text{DGsimplify}(\text{evalDG}(\text{subs}(\{ \text{xi}[2, 2] = \Lambda_1, \text{xi}[2, 3] = \Lambda_2, \text{xi}[2, 4] = \Lambda_3, \text{xi}[3, 3] = \Lambda_4, \text{xi}[3, 4] \\
& \quad = \Lambda_5, \text{xi}[4, 4] = \Lambda_6 \}, \text{subs}(\text{solve}(\{ \text{coeffs}(\text{ContractIndices}(\text{QuT}, \text{Vt0} \& \text{Vt0}), [[1, 1], [2, \\
& \quad 2]]), \{ q_{[]} \} \}), \{ \text{seq}(\text{seq}(\text{xi}[i, j], j = i .. 5), i = 1 .. 5) \}), \text{QuT}))) \\
& - \left(\left(\frac{4 \Lambda_1}{3} dx \right) dt \right) - ((3 \Lambda_2 dx) dz) - (((20 \Lambda_1 u_1 + 8 \Lambda_3 + 18 \Lambda_4) dx) ds) \tag{5.45} \\
& + (\Lambda_1 dy) dy + \left(\frac{\Lambda_2}{2} dy \right) dt + \left(\frac{\Lambda_3}{2} dy \right) dz - \left(\left(\left(\frac{21 \Lambda_2 u_1}{2} + 6 \Lambda_1 u_2 \right. \right. \right. \\
& \left. \left. + 3 \Lambda_5 \right) dy \right) ds \right) - \left(\left(\left(\frac{4 \Lambda_1}{3} dt \right) dx \right) + \left(\frac{\Lambda_2}{2} dt \right) dy + (\Lambda_4 dt) dt + \left(\frac{\Lambda_5}{2} dt \right) dz \right. \\
& \left. - \left(\left(\left(\frac{56}{3} \Lambda_1 u_1^2 + \frac{4}{3} \Lambda_1 u_3 + \frac{16}{3} \Lambda_3 u_1 + 15 u_1 \Lambda_4 + 7 \Lambda_2 u_2 + \frac{4}{3} \Lambda_6 \right) dt \right) ds \right) \right. \\
& \left. - ((3 \Lambda_2 dz) dx) + \left(\frac{\Lambda_3}{2} dz \right) dy + \left(\frac{\Lambda_5}{2} dz \right) dt + (\Lambda_6 dz) dz - \left(\left(\left(18 \Lambda_2 u_1^2 \right. \right. \right. \right. \\
& \left. \left. + 32 \Lambda_1 u_1 u_2 + \frac{9}{2} u_1 \Lambda_5 + 7 \Lambda_3 u_2 + 18 u_2 \Lambda_4 + 3 u_3 \Lambda_2 \right) dz \right) ds \right) - (((20 \Lambda_1 u_1 \\
& + 8 \Lambda_3 + 18 \Lambda_4) ds) dx) - \left(\left(\left(\left(\frac{21 \Lambda_2 u_1}{2} + 6 \Lambda_1 u_2 + 3 \Lambda_5 \right) ds \right) dy \right) - \left(\left(\left(\frac{56}{3} \Lambda_1 \right. \right. \right. \right. \\
& \left. \left. u_1^2 + \frac{4}{3} \Lambda_1 u_3 + \frac{16}{3} \Lambda_3 u_1 + 15 u_1 \Lambda_4 + 7 \Lambda_2 u_2 + \frac{4}{3} \Lambda_6 \right) ds \right) dt \right) - \left(\left(\left(18 \Lambda_2 u_1^2 \right. \right. \right. \right.
\end{aligned}$$

$$+ 32 \Lambda_1 u_1 u_2 + \frac{9}{2} u_1 \Lambda_5 + 7 \Lambda_3 u_2 + 18 u_2 \Lambda_4 + 3 u_3 \Lambda_2 \Big) ds \Big) dz \Big) - \Big(\Big((48 \Lambda_1 u_1^3 + 40 \Lambda_1 u_1 u_3 + 28 \Lambda_1 u_2^2 + 90 \Lambda_2 u_1 u_2 + 27 \Lambda_4 u_1^2 + 16 \Lambda_3 u_3 + 36 \Lambda_4 u_3 + 12 \Lambda_5 u_2 + 8 \Lambda_6 u_1) ds \Big) ds \Big)$$

> eliminate(DGinfo(DGsimply(evalDG(subs(S3,subs({xi[2,2]=Λ₁,xi[2,3]=Λ₂,xi[2,4]=Λ₃,xi[3,3]=Λ₄,xi[3,4]=Λ₅,xi[4,4]=Λ₆}),subs(solve({coeffs(ContractIndices(QuT,Vt0&t Vt0,[[1,1],[2,2]]),{q[_]}),{seq(seq(xi[i,j],j=i..5),i=1..5)}),QuT))-RicCcSymm)), "CoefficientSet"),{Λ₁,Λ₂,Λ₃,Λ₄,Λ₅,Λ₆})

$$\left\{ \Lambda_1 = \frac{3}{4} u_{1,1,1}, \Lambda_2 = -\frac{1}{2} \frac{1}{(3u_1^3 + 3u_1u_3 - u_2^2)u_2} (3u_1^5 u_{1,1,1} + 12u_1^4 u_{1,1}^2 + 6u_1^4 u_{1,2,2} \right. \quad (5.46)$$

$$+ 6u_1^3 u_{1,1,1} + 17u_1^3 u_{1,1}u_{2,2} + 6u_1^3 u_{1,2}^2 - u_1^2 u_2^2 u_{1,1,1} + 10u_1^2 u_2 u_{1,1}u_{1,2} + 19$$

$$u_1^2 u_3 u_{1,1}^2 - 4u_1 u_2^2 u_{1,1}^2 + 3u_1^3 u_{2,2,3} + 6u_1^2 u_3 u_{1,2,2} + 7u_1^2 u_{1,1}u_{3,3} + 2u_1^2 u_{2,2}^2 - 2u_1$$

$$u_2^2 u_{1,2,2} + 3u_1 u_3^2 u_{1,1,1} + 16u_1 u_3 u_{1,1}u_{2,2} + 6u_1 u_3 u_{1,2}^2 - u_2^2 u_3 u_{1,1,1} - 6$$

$$u_2^2 u_{1,1}u_{2,2} - 6u_2^2 u_{1,2}^2 - 2u_2 u_3 u_{1,1}u_{1,2} + u_3^2 u_{1,1}^2 + 2u_1^2 u_{2,2,5} + 3u_1 u_3 u_{2,2,3}$$

$$+ 3u_1 u_{1,1}u_{4,4} + 2u_1 u_{2,2}u_{3,3} - u_2^2 u_{2,2,3} - 2u_2 u_{1,1}u_{3,4} - 2u_2 u_{1,2}u_{3,3}$$

$$+ u_3 u_{1,1}u_{3,3} + u_3 u_{2,2}^2 - 2u_1 u_{1,5,5} + 2u_1 u_{2,4,5} + u_3 u_{2,2,5} + u_{2,2}u_{4,4} - u_{3,5,5}$$

$$+ u_{4,4,5}), \Lambda_3 = 3u_1 u_{1,1,1} + \frac{3}{4} u_{1,1}^2 - \frac{3}{2} u_{1,2,2}, \Lambda_4 = -2u_1 u_{1,1,1} + u_{1,2,2}, \Lambda_5$$

$$= \frac{1}{2} \frac{1}{(3u_1^3 + 3u_1u_3 - u_2^2)u_2} (12u_1^6 u_{1,1,1} + 48u_1^5 u_{1,1}^2 + 24u_1^5 u_{1,2,2} + 24$$

$$u_1^4 u_3 u_{1,1,1} + 68u_1^4 u_{1,1}u_{2,2} + 24u_1^4 u_{1,2}^2 - 16u_1^3 u_2^2 u_{1,1,1} + 49u_1^3 u_2 u_{1,1}u_{1,2} + 76$$

$$u_1^3 u_3 u_{1,1}^2 - 16u_1^2 u_2^2 u_{1,1}^2 + 12u_1^4 u_{2,2,3} + 6u_1^3 u_2 u_{2,2,2} + 24u_1^3 u_3 u_{1,2,2} + 28$$

$$\begin{aligned}
& u_1^3 u_{1,1} u_{3,3} + 8 u_1^3 u_{2,2}^2 - 8 u_1^2 u_2^2 u_{1,2,2} + 12 u_1^2 u_3^2 u_{1,1,1} + 64 u_1^2 u_3 u_{1,1} u_{2,2} + 24 u_1^2 u_3 \\
& u_{1,2}^2 - 16 u_1 u_2^2 u_3 u_{1,1,1} - 24 u_1 u_2^2 u_{1,1} u_{2,2} - 24 u_1 u_2^2 u_{1,2}^2 + u_1 u_2 u_3 u_{1,1} u_{1,2} + 4 u_1 \\
& u_3^2 u_{1,1}^2 + 4 u_2^4 u_{1,1,1} - 3 u_2^3 u_{1,1} u_{1,2} + 8 u_1^3 u_{2,2,5} + 12 u_1^2 u_3 u_{2,2,3} + 12 u_1^2 u_{1,1} u_{4,4} \\
& + 8 u_1^2 u_{2,2} u_{3,3} - 4 u_1 u_2^2 u_{2,2,3} + 6 u_1 u_2 u_3 u_{2,2,2} - 8 u_1 u_2 u_{1,1} u_{3,4} - 8 u_1 u_2 u_{1,2} u_{3,3} \\
& + 4 u_1 u_3 u_{1,1} u_{3,3} + 4 u_1 u_3 u_{2,2}^2 - 2 u_2^3 u_{2,2,2} - 8 u_1^2 u_{1,5,5} + 8 u_1^2 u_{2,4,5} \\
& + 4 u_1 u_3 u_{2,2,5} + 4 u_1 u_{2,2} u_{4,4} - 4 u_1 u_{3,5,5} + 4 u_1 u_{4,4,5}), \Lambda_6 \\
& = \frac{3}{4} \frac{1}{3 u_1^3 + 3 u_1 u_3 - u_2^2} (24 u_1^5 u_{1,1,1} + 54 u_1^4 u_{1,1}^2 + 9 u_1^4 u_{1,2,2} + 36 u_1^3 u_3 u_{1,1,1} \\
& + 68 u_1^3 u_{1,1} u_{2,2} + 30 u_1^3 u_{1,2}^2 - 8 u_1^2 u_2^2 u_{1,1,1} + 40 u_1^2 u_2 u_{1,1} u_{1,2} + 82 u_1^2 u_3 u_{1,1}^2 \\
& - 18 u_1 u_2^2 u_{1,1}^2 + 15 u_1^3 u_{2,2,3} + 9 u_1^2 u_3 u_{1,2,2} + 28 u_1^2 u_{1,1} u_{3,3} + 8 u_1^2 u_{2,2}^2 - 3 u_1 \\
& u_2^2 u_{1,2,2} + 12 u_1 u_3^2 u_{1,1,1} + 64 u_1 u_3 u_{1,1} u_{2,2} + 30 u_1 u_3 u_{1,2}^2 - 4 u_2^2 u_3 u_{1,1,1} - 24 \\
& u_2^2 u_{1,1} u_{2,2} - 26 u_2^2 u_{1,2}^2 - 8 u_2 u_3 u_{1,1} u_{1,2} + 4 u_3^2 u_{1,1}^2 + 8 u_1^2 u_{2,2,5} + 15 u_1 u_3 u_{2,2,3} \\
& + 12 u_1 u_{1,1} u_{4,4} + 8 u_1 u_{2,2} u_{3,3} - 5 u_2^2 u_{2,2,3} - 8 u_2 u_{1,1} u_{3,4} - 8 u_2 u_{1,2} u_{3,3} \\
& + 4 u_3 u_{1,1} u_{3,3} + 4 u_3 u_{2,2}^2 - 8 u_1 u_{1,5,5} + 8 u_1 u_{2,4,5} + 4 u_3 u_{2,2,5} + 4 u_{2,2} u_{4,4} \\
& - 4 u_{3,5,5} + 4 u_{4,4,5}) \}, \{ \}]
\end{aligned}$$

This proves that symmetrized Ricci tensor of the canonical GL2-connection is a section of the bundle of quadrics annihilating the dual (tangent) to the canonical (cotangent) rational normal curve!

Here is the connection:

> CC

(5.47)

>

>

>

II. Let us check if there exists a unique symmetric connection preserving GL2-structure

> ChrSeq := {seq(seq(seq(((a||i)||j)||k), k=1..5), j=i..5), i=1..5)}: nops(ChrSeq);
CsC := Connection(subs({seq(seq(seq(((a||i)||j)||k)=(((a||j)||i)||k), k=1..5), j=1..(i-1)), i=1..5}), add(add(add(((a||i)||j)||k)·vM[k]&t dM[i]&t dM[j], k=1..5), j=1..5), i=1..5)): TorsionTensor(CsC)

75

((0 D_x) dx) dx

(5.48)

```

> RNE := { } :for i to 5 do for j to 6 do RNE := {op(RNE),
      coeffs( expand( numer( simplify( subs( S2, subs( Qsyst,
      ToJet( ContractIndices( DirectionalCovariantDerivative(vM[i], Quadu[j], CsC), ω
      &t ω, [[1, 1], [2, 2]]), {u(x, y, t, z, s)}, notation = jetnumbers) ) ) ) ), {q[_]} ) } :od:od:
> nops(RNE)

```

227

(5.49)

```

> elRN := eliminate(RNE, ChrSeq) :
> elRN[2]

```

$$\{u_{1,1}, u_{1,2}, u_1 u_{1,1}, u_1 u_{1,2}, u_1 (3 u_1 u_{1,2} - 2 u_2 u_{1,1}), u_1 (3 u_1 u_{1,2} + u_2 u_{1,1}), u_1 (6 u_1 u_{1,2} - u_2 u_{1,1}), u_2 u_{1,1}, u_2 u_{1,2}, u_3 u_{1,1}, u_{1,1} (4 u_1^2 + u_3), u_1 u_2 u_{1,1}, u_1 u_3 u_{1,1}, u_1 u_{1,1} (3 u_1^2 + u_3), -2 u_1 u_3 u_{1,1}, 2 u_1 u_3 u_{1,1}, 3 u_1 u_{1,2} + u_2 u_{1,1}, u_1^2 u_{1,1} - 3 u_2 u_{1,2}, u_1^2 u_{1,1} + 3 u_2 u_{1,2}, u_1^3 u_{1,1} + 3 u_1 u_2 u_{1,2} - u_2^2 u_{1,1}, 12 u_1^2 u_{1,2} - u_1 u_2 u_{1,1} + 6 u_3 u_{1,2}, 6 u_1^3 u_{1,2} - u_1^2 u_2 u_{1,1} + u_2 u_3 u_{1,1}, 12 u_1^2 u_2 u_{1,2} - 3 u_1^2 u_3 u_{1,1} - 2 u_1 u_2^2 u_{1,1} - u_3^2 u_{1,1}, 6 u_1^3 u_{1,2} + 5 u_1^2 u_2 u_{1,1} + 6 u_2^2 u_{1,2} + u_2 u_3 u_{1,1}, 3 u_1^3 u_{1,2} - u_1^2 u_2 u_{1,1} - 3 u_1 u_3 u_{1,2} + u_2 u_3 u_{1,1}\} \quad (5.50)$$

$$\begin{aligned} & \text{simplify}(\text{subs}(\{u_{1,1}=0\}, \{u_{1,1}, u_{1,2}, u_1 u_{1,1}, u_1 u_{1,2}, u_1 (3 u_1 u_{1,2} - 2 u_2 u_{1,1}), u_1 (3 u_1 u_{1,2} + u_2 u_{1,1}), u_1 (6 u_1 u_{1,2} - u_2 u_{1,1}), u_2 u_{1,1}, u_2 u_{1,2}, u_3 u_{1,1}, u_{1,1} (4 u_1^2 + u_3), u_1 u_2 u_{1,1}, u_1 u_3 u_{1,1}, u_1 u_{1,1} (3 u_1^2 + u_3), -2 u_1 u_3 u_{1,1}, 2 u_1 u_3 u_{1,1}, 3 u_1 u_{1,2} + u_2 u_{1,1}, u_1^2 u_{1,1} - 3 u_2 u_{1,2}, u_1^2 u_{1,1} + 3 u_2 u_{1,2}, u_1^3 u_{1,1} + 3 u_1 u_2 u_{1,2} - u_2^2 u_{1,1}, 12 u_1^2 u_{1,2} - u_1 u_2 u_{1,1} + 6 u_3 u_{1,2}, 6 u_1^3 u_{1,2} - u_1^2 u_2 u_{1,1} + u_2 u_3 u_{1,1}, 12 u_1^2 u_2 u_{1,2} - 3 u_1^2 u_3 u_{1,1} - 2 u_1 u_2^2 u_{1,1} - u_3^2 u_{1,1}, 6 u_1^3 u_{1,2} + 5 u_1^2 u_2 u_{1,1} + 6 u_2^2 u_{1,2} + u_2 u_3 u_{1,1}, 3 u_1^3 u_{1,2} - u_1^2 u_2 u_{1,1} - 3 u_1 u_3 u_{1,2} + u_2 u_3 u_{1,1}\})) \\ & \{0, u_{1,2}, u_1 u_{1,2}, u_2 u_{1,2}, 3 u_1 u_{1,2}, 3 u_1^2 u_{1,2}, 6 u_1^2 u_{1,2}, 6 u_1^3 u_{1,2}, -3 u_2 u_{1,2}, 3 u_2 u_{1,2}, 3 u_1 u_2 u_{1,2}, 12 u_1^2 u_2 u_{1,2}, 6 u_1^3 u_{1,2} + 6 u_2^2 u_{1,2}, 12 u_1^2 u_{1,2} + 6 u_3 u_{1,2}, 3 u_1^3 u_{1,2} - 3 u_1 u_3 u_{1,2}\} \end{aligned} \quad (5.51)$$

$$\begin{aligned} & \text{simplify}(\text{subs}(\{u_{1,1}=0\}, \{u_{1,1}, u_1 u_{1,1}, u_2 u_{1,1}, (u_1^2 + u_3) u_{1,1}, u_1 u_2 u_{1,1}, u_1 (u_1^2 + u_3) u_{1,1}, u_2 (u_1^2 + u_3) u_{1,1}, -8 u_1 u_{1,1}, -7 u_1 u_{1,1}, -6 u_1 u_{1,1}, -5 u_1 u_{1,1}, -3 u_1 u_{1,1}, -u_1 u_{1,1}, -5 u_1^2 u_{1,1}, -u_1^2 u_{1,1}, -5 u_1 u_2 u_{1,1}, -u_1 u_{1,1} (6 u_1^2 + u_3), -5 u_1^2 u_2 u_{1,1}, -u_2 u_{1,1} (6 u_1^2 + u_3), -16 u_1^2 u_{1,1} - u_3 u_{1,1}, -15 u_1^2 u_{1,1} - u_3 u_{1,1}, -12 u_1^2 u_{1,1} - u_3 u_{1,1}, -9 u_1^2 u_{1,1} - u_3 u_{1,1}, -6 u_1^2 u_{1,1} - u_3 u_{1,1}, -5 u_1^3 u_{1,1} - 5 u_1 u_3 u_{1,1}, -6 u_1^4 u_{1,1} - 7 u_1^2 u_3 u_{1,1} - u_3^2 u_{1,1}, -21 u_1^3 u_{1,1} - 6 u_1 u_3 u_{1,1} + 4 u_2^2 u_{1,1}, -6 u_1^3 u_{1,1} - u_1 u_3 u_{1,1} + u_2^2 u_{1,1}, -u_1^3 u_{1,1} - u_1 u_3 u_{1,1} + 2 u_2^2 u_{1,1}\})) \\ & \{0\} \end{aligned} \quad (5.52)$$

```

> Sss1 := solve(RNE, ChrSeq)

```

Sss1 :=

(5.53)

So it does not exist in 5D (for dKP) as expected ...

>

III. Finally let us compute the canonical projective totally geodesic connection

> CsC := Connection(subs({seq(seq(seq(((a||i)||j)||k)=(((a||j)||i)||k), k=1..5), j=1..(i-1)), i=1..5}), add(add(add(((a||i)||j)||k)·vM[k]&t dM[i]&t dM[j], k=1..5), j=1..5), i=1..5)):

> DD := {YY, ZZ, TT, SS} : ChrCoe := {seq(seq(seq(((a||i)||j)||k), k=1..5), j=i..5), i=1..5):

TGE := { } :for i to 4 do for j to 4 do TGE := {op(TGE), coeffs(expand(subs(S2, subs(Qsyst, ToJet(Hook(convert(DirectionalCovariantDerivative(DD[i], DD[j], CsC), 'DGvector'), ωω), {u(x, y, t, z, s), q(x, y, t, z, s)}, notation=jetnumbers))), {q_[j], q₁})} :od:od:
[nops(ChrCoe), nops(TGE)]

[75, 84]

(5.54)

> Sss2 := solve(TGE, ChrCoe\{a155, a255, a355, a455, a555}) : nops(Sss2)

70

(5.55)

> tmp := { } :for i to nops(Ss2) do if lhs(Sss2[i]) = rhs(Sss2[i]) then tmp := {op(tmp), lhs(Sss2[i])} :fi:od: tmp

{ }

(5.56)

> Sss2

{ a111 = 2 a155, a112 = 0, a113 = 0, a114 = 0, a115 = 0, a121 = a255, a122 = a155, a123

(5.57)

$$= 0, a124 = 0, a125 = 0, a131 = a355 - \frac{1}{2} u_{1,1}, a132 = 0, a133 = a155, a134 = 0, a135$$

$$= 0, a141 = a455 - u_{1,2}, a142 = -\frac{1}{2} u_{1,1}, a143 = 0, a144 = a155, a145 = 0, a151 =$$

$$-u_1 u_{1,1} + \frac{1}{2} a555 - \frac{3}{2} u_{2,2}, a152 = -u_{1,2}, a153 = -\frac{1}{2} u_{1,1}, a154 = 0, a221 = u_{1,1},$$

$$a222 = 2 a255, a223 = 0, a224 = 0, a225 = 0, a231 = u_{1,2}, a232 = \frac{1}{2} u_{1,1} + a355, a233$$

$$= a255, a234 = 0, a235 = 0, a241 = \frac{3}{2} u_1 u_{1,1} + u_{2,2}, a242 = a455, a243 = \frac{1}{2} u_{1,1},$$

$$a244 = a255, a245 = 0, a251 = 2 u_1 u_{1,2} + \frac{3}{2} u_2 u_{1,1} + u_{2,3}, a252 = \frac{1}{2} u_1 u_{1,1}$$

$$+ \frac{1}{2} a555 - \frac{1}{2} u_{2,2}, a253 = 0, a254 = \frac{1}{2} u_{1,1}, a331 = -u_1 u_{1,1} + u_{2,2}, a332 = 2 u_{1,2},$$

$$a333 = 2 a355, a334 = 0, a335 = 0, a341 = -u_1 u_{1,2} - \frac{1}{2} u_2 u_{1,1} + u_{2,3}, a342$$

$$= \frac{1}{2} u_1 u_{1,1} + 2 u_{2,2}, a343 = a455 + u_{1,2}, a344 = a355, a345 = 0, a351 = -\frac{5}{2} u_1^2 u_{1,1}$$

$$-u_1 u_{2,2} + u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + u_{3,3}, a352 = u_1 u_{1,2} + u_2 u_{1,1} + 2 u_{2,3}, a353 =$$

$$\begin{aligned}
& -\frac{1}{2} u_1 u_{1,1} + \frac{1}{2} a555 + \frac{1}{2} u_{2,2}, a354 = u_{1,2}, a441 = u_1^2 u_{1,1} - u_1 u_{2,2} - 2 u_2 u_{1,2} \\
& + u_{3,3}, a442 = -2 u_1 u_{1,2} - u_2 u_{1,1} + 2 u_{2,3}, a443 = 2 u_1 u_{1,1} + 3 u_{2,2}, a444 = 2 a455, \\
& a445 = 0, a451 = u_1^2 u_{1,2} - u_1 u_{2,3} - 2 u_2 u_{2,2} - u_3 u_{1,2} + u_{3,4}, a452 = -\frac{3}{2} u_1^2 u_{1,1} \\
& - 2 u_1 u_{2,2} - u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + 2 u_{3,3}, a453 = \frac{1}{2} u_2 u_{1,1} + 3 u_{2,3}, a454 = u_1 u_{1,1} \\
& + \frac{1}{2} a555 + \frac{3}{2} u_{2,2}, a551 = -u_1^3 u_{1,1} + u_1^2 u_{2,2} + 4 u_1 u_2 u_{1,2} - 2 u_1 u_3 u_{1,1} + u_2^2 u_{1,1} \\
& - u_1 u_{3,3} - 2 u_2 u_{2,3} - 3 u_3 u_{2,2} + u_{4,4}, a552 = 2 u_1^2 u_{1,2} - 2 u_1 u_{2,3} - 4 u_2 u_{2,2} \\
& - 2 u_3 u_{1,2} + 2 u_{3,4}, a553 = -4 u_1^2 u_{1,1} - 3 u_1 u_{2,2} - u_3 u_{1,1} + 3 u_{3,3}, a554 = 2 u_1 u_{1,2} \\
& + 2 u_2 u_{1,1} + 4 u_{2,3} \}
\end{aligned}$$

$$\triangleright pt := \{a155=0, a255=0, a355=0, a455=0, a555=0\} :$$

$$\triangleright Sss20 := \{op(subs(pt, Sss2)), op(pt)\}$$

$$Sss20 := \left\{ a111=0, a112=0, a113=0, a114=0, a115=0, a121=0, a122=0, a123=0, \quad (5.58) \right.$$

$$\begin{aligned}
& a124=0, a125=0, a131 = -\frac{1}{2} u_{1,1}, a132=0, a133=0, a134=0, a135=0, a141 = \\
& -u_{1,2}, a142 = -\frac{1}{2} u_{1,1}, a143=0, a144=0, a145=0, a151 = -u_1 u_{1,1} - \frac{3}{2} u_{2,2}, a152 \\
& = -u_{1,2}, a153 = -\frac{1}{2} u_{1,1}, a154=0, a155=0, a221 = u_{1,1}, a222=0, a223=0, a224 \\
& = 0, a225=0, a231 = u_{1,2}, a232 = \frac{1}{2} u_{1,1}, a233=0, a234=0, a235=0, a241 \\
& = \frac{3}{2} u_1 u_{1,1} + u_{2,2}, a242=0, a243 = \frac{1}{2} u_{1,1}, a244=0, a245=0, a251 = 2 u_1 u_{1,2} \\
& + \frac{3}{2} u_2 u_{1,1} + u_{2,3}, a252 = \frac{1}{2} u_1 u_{1,1} - \frac{1}{2} u_{2,2}, a253=0, a254 = \frac{1}{2} u_{1,1}, a255=0, \\
& a331 = -u_1 u_{1,1} + u_{2,2}, a332 = 2 u_{1,2}, a333=0, a334=0, a335=0, a341 = -u_1 u_{1,2} \\
& - \frac{1}{2} u_2 u_{1,1} + u_{2,3}, a342 = \frac{1}{2} u_1 u_{1,1} + 2 u_{2,2}, a343 = u_{1,2}, a344=0, a345=0, a351 \\
& = -\frac{5}{2} u_1^2 u_{1,1} - u_1 u_{2,2} + u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + u_{3,3}, a352 = u_1 u_{1,2} + u_2 u_{1,1} \\
& + 2 u_{2,3}, a353 = -\frac{1}{2} u_1 u_{1,1} + \frac{1}{2} u_{2,2}, a354 = u_{1,2}, a355=0, a441 = u_1^2 u_{1,1} - u_1 u_{2,2} \\
& - 2 u_2 u_{1,2} + u_{3,3}, a442 = -2 u_1 u_{1,2} - u_2 u_{1,1} + 2 u_{2,3}, a443 = 2 u_1 u_{1,1} + 3 u_{2,2}, a444 \\
& = 0, a445=0, a451 = u_1^2 u_{1,2} - u_1 u_{2,3} - 2 u_2 u_{2,2} - u_3 u_{1,2} + u_{3,4}, a452 = -\frac{3}{2} u_1^2 u_{1,1}
\end{aligned}$$

$$\begin{aligned}
& -2 u_1 u_{2,2} - u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + 2 u_{3,3}, a453 = \frac{1}{2} u_2 u_{1,1} + 3 u_{2,3}, a454 = u_1 u_{1,1} \\
& + \frac{3}{2} u_{2,2}, a455 = 0, a551 = -u_1^3 u_{1,1} + u_1^2 u_{2,2} + 4 u_1 u_2 u_{1,2} - 2 u_1 u_3 u_{1,1} + u_2^2 u_{1,1} \\
& - u_1 u_{3,3} - 2 u_2 u_{2,3} - 3 u_3 u_{2,2} + u_{4,4}, a552 = 2 u_1^2 u_{1,2} - 2 u_1 u_{2,3} - 4 u_2 u_{2,2} \\
& - 2 u_3 u_{1,2} + 2 u_{3,4}, a553 = -4 u_1^2 u_{1,1} - 3 u_1 u_{2,2} - u_3 u_{1,1} + 3 u_{3,3}, a554 = 2 u_1 u_{1,2} \\
& + 2 u_2 u_{1,1} + 4 u_{2,3}, a555 = 0 \}
\end{aligned}$$

> CsC0 := subs(Sss20, CsC) :

> DGsimplify(CsC0)

$$\begin{aligned}
& - \left(\left(\left(\frac{u_{1,1}}{2} D_{-y} \right) dx \right) dz \right) + \left(\left(\left(u_1 u_{1,1} + \frac{3 u_{2,2}}{2} \right) D_{-z} \right) ds \right) dz + \left((u_{1,2} D_{-z}) ds \right) dt \quad (5.59) \\
& + \left(\left(\left(2 u_1 u_{1,2} + \frac{3 u_2 u_{1,1}}{2} + u_{2,3} \right) D_{-x} \right) ds \right) dy + \left((u_{1,2} D_{-x}) dt \right) dy + \left(\left((u_1^2 u_{1,1} \right. \right. \\
& \left. \left. - u_1 u_{2,2} - 2 u_2 u_{1,2} + u_{3,3} \right) D_{-x} \right) dz \right) dz - \left((u_{1,2} D_{-x}) dz \right) dx \\
& - \left((u_{1,2} D_{-y}) ds \right) dx + \left(\left(\left(\frac{3 u_1 u_{1,1}}{2} + u_{2,2} \right) D_{-x} \right) dz \right) dy + \left(\left(\left(\frac{u_1 u_{1,1}}{2} \right. \right. \right. \\
& \left. \left. - \frac{u_{2,2}}{2} \right) D_{-y} \right) dy \right) ds + \left((2 u_1 u_{1,2} + 2 u_2 u_{1,1} + 4 u_{2,3}) D_{-z} \right) ds \right) ds \\
& + \left(\left(\left(\left(\frac{u_2 u_{1,1}}{2} + 3 u_{2,3} \right) D_{-t} \right) dz \right) ds - \left(\left(\left(\frac{3}{2} u_1^2 u_{1,1} + 2 u_1 u_{2,2} + u_2 u_{1,2} \right. \right. \right. \\
& \left. \left. + \frac{1}{2} u_3 u_{1,1} - 2 u_{3,3} \right) D_{-y} \right) dz \right) ds + \left(\left(\left(\frac{3 u_1 u_{1,1}}{2} + u_{2,2} \right) D_{-x} \right) dy \right) dz \\
& - \left(\left(\left(\left(\frac{5}{2} u_1^2 u_{1,1} + u_1 u_{2,2} - u_2 u_{1,2} + \frac{1}{2} u_3 u_{1,1} - u_{3,3} \right) D_{-x} \right) ds \right) dt \right) \\
& + \left((u_1 u_{1,2} + u_2 u_{1,1} + 2 u_{2,3}) D_{-y} \right) ds \right) dt - \left(\left(\left(\frac{u_{1,1}}{2} D_{-x} \right) dx \right) dt \right) \\
& - \left(\left(\left(\left(\frac{u_1 u_{1,1}}{2} - \frac{u_{2,2}}{2} \right) D_{-t} \right) dt \right) ds \right) + \left(\left(\left(\frac{u_2 u_{1,1}}{2} + 3 u_{2,3} \right) D_{-t} \right) ds \right) dz \\
& + \left(\left((2 u_1^2 u_{1,2} - 2 u_1 u_{2,3} - 4 u_2 u_{2,2} - 2 u_3 u_{1,2} + 2 u_{3,4}) D_{-y} \right) ds \right) ds \\
& - \left(\left(\left(\frac{u_{1,1}}{2} D_{-t} \right) ds \right) dx \right) + \left(\left((u_1^2 u_{1,2} - u_1 u_{2,3} - 2 u_2 u_{2,2} - u_3 u_{1,2} \right. \right. \right. \\
& \left. \left. + u_{3,4} \right) D_{-x} \right) dz \right) ds - \left(\left((2 u_1 u_{1,2} + u_2 u_{1,1} - 2 u_{2,3}) D_{-y} \right) dz \right) dz + \left((u_1 u_{1,2} \right. \\
& \left. + u_2 u_{1,1} + 2 u_{2,3}) D_{-y} \right) dt \right) ds + \left((u_{1,2} D_{-t}) dt \right) dz - \left((u_{1,2} D_{-x}) dx \right) dz \\
& + \left((u_{1,2} D_{-z}) dt \right) ds + \left(\left(\frac{u_{1,1}}{2} D_{-y} \right) dy \right) dt + \left(\left(\left(\frac{u_1 u_{1,1}}{2} + 2 u_{2,2} \right) D_{-y} \right) dz \right) dt \\
& + \left((2 u_{1,2} D_{-y}) dt \right) dt + \left(\left(\left(u_1 u_{1,1} + \frac{3 u_{2,2}}{2} \right) D_{-z} \right) dz \right) ds + \left(\left(\left(\frac{u_1 u_{1,1}}{2} \right. \right. \right.
\end{aligned}$$

$$\begin{aligned}
& -\frac{u_{2,2}}{2} \Big) D_{-y} \Big) ds \Big) dy + \Big(\Big(\Big(\frac{u_{1,1}}{2} D_{-t} \Big) dy \Big) dz - \Big(\Big(\Big(\frac{u_{1,1}}{2} D_{-y} \Big) dz \Big) dx \Big) \\
& - \Big(\Big(\Big(\Big(u_1 u_{1,2} + \frac{u_2 u_{1,1}}{2} - u_{2,3} \Big) D_{-x} \Big) dz \Big) dt \Big) - \Big(\Big(\Big(\Big(u_1 u_{1,2} + \frac{u_2 u_{1,1}}{2} \\
& - u_{2,3} \Big) D_{-x} \Big) dt \Big) dz \Big) - \Big(\Big(\Big(\Big(u_1^3 u_{1,1} - u_1^2 u_{2,2} - 4 u_1 u_2 u_{1,2} + 2 u_1 u_3 u_{1,1} - u_2^2 u_{1,1} \\
& + u_1 u_{3,3} + 2 u_2 u_{2,3} + 3 u_3 u_{2,2} - u_{4,4} \Big) D_{-x} \Big) ds \Big) ds \Big) - \Big(\Big(\Big(\Big(u_1 u_{1,1} \\
& + \frac{3 u_{2,2}}{2} \Big) D_{-x} \Big) ds \Big) dx \Big) + \Big(\Big(\Big(\frac{u_{1,1}}{2} D_{-z} \Big) ds \Big) dy - \Big(\Big(\Big(\Big(\frac{u_1 u_{1,1}}{2} \\
& - \frac{u_{2,2}}{2} \Big) D_{-t} \Big) ds \Big) dt \Big) + \Big((u_{1,1} D_{-x}) dy \Big) dy - \Big(\Big(\Big(\Big(\frac{3}{2} u_1^2 u_{1,1} + 2 u_1 u_{2,2} \\
& + u_2 u_{1,2} + \frac{1}{2} u_3 u_{1,1} - 2 u_{3,3} \Big) D_{-y} \Big) ds \Big) dz \Big) + \Big(\Big(\Big(\frac{u_{1,1}}{2} D_{-t} \Big) dz \Big) dy \\
& + \Big((u_{1,2} D_{-x}) dy \Big) dt - \Big(\Big(\Big(\Big(u_1 u_{1,1} + \frac{3 u_{2,2}}{2} \Big) D_{-x} \Big) dx \Big) ds \Big) + \Big(\Big(\Big(\Big(2 u_1 u_{1,2} \\
& + \frac{3 u_2 u_{1,1}}{2} + u_{2,3} \Big) D_{-x} \Big) dy \Big) ds - \Big(\Big(\Big(\Big(\frac{5}{2} u_1^2 u_{1,1} + u_1 u_{2,2} - u_2 u_{1,2} + \frac{1}{2} u_3 u_{1,1} \\
& - u_{3,3} \Big) D_{-x} \Big) dt \Big) ds \Big) + \Big(\Big(\Big(\frac{u_{1,1}}{2} D_{-y} \Big) dt \Big) dy + \Big((u_{1,2} D_{-t}) dz \Big) dt - \Big(\Big(\Big((u_1 u_{1,1} \\
& - u_{2,2}) D_{-x} \Big) dt \Big) dt \Big) - \Big(\Big((u_{1,2} D_{-y}) dx \Big) ds \Big) - \Big(\Big(\Big(\Big(\frac{u_{1,1}}{2} D_{-t} \Big) dx \Big) ds \Big) + \Big(\Big(\Big(\\
& u_1^2 u_{1,2} - u_1 u_{2,3} - 2 u_2 u_{2,2} - u_3 u_{1,2} + u_{3,4} \Big) D_{-x} \Big) ds \Big) dz + \Big(\Big(\Big(2 u_1 u_{1,1} \\
& + 3 u_{2,2} \Big) D_{-t} \Big) dz \Big) dz - \Big(\Big(\Big(\Big(4 u_1^2 u_{1,1} + 3 u_1 u_{2,2} + u_3 u_{1,1} - 3 u_{3,3} \Big) D_{-t} \Big) ds \Big) ds \Big) \\
& + \Big(\Big(\Big(\frac{u_{1,1}}{2} D_{-z} \Big) dy \Big) ds + \Big(\Big(\Big(\Big(\frac{u_1 u_{1,1}}{2} + 2 u_{2,2} \Big) D_{-y} \Big) dt \Big) dz \\
& - \Big(\Big(\Big(\Big(\frac{u_{1,1}}{2} D_{-x} \Big) dt \Big) dx \Big)
\end{aligned}$$

$$> \Pi_{31} = -\frac{1}{2} u_{1,1},$$

$$\Pi_{41} = -u_{1,2}, \Pi_{42} = -\frac{1}{2} u_{1,1},$$

$$\Pi_{51} = -u_1 u_{1,1} - \frac{3}{2} u_{2,2}, \Pi_{52} = -u_{1,2}, \Pi_{53} = -\frac{1}{2} u_{1,1},$$

$$\Pi_{21} = u_{1,1},$$

$$\Pi_{31} = u_{1,2}, \Pi_{32} = \frac{1}{2} u_{1,1},$$

$$\Pi_{41} = \frac{3}{2} u_1 u_{1,1} + u_{2,2}, \Pi_{43} = \frac{1}{2} u_{1,1},$$

$$\begin{aligned}
I251 &= 2 u_1 u_{1,2} + \frac{3}{2} u_2 u_{1,1} + u_{2,3}, I252 = \frac{1}{2} u_1 u_{1,1} - \frac{1}{2} u_{2,2}, I254 = \frac{1}{2} u_{1,1}, \\
I331 &= -u_1 u_{1,1} + u_{2,2}, I332 = 2 u_{1,2}, \\
I341 &= -u_1 u_{1,2} - \frac{1}{2} u_2 u_{1,1} + u_{2,3}, I342 = \frac{1}{2} u_1 u_{1,1} + 2 u_{2,2}, I343 = u_{1,2}, \\
I351 &= -\frac{5}{2} u_1^2 u_{1,1} - u_1 u_{2,2} + u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + u_{3,3}, I352 = u_1 u_{1,2} + u_2 u_{1,1} \\
&\quad + 2 u_{2,3}, I353 = -\frac{1}{2} u_1 u_{1,1} + \frac{1}{2} u_{2,2}, I354 = u_{1,2}, \\
I441 &= u_1^2 u_{1,1} - u_1 u_{2,2} - 2 u_2 u_{1,2} + u_{3,3}, I442 = -2 u_1 u_{1,2} - u_2 u_{1,1} + 2 u_{2,3}, I443 \\
&= 2 u_1 u_{1,1} + 3 u_{2,2}, \\
I451 &= u_1^2 u_{1,2} - u_1 u_{2,3} - 2 u_2 u_{2,2} - u_3 u_{1,2} + u_{3,4}, I452 = -\frac{3}{2} u_1^2 u_{1,1} - 2 u_1 u_{2,2} \\
&\quad - u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + 2 u_{3,3}, I453 = \frac{1}{2} u_2 u_{1,1} + 3 u_{2,3}, I454 = u_1 u_{1,1} + \frac{3}{2} u_{2,2}, \\
I551 &= -u_1^3 u_{1,1} + u_1^2 u_{2,2} + 4 u_1 u_2 u_{1,2} - 2 u_1 u_3 u_{1,1} + u_2^2 u_{1,1} - u_1 u_{3,3} - 2 u_2 u_{2,3} \\
&\quad - 3 u_3 u_{2,2} + u_{4,4}, I552 = 2 u_1^2 u_{1,2} - 2 u_1 u_{2,3} - 4 u_2 u_{2,2} - 2 u_3 u_{1,2} + 2 u_{3,4}, I553 = \\
&\quad -4 u_1^2 u_{1,1} - 3 u_1 u_{2,2} - u_3 u_{1,1} + 3 u_{3,3}, I554 = 2 u_1 u_{1,2} + 2 u_2 u_{1,1} + 4 u_{2,3}
\end{aligned}$$

Veronese web hierarchy 5D

$$\begin{aligned}
&> DGsetup([x1, x2, x3, x4, x5], [u], Ju, 3) : \\
&> X2f := f \rightarrow TotalDiff(f, 2) - \frac{(q - a_1) u_2}{(q - a_2) u_1} \cdot TotalDiff(f, 1) : \\
&X3f := f \rightarrow TotalDiff(f, 3) - \frac{(q - a_1) u_3}{(q - a_3) u_1} \cdot TotalDiff(f, 1) : \\
&X4f := f \rightarrow TotalDiff(f, 4) - \frac{(q - a_1) u_4}{(q - a_4) u_1} \cdot TotalDiff(f, 1) : \\
&X5f := f \rightarrow TotalDiff(f, 5) - \frac{(q - a_1) u_5}{(q - a_5) u_1} \cdot TotalDiff(f, 1) : \\
&> numer(X2f(X3f(x1)) - X3f(X2f(x1))); numer(X2f(X4f(x1)) - X4f(X2f(x1))); \\
&\quad numer(X3f(X4f(x1)) - X4f(X3f(x1))); numer(X2f(X5f(x1)) - X5f(X2f(x1))); \\
&\quad numer(X3f(X5f(x1)) - X5f(X3f(x1))); numer(X4f(X5f(x1)) - X5f(X4f(x1))) \\
&\quad - (q - a_1) u_1 (a_1 u_2 u_{1,3} - a_1 u_3 u_{1,2} - a_2 u_1 u_{2,3} + a_2 u_3 u_{1,2} + a_3 u_1 u_{2,3} - a_3 u_2 u_{1,3}) \\
&\quad - (q - a_1) u_1 (a_1 u_2 u_{1,4} - a_1 u_4 u_{1,2} - a_2 u_1 u_{2,4} + a_2 u_4 u_{1,2} + a_4 u_1 u_{2,4} - a_4 u_2 u_{1,4}) \\
&\quad - (q - a_1) u_1 (a_1 u_3 u_{1,4} - a_1 u_4 u_{1,3} - a_3 u_1 u_{3,4} + a_3 u_4 u_{1,3} + a_4 u_1 u_{3,4} - a_4 u_3 u_{1,4}) \\
&\quad - (q - a_1) u_1 (a_1 u_2 u_{1,5} - a_1 u_5 u_{1,2} - a_2 u_1 u_{2,5} + a_2 u_5 u_{1,2} + a_5 u_1 u_{2,5} - a_5 u_2 u_{1,5}) \\
&\quad - (q - a_1) u_1 (a_1 u_3 u_{1,5} - a_1 u_5 u_{1,3} - a_3 u_1 u_{3,5} + a_3 u_5 u_{1,3} + a_5 u_1 u_{3,5} - a_5 u_3 u_{1,5})
\end{aligned}$$

$$-(q - a_1) u_1 (a_1 u_4 u_{1,5} - a_1 u_5 u_{1,4} - a_4 u_1 u_{4,5} + a_4 u_5 u_{1,4} + a_5 u_1 u_{4,5} - a_5 u_4 u_{1,5}) \quad (6.1)$$

$$\begin{aligned} & \text{Eq1} := \text{collect}(a_1 u_2 u_{1,3} - a_1 u_3 u_{1,2} - a_2 u_1 u_{2,3} + a_2 u_3 u_{1,2} + a_3 u_1 u_{2,3} - a_3 u_2 u_{1,3}, \\ & \quad \{u_{1,2}, u_{1,3}, u_{1,4}, u_{1,5}, u_{2,3}, u_{2,4}, u_{2,5}, u_{3,4}, u_{3,5}, u_{4,5}\}); \\ & \text{Eq2} := \text{collect}(a_1 u_2 u_{1,4} - a_1 u_4 u_{1,2} - a_2 u_1 u_{2,4} + a_2 u_4 u_{1,2} + a_4 u_1 u_{2,4} - a_4 u_2 u_{1,4}, \\ & \quad \{u_{1,2}, u_{1,3}, u_{1,4}, u_{1,5}, u_{2,3}, u_{2,4}, u_{2,5}, u_{3,4}, u_{3,5}, u_{4,5}\}); \\ & \text{Eq3} := \text{collect}(a_1 u_3 u_{1,4} - a_1 u_4 u_{1,3} - a_3 u_1 u_{3,4} + a_3 u_4 u_{1,3} + a_4 u_1 u_{3,4} - a_4 u_3 u_{1,4}, \\ & \quad \{u_{1,2}, u_{1,3}, u_{1,4}, u_{1,5}, u_{2,3}, u_{2,4}, u_{2,5}, u_{3,4}, u_{3,5}, u_{4,5}\}); \\ & \text{Eq4} := \text{collect}(a_1 u_2 u_{1,5} - a_1 u_5 u_{1,2} - a_2 u_1 u_{2,5} + a_2 u_5 u_{1,2} + a_5 u_1 u_{2,5} - a_5 u_2 u_{1,5}, \\ & \quad \{u_{1,2}, u_{1,3}, u_{1,4}, u_{1,5}, u_{2,3}, u_{2,4}, u_{2,5}, u_{3,4}, u_{3,5}, u_{4,5}\}); \\ & \text{Eq5} := \text{collect}(a_1 u_3 u_{1,5} - a_1 u_5 u_{1,3} - a_3 u_1 u_{3,5} + a_3 u_5 u_{1,3} + a_5 u_1 u_{3,5} - a_5 u_3 u_{1,5}, \\ & \quad \{u_{1,2}, u_{1,3}, u_{1,4}, u_{1,5}, u_{2,3}, u_{2,4}, u_{2,5}, u_{3,4}, u_{3,5}, u_{4,5}\}); \\ & \text{Eq6} := \text{collect}(a_1 u_4 u_{1,5} - a_1 u_5 u_{1,4} - a_4 u_1 u_{4,5} + a_4 u_5 u_{1,4} + a_5 u_1 u_{4,5} - a_5 u_4 u_{1,5}, \\ & \quad \{u_{1,2}, u_{1,3}, u_{1,4}, u_{1,5}, u_{2,3}, u_{2,4}, u_{2,5}, u_{3,4}, u_{3,5}, u_{4,5}\}); \\ & \text{Eq1} := (-a_1 u_3 + a_2 u_3) u_{1,2} + (a_1 u_2 - a_3 u_2) u_{1,3} + (-a_2 u_1 + a_3 u_1) u_{2,3} \\ & \text{Eq2} := (-a_1 u_4 + a_2 u_4) u_{1,2} + (a_1 u_2 - a_4 u_2) u_{1,4} + (-a_2 u_1 + a_4 u_1) u_{2,4} \\ & \text{Eq3} := (-a_1 u_4 + a_3 u_4) u_{1,3} + (a_1 u_3 - a_4 u_3) u_{1,4} + (-a_3 u_1 + a_4 u_1) u_{3,4} \\ & \text{Eq4} := (-a_1 u_5 + a_2 u_5) u_{1,2} + (a_1 u_2 - a_5 u_2) u_{1,5} + (-a_2 u_1 + a_5 u_1) u_{2,5} \\ & \text{Eq5} := (-a_1 u_5 + a_3 u_5) u_{1,3} + (a_1 u_3 - a_5 u_3) u_{1,5} + (-a_3 u_1 + a_5 u_1) u_{3,5} \\ & \text{Eq6} := (-a_1 u_5 + a_4 u_5) u_{1,4} + (a_1 u_4 - a_5 u_4) u_{1,5} + (-a_4 u_1 + a_5 u_1) u_{4,5} \end{aligned} \quad (6.2)$$

$$\text{S2} := \text{solve}(\{\text{Eq1}, \text{Eq2}, \text{Eq3}, \text{Eq4}, \text{Eq5}, \text{Eq6}\}, \{u_{2,3}, u_{2,4}, u_{2,5}, u_{3,4}, u_{3,5}, u_{4,5}\})$$

$$\begin{aligned} \text{S2} &:= \left\{ u_{2,3} = \frac{a_1 u_2 u_{1,3} - a_1 u_3 u_{1,2} + a_2 u_3 u_{1,2} - a_3 u_2 u_{1,3}}{u_1 (a_2 - a_3)}, u_{2,4} \right. \\ &= \frac{a_1 u_2 u_{1,4} - a_1 u_4 u_{1,2} + a_2 u_4 u_{1,2} - a_4 u_2 u_{1,4}}{u_1 (a_2 - a_4)}, u_{2,5} \\ &= \frac{a_1 u_2 u_{1,5} - a_1 u_5 u_{1,2} + a_2 u_5 u_{1,2} - a_5 u_2 u_{1,5}}{u_1 (a_2 - a_5)}, u_{3,4} \\ &= \frac{a_1 u_3 u_{1,4} - a_1 u_4 u_{1,3} + a_3 u_4 u_{1,3} - a_4 u_3 u_{1,4}}{u_1 (a_3 - a_4)}, u_{3,5} \\ &= \frac{a_1 u_3 u_{1,5} - a_1 u_5 u_{1,3} + a_3 u_5 u_{1,3} - a_5 u_3 u_{1,5}}{u_1 (a_3 - a_5)}, u_{4,5} \\ &= \left. \frac{a_1 u_4 u_{1,5} - a_1 u_5 u_{1,4} + a_4 u_5 u_{1,4} - a_5 u_4 u_{1,5}}{u_1 (a_4 - a_5)} \right\} \end{aligned} \quad (6.3)$$

$$\begin{aligned} & \text{simplify}(\text{subs}(S2, u_4 \cdot \text{TotalDiff}(\text{Eq1}, 1) - u_1 \cdot \text{TotalDiff}(\text{Eq1}, 4) - u_3 \cdot \text{TotalDiff}(\text{Eq2}, 1) \\ & \quad + u_1 \cdot \text{TotalDiff}(\text{Eq2}, 3) + u_2 \cdot \text{TotalDiff}(\text{Eq3}, 1) - u_1 \cdot \text{TotalDiff}(\text{Eq3}, 2))) \\ & \text{simplify}(\text{subs}(VsS2, (a_1 - a_4) \cdot \text{TotalDiff}(\text{Eq1}, 4) + (a_3 - a_1) \cdot \text{TotalDiff}(\text{Eq2}, 3) + (a_1 \\ & \quad - a_2) \cdot \text{TotalDiff}(\text{Eq3}, 2))) \end{aligned}$$

> Vss3 := {Eq1, Eq2, Eq3, Eq4, Eq5, Eq6} :for i to 5 do for j to 6 do Vss3 := {op(Vss3),
TotalDiff(Eq||j, i)} :od:od:
tmp := eliminate(Vss3, {u_{2,3}, u_{2,4}, u_{2,5}, u_{3,4}, u_{3,5}, u_{4,5}, u_{1,2}, u_{1,3}, u_{1,4}, u_{1,5}, u_{1,3,4},
u_{1,3,5}, u_{1,4,5}, u_{2,2,3}, u_{2,2,4}, u_{2,2,5}, u_{2,3,3}, u_{2,3,4}, u_{2,3,5}, u_{2,4,4}, u_{2,4,5}, u_{2,5,5}, u_{3,3,4},
u_{3,3,5}, u_{3,4,4}, u_{3,4,5}, u_{3,5,5}, u_{4,4,5}, u_{4,5,5}}) : [nops(tmp[1]), nops(tmp[2])];
[28, 0]

(6.5)

> S3 := solve(Vss3, {u_{2,3}, u_{2,4}, u_{2,5}, u_{3,4}, u_{3,5}, u_{4,5}, u_{1,2}, u_{1,3}, u_{1,4}, u_{1,5}, u_{1,3,4}, u_{1,3,5},
u_{1,4,5}, u_{2,2,3}, u_{2,2,4}, u_{2,2,5}, u_{2,3,3}, u_{2,3,4}, u_{2,3,5}, u_{2,4,4}, u_{2,4,5}, u_{2,5,5}, u_{3,3,4}, u_{3,3,5},
u_{3,4,4}, u_{3,4,5}, u_{3,5,5}, u_{4,4,5}, u_{4,5,5}}) :

> nops(%)

28

(6.6)

> # ... #

> SymbE := k → add(add(diff(Eq||k, u_{i,j})·p_i·p_j, j=i..5), i=1..5) :

> simplify(subs({seq(p_k = $\frac{u[k]}{q - a[k]}$, k=1..5)}, {seq(SymbE(k), k=1..6)}))
{0}

(6.7)

> tmp := solve({coeffs(expand(add(add(a[k, l]·p_k, k=1..5)·SymbE(l), l=1..6)), {p₁, p₂,
p₃, p₄, p₅}}), {seq(seq(a[k, l], k=1..5), l=1..6)})

tmp := $\left\{ a_{1,1} = \frac{1}{(a_3 - a_5)(a_3 - a_4)} u_1 (a_1 a_3 a_{3,2} u_4 + a_1 a_3 a_{3,4} u_5 + a_1 a_3 a_{4,1} u_4 \right.$

(6.8)

$+ a_1 a_3 a_{5,1} u_5 - a_1 a_4 a_{3,4} u_5 - a_1 a_4 a_{5,1} u_5 - a_1 a_5 a_{3,2} u_4 - a_1 a_5 a_{4,1} u_4 - a_3^2 a_{4,1} u_4$

$- a_3^2 a_{5,1} u_5 - a_3 a_4 a_{3,2} u_4 + a_3 a_4 a_{5,1} u_5 - a_3 a_5 a_{3,4} u_5 + a_3 a_5 a_{4,1} u_4 + a_4 a_5 a_{3,2} u_4$

$+ a_4 a_5 a_{3,4} u_5), a_{1,2} = \frac{1}{(a_4 - a_5)(a_3 - a_4)} u_1 (a_1 a_3 a_{4,4} u_5 + a_1 a_3 a_{5,2} u_5$

$- a_1 a_4 a_{3,2} u_3 - a_1 a_4 a_{4,1} u_3 - a_1 a_4 a_{4,4} u_5 - a_1 a_4 a_{5,2} u_5 + a_1 a_5 a_{3,2} u_3$

$+ a_1 a_5 a_{4,1} u_3 + a_3 a_4 a_{4,1} u_3 - a_3 a_4 a_{5,2} u_5 - a_3 a_5 a_{4,1} u_3 - a_3 a_5 a_{4,4} u_5 + a_4^2 a_{3,2} u_3$

$+ a_4^2 a_{5,2} u_5 - a_4 a_5 a_{3,2} u_3 + a_4 a_5 a_{4,4} u_5), a_{1,3}$

$= \frac{1}{u_1 (a_3 a_4 - a_3 a_5 - a_4^2 + a_4 a_5)} (a_1 a_3 a_{4,5} u_5 + a_1 a_3 a_{5,3} u_5 + a_1 a_4 a_{3,2} u_2$

$$\begin{aligned}
& + a_1 a_4 a_{4,1} u_2 - a_1 a_4 a_{4,5} u_5 - a_1 a_4 a_{5,3} u_5 - a_1 a_5 a_{3,2} u_2 - a_1 a_5 a_{4,1} u_2 \\
& - a_3 a_4 a_{4,1} u_2 - a_3 a_4 a_{5,3} u_5 + a_3 a_5 a_{4,1} u_2 - a_3 a_5 a_{4,5} u_5 - a_4^2 a_{3,2} u_2 + a_4^2 a_{5,3} u_5 \\
& + a_4 a_5 a_{3,2} u_2 + a_4 a_5 a_{4,5} u_5), a_{1,4} = - \frac{1}{(a_4 - a_5)(a_3 - a_5)u_1} (a_1 a_3 a_{4,4} u_4 \\
& + a_1 a_3 a_{5,2} u_4 + a_1 a_4 a_{3,4} u_3 + a_1 a_4 a_{5,1} u_3 - a_1 a_5 a_{3,4} u_3 - a_1 a_5 a_{4,4} u_4 \\
& - a_1 a_5 a_{5,1} u_3 - a_1 a_5 a_{5,2} u_4 - a_3 a_4 a_{5,1} u_3 - a_3 a_4 a_{5,2} u_4 - a_3 a_5 a_{4,4} u_4 \\
& + a_3 a_5 a_{5,1} u_3 - a_4 a_5 a_{3,4} u_3 + a_4 a_5 a_{5,2} u_4 + a_5^2 a_{3,4} u_3 + a_5^2 a_{4,4} u_4), a_{1,5} = \\
& - \frac{1}{u_1 (a_3 a_4 - a_3 a_5 - a_4 a_5 + a_5^2)} (a_1 a_3 a_{4,5} u_4 + a_1 a_3 a_{5,3} u_4 - a_1 a_4 a_{3,4} u_2 \\
& - a_1 a_4 a_{5,1} u_2 + a_1 a_5 a_{3,4} u_2 - a_1 a_5 a_{4,5} u_4 + a_1 a_5 a_{5,1} u_2 - a_1 a_5 a_{5,3} u_4 \\
& + a_3 a_4 a_{5,1} u_2 - a_3 a_4 a_{5,3} u_4 - a_3 a_5 a_{4,5} u_4 - a_3 a_5 a_{5,1} u_2 + a_4 a_5 a_{3,4} u_2 \\
& + a_4 a_5 a_{5,3} u_4 - a_5^2 a_{3,4} u_2 + a_5^2 a_{4,5} u_4), a_{1,6} = \frac{1}{u_1 (a_4 - a_5)} (a_1 a_{4,4} u_2 \\
& + a_1 a_{4,5} u_3 + a_1 a_{5,2} u_2 + a_1 a_{5,3} u_3 - a_4 a_{5,2} u_2 - a_4 a_{5,3} u_3 - a_5 a_{4,4} u_2 - a_5 a_{4,5} u_3) \\
& , a_{2,1} = 0, a_{2,2} = 0, a_{2,3} = - \frac{a_2 a_{3,2} + a_2 a_{4,1} - a_3 a_{4,1} - a_4 a_{3,2}}{a_3 - a_4}, a_{2,4} = 0, a_{2,5} = \\
& - \frac{a_2 a_{3,4} + a_2 a_{5,1} - a_3 a_{5,1} - a_5 a_{3,4}}{a_3 - a_5}, a_{2,6} = - \frac{a_2 a_{4,4} + a_2 a_{5,2} - a_4 a_{5,2} - a_5 a_{4,4}}{a_4 - a_5}, \\
& a_{3,1} = 0, a_{3,2} = a_{3,2}, a_{3,3} = 0, a_{3,4} = a_{3,4}, a_{3,5} = 0, a_{3,6} = \\
& - \frac{a_3 a_{4,5} + a_3 a_{5,3} - a_4 a_{5,3} - a_5 a_{4,5}}{a_4 - a_5}, a_{4,1} = a_{4,1}, a_{4,2} = 0, a_{4,3} = 0, a_{4,4} = a_{4,4}, a_{4,5} \\
& = a_{4,5}, a_{4,6} = 0, a_{5,1} = a_{5,1}, a_{5,2} = a_{5,2}, a_{5,3} = a_{5,3}, a_{5,4} = 0, a_{5,5} = 0, a_{5,6} = 0 \Bigg\}
\end{aligned}$$

so 8 compatibility conditions - we check them at one step

$$\text{> simplify(subs(S2, subs(tmp, add(add(a[k, l] \cdot \text{TotalDiff}(Eq \| l, k), k = 1 \dots 5), l = 1 \dots 6))) \quad (6.9)$$

This shows that the Veronese system is compatible and the Hilbert polynomial is $\dim(g[k])=1+4k$, whence the general solution depends on 4 functions of 1 variable.

This is also what is expected from the characteristic variety (rational normal curve in our case) that has order 4 and dimension 1.

$$\begin{aligned} &\text{> DGsetup}([x1, x2, x3, x4, x5], M) : \\ &\text{> } vM := [D_x1, D_x2, D_x3, D_x4, D_x5] : dM := [dx1, dx2, dx3, dx4, dx5] : \\ &\text{> } C := \text{Connection}(add(add(add(((a \| i) \| j) \| k) \cdot vM[k] \&t dM[i] \&t dM[j], k = 1 \dots 5), j = 1 \dots 5), i = 1 \dots 5)) : \\ &\text{> } X2 := \text{evalDG}\left(D_x2 - \frac{(q - a_1) u_2}{(q - a_2) u_1} \cdot D_x1\right) : \\ &\quad X3 := \text{evalDG}\left(D_x3 - \frac{(q - a_1) u_3}{(q - a_3) u_1} \cdot D_x1\right) : \\ &\quad X4 := \text{evalDG}\left(D_x4 - \frac{(q - a_1) u_4}{(q - a_4) u_1} \cdot D_x1\right) : \\ &\quad X5 := \text{evalDG}\left(D_x5 - \frac{(q - a_1) u_5}{(q - a_5) u_1} \cdot D_x1\right) : \\ &\quad \omega := \text{evalDG}\left(\frac{u_1}{q - a_1} \cdot dx1 + \frac{u_2}{q - a_2} \cdot dx2 + \frac{u_3}{q - a_3} \cdot dx3 + \frac{u_4}{q - a_4} \cdot dx4 + \frac{u_5}{q - a_5} \cdot dx5\right) : \\ &\quad [\text{Hook}(X2, \omega), \text{Hook}(X3, \omega), \text{Hook}(X4, \omega), \text{Hook}(X5, \omega)] \\ &\quad [0, 0, 0, 0] \quad (6.10) \end{aligned}$$

$$\begin{aligned} &\text{> } XX2 := \text{FromJet}(X2, \{u(x1, x2, x3, x4, x5)\}) : XX3 := \text{FromJet}(X3, \{u(x1, x2, x3, x4, x5)\}) : XX4 := \text{FromJet}(X4, \{u(x1, x2, x3, x4, x5)\}) : XX5 := \text{FromJet}(X5, \{u(x1, x2, x3, x4, x5)\}) : \\ &\quad \omega\omega := \text{FromJet}(\omega, \{u(x1, x2, x3, x4, x5)\}) : \\ &\text{> } DD := \{XX2, XX3, XX4, XX5\} : \text{ChrCoe} := \{\text{seq}(\text{seq}(\text{seq}(((a \| i) \| j) \| k), i = 1 \dots 5), j = 1 \dots 5), k = 1 \dots 5)\} : \\ &\quad TGE := \{\} : \text{for } i \text{ to } 4 \text{ do for } j \text{ to } 4 \text{ do } TGE := \{\text{op}(TGE), \text{coeffs}(\text{expand}(\text{numer}(\text{subs}(VsS2, \text{ToJet}(\text{Hook}(\text{convert}(\text{DirectionalCovariantDerivative}(DD[i], DD[j], C), 'DGvector'), \omega\omega), \{u(x1, x2, x3, x4, x5)\}, \text{notation} = \text{jetnumbers})) \&t dM[i] \&t dM[j], C), \{q\})\} : \text{od} : \text{od} : \\ &\quad [\text{nops}(\text{ChrCoe}), \text{nops}(TGE)] \\ &\quad [125, 112] \quad (6.11) \end{aligned}$$

$$\begin{aligned} &\text{> } Ck := \text{Connection}\left(\text{evalDG}\left(\text{add}\left(\text{add}\left(\left(\frac{u_{i,j}}{u_i} + \alpha_j\right) \cdot vM[i] \&t dM[i] \&t dM[j], j = i \dots 5\right), i = 1 \dots 5\right) + \text{add}\left(\text{add}\left(\left(\frac{u_{j,i}}{u_i} + \alpha_j\right) \cdot vM[i] \&t dM[i] \&t dM[j], j = 1 \dots (i - 1)\right), i = 1 \dots 5\right)\right) : CK := \text{FromJet}(Ck, \{u(x1, x2, x3, x4, x5)\}) : \\ &\text{> } \text{ToJet}(\text{DirectionalCovariantDerivative}(vM[1], vM[4], CK), \{u(x1, x2, x3, x4, x5)\}, \text{notation} = \text{jetnumbers}) \end{aligned}$$

$$\frac{\alpha_1 u_4 + u_{1,4}}{u_4} D_{x4} \quad (6.12)$$

$$\begin{aligned} &> \{ \text{seq}(\text{evalDG}(\text{convert}(\text{DirectionalCovariantDerivative}(\text{vM}[i], \omega\omega, CK), 'DGform') \\ &\quad \&w \omega\omega), i = 1 \dots 5) \} \\ &\quad \{ (0 \, dx1) \wedge dx2 \} \end{aligned} \quad (6.13)$$

$$\begin{aligned} &> \text{Sconn} := \text{solve}(\text{TGE}, \text{ChrCoe}) : \\ &> \text{FreePar} := \{ \} : \text{for } i \text{ to } \text{nops}(\text{Sconn}) \text{ do if } \text{lhs}(\text{Sconn}[i]) = \text{rhs}(\text{Sconn}[i]) \text{ then } \text{FreePar} \\ &\quad := \{ \text{op}(\text{FreePar}), \text{lhs}(\text{Sconn}[i]) \} \text{ fi:od: } \text{FreePar}, \text{nops}(\%) \\ &\quad \{ a121, a144, a222, a255, a333, a344, a424, a455, a515, a535, a544, a545, a555 \}, 13 \end{aligned} \quad (6.14)$$

$$\begin{aligned} &> \text{ChrCoe1} := \text{ChrCoe} \setminus \text{FreePar}, \text{nops}(\%) \\ \text{ChrCoe1} &:= \{ a111, a112, a113, a114, a115, a122, a123, a124, a125, a131, a132, a133, \\ &\quad a134, a135, a141, a142, a143, a145, a151, a152, a153, a154, a155, a211, a212, a213, \\ &\quad a214, a215, a221, a223, a224, a225, a231, a232, a233, a234, a235, a241, a242, a243, \\ &\quad a244, a245, a251, a252, a253, a254, a311, a312, a313, a314, a315, a321, a322, a323, \\ &\quad a324, a325, a331, a332, a334, a335, a341, a342, a343, a345, a351, a352, a353, a354, \\ &\quad a355, a411, a412, a413, a414, a415, a421, a422, a423, a425, a431, a432, a433, a434, \\ &\quad a435, a441, a442, a443, a444, a445, a451, a452, a453, a454, a511, a512, a513, a514, \\ &\quad a521, a522, a523, a524, a525, a531, a532, a533, a534, a541, a542, a543, a551, a552, \\ &\quad a553, a554 \}, 112 \end{aligned} \quad (6.15)$$

$$\begin{aligned} &> \text{ChrCoe1} := \{ a111, a112, a113, a114, a115, a122, a123, a124, a125, a131, a132, a133, \\ &\quad a134, a135, a141, a142, a143, a145, a151, a152, a153, a154, a155, a211, a212, a213, \\ &\quad a214, a215, a221, a223, a224, a225, a231, a232, a233, a234, a235, a241, a242, a243, \\ &\quad a244, a245, a251, a252, a253, a254, a311, a312, a313, a314, a315, a321, a322, a323, \\ &\quad a324, a325, a331, a332, a334, a335, a341, a342, a343, a345, a351, a352, a353, a354, \\ &\quad a355, a411, a412, a413, a414, a415, a421, a422, a423, a425, a431, a432, a433, a434, \\ &\quad a435, a441, a442, a443, a444, a445, a451, a452, a453, a454, a511, a512, a513, a514, \\ &\quad a521, a522, a523, a524, a525, a531, a532, a533, a534, a541, a542, a543, a551, a552, \\ &\quad a553, a554 \} : \\ \text{FreePar} &:= \{ a121, a144, a222, a255, a333, a344, a424, a455, a515, a535, a544, a545, \\ &\quad a555 \} : \\ &> \text{Sconn} := \text{solve}(\text{TGE}, \text{ChrCoe1}) : \\ &> \text{CC} := \text{DGsimplify}(\text{subs}(\text{Sconn}, C)) : \\ &> \text{Quad} := k \rightarrow \text{evalDG}(\text{add}(\text{add}(\text{diff}(\text{Eq} \| k, u_{i,j}) \cdot \text{vM}[i] \&s \text{vM}[j], j = i \dots 5), i = 1 \dots 5)) : \\ &\quad \text{Quadu} := k \rightarrow \text{FromJet}(\text{Quad}(k), \{ u(x1, x2, x3, x4, x5) \}) : \\ &> \{ \text{seq}(\text{ContractIndices}(\text{Quad}(i), \omega \&t \omega, [[1, 1], [2, 2]]), i = 1 \dots 6) \} \\ &\quad \{ 0 \} \end{aligned} \quad (6.16)$$

$$\begin{aligned} &> \text{RNE} := \{ \} : \text{for } i \text{ to } 5 \text{ do for } j \text{ to } 6 \text{ do } \text{RNE} := \{ \text{op}(\text{RNE}), \\ &\quad \text{coeffs}(\text{expand}(\text{numer}(\text{simplify}(\text{subs}(\text{S2}, \\ &\quad \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(\text{vM}[i], \text{Quadu}(j), \text{CC}), \omega\omega \\ &\quad \&t \omega\omega, [[1, 1], [2, 2]]), \{ u(x1, x2, x3, x4, x5) \}, \text{notation} = \text{jetnumbers}))) \}, \{ q \} \} : \\ &\quad \text{od:od: } \text{nops}(\text{RNE}) \end{aligned} \quad (6.17)$$

$$> \text{sRNE} := \text{solve}(\text{RNE}, \text{FreePar})$$

$$sRNE := \left\{ a_{121} \right. \quad (6.18)$$

$$= \frac{a_{424} a_2 u_1 u_4 - a_{424} a_4 u_1 u_4 - a_1 u_2 u_{1,4} + a_1 u_4 u_{1,2} + a_4 u_2 u_{1,4} - a_4 u_4 u_{1,2}}{u_1 u_4 (a_2 - a_4)},$$

$$a_{144} = 0, a_{222} = \frac{1}{(a_2 - a_4) u_1 u_2 u_4} (a_{424} a_2 u_1 u_2 u_4 - a_{424} a_4 u_1 u_2 u_4 - a_1$$

$$u_2^2 u_{1,4} + a_1 u_2 u_4 u_{1,2} + a_2 u_1 u_4 u_{2,2} - a_2 u_2 u_4 u_{1,2} - a_4 u_1 u_4 u_{2,2} + a_4 u_2^2 u_{1,4}), a_{255}$$

$$= 0, a_{333} = \frac{1}{(a_3 - a_5) u_1 u_3 u_5} (a_{535} a_3 u_1 u_3 u_5 - a_{535} a_5 u_1 u_3 u_5 - a_1 u_3^2 u_{1,5}$$

$$+ a_1 u_3 u_5 u_{1,3} + a_3 u_1 u_5 u_{3,3} - a_3 u_3 u_5 u_{1,3} - a_5 u_1 u_5 u_{3,3} + a_5 u_3^2 u_{1,5}), a_{344} = 0,$$

$$a_{424} = a_{424}, a_{455} = 0, a_{515} = a_{515}, a_{535} = a_{535}, a_{544} = 0, a_{545} = a_{545}, a_{555} = a_{555} \}$$

$$\begin{aligned} &> CT := DGsimplify \left(evalDG \left(subs \left(\left\{ a_{515} = \frac{u_{1,5}}{u_5} + \alpha_1, a_{424} = \frac{u_{2,4}}{u_4} + \alpha_2, a_{535} = \frac{u_{3,5}}{u_5} \right. \right. \right. \right. \\ &\quad \left. \left. \left. + \alpha_3, a_{545} = \frac{u_{4,5}}{u_5} + \alpha_4, a_{555} = \frac{u_{5,5}}{u_5} + \alpha_5 \right\}, subs(sRNE, CC) \right) \right) : \end{aligned}$$

$$\begin{aligned} &> \{ seq(seq(simplify(subs(S2, evalDG(DirectionalCovariantDerivative(vM[i], vM[j], CT) \\ &\quad - DirectionalCovariantDerivative(vM[i], vM[j], Ck))))), i = 1 .. 5), j = 1 .. 5) \} \\ &\quad \{0, 0 D_{x1}, 0 D_{x2}, 0 D_{x3}, 0 D_{x4}, 0 D_{x5} \} \end{aligned} \quad (6.19)$$

$$\begin{aligned} &> simplify \left(subs \left(S2, evalDG \left(convert(DirectionalCovariantDerivative(D_{x1}, D_{x4}, CT), \right. \right. \right. \\ &\quad \left. \left. \left. 'DGvector') - \frac{\alpha_1 u_4 + u_{1,4}}{u_4} \cdot D_{x4} \right) \right) \right), simplify \left(subs \left(S2, \right. \right. \\ &\quad evalDG \left(convert(DirectionalCovariantDerivative(D_{x4}, D_{x4}, CT), 'DGvector') \right. \\ &\quad \left. \left. - \frac{\alpha_4 u_4 + u_{4,4}}{u_4} \cdot D_{x4} \right) \right) \right); \end{aligned}$$

$$0 D_{x1}, 0 D_{x4} \quad (6.20)$$

$$> \# TorsionTensor(CT)$$

$$\begin{aligned} &> sCTa := solve(DGinfo(ContractIndices(TorsionTensor(CT), [[1, 2]]), "CoefficientSet"), \\ &\quad \{ \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5 \}) : CTa := DGsimplify(evalDG(subs(sCTa, CT))) : \end{aligned}$$

$$> simplify \left(subs \left(S2, subs \left(sCTa, \left\{ 4 \alpha_1 + \frac{u_{1,2}}{u_2} + \frac{u_{1,3}}{u_3} + \frac{u_{1,4}}{u_4} + \frac{u_{1,5}}{u_5}, 4 \alpha_2 + \frac{u_{1,2}}{u_1} \right. \right. \right. \right.$$

$$\begin{aligned}
& + \frac{u_{2,3}}{u_3} + \frac{u_{2,4}}{u_4} + \frac{u_{2,5}}{u_5}, 4 \alpha_3 + \frac{u_{1,3}}{u_1} + \frac{u_{2,3}}{u_2} + \frac{u_{3,4}}{u_4} + \frac{u_{3,5}}{u_5}, 4 \alpha_4 + \frac{u_{1,4}}{u_1} + \frac{u_{2,4}}{u_2} \\
& + \frac{u_{3,4}}{u_3} + \frac{u_{4,5}}{u_5}, 4 \alpha_5 + \frac{u_{1,5}}{u_1} + \frac{u_{2,5}}{u_2} + \frac{u_{3,5}}{u_3} + \frac{u_{4,5}}{u_4} \} \} \} \} \} \\
& \{0\}
\end{aligned} \tag{6.21}$$

$$\begin{aligned}
& \text{DirectionalCovariantDerivative}(D_x1, D_x4, CTa); \text{DirectionalCovariantDerivative}(D_x1, \\
& D_x1, CTa) \\
& - \left(\frac{u_{1,5} u_4 u_3 u_2 - 3 u_{1,4} u_5 u_3 u_2 + u_2 u_4 u_5 u_{1,3} + u_3 u_4 u_5 u_{1,2}}{4 u_2 u_5 u_4 u_3} D_x4 \right) \\
& - \left(\frac{u_1 u_2 u_3 u_4 u_{1,5} + u_1 u_2 u_3 u_5 u_{1,4} + u_1 u_2 u_4 u_5 u_{1,3} + u_1 u_3 u_4 u_5 u_{1,2} - 4 u_{1,1} u_2 u_5 u_4 u_3}{4 u_2 u_5 u_4 u_3 u_1} \right. \\
& \left. D_x1 \right)
\end{aligned} \tag{6.22}$$

This gives a totally geodesic connection.

I. Let us recompute it via a different normalization of the connection (without referring to totally geodesic) and check generalized EW property.

>
>
>
>
>
>
>
>

$$\begin{aligned}
& \text{RNE} := \{ \} : \text{for } i \text{ to } 5 \text{ do for } j \text{ to } 6 \text{ do RNE} := \{ \text{op}(\text{RNE}), \\
& \text{coeffs}(\text{expand}(\text{numer}(\text{simplify}(\text{subs}(S2, \\
& \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(vM[i], \text{Quadu}(j), C), \omega \& \omega, \\
& [[1, 1], [2, 2]]), \{u(x1, x2, x3, x4, x5)\}, \text{notation} = \text{jetnumbers})) \} \} : \text{od:od:} \\
& \text{nops}(\text{RNE}) \\
& 210
\end{aligned} \tag{6.23}$$

$$\begin{aligned}
& \text{Sconn} := \text{solve}(\text{RNE}, \text{ChrCoe}) : \\
& \text{FreePar} := \{ \} : \text{for } i \text{ to } \text{nops}(\text{Sconn}) \text{ do if } \text{lhs}(\text{Sconn}[i]) = \text{rhs}(\text{Sconn}[i]) \text{ then FreePar} \\
& := \{ \text{op}(\text{FreePar}), \text{lhs}(\text{Sconn}[i]) \} \text{ fi:od: FreePar, nops}(\%) \\
& \{a222, a225, a232, a235, a444, a452, a455, a511, a512, a514, a515, a523, a525, a534, \\
& a535, a542, a543, a544, a553, a555\}, 20
\end{aligned} \tag{6.24}$$

$$\begin{aligned}
& \text{ChrCoe1} := \text{ChrCoe} \setminus \text{FreePar} \\
& \text{ChrCoe1} := \{a111, a112, a113, a114, a115, a121, a122, a123, a124, a125, a131, a132, \\
& a133, a134, a135, a141, a142, a143, a144, a145, a151, a152, a153, a154, a155, a211, \\
& a212, a213, a214, a215, a221, a223, a224, a231, a233, a234, a241, a242, a243, a244, \\
& a245, a251, a252, a253, a254, a255, a311, a312, a313, a314, a315, a321, a322, a323, \\
& a324, a325, a331, a332, a333, a334, a335, a341, a342, a343, a344, a345, a351, a352, \\
& a353, a354, a355, a411, a412, a413, a414, a415, a421, a422, a423, a424, a425, a431,
\end{aligned} \tag{6.25}$$

$a432, a433, a434, a435, a441, a442, a443, a445, a451, a453, a454, a513, a521, a522, a524, a531, a532, a533, a541, a545, a551, a552, a554\}$

$$\begin{aligned} &> [nops(ChrCoe), nops(FreePar), nops(ChrCoe1)] \\ &\quad [125, 20, 105] \end{aligned} \quad (6.26)$$

$$\begin{aligned} &> FreePar := \{a222, a225, a232, a235, a444, a452, a455, a511, a512, a514, a515, a523, \\ &\quad a525, a534, a535, a542, a543, a544, a553, a555\} : \end{aligned}$$

$$\begin{aligned} &> ChrCoe1 := \{a111, a112, a113, a114, a115, a121, a122, a123, a124, a125, a131, a132, \\ &\quad a133, a134, a135, a141, a142, a143, a144, a145, a151, a152, a153, a154, a155, a211, \\ &\quad a212, a213, a214, a215, a221, a223, a224, a231, a233, a234, a241, a242, a243, a244, \\ &\quad a245, a251, a252, a253, a254, a255, a311, a312, a313, a314, a315, a321, a322, a323, \\ &\quad a324, a325, a331, a332, a333, a334, a335, a341, a342, a343, a344, a345, a351, a352, \\ &\quad a353, a354, a355, a411, a412, a413, a414, a415, a421, a422, a423, a424, a425, a431, \\ &\quad a432, a433, a434, a435, a441, a442, a443, a445, a451, a453, a454, a513, a521, a522, \\ &\quad a524, a531, a532, a533, a541, a545, a551, a552, a554\} : \end{aligned}$$

$$\begin{aligned} &> Sconn := solve(RNE, ChrCoe1) : \end{aligned}$$

$$\begin{aligned} &> CC := DGsimplify(subs(Sconn, C)) : \end{aligned}$$

$$\begin{aligned} &> TorC := TorsionTensor(CC) : \end{aligned}$$

$$\begin{aligned} REN := \{ &seq(seq(coeffs(expand(numer(ToJet(Hook(convert(ContractIndices(DD[i] \\ &\quad \&t DD[j], TorC, [[1, 2], [2, 3]]), 'DGvector'), \omega), \{u(x, y, t, z, s), q(x, y, t, z, s)\}, \\ &\quad notation = jetnumbers))), \{q_{[j]}\}), j = i + 1 .. 4), i = 1 .. 4) \} : nops(REN) \end{aligned}$$

38

(6.27)

$$\begin{aligned} &> tmp := eliminate(REN \text{ union } DGinfo(ContractIndices(TorsionTensor(CC), [[1, 2]]), \\ &\quad "CoefficientSet"), FreePar) \end{aligned}$$

$$tmp := \left[\left\{ a413 = 0, a414 = 0, a415 = 0, a423 = \frac{9}{4} u_{1,1}, a424 = 0, a425 = 0, a433 = 3 u_{1,2}, \right. \right. \quad (6.28)$$

$$a434 = u_{1,1}, a435 = 0, a443 = \frac{15}{4} u_{1,1} + 3 u_{2,2}, a444 = \frac{5}{2} u_{1,2}, a445 = -\frac{1}{4} u_{1,1},$$

$$a453 = 3 u_{1,1} + \frac{9}{2} u_{2,1} + 3 u_{2,3}, a454 = 3 u_{1,1} + 4 u_{2,2}, a455 = -u_{1,2}, a515$$

$$= 0, a525 = 0, a535 = \frac{3}{2} u_{1,1}, a545 = \frac{7}{2} u_{1,2}, a555 = \frac{9}{2} u_{1,1} + 5 u_{2,2} \Big\}, \{ \} \Big]$$

$$\begin{aligned} &> Sconn2 := tmp[1] : CC := DGsimplify(subs(Sconn2, CC)) : \end{aligned}$$

$$\begin{aligned} &> \{seq(seq(coeffs(expand(numer(simplify(subs(S2, subs(Qsyst, \\ &\quad ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CC), \omega \\ &\quad \&t \omega, [[1, 1], [2, 2]]), \{u(x, y, t, z, s)\}, notation = jetnumbers))))), \{q_{[j]}\}), i = 1 \\ &\quad .. 5), j = 1 .. 6) \} \text{ union } DGinfo(ContractIndices(TorsionTensor(CC), [[1, 2]]), \\ &\quad "CoefficientSet") \end{aligned}$$

{0}

(6.29)

$$\begin{aligned} &> coeffs(expand(subs(S2, subs(Qsyst, \\ &\quad ToJet(Hook(convert(DirectionalCovariantDerivative(DD[3], DD[1], CC), 'DGvector \\ &\quad '), \omega\omega), \{u(x, y, t, z, s), q(x, y, t, z, s)\}, notation = jetnumbers))))), \{q_{[1]}, q_1\}) \end{aligned}$$

$$-\frac{1}{2} u_{1,1} - \frac{3}{4} u_{1,1}$$

(6.30)

```

> RicCC := RicciTensor(FromJet(CC, {u(x, y, t, z, s)})) : RicCc := DGsimplify(subs(S3,
  ToJet(RicCC, {u(x, y, t, z, s)}, notation = jetnumbers))) :
> RicCcSkew := SymmetrizeIndices(RicCc, [1, 2], "SkewSymmetric") :
  RicCcSymm := SymmetrizeIndices(RicCc, [1, 2], "Symmetric") :
> ω
dx + q[ ] dy + (q[ ]^2 + u1) dt + (q[ ]^3 + 2 q[ ] u1 + u2) dz + (q[ ]^4 + 3 q[ ]^2 u1
  + 2 q[ ] u2 + u12 + u3) ds

```

(6.31)

```

> ω0 := ω : ω0b := DGsimplify(evalDG(subs(q[ ] = 0, ω)));
ω1a := evalDG(1/ε · (subs(q[ ] = q[ ] + ε, ω) - ω)) : ω1 := evalDG(subs(ε = 0, ω1a)) :
  ω1b := DGsimplify(evalDG(subs(q[ ] = 0, ω1)));
ω2a := evalDG(1/ε · (subs(q[ ] = q[ ] + ε, ω1) - ω1)) : ω2 := evalDG(subs(ε = 0,
  ω2a)) : ω2b := DGsimplify(evalDG(1/2 · subs(q[ ] = 0, ω2)));
ω3a := evalDG(1/ε · (subs(q[ ] = q[ ] + ε, ω2) - ω2)) : ω3 := evalDG(subs(ε = 0,
  ω3a)) : ω3b := DGsimplify(evalDG(1/6 · subs(q[ ] = 0, ω3)));
ω4a := evalDG(1/ε · (subs(q[ ] = q[ ] + ε, ω3) - ω3)) : ω4 := evalDG(subs(ε = 0,
  ω4a)) : ω4b := DGsimplify(evalDG(1/24 · subs(q[ ] = 0, ω4)));
ω0b := dx + u1 dt + u2 dz + (u12 + u3) ds
ω1b := dy + 2 u1 dz + 2 u2 ds
ω2b := dt + 3 u1 ds
ω3b := dz
ω4b := ds

```

(6.32)

```

> evalDG(ω - ω0b - ω1b · q[ ] - ω2b · q[ ]2 - ω3b · q[ ]3 - ω4b · q[ ]4)
  0 dx

```

(6.33)

```

> Quaω := evalDG([ω0b &s ω0b, ω0b &s ω1b, 2 ω0b &s ω2b + ω1b &s ω1b, ω0b &s ω3b
  + ω1b &s ω2b, 2 ω0b &s ω4b + 2 ω1b &s ω3b + ω2b &s ω2b, ω1b &s ω4b + ω2b
  &s ω3b, 2 ω2b &s ω4b + ω3b &s ω3b, ω3b &s ω4b, ω4b &s ω4b]) : tmpΩ := { } :for i
  to nops(Quaω) do for j to 6 do tmpΩ := {op(tmpΩ), ContractIndices(Quad|| j,
  Quaω[i], [[1, 1], [2, 2]])} :od:od: tmpΩ
  {0}

```

(6.34)

```

> eliminate(simplify(subs(S3, DGinfo(evalDG(RicCcSymm - add(Λ[i] · Quaω[i], i = 1
  ..nops(Quaω))), "CoefficientSet")), {seq(Λ[i], i = 1 ..nops(Quaω))})[2]
{u1, 1, 1, 6 u1, 12 + 7 u1, 2, 2, 28 u1 u1, 1 + 3 u1, 12 - 14 u1, 2, 2, 3 u15 u1, 1 + 12 u14 u1, 12 + 6

```

(6.35)

$$\begin{aligned}
& u_1^4 u_{1,2,2} + 6 u_1^3 u_3 u_{1,1,1} + 17 u_1^3 u_{1,1} u_{2,2} + 6 u_1^3 u_{1,2}^2 - u_1^2 u_2^2 u_{1,1,1} + 10 u_1^2 u_2 u_{1,1} u_{1,2} \\
& + 19 u_1^2 u_3 u_{1,1}^2 - 4 u_1 u_2^2 u_{1,1}^2 + 3 u_1^3 u_{2,2,3} + 6 u_1^2 u_3 u_{1,2,2} + 7 u_1^2 u_{1,1} u_{3,3} + 2 u_1^2 u_{2,2}^2 \\
& - 2 u_1 u_2^2 u_{1,2,2} + 3 u_1 u_3^2 u_{1,1,1} + 16 u_1 u_3 u_{1,1} u_{2,2} + 6 u_1 u_3 u_{1,2}^2 - u_2^2 u_3 u_{1,1,1} - 6 \\
& u_2^2 u_{1,1} u_{2,2} - 6 u_2^2 u_{1,2}^2 - 2 u_2 u_3 u_{1,1} u_{1,2} + u_3^2 u_{1,1}^2 + 2 u_1^2 u_{2,2,5} + 3 u_1 u_3 u_{2,2,3} \\
& + 3 u_1 u_{1,1} u_{4,4} + 2 u_1 u_{2,2} u_{3,3} - u_2^2 u_{2,2,3} - 2 u_2 u_{1,1} u_{3,4} - 2 u_2 u_{1,2} u_{3,3} \\
& + u_3 u_{1,1} u_{3,3} + u_3 u_{2,2}^2 - 2 u_1 u_{1,5,5} + 2 u_1 u_{2,4,5} + u_3 u_{2,2,5} + u_{2,2} u_{4,4} - u_{3,5,5} \\
& + u_{4,4,5}, 126 u_1^5 u_{1,1,1} + 258 u_1^4 u_{1,1}^2 + 63 u_1^4 u_{1,2,2} + 168 u_1^3 u_3 u_{1,1,1} + 238 u_1^3 u_{1,1} u_{2,2} \\
& + 126 u_1^3 u_{1,2}^2 - 42 u_1^2 u_2^2 u_{1,1,1} + 140 u_1^2 u_2 u_{1,1} u_{1,2} + 356 u_1^2 u_3 u_{1,1}^2 - 86 u_1 u_2^2 u_{1,1}^2 \\
& + 63 u_1^3 u_{2,2,3} + 63 u_1^2 u_3 u_{1,2,2} + 98 u_1^2 u_{1,1} u_{3,3} + 28 u_1^2 u_{2,2}^2 - 21 u_1 u_2^2 u_{1,2,2} + 42 u_1 \\
& u_3^2 u_{1,1,1} + 224 u_1 u_3 u_{1,1} u_{2,2} + 126 u_1 u_3 u_{1,2}^2 - 14 u_2^2 u_3 u_{1,1,1} - 84 u_2^2 u_{1,1} u_{2,2} - 98 \\
& u_2^2 u_{1,2}^2 - 28 u_2 u_3 u_{1,1} u_{1,2} + 14 u_3^2 u_{1,1}^2 + 28 u_1^2 u_{2,2,5} + 63 u_1 u_3 u_{2,2,3} + 42 u_1 u_{1,1} u_{4,4} \\
& + 28 u_1 u_{2,2} u_{3,3} - 21 u_2^2 u_{2,2,3} - 28 u_2 u_{1,1} u_{3,4} - 28 u_2 u_{1,2} u_{3,3} + 14 u_3 u_{1,1} u_{3,3} \\
& + 14 u_3 u_{2,2}^2 - 28 u_1 u_{1,5,5} + 28 u_1 u_{2,4,5} + 14 u_3 u_{2,2,5} + 14 u_{2,2} u_{4,4} - 14 u_{3,5,5} \\
& + 14 u_{4,4,5}, 3 u_1^6 u_{1,1,1} + 12 u_1^5 u_{1,1}^2 + 6 u_1^5 u_{1,2,2} + 6 u_1^4 u_3 u_{1,1,1} + 17 u_1^4 u_{1,1} u_{2,2} + 6 \\
& u_1^4 u_{1,2}^2 - 7 u_1^3 u_2^2 u_{1,1,1} + 19 u_1^3 u_2 u_{1,1} u_{1,2} + 19 u_1^3 u_3 u_{1,1}^2 - 4 u_1^2 u_2^2 u_{1,1}^2 + 3 u_1^4 u_{2,2,3} \\
& + 6 u_1^3 u_2 u_{2,2,2} + 6 u_1^3 u_3 u_{1,2,2} + 7 u_1^3 u_{1,1} u_{3,3} + 2 u_1^3 u_{2,2}^2 - 2 u_1^2 u_2^2 u_{1,2,2} + 3 u_1^2 \\
& u_3^2 u_{1,1,1} + 16 u_1^2 u_3 u_{1,1} u_{2,2} + 6 u_1^2 u_3 u_{1,2}^2 - 7 u_1 u_2^2 u_3 u_{1,1,1} - 6 u_1 u_2^2 u_{1,1} u_{2,2} - 6 u_1 \\
& u_2^2 u_{1,2}^2 + 7 u_1 u_2 u_3 u_{1,1} u_{1,2} + u_1 u_3^2 u_{1,1}^2 + 2 u_2^4 u_{1,1,1} - 3 u_2^3 u_{1,1} u_{1,2} + 2 u_1^3 u_{2,2,5} + 3 \\
& u_1^2 u_3 u_{2,2,3} + 3 u_1^2 u_{1,1} u_{4,4} + 2 u_1^2 u_{2,2} u_{3,3} - u_1 u_2^2 u_{2,2,3} + 6 u_1 u_2 u_3 u_{2,2,2} \\
& - 2 u_1 u_2 u_{1,1} u_{3,4} - 2 u_1 u_2 u_{1,2} u_{3,3} + u_1 u_3 u_{1,1} u_{3,3} + u_1 u_3 u_{2,2}^2 - 2 u_2^3 u_{2,2,2} - 2 \\
& u_1^2 u_{1,5,5} + 2 u_1^2 u_{2,4,5} + u_1 u_3 u_{2,2,5} + u_1 u_{2,2} u_{4,4} - u_1 u_{3,5,5} + u_1 u_{4,4,5} \}
\end{aligned}$$

Thus the condition that $\text{sym}(\text{Ric}(\nabla))$ belongs to the span of quadrics defining the GL2 structure is non-vacous. It is another differential constraint, giving exact solutions.

$$\begin{aligned}
& \triangleright \text{Vt0} := \text{Annihilator}([\omega 0, \omega 1, \omega 2, \omega 3])[1]; \text{Vt1} := \text{Annihilator}([\omega 0, \omega 1, \omega 2])[2]; \text{Vt2} \\
& \quad := \text{Annihilator}([\omega 0, \omega 1])[3]; \text{Vt3} := \text{Annihilator}([\omega 0])[4]; \\
& \text{Vt01} := \text{ContractIndices}(\text{TorsionTensor}(\text{CC}), \text{evalDG}(\text{Vt0} \& \text{Vt1}), [[2, 1], [3, 2]]): \\
& \text{Vt0} := (q[]^4 - 6 q[]^2 u_1 + 4 q[] u_2 + 2 u_1^2 - u_3) D_x - (4 q[]^3 - 8 q[] u_1 \\
& \quad + 2 u_2) D_y + (6 q[]^2 - 3 u_1) D_t - (4 q[] D_z) + D_s \\
& \text{Vt1} := -((q[]^3 - 3 q[] u_1 + u_2) D_x) + (3 q[]^2 - 2 u_1) D_y - (3 q[] D_t) + D_z \\
& \text{Vt2} := (3 q[]^4 + 3 q[]^2 u_1 - u_1^2 - u_3) D_x - ((4 q[]^3 + 6 q[] u_1 + 2 u_2) D_y) + D_s \\
& \text{Vt3} := -(q[] D_x) + D_y
\end{aligned} \tag{6.36}$$

Thus there exist GL2-connections, with respect to which the null Lax surfaces are totally geodesic.

The 4 parameters can be fixed uniquely by the condition $\text{Trace}(\text{Torsion})=0$.

$$\begin{aligned} & \text{seq}(\text{ContractIndices}(\text{Qua}\omega[i], \text{evalDG}(\text{Vt0} \& \text{Vt0}), [[1, 1], [2, 2]]), i = 1 \dots \text{nops}(\text{Qua}\omega)) \\ & \quad q_{[]}^8, -4 q_{[]}^7, 28 q_{[]}^6, -28 q_{[]}^5, 70 q_{[]}^4, -28 q_{[]}^3, 28 q_{[]}^2, -4 q_{[]}, 1 \end{aligned} \quad (6.37)$$

$$\begin{aligned} & \text{QuT} := \text{evalDG}(\text{add}(\text{add}(\text{xi}[i, j] \cdot dM[i] \& s dM[j], j = i \dots 5), i = 1 \dots 5)) : \\ & \text{DGsimplify}(\text{evalDG}(\text{subs}(\{\text{xi}[2, 2] = \Lambda_1, \text{xi}[2, 3] = \Lambda_2, \text{xi}[2, 4] = \Lambda_3, \text{xi}[3, 3] = \Lambda_4, \text{xi}[3, 4] \\ & \quad = \Lambda_5, \text{xi}[4, 4] = \Lambda_6\}, \text{subs}(\text{solve}(\{\text{coeffs}(\text{ContractIndices}(\text{QuT}, \text{Vt0} \& \text{Vt0}), [[1, 1], [2, \\ & \quad 2]]), \{q_{[]}\}\}, \{\text{seq}(\text{seq}(\text{xi}[i, j], j = i \dots 5), i = 1 \dots 5)\}), \text{QuT}))) \end{aligned}$$

$$- \left(\left(\frac{4 \Lambda_1}{3} dx \right) dt \right) - \left((3 \Lambda_2 dx) dz \right) - \left(((20 \Lambda_1 u_1 + 8 \Lambda_3 + 18 \Lambda_4) dx) ds \right) \quad (6.38)$$

$$\begin{aligned} & + (\Lambda_1 dy) dy + \left(\frac{\Lambda_2}{2} dy \right) dt + \left(\frac{\Lambda_3}{2} dy \right) dz - \left(\left(\frac{21 \Lambda_2 u_1}{2} + 6 \Lambda_1 u_2 \right. \right. \\ & \left. \left. + 3 \Lambda_5 \right) dy \right) ds \Big) - \left(\left(\frac{4 \Lambda_1}{3} dt \right) dx \right) + \left(\frac{\Lambda_2}{2} dt \right) dy + (\Lambda_4 dt) dt + \left(\frac{\Lambda_5}{2} dt \right) dz \\ & - \left(\left(\left(\frac{56}{3} \Lambda_1 u_1^2 + \frac{4}{3} \Lambda_1 u_3 + \frac{16}{3} \Lambda_3 u_1 + 15 u_1 \Lambda_4 + 7 \Lambda_2 u_2 + \frac{4}{3} \Lambda_6 \right) dt \right) ds \right) \\ & - \left((3 \Lambda_2 dz) dx \right) + \left(\frac{\Lambda_3}{2} dz \right) dy + \left(\frac{\Lambda_5}{2} dz \right) dt + (\Lambda_6 dz) dz - \left(\left(\left(18 \Lambda_2 u_1^2 \right. \right. \right. \\ & \left. \left. + 32 \Lambda_1 u_1 u_2 + \frac{9}{2} u_1 \Lambda_5 + 7 \Lambda_3 u_2 + 18 u_2 \Lambda_4 + 3 u_3 \Lambda_2 \right) dz \right) ds \Big) - \left(((20 \Lambda_1 u_1 \right. \\ & \left. + 8 \Lambda_3 + 18 \Lambda_4) ds) dx \right) - \left(\left(\left(\frac{21 \Lambda_2 u_1}{2} + 6 \Lambda_1 u_2 + 3 \Lambda_5 \right) ds \right) dy \right) - \left(\left(\left(\frac{56}{3} \Lambda_1 \right. \right. \right. \\ & \left. \left. u_1^2 + \frac{4}{3} \Lambda_1 u_3 + \frac{16}{3} \Lambda_3 u_1 + 15 u_1 \Lambda_4 + 7 \Lambda_2 u_2 + \frac{4}{3} \Lambda_6 \right) ds \right) dt \Big) - \left(\left(\left(18 \Lambda_2 u_1^2 \right. \right. \right. \\ & \left. \left. + 32 \Lambda_1 u_1 u_2 + \frac{9}{2} u_1 \Lambda_5 + 7 \Lambda_3 u_2 + 18 u_2 \Lambda_4 + 3 u_3 \Lambda_2 \right) ds \right) dz \Big) - \left(((48 \Lambda_1 u_1^3 \right. \\ & \left. + 40 \Lambda_1 u_1 u_3 + 28 \Lambda_1 u_2^2 + 90 \Lambda_2 u_1 u_2 + 27 \Lambda_4 u_1^2 + 16 \Lambda_3 u_3 + 36 \Lambda_4 u_3 + 12 \Lambda_5 u_2 \right. \\ & \left. + 8 \Lambda_6 u_1) ds) ds \right) \end{aligned}$$

$$\begin{aligned} & \text{eliminate}(\text{DGinfo}(\text{DGsimplify}(\text{evalDG}(\text{subs}(\text{S3}, \text{subs}(\{\text{xi}[2, 2] = \Lambda_1, \text{xi}[2, 3] = \Lambda_2, \text{xi}[2, 4] \\ & \quad = \Lambda_3, \text{xi}[3, 3] = \Lambda_4, \text{xi}[3, 4] = \Lambda_5, \text{xi}[4, 4] = \Lambda_6\}, \\ & \quad \text{subs}(\text{solve}(\{\text{coeffs}(\text{ContractIndices}(\text{QuT}, \text{Vt0} \& \text{Vt0}), [[1, 1], [2, 2]]), \{q_{[]}\}\}, \\ & \quad \{\text{seq}(\text{seq}(\text{xi}[i, j], j = i \dots 5), i = 1 \dots 5)\}), \text{QuT})) - \text{RicCcSymm})), \text{"CoefficientSet"}), \\ & \quad \{\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4, \Lambda_5, \Lambda_6\}) \end{aligned}$$

$$\left[\left\{ \Lambda_1 = \frac{3}{4} u_{1,1,1}, \Lambda_2 = -\frac{1}{2} \frac{1}{(3 u_1^3 + 3 u_1 u_3 - u_2^2) u_2} (3 u_1^5 u_{1,1,1} + 12 u_1^4 u_{1,1}^2 + 6 u_1^4 u_{1,2,2} \right. \right. \quad (6.39)$$

$$\left. + 6 u_1^3 u_3 u_{1,1,1} + 17 u_1^3 u_{1,1} u_{2,2} + 6 u_1^3 u_{1,2}^2 - u_1^2 u_2^2 u_{1,1,1} + 10 u_1^2 u_2 u_{1,1} u_{1,2} + 19 \right.$$

$$u_1^2 u_3 u_{1,1}^2 - 4 u_1 u_2^2 u_{1,1}^2 + 3 u_1^3 u_{2,2,3} + 6 u_1^2 u_3 u_{1,2,2} + 7 u_1^2 u_{1,1} u_{3,3} + 2 u_1^2 u_{2,2}^2 - 2 u_1$$

$$u_2^2 u_{1,2,2} + 3 u_1 u_3^2 u_{1,1,1} + 16 u_1 u_3 u_{1,1} u_{2,2} + 6 u_1 u_3 u_{1,2}^2 - u_2^2 u_3 u_{1,1,1} - 6$$

$$u_2^2 u_{1,1} u_{2,2} - 6 u_2^2 u_{1,2}^2 - 2 u_2 u_3 u_{1,1} u_{1,2} + u_3^2 u_{1,1}^2 + 2 u_1^2 u_{2,2,5} + 3 u_1 u_3 u_{2,2,3}$$

$$+ 3 u_1 u_{1,1} u_{4,4} + 2 u_1 u_{2,2} u_{3,3} - u_2^2 u_{2,2,3} - 2 u_2 u_{1,1} u_{3,4} - 2 u_2 u_{1,2} u_{3,3}$$

$$+ u_3 u_{1,1} u_{3,3} + u_3 u_{2,2}^2 - 2 u_1 u_{1,5,5} + 2 u_1 u_{2,4,5} + u_3 u_{2,2,5} + u_{2,2} u_{4,4} - u_{3,5,5}$$

$$+ u_{4,4,5}), \Lambda_3 = 3 u_1 u_{1,1,1} + \frac{3}{4} u_{1,1}^2 - \frac{3}{2} u_{1,2,2}, \Lambda_4 = -2 u_1 u_{1,1,1} + u_{1,2,2}, \Lambda_5$$

$$= \frac{1}{2} \frac{1}{(3 u_1^3 + 3 u_1 u_3 - u_2^2) u_2} (12 u_1^6 u_{1,1,1} + 48 u_1^5 u_{1,1}^2 + 24 u_1^5 u_{1,2,2} + 24$$

$$u_1^4 u_3 u_{1,1,1} + 68 u_1^4 u_{1,1} u_{2,2} + 24 u_1^4 u_{1,2}^2 - 16 u_1^3 u_2^2 u_{1,1,1} + 49 u_1^3 u_2 u_{1,1} u_{1,2} + 76$$

$$u_1^3 u_3 u_{1,1}^2 - 16 u_1^2 u_2^2 u_{1,1}^2 + 12 u_1^4 u_{2,2,3} + 6 u_1^3 u_2 u_{2,2,2} + 24 u_1^3 u_3 u_{1,2,2} + 28$$

$$u_1^3 u_{1,1} u_{3,3} + 8 u_1^3 u_{2,2}^2 - 8 u_1^2 u_2^2 u_{1,2,2} + 12 u_1^2 u_3^2 u_{1,1,1} + 64 u_1^2 u_3 u_{1,1} u_{2,2} + 24 u_1^2 u_3$$

$$u_{1,2}^2 - 16 u_1 u_2^2 u_3 u_{1,1,1} - 24 u_1 u_2^2 u_{1,1} u_{2,2} - 24 u_1 u_2^2 u_{1,2}^2 + u_1 u_2 u_3 u_{1,1} u_{1,2} + 4 u_1$$

$$u_3^2 u_{1,1}^2 + 4 u_2^4 u_{1,1,1} - 3 u_2^3 u_{1,1} u_{1,2} + 8 u_1^3 u_{2,2,5} + 12 u_1^2 u_3 u_{2,2,3} + 12 u_1^2 u_{1,1} u_{4,4}$$

$$+ 8 u_1^2 u_{2,2} u_{3,3} - 4 u_1 u_2^2 u_{2,2,3} + 6 u_1 u_2 u_3 u_{2,2,2} - 8 u_1 u_2 u_{1,1} u_{3,4} - 8 u_1 u_2 u_{1,2} u_{3,3}$$

$$+ 4 u_1 u_3 u_{1,1} u_{3,3} + 4 u_1 u_3 u_{2,2}^2 - 2 u_2^3 u_{2,2,2} - 8 u_1^2 u_{1,5,5} + 8 u_1^2 u_{2,4,5}$$

$$+ 4 u_1 u_3 u_{2,2,5} + 4 u_1 u_{2,2} u_{4,4} - 4 u_1 u_{3,5,5} + 4 u_1 u_{4,4,5}), \Lambda_6$$

$$\begin{aligned}
&= \frac{3}{4} \frac{1}{3 u_1^3 + 3 u_1 u_3 - u_2^2} (24 u_1^5 u_{1,1,1} + 54 u_1^4 u_{1,1}^2 + 9 u_1^4 u_{1,2,2} + 36 u_1^3 u_3 u_{1,1,1} \\
&+ 68 u_1^3 u_{1,1} u_{2,2} + 30 u_1^3 u_{1,2}^2 - 8 u_1^2 u_{2,1,1}^2 + 40 u_1^2 u_2 u_{1,1} u_{1,2} + 82 u_1^2 u_3 u_{1,1}^2 \\
&- 18 u_1 u_2^2 u_{1,1}^2 + 15 u_1^3 u_{2,2,3} + 9 u_1^2 u_3 u_{1,2,2} + 28 u_1^2 u_{1,1} u_{3,3} + 8 u_1^2 u_{2,2}^2 - 3 u_1 \\
&u_2^2 u_{1,2,2} + 12 u_1 u_3^2 u_{1,1,1} + 64 u_1 u_3 u_{1,1} u_{2,2} + 30 u_1 u_3 u_{1,2}^2 - 4 u_2^2 u_3 u_{1,1,1} - 24 \\
&u_2^2 u_{1,1} u_{2,2} - 26 u_2^2 u_{1,2}^2 - 8 u_2 u_3 u_{1,1} u_{1,2} + 4 u_3^2 u_{1,1}^2 + 8 u_1^2 u_{2,2,5} + 15 u_1 u_3 u_{2,2,3} \\
&+ 12 u_1 u_{1,1} u_{4,4} + 8 u_1 u_{2,2} u_{3,3} - 5 u_2^2 u_{2,2,3} - 8 u_2 u_{1,1} u_{3,4} - 8 u_2 u_{1,2} u_{3,3} \\
&+ 4 u_3 u_{1,1} u_{3,3} + 4 u_3 u_{2,2}^2 - 8 u_1 u_{1,5,5} + 8 u_1 u_{2,4,5} + 4 u_3 u_{2,2,5} + 4 u_{2,2} u_{4,4} \\
&- 4 u_{3,5,5} + 4 u_{4,4,5}) \}, \{ \}]
\end{aligned}$$

This proves that symmetrized Ricci tensor of the canonical GL2-connection is a section of the bundle of quadrics annihilating the dual (tangent) to the canonical (cotangent) rational normal curve!
Here is the connection:

> CC

(6.40)

>
>
>
>

II. Let us check if there exists a unique symmetric connection preserving GL2-structure

> ChrSeq := {seq(seq(seq(((a||i)||j)||k), k=1..5), j=i..5), i=1..5)}: nops(ChrSeq);
CsC := Connection(subs({seq(seq(seq(((a||i)||j)||k)=(((a||j)||i)||k), k=1..5), j=1..(i-1)), i=1..5}), add(add(add(((a||i)||j)||k)·vM[k]&t dM[i]&t dM[j], k=1..5), j=1..5), i=1..5)): TorsionTensor(CsC)

75

((0 D_x) dx) dx

(6.41)

> RNE := { } :for i to 5 do for j to 6 do RNE := {op(RNE),
coeffs(expand(numer(simplify(subs(S2, subs(Qsyst,
ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CsC), ω
&t ω, [[1, 1], [2, 2]]), {u(x, y, t, z, s)}, notation = jetnumbers)))))}, {q[]})} :od:od:

> nops(RNE)

227

(6.42)

> elRN := eliminate(RNE, ChrSeq) :

> elRN[2]

$$\begin{aligned}
&\{u_{1,1}, u_{1,2}, u_1 u_{1,1}, u_1 u_{1,2}, u_1 (3 u_1 u_{1,2} - 2 u_2 u_{1,1}), u_1 (3 u_1 u_{1,2} + u_2 u_{1,1}), u_1 (6 u_1 u_{1,2} \\
&- u_2 u_{1,1}), u_2 u_{1,1}, u_2 u_{1,2}, u_3 u_{1,1}, u_{1,1} (4 u_1^2 + u_3), u_1 u_2 u_{1,1}, u_1 u_3 u_{1,1}, u_1 u_{1,1} (3 u_1^2 \\
&+ u_3), -2 u_1 u_3 u_{1,1}, 2 u_1 u_3 u_{1,1}, 3 u_1 u_{1,2} + u_2 u_{1,1}, u_1^2 u_{1,1} - 3 u_2 u_{1,2}, u_1^2 u_{1,1} \\
&+ 3 u_2 u_{1,2}, u_1^3 u_{1,1} + 3 u_1 u_2 u_{1,2} - u_2^2 u_{1,1}, 12 u_1^2 u_{1,2} - u_1 u_2 u_{1,1} + 6 u_3 u_{1,2}, 6 u_1^3 u_{1,2} \\
&- u_1^2 u_2 u_{1,1} + u_2 u_3 u_{1,1}, 12 u_1^2 u_2 u_{1,2} - 3 u_1^2 u_3 u_{1,1} - 2 u_1 u_2^2 u_{1,1} - u_3^2 u_{1,1}, 6 u_1^3 u_{1,2}
\end{aligned}$$

(6.43)

$$+ 5 u_1^2 u_2 u_{1,1} + 6 u_2^2 u_{1,2} + u_2 u_3 u_{1,1}, 3 u_1^3 u_{1,2} - u_1^2 u_2 u_{1,1} - 3 u_1 u_3 u_{1,2} + u_2 u_3 u_{1,1} \}$$

$$\begin{aligned} & \text{> simplify(subs(\{u_{1,1}=0\}, \{u_{1,1}, u_{1,2}, u_1 u_{1,1}, u_1 u_{1,2}, u_1 (3 u_1 u_{1,2} - 2 u_2 u_{1,1}), u_1 (3 u_1 u_{1,2} \\ & + u_2 u_{1,1}), u_1 (6 u_1 u_{1,2} - u_2 u_{1,1}), u_2 u_{1,1}, u_2 u_{1,2}, u_3 u_{1,1}, u_{1,1} (4 u_1^2 + u_3), u_1 u_2 u_{1,1}, \\ & u_1 u_3 u_{1,1}, u_1 u_{1,1} (3 u_1^2 + u_3), -2 u_1 u_3 u_{1,1}, 2 u_1 u_3 u_{1,1}, 3 u_1 u_{1,2} + u_2 u_{1,1}, u_1^2 u_{1,1} \\ & - 3 u_2 u_{1,2}, u_1^2 u_{1,1} + 3 u_2 u_{1,2}, u_1^3 u_{1,1} + 3 u_1 u_2 u_{1,2} - u_2^2 u_{1,1}, 12 u_1^2 u_{1,2} - u_1 u_2 u_{1,1} \\ & + 6 u_3 u_{1,2}, 6 u_1^3 u_{1,2} - u_1^2 u_2 u_{1,1} + u_2 u_3 u_{1,1}, 12 u_1^2 u_2 u_{1,2} - 3 u_1^2 u_3 u_{1,1} - 2 u_1 u_2^2 u_{1,1} \\ & - u_3^2 u_{1,1}, 6 u_1^3 u_{1,2} + 5 u_1^2 u_2 u_{1,1} + 6 u_2^2 u_{1,2} + u_2 u_3 u_{1,1}, 3 u_1^3 u_{1,2} - u_1^2 u_2 u_{1,1} \\ & - 3 u_1 u_3 u_{1,2} + u_2 u_3 u_{1,1}\})) \\ & \{0, u_{1,2}, u_1 u_{1,2}, u_2 u_{1,2}, 3 u_1 u_{1,2}, 3 u_1^2 u_{1,2}, 6 u_1^3 u_{1,2}, 6 u_1^3 u_{1,2}, -3 u_2 u_{1,2}, 3 u_2 u_{1,2}, \\ & 3 u_1 u_2 u_{1,2}, 12 u_1^2 u_2 u_{1,2}, 6 u_1^3 u_{1,2} + 6 u_2^2 u_{1,2}, 12 u_1^2 u_{1,2} + 6 u_3 u_{1,2}, 3 u_1^3 u_{1,2} \\ & - 3 u_1 u_3 u_{1,2}\} \end{aligned} \quad (6.44)$$

$$\begin{aligned} & \text{> simplify(subs(\{u_{1,1}=0\}, \{u_{1,1}, u_1 u_{1,1}, u_2 u_{1,1}, (u_1^2 + u_3) u_{1,1}, u_1 u_2 u_{1,1}, u_1 (u_1^2 + u_3) u_{1,1}, \\ & u_2 (u_1^2 + u_3) u_{1,1}, -8 u_1 u_{1,1}, -7 u_1 u_{1,1}, -6 u_1 u_{1,1}, -5 u_1 u_{1,1}, -3 u_1 u_{1,1}, -u_1 u_{1,1}, -5 \\ & u_1^2 u_{1,1}, -u_1^2 u_{1,1}, -5 u_1 u_2 u_{1,1}, -u_1 u_{1,1} (6 u_1^2 + u_3), -5 u_1^2 u_2 u_{1,1}, -u_2 u_{1,1} (6 u_1^2 + u_3), \\ & -16 u_1^2 u_{1,1} - u_3 u_{1,1}, -15 u_1^2 u_{1,1} - u_3 u_{1,1}, -12 u_1^2 u_{1,1} - u_3 u_{1,1}, -9 u_1^2 u_{1,1} - u_3 u_{1,1}, \\ & -6 u_1^2 u_{1,1} - u_3 u_{1,1}, -5 u_1^3 u_{1,1} - 5 u_1 u_3 u_{1,1}, -6 u_1^4 u_{1,1} - 7 u_1^2 u_3 u_{1,1} - u_3^2 u_{1,1}, -21 \\ & u_1^3 u_{1,1} - 6 u_1 u_3 u_{1,1} + 4 u_2^2 u_{1,1}, -6 u_1^3 u_{1,1} - u_1 u_3 u_{1,1} + u_2^2 u_{1,1}, -u_1^3 u_{1,1} - u_1 u_3 u_{1,1} \\ & + 2 u_2^2 u_{1,1}\})) \\ & \{0\} \end{aligned} \quad (6.45)$$

$$\begin{aligned} & \text{> Sss1 := solve(RNE, ChrSeq)} \\ & \text{Sss1 :=} \end{aligned} \quad (6.46)$$

So it does not exist in 5D (for dKP) as expected ...

>

III. Finally let us compute the canonical projective totally geodesic connection

$$\begin{aligned} & \text{> CsC := Connection(subs(\{seq(seq(seq(((a||i)||j)||k)=(((a||j)||i)||k), k=1..5), j=1 \\ & ..(i-1)), i=1..5\}), add(add(add(((a||i)||j)||k) \cdot vM[k] \&t dM[i] \&t dM[j], k=1 \\ & ..5), j=1..5), i=1..5))) \\ & \text{> DD := \{YY, ZZ, TT, SS\} : ChrCoe := \{seq(seq(seq(((a||i)||j)||k), k=1..5), j=i..5), i \\ & =1..5\} : \\ & \text{TGE := \{ \} : for i to 4 do for j to 4 do TGE := \{op(TGE), coeffs(expand(subs(S2, \\ & subs(Qsyst, ToJet(Hook(convert(DirectionalCovariantDerivative(DD[i], DD[j], CsC), \\ & 'DGvector'), \omega\omega), \{u(x,y,t,z,s), q(x,y,t,z,s)\}, notation=jetnumbers))\}), \{q_{[}, q_1\}) \\ & \} : od:od: \\ & [nops(ChrCoe), nops(TGE)] \end{aligned} \quad [75, 84] \quad (6.47)$$

$$\begin{aligned} & \text{> Sss2 := solve(TGE, ChrCoe \setminus \{a155, a255, a355, a455, a555\}) : nops(Sss2)} \\ & \end{aligned} \quad (6.48)$$

$\triangleright tmp := \{ \} : \text{for } i \text{ to } nops(Ss2) \text{ do if } lhs(Sss2[i]) = rhs(Sss2[i]) \text{ then } tmp := \{ op(tmp),$
 $lhs(Sss2[i]) \} : \text{fi. od: } tmp$

$\{ \}$

(6.49)

$\triangleright Sss2$

$\{ a111 = 2 a155, a112 = 0, a113 = 0, a114 = 0, a115 = 0, a121 = a255, a122 = a155, a123$ (6.50)

$$= 0, a124 = 0, a125 = 0, a131 = a355 - \frac{1}{2} u_{1,1}, a132 = 0, a133 = a155, a134 = 0, a135$$

$$= 0, a141 = a455 - u_{1,2}, a142 = -\frac{1}{2} u_{1,1}, a143 = 0, a144 = a155, a145 = 0, a151 =$$

$$-u_1 u_{1,1} + \frac{1}{2} a555 - \frac{3}{2} u_{2,2}, a152 = -u_{1,2}, a153 = -\frac{1}{2} u_{1,1}, a154 = 0, a221 = u_{1,1},$$

$$a222 = 2 a255, a223 = 0, a224 = 0, a225 = 0, a231 = u_{1,2}, a232 = \frac{1}{2} u_{1,1} + a355, a233$$

$$= a255, a234 = 0, a235 = 0, a241 = \frac{3}{2} u_1 u_{1,1} + u_{2,2}, a242 = a455, a243 = \frac{1}{2} u_{1,1},$$

$$a244 = a255, a245 = 0, a251 = 2 u_1 u_{1,2} + \frac{3}{2} u_2 u_{1,1} + u_{2,3}, a252 = \frac{1}{2} u_1 u_{1,1}$$

$$+ \frac{1}{2} a555 - \frac{1}{2} u_{2,2}, a253 = 0, a254 = \frac{1}{2} u_{1,1}, a331 = -u_1 u_{1,1} + u_{2,2}, a332 = 2 u_{1,2},$$

$$a333 = 2 a355, a334 = 0, a335 = 0, a341 = -u_1 u_{1,2} - \frac{1}{2} u_2 u_{1,1} + u_{2,3}, a342$$

$$= \frac{1}{2} u_1 u_{1,1} + 2 u_{2,2}, a343 = a455 + u_{1,2}, a344 = a355, a345 = 0, a351 = -\frac{5}{2} u_1^2 u_{1,1}$$

$$-u_1 u_{2,2} + u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + u_{3,3}, a352 = u_1 u_{1,2} + u_2 u_{1,1} + 2 u_{2,3}, a353 =$$

$$-\frac{1}{2} u_1 u_{1,1} + \frac{1}{2} a555 + \frac{1}{2} u_{2,2}, a354 = u_{1,2}, a441 = u_1^2 u_{1,1} - u_1 u_{2,2} - 2 u_2 u_{1,2}$$

$$+ u_{3,3}, a442 = -2 u_1 u_{1,2} - u_2 u_{1,1} + 2 u_{2,3}, a443 = 2 u_1 u_{1,1} + 3 u_{2,2}, a444 = 2 a455,$$

$$a445 = 0, a451 = u_1^2 u_{1,2} - u_1 u_{2,3} - 2 u_2 u_{2,2} - u_3 u_{1,2} + u_{3,4}, a452 = -\frac{3}{2} u_1^2 u_{1,1}$$

$$-2 u_1 u_{2,2} - u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + 2 u_{3,3}, a453 = \frac{1}{2} u_2 u_{1,1} + 3 u_{2,3}, a454 = u_1 u_{1,1}$$

$$+ \frac{1}{2} a555 + \frac{3}{2} u_{2,2}, a551 = -u_1^3 u_{1,1} + u_1^2 u_{2,2} + 4 u_1 u_2 u_{1,2} - 2 u_1 u_3 u_{1,1} + u_2^2 u_{1,1}$$

$$-u_1 u_{3,3} - 2 u_2 u_{2,3} - 3 u_3 u_{2,2} + u_{4,4}, a552 = 2 u_1^2 u_{1,2} - 2 u_1 u_{2,3} - 4 u_2 u_{2,2}$$

$$-2 u_3 u_{1,2} + 2 u_{3,4}, a553 = -4 u_1^2 u_{1,1} - 3 u_1 u_{2,2} - u_3 u_{1,1} + 3 u_{3,3}, a554 = 2 u_1 u_{1,2}$$

$$+ 2 u_2 u_{1,1} + 4 u_{2,3} \}$$

$\triangleright pt := \{ a155 = 0, a255 = 0, a355 = 0, a455 = 0, a555 = 0 \} :$

$\triangleright Sss20 := \{ op(subs(pt, Sss2)), op(pt) \}$

$$Sss20 := \left\{ a111=0, a112=0, a113=0, a114=0, a115=0, a121=0, a122=0, a123=0, \quad (6.51) \right.$$

$$\begin{aligned} & a124=0, a125=0, a131=-\frac{1}{2} u_{1,1}, a132=0, a133=0, a134=0, a135=0, a141= \\ & -u_{1,2}, a142=-\frac{1}{2} u_{1,1}, a143=0, a144=0, a145=0, a151=-u_1 u_{1,1}-\frac{3}{2} u_{2,2}, a152 \\ & =-u_{1,2}, a153=-\frac{1}{2} u_{1,1}, a154=0, a155=0, a221=u_{1,1}, a222=0, a223=0, a224 \\ & =0, a225=0, a231=u_{1,2}, a232=\frac{1}{2} u_{1,1}, a233=0, a234=0, a235=0, a241 \\ & =\frac{3}{2} u_1 u_{1,1}+u_{2,2}, a242=0, a243=\frac{1}{2} u_{1,1}, a244=0, a245=0, a251=2 u_1 u_{1,2} \\ & +\frac{3}{2} u_2 u_{1,1}+u_{2,3}, a252=\frac{1}{2} u_1 u_{1,1}-\frac{1}{2} u_{2,2}, a253=0, a254=\frac{1}{2} u_{1,1}, a255=0, \\ & a331=-u_1 u_{1,1}+u_{2,2}, a332=2 u_{1,2}, a333=0, a334=0, a335=0, a341=-u_1 u_{1,2} \\ & -\frac{1}{2} u_2 u_{1,1}+u_{2,3}, a342=\frac{1}{2} u_1 u_{1,1}+2 u_{2,2}, a343=u_{1,2}, a344=0, a345=0, a351 \\ & =-\frac{5}{2} u_1^2 u_{1,1}-u_1 u_{2,2}+u_2 u_{1,2}-\frac{1}{2} u_3 u_{1,1}+u_{3,3}, a352=u_1 u_{1,2}+u_2 u_{1,1} \\ & +2 u_{2,3}, a353=-\frac{1}{2} u_1 u_{1,1}+\frac{1}{2} u_{2,2}, a354=u_{1,2}, a355=0, a441=u_1^2 u_{1,1}-u_1 u_{2,2} \\ & -2 u_2 u_{1,2}+u_{3,3}, a442=-2 u_1 u_{1,2}-u_2 u_{1,1}+2 u_{2,3}, a443=2 u_1 u_{1,1}+3 u_{2,2}, a444 \\ & =0, a445=0, a451=u_1^2 u_{1,2}-u_1 u_{2,3}-2 u_2 u_{2,2}-u_3 u_{1,2}+u_{3,4}, a452=-\frac{3}{2} u_1^2 u_{1,1} \\ & -2 u_1 u_{2,2}-u_2 u_{1,2}-\frac{1}{2} u_3 u_{1,1}+2 u_{3,3}, a453=\frac{1}{2} u_2 u_{1,1}+3 u_{2,3}, a454=u_1 u_{1,1} \\ & +\frac{3}{2} u_{2,2}, a455=0, a551=-u_1^3 u_{1,1}+u_1^2 u_{2,2}+4 u_1 u_2 u_{1,2}-2 u_1 u_3 u_{1,1}+u_2^2 u_{1,1} \\ & -u_1 u_{3,3}-2 u_2 u_{2,3}-3 u_3 u_{2,2}+u_{4,4}, a552=2 u_1^2 u_{1,2}-2 u_1 u_{2,3}-4 u_2 u_{2,2} \\ & -2 u_3 u_{1,2}+2 u_{3,4}, a553=-4 u_1^2 u_{1,1}-3 u_1 u_{2,2}-u_3 u_{1,1}+3 u_{3,3}, a554=2 u_1 u_{1,2} \\ & +2 u_2 u_{1,1}+4 u_{2,3}, a555=0 \} \end{aligned}$$

> CsC0 := subs(Sss20, CsC) :

> DGsimplify(CsC0)

(6.52)

$$> \Pi_{131} = -\frac{1}{2} u_{1,1},$$

$$\Pi_{141} = -u_{1,2}, \Pi_{142} = -\frac{1}{2} u_{1,1},$$

$$\Pi_{151} = -u_1 u_{1,1} - \frac{3}{2} u_{2,2}, \Pi_{152} = -u_{1,2}, \Pi_{153} = -\frac{1}{2} u_{1,1},$$

$$\Gamma_{221} = u_{1,1},$$

$$\Gamma_{231} = u_{1,2}, \Gamma_{232} = \frac{1}{2} u_{1,1},$$

$$\Gamma_{241} = \frac{3}{2} u_1 u_{1,1} + u_{2,2}, \Gamma_{243} = \frac{1}{2} u_{1,1},$$

$$\Gamma_{251} = 2 u_1 u_{1,2} + \frac{3}{2} u_2 u_{1,1} + u_{2,3}, \Gamma_{252} = \frac{1}{2} u_1 u_{1,1} - \frac{1}{2} u_{2,2}, \Gamma_{254} = \frac{1}{2} u_{1,1},$$

$$\Gamma_{331} = -u_1 u_{1,1} + u_{2,2}, \Gamma_{332} = 2 u_{1,2},$$

$$\Gamma_{341} = -u_1 u_{1,2} - \frac{1}{2} u_2 u_{1,1} + u_{2,3}, \Gamma_{342} = \frac{1}{2} u_1 u_{1,1} + 2 u_{2,2}, \Gamma_{343} = u_{1,2},$$

$$\Gamma_{351} = -\frac{5}{2} u_1^2 u_{1,1} - u_1 u_{2,2} + u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + u_{3,3}, \Gamma_{352} = u_1 u_{1,2} + u_2 u_{1,1} + 2 u_{2,3}, \Gamma_{353} = -\frac{1}{2} u_1 u_{1,1} + \frac{1}{2} u_{2,2}, \Gamma_{354} = u_{1,2},$$

$$\Gamma_{441} = u_1^2 u_{1,1} - u_1 u_{2,2} - 2 u_2 u_{1,2} + u_{3,3}, \Gamma_{442} = -2 u_1 u_{1,2} - u_2 u_{1,1} + 2 u_{2,3}, \Gamma_{443} = 2 u_1 u_{1,1} + 3 u_{2,2},$$

$$\Gamma_{451} = u_1^2 u_{1,2} - u_1 u_{2,3} - 2 u_2 u_{2,2} - u_3 u_{1,2} + u_{3,4}, \Gamma_{452} = -\frac{3}{2} u_1^2 u_{1,1} - 2 u_1 u_{2,2} - u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + 2 u_{3,3}, \Gamma_{453} = \frac{1}{2} u_2 u_{1,1} + 3 u_{2,3}, \Gamma_{454} = u_1 u_{1,1} + \frac{3}{2} u_{2,2},$$

$$\Gamma_{551} = -u_1^3 u_{1,1} + u_1^2 u_{2,2} + 4 u_1 u_2 u_{1,2} - 2 u_1 u_3 u_{1,1} + u_2^2 u_{1,1} - u_1 u_{3,3} - 2 u_2 u_{2,3} - 3 u_3 u_{2,2} + u_{4,4}, \Gamma_{552} = 2 u_1^2 u_{1,2} - 2 u_1 u_{2,3} - 4 u_2 u_{2,2} - 2 u_3 u_{1,2} + 2 u_{3,4}, \Gamma_{553} = -4 u_1^2 u_{1,1} - 3 u_1 u_{2,2} - u_3 u_{1,1} + 3 u_{3,3}, \Gamma_{554} = 2 u_1 u_{1,2} + 2 u_2 u_{1,1} + 4 u_{2,3}$$

Adler-Shabat hierarchy 5D

> DGsetup([x1, x2, x3, x4, x5], [u], J, 4) :

> Eq := (i, j) → (u_i - u_j) · u_{i,j} - u_{1,i} + u_{1,j} :

EQ := {Eq(2, 3), Eq(2, 4), Eq(2, 5), Eq(3, 4), Eq(3, 5), Eq(4, 5)} :

> S2 := solve(EQ, {u_{2,3}, u_{2,4}, u_{2,5}, u_{3,4}, u_{3,5}, u_{4,5}})

$$S2 := \left\{ u_{2,3} = \frac{u_{1,2} - u_{1,3}}{u_2 - u_3}, u_{2,4} = \frac{u_{1,2} - u_{1,4}}{u_2 - u_4}, u_{2,5} = \frac{u_{1,2} - u_{1,5}}{u_2 - u_5}, u_{3,4} = \frac{u_{1,3} - u_{1,4}}{u_3 - u_4}, u_{3,5} = \frac{u_{1,3} - u_{1,5}}{u_3 - u_5}, u_{4,5} = \frac{u_{1,4} - u_{1,5}}{u_4 - u_5} \right\} \quad (7.1)$$

> Symb := k → add(add(diff(EQ[k], u_{i,j}) · p_i · p_j, j = 1..5), i = 1..5) :

$$\text{simplify}\left(\text{subs}\left(\left\{p_1 = 1, \text{seq}\left(p_i = \frac{1}{q - u_i}, i = 2..5\right)\right\}, \left\{\text{seq}(\text{Symb}(k), k = 1..6)\right\}\right)\right) \quad \{0\} \quad (7.2)$$

> Vss3 := EQ :for i to 5 do for j to 6 do Vss3 := {op(Vss3), TotalDiff(EQ[j], i)} :od:od:

$$tmp := eliminate(Vss3, \{u_{2,3}, u_{2,4}, u_{2,5}, u_{3,4}, u_{3,5}, u_{4,5}, u_{1,2,3}, u_{1,2,4}, u_{1,3,4}, u_{1,2,5}, u_{1,3,4}, u_{1,3,5}, u_{1,4,5}, u_{2,2,3}, u_{2,2,4}, u_{2,2,5}, u_{2,3,3}, u_{2,3,4}, u_{2,3,5}, u_{2,4,4}, u_{2,4,5}, u_{2,5,5}, u_{3,3,4}, u_{3,3,5}, u_{3,4,4}, u_{3,4,5}, u_{3,5,5}, u_{4,4,5}, u_{4,5,5}\}) : [nops(tmp[1]), nops(tmp[2])];$$

(7.3)

$$> S3 := solve(Vss3, \{u_{2,3}, u_{2,4}, u_{2,5}, u_{3,4}, u_{3,5}, u_{4,5}, u_{1,2,3}, u_{1,2,4}, u_{1,3,4}, u_{1,2,5}, u_{1,3,4}, u_{1,3,5}, u_{1,4,5}, u_{2,2,3}, u_{2,2,4}, u_{2,2,5}, u_{2,3,3}, u_{2,3,4}, u_{2,3,5}, u_{2,4,4}, u_{2,4,5}, u_{2,5,5}, u_{3,3,4}, u_{3,3,5}, u_{3,4,4}, u_{3,4,5}, u_{3,5,5}, u_{4,4,5}, u_{4,5,5}\}) :$$

$$> tmp := solve(\{coeffs(expand(add(add(a[k,l] \cdot p_k, k=1..5) \cdot Symb(l), l=1..6)), \{p_1, p_2, p_3, p_4, p_5\})\}, \{seq(seq(a[k,l], k=1..5), l=1..6)\})$$

$$tmp := \left\{ a_{1,1} = a_{1,4} + a_{1,5}, a_{1,2} = -a_{1,4} + a_{1,6}, a_{1,3} = -a_{1,5} - a_{1,6}, a_{1,4} = a_{1,4}, a_{1,5} = a_{1,5}, \right.$$

(7.4)

$$a_{1,6} = a_{1,6}, a_{2,1} = 0, a_{2,2} = 0, a_{2,3} = 0, a_{2,4} = \frac{1}{u_4 - u_5} (a_{1,4} u_2 u_4 - a_{1,4} u_2 u_5 - a_{1,4}$$

$$u_4^2 + a_{1,4} u_4 u_5 + a_{4,1} u_4 - a_{4,1} u_5 + a_{4,5} u_2 - a_{4,5} u_4 + a_{5,4} u_2 - a_{5,4} u_4), a_{2,5}$$

$$= \frac{1}{u_4 - u_5} (a_{1,5} u_2 u_4 - a_{1,5} u_2 u_5 - a_{1,5} u_4 u_5 + a_{1,5} u_5^2 - a_{4,5} u_2 + a_{4,5} u_5 + a_{5,1} u_4$$

$$- a_{5,1} u_5 - a_{5,4} u_2 + a_{5,4} u_5), a_{2,6} = \frac{1}{u_4 - u_5} (a_{1,6} u_2 u_4 - a_{1,6} u_2 u_5 - a_{1,6} u_4 u_5$$

$$+ a_{1,6} u_5^2 + a_{4,5} u_2 - a_{4,5} u_5 + a_{5,2} u_4 - a_{5,2} u_5 + a_{5,4} u_2 - a_{5,4} u_5), a_{3,1} = 0, a_{3,2}$$

$$= -\frac{1}{u_4 - u_5} (a_{1,4} u_3 u_4 - a_{1,4} u_3 u_5 - a_{1,4} u_4^2 + a_{1,4} u_4 u_5 + a_{4,1} u_4 - a_{4,1} u_5 + a_{4,5} u_3$$

$$- a_{4,5} u_4 + a_{5,4} u_3 - a_{5,4} u_4), a_{3,3} = -\frac{1}{u_4 - u_5} (a_{1,5} u_3 u_4 - a_{1,5} u_3 u_5 - a_{1,5} u_4 u_5$$

$$+ a_{1,5} u_5^2 - a_{4,5} u_3 + a_{4,5} u_5 + a_{5,1} u_4 - a_{5,1} u_5 - a_{5,4} u_3 + a_{5,4} u_5), a_{3,4} = 0, a_{3,5} = 0,$$

$$a_{3,6} = -\frac{a_{4,5} u_3 - a_{4,5} u_5 + a_{5,4} u_3 - a_{5,4} u_4}{u_4 - u_5}, a_{4,1} = a_{4,1}, a_{4,2} = 0, a_{4,3} = -a_{1,6} u_4$$

$$+ a_{1,6} u_5 - a_{4,5} - a_{5,2} - a_{5,4}, a_{4,4} = 0, a_{4,5} = a_{4,5}, a_{4,6} = 0, a_{5,1} = a_{5,1}, a_{5,2} = a_{5,2}, a_{5,3}$$

$$= 0, a_{5,4} = a_{5,4}, a_{5,5} = 0, a_{5,6} = 0\}$$

so 8 compatibility conditions parametrized by (c1,c2,e1,e2,e3,f1,f2,f3) - we check them at one step

$$> simplify(subs(S2, subs(tmp, add(add(a[i,j] \cdot TotalDiff(EQ[j], i), i=1..5), j=1..6))))$$

0

(7.5)

This shows that dimensions of the symbol spaces are given by $\dim(g[k])=1+4k$, whence the general solution depends on 4 functions of 1 variable.

This is also what is expected from the characteristic variety (rational normal curve in our case) that has order 4 and dimension 1.

```

>
> DGsetup( [x1, x2, x3, x4, x5], [u, q], Jq, 3) :
> Qsyst := { seq( q_i = \frac{1}{q_{[ ]} - u_i} q_1 - \frac{u_{1,i}}{q_{[ ]} - u_i}, i = 2 .. 5 ) } :
> simplify( subs( S3, expand( subs( Qsyst, subs( { seq( q_{1,i} = TotalDiff( subs( Qsyst, q_i ), 1 ), i = 2
..5 ), { seq( seq( TotalDiff( subs( Qsyst, q_i ), j ) - TotalDiff( subs( Qsyst, q_j ), i ), j = i + 1
..5 ), i = 2 .. 4 ) } ) ) ) ) ) )
{0}
(7.6)

```

```

> DGsetup( [x1, x2, x3, x4, x5], [u], Jqu, 3) :
> Xf := k → ( f → TotalDiff( f, k ) - \frac{1}{q - u_k} · TotalDiff( f, 1 ) - \frac{u_{1,k}}{q - u_k} · diff( f, q ) ) :
> simplify( subs( S3, { seq( seq( simplify( Xf( i ) ( Xf( j ) ( x1 ) ) ) - Xf( j ) ( Xf( i ) ( x1 ) ) ), j = i + 1 .. 5 ),
i = 2 .. 4 ), seq( seq( simplify( Xf( i ) ( Xf( j ) ( q ) ) ) - Xf( j ) ( Xf( i ) ( q ) ) ), j = i + 1 .. 5 ), i = 2
..4 ) } ) ) )
{0}
(7.7)

```

I. We first look for a connection (with torsion) preserving the field of GL2-curves and having α -surfaces totally geodesic + trace of torsion zero.

```

> DGsetup( [x1, x2, x3, x4, x5], M ) :
> vM := [ seq( D_x || i, i = 1 .. 5 ) ] : dM := [ seq( dx || i, i = 1 .. 5 ) ] : var := seq( x || i, i = 1 .. 5 ) :
> C := Connection( add( add( add( ( ( ( a || i ) || j ) || k ) · vM[k] &t dM[i] &t dM[j], k = 1 .. 5 ), j = 1
..5 ), i = 1 .. 5 ) ) :
> X := k → evalDG( vM[k] - \frac{1}{q_{[ ]} - u_k} · vM[1] ) :
ω := evalDG( dM[1] + add( \frac{1}{q_{[ ]} - u_k} · dM[k], k = 2 .. 5 ) ) :
[ seq( Hook( X( i ), ω ), i = 2 .. 5 ) ]
[0, 0, 0, 0]
(7.8)

```

```

> FromJet( u_1, { u( var ), q( var ) } )
\frac{\partial}{\partial x1} u(x1, x2, x3, x4, x5)
(7.9)

```

```

> Y := k → FromJet( X( k ), { u( var ), q( var ) } ) : ωω := FromJet( ω, { u( var ), q( var ) } ) :
> DD := { seq( Y( k ), k = 2 .. 5 ) } : ChrCoe := { seq( seq( seq( ( ( ( a || i ) || j ) || k ), i = 1 .. 5 ), j = 1
..5 ), k = 1 .. 5 ) } :
TGE := { } : for i to 4 do for j to 4 do TGE := { op( TGE ), coeffs( expand( numer( subs( S2,
subs( Qsyst, ToJet( Hook( convert( DirectionalCovariantDerivative( DD[i], DD[j], C ),
'DGvector' ), ωω ), { u( var ), q( var ) }, notation = jetnumbers ) ) ) ), { q_{[ ]}, q_1 } ) } : od: od:
[ nops( ChrCoe ), nops( TGE ) ]
[125, 116]
(7.10)

```

```

> Sconn := solve( TGE, ChrCoe ) :
> FreePar := { } : for i to nops( Sconn ) do if lhs( Sconn[i] ) = rhs( Sconn[i] ) then FreePar

```

$$:= \{op(\text{FreePar}), lhs(\text{Sconn}[i])\} \text{ fi od: FreePar} \\ \{a155, a244, a255, a323, a355, a455, a511, a515, a525, a535, a544, a545, a555\} \quad (7.11)$$

$$\begin{aligned} &> \text{ChrCoe1} := \text{ChrCoe} \setminus \text{FreePar} \\ \text{ChrCoe1} &:= \{a111, a112, a113, a114, a115, a121, a122, a123, a124, a125, a131, a132, \\ &a133, a134, a135, a141, a142, a143, a144, a145, a151, a152, a153, a154, a211, a212, \\ &a213, a214, a215, a221, a222, a223, a224, a225, a231, a232, a233, a234, a235, a241, \\ &a242, a243, a245, a251, a252, a253, a254, a311, a312, a313, a314, a315, a321, a322, \\ &a324, a325, a331, a332, a333, a334, a335, a341, a342, a343, a344, a345, a351, a352, \\ &a353, a354, a411, a412, a413, a414, a415, a421, a422, a423, a424, a425, a431, a432, \\ &a433, a434, a435, a441, a442, a443, a444, a445, a451, a452, a453, a454, a512, a513, \\ &a514, a521, a522, a523, a524, a531, a532, a533, a534, a541, a542, a543, a551, a552, \\ &a553, a554\} \end{aligned} \quad (7.12)$$

$$\begin{aligned} &> [\text{nops}(\text{ChrCoe}), \text{nops}(\text{FreePar}), \text{nops}(\text{ChrCoe1})] \\ &\quad [125, 13, 112] \end{aligned} \quad (7.13)$$

$$\begin{aligned} &> \text{FreePar} := \{a155, a244, a255, a323, a355, a455, a511, a515, a525, a535, a544, a545, \\ &a555\} : \end{aligned}$$

$$\begin{aligned} &> \text{ChrCoe1} := \{a111, a112, a113, a114, a115, a121, a122, a123, a124, a125, a131, a132, \\ &a133, a134, a135, a141, a142, a143, a144, a145, a151, a152, a153, a154, a211, a212, \\ &a213, a214, a215, a221, a222, a223, a224, a225, a231, a232, a233, a234, a235, a241, \\ &a242, a243, a245, a251, a252, a253, a254, a311, a312, a313, a314, a315, a321, a322, \\ &a324, a325, a331, a332, a333, a334, a335, a341, a342, a343, a344, a345, a351, a352, \\ &a353, a354, a411, a412, a413, a414, a415, a421, a422, a423, a424, a425, a431, a432, \\ &a433, a434, a435, a441, a442, a443, a444, a445, a451, a452, a453, a454, a512, a513, \\ &a514, a521, a522, a523, a524, a531, a532, a533, a534, a541, a542, a543, a551, a552, \\ &a553, a554\} : \end{aligned}$$

$$\begin{aligned} &> \text{Sconn} := \text{solve}(\text{TGE}, \text{ChrCoe1}) : \end{aligned}$$

$$\begin{aligned} &> \text{CC} := \text{DGsimplify}(\text{subs}(\text{Sconn}, \text{C})) : \end{aligned}$$

$$\begin{aligned} &> \text{for } k \text{ to } 6 \text{ do } \text{Quad} \parallel k := \text{evalDG}(\text{add}(\text{add}(\text{diff}(\text{EQ}[k], u_{i,j}) \cdot vM[i] \&s vM[j], j = i \dots 5), i \\ &\quad = 1 \dots 5)) \text{ od:} \end{aligned}$$

$$\begin{aligned} &\text{for } k \text{ to } 6 \text{ do } \text{Quadu} \parallel k := \text{FromJet}(\text{Quad} \parallel k, \{u(\text{var}), q(\text{var})\}, \text{notation} = \text{jetnumbers}) \text{ od:} \\ &\text{Quadu} := [\text{seq}(\text{Quadu} \parallel k, k = 1 \dots 6)] : \end{aligned}$$

$$\begin{aligned} &> \{\text{seq}(\text{ContractIndices}(\text{Quad} \parallel i, \omega \&t \omega, [[1, 1], [2, 2]]), i = 1 \dots 6)\} \\ &\quad \{0\} \end{aligned} \quad (7.14)$$

$$\begin{aligned} &> \text{RNE} := \{ \} : \text{for } i \text{ to } 5 \text{ do for } j \text{ to } 6 \text{ do } \text{RNE} := \{op(\text{RNE}), \\ &\quad \text{coeffs}(\text{expand}(\text{numer}(\text{simplify}(\text{subs}(\text{S2}, \text{subs}(\text{Qsyst}, \\ &\quad \text{ToJet}(\text{ContractIndices}(\text{DirectionalCovariantDerivative}(vM[i], \text{Quadu}[j], \text{CC}), \omega \\ &\quad \&t \omega, [[1, 1], [2, 2]]), \{u(\text{var})\}, \text{notation} = \text{jetnumbers}))) \&s \{q[\]\}) \} \text{ od od:} \\ &\quad \text{nops}(\text{RNE}) \end{aligned}$$

$$18 \quad (7.15)$$

$$\begin{aligned} &> \text{tmp} := \text{eliminate}(\text{RNE}, \text{FreePar}) \end{aligned}$$

$$\text{tmp} := \left[\left\{ a155 = - \frac{1}{(u_4 - u_5)(u_2^2 - u_2 u_4 - u_2 u_5 + u_4 u_5)} (u_2^2 u_4 u_{5,5} - u_2^2 u_5 u_{5,5} - u_2 \right. \right. \quad (7.16)$$

$$\begin{aligned}
& u_4^2 u_{5,5} + u_2 u_5^2 u_{5,5} + u_4^2 u_5 u_{5,5} - u_4 u_5^2 u_{5,5} - u_2^2 u_{1,4} + u_2^2 u_{1,5} + 2 u_2 u_5 u_{1,4} \\
& - 2 u_2 u_5 u_{1,5} + u_4^2 u_{1,2} - u_4^2 u_{1,5} - 2 u_4 u_5 u_{1,2} + 2 u_4 u_5 u_{1,5} + u_5^2 u_{1,2} - u_5^2 u_{1,4}), \\
& a_{244} = (u_2^2 u_4 u_{4,4} - u_2^2 u_5 u_{4,4} - u_2 u_4^2 u_{4,4} + u_2 u_5^2 u_{4,4} + u_4^2 u_5 u_{4,4} - u_4 u_5^2 u_{4,4} - \\
& u_2^2 u_{1,4} + u_2^2 u_{1,5} + 2 u_2 u_4 u_{1,4} - 2 u_2 u_4 u_{1,5} - u_4^2 u_{1,2} + u_4^2 u_{1,5} + 2 u_4 u_5 u_{1,2} \\
& - 2 u_4 u_5 u_{1,4} - u_5^2 u_{1,2} + u_5^2 u_{1,4}) / (u_2^3 u_4 - u_2^3 u_5 - 2 u_2^2 u_4^2 + u_2^2 u_4 u_5 + u_2^2 u_5^2 + u_2 u_4^3 \\
& + u_2 u_4^2 u_5 - 2 u_2 u_4 u_5^2 - u_4^3 u_5 + u_4^2 u_5^2), a_{255} = (u_2^2 u_4 u_{5,5} - u_2^2 u_5 u_{5,5} - u_2 u_4^2 u_{5,5} \\
& + u_2 u_5^2 u_{5,5} + u_4^2 u_5 u_{5,5} - u_4 u_5^2 u_{5,5} - u_2^2 u_{1,4} + u_2^2 u_{1,5} + 2 u_2 u_5 u_{1,4} - 2 u_2 u_5 u_{1,5} + \\
& u_4^2 u_{1,2} - u_4^2 u_{1,5} - 2 u_4 u_5 u_{1,2} + 2 u_4 u_5 u_{1,5} + u_5^2 u_{1,2} - u_5^2 u_{1,4}) / ((u_2^2 u_4 - u_2^2 u_5 \\
& - 2 u_2 u_4 u_5 + 2 u_2 u_5^2 + u_4 u_5^2 - u_5^3) (u_2 - u_4)), a_{323} = (a_{525} u_2^4 - 2 a_{525} u_2^3 u_3 \\
& - 2 a_{525} u_2^3 u_5 + a_{525} u_2^2 u_3^2 + 4 a_{525} u_2^2 u_3 u_5 + a_{525} u_2^2 u_5^2 - 2 a_{525} u_2 u_3^2 u_5 \\
& - 2 a_{525} u_2 u_3 u_5^2 + a_{525} u_3^2 u_5^2 + u_2^2 u_3 u_{2,2} - u_2^2 u_5 u_{2,2} - u_2 u_3^2 u_{2,2} + u_2 u_5^2 u_{2,2} + \\
& u_3^2 u_5 u_{2,2} - u_3 u_5^2 u_{2,2} + u_2^2 u_{1,3} - u_2^2 u_{1,5} - 2 u_2 u_3 u_{1,2} + 2 u_2 u_3 u_{1,5} + 2 u_2 u_5 u_{1,2} \\
& - 2 u_2 u_5 u_{1,3} + u_3^2 u_{1,2} - u_3^2 u_{1,5} - u_5^2 u_{1,2} + u_5^2 u_{1,3}) / (u_2^4 - 2 u_2^3 u_3 - 2 u_2^3 u_5 + u_2^2 u_3^2 \\
& + 4 u_2^2 u_3 u_5 + u_2^2 u_5^2 - 2 u_2 u_3^2 u_5 - 2 u_2 u_3 u_5^2 + u_3^2 u_5^2), a_{355} = (u_2^2 u_3 u_4 u_{5,5} - \\
& u_2^2 u_3 u_5 u_{5,5} - u_2^2 u_4 u_5 u_{5,5} + u_2^2 u_5^2 u_{5,5} - u_2 u_3 u_4^2 u_{5,5} + u_2 u_3 u_5^2 u_{5,5} + u_2 u_4^2 u_5 u_{5,5}
\end{aligned}$$

$$\begin{aligned}
& -u_2 u_5^3 u_{5,5} + u_3 u_4^2 u_5 u_{5,5} - u_3 u_4 u_5^2 u_{5,5} - u_4^2 u_5^2 u_{5,5} + u_4 u_5^3 u_{5,5} - u_2^2 u_4 u_{1,3} + \\
& u_2^2 u_4 u_{1,5} + u_2^2 u_5 u_{1,3} - u_2^2 u_5 u_{1,5} - u_2 u_3^2 u_{1,4} + u_2 u_3^2 u_{1,5} + 2 u_2 u_3 u_5 u_{1,4} \\
& - 2 u_2 u_3 u_5 u_{1,5} + u_2 u_4^2 u_{1,3} - u_2 u_4^2 u_{1,5} - u_2 u_5^2 u_{1,3} - u_2 u_5^2 u_{1,4} + 2 u_2 u_5^2 u_{1,5} + \\
& u_3^2 u_4 u_{1,2} - u_3^2 u_4 u_{1,5} - u_3^2 u_5 u_{1,2} + u_3^2 u_5 u_{1,4} - 2 u_3 u_4 u_5 u_{1,2} + 2 u_3 u_4 u_5 u_{1,5} + 2 u_3 \\
& u_5^2 u_{1,2} - 2 u_3 u_5^2 u_{1,4} - u_4^2 u_5 u_{1,3} + u_4^2 u_5 u_{1,5} + u_4 u_5^2 u_{1,2} + u_4 u_5^2 u_{1,3} - 2 u_4 u_5^2 u_{1,5} - \\
& u_5^3 u_{1,2} + u_5^3 u_{1,4}) / (u_2^2 u_3^2 u_4 - u_2^2 u_3^2 u_5 - 2 u_2^2 u_3 u_4 u_5 + 2 u_2^2 u_3 u_5^2 + u_2^2 u_4 u_5^2 - u_2^2 u_5^3 \\
& - u_2 u_3^2 u_4^2 + u_2 u_3^2 u_5^2 + 2 u_2 u_3 u_4^2 u_5 - 2 u_2 u_3 u_5^3 - u_2 u_4^2 u_5^2 + u_2 u_5^4 + u_3^2 u_4^2 u_5 - u_3^2 u_4 u_5^2 \\
& - 2 u_3 u_4^2 u_5^2 + 2 u_3 u_4 u_5^3 + u_4^2 u_5^3 - u_4 u_5^4), a455 = (u_2^2 u_4 u_{5,5} - u_2^2 u_5 u_{5,5} - u_2 \\
& u_4^2 u_{5,5} + u_2 u_5^2 u_{5,5} + u_4^2 u_5 u_{5,5} - u_4 u_5^2 u_{5,5} - u_2^2 u_{1,4} + u_2^2 u_{1,5} + 2 u_2 u_5 u_{1,4} \\
& - 2 u_2 u_5 u_{1,5} + u_4^2 u_{1,2} - u_4^2 u_{1,5} - 2 u_4 u_5 u_{1,2} + 2 u_4 u_5 u_{1,5} + u_5^2 u_{1,2} - u_5^2 u_{1,4}) / ((\\
& u_2^2 - u_2 u_4 - u_2 u_5 + u_4 u_5) (u_4^2 - 2 u_4 u_5 + u_5^2)), a511 \\
& = \frac{u_2 u_{1,4} - u_2 u_{1,5} - u_4 u_{1,2} + u_4 u_{1,5} + u_5 u_{1,2} - u_5 u_{1,4}}{(u_4 - u_5) (u_2^2 - u_2 u_4 - u_2 u_5 + u_4 u_5)}, a515 = a515, a525 = a525, \\
& a535 = a535, a544 = - (u_2^2 u_4 u_{4,4} - u_2^2 u_5 u_{4,4} - u_2 u_4^2 u_{4,4} + u_2 u_5^2 u_{4,4} + u_4^2 u_5 u_{4,4} \\
& - u_4 u_5^2 u_{4,4} - u_2^2 u_{1,4} + u_2^2 u_{1,5} + 2 u_2 u_4 u_{1,4} - 2 u_2 u_4 u_{1,5} - u_4^2 u_{1,2} + u_4^2 u_{1,5} \\
& + 2 u_4 u_5 u_{1,2} - 2 u_4 u_5 u_{1,4} - u_5^2 u_{1,2} + u_5^2 u_{1,4}) / (u_2^2 u_4^2 - 2 u_2^2 u_4 u_5 + u_2^2 u_5^2 - u_2 u_4^3 \\
& + u_2 u_4^2 u_5 + u_2 u_4 u_5^2 - u_2 u_5^3 + u_4^3 u_5 - 2 u_4^2 u_5^2 + u_4 u_5^3), a545 = a545, a555 = a555\}, \{ \\
& u_2^4 u_3^3 u_4 u_{1,4} - u_2^4 u_3^3 u_4 u_{1,5} - u_2^4 u_3^3 u_5 u_{1,4} + u_2^4 u_3^3 u_5 u_{1,5} - u_2^4 u_3^2 u_4^2 u_{1,3} + u_2^4 u_3^2 u_4^2 u_{1,5} \\
& + 2 u_2^4 u_3^2 u_4 u_5 u_{1,3} - 3 u_2^4 u_3^2 u_4 u_5 u_{1,4} + u_2^4 u_3^2 u_4 u_5 u_{1,5} - u_2^4 u_3^2 u_5^2 u_{1,3} + 3 u_2^4 u_3^2
\end{aligned}$$

$$\begin{aligned}
& u_2^2 u_{1,4} - 2 u_2^4 u_3^2 u_5^2 u_{1,5} + 2 u_2^4 u_3 u_4^2 u_5 u_{1,3} - 2 u_2^4 u_3 u_4^2 u_5 u_{1,5} - 4 u_2^4 u_3 u_4 u_5^2 u_{1,3} + 3 \\
& u_2^4 u_3 u_4 u_5^2 u_{1,4} + u_2^4 u_3 u_4 u_5^2 u_{1,5} + 2 u_2^4 u_3 u_5^3 u_{1,3} - 3 u_2^4 u_3 u_5^3 u_{1,4} + u_2^4 u_3 u_5^3 u_{1,5} - u_2^4 \\
& u_4^2 u_5^2 u_{1,3} + u_2^4 u_4^2 u_5^2 u_{1,5} + 2 u_2^4 u_4 u_5^3 u_{1,3} - u_2^4 u_4 u_5^3 u_{1,4} - u_2^4 u_4 u_5^3 u_{1,5} - u_2^4 u_5^4 u_{1,3} + \\
& u_2^4 u_5^4 u_{1,4} - 2 u_2^3 u_3^4 u_4 u_{1,4} + 2 u_2^3 u_3^4 u_4 u_{1,5} + 2 u_2^3 u_3^4 u_5 u_{1,4} - 2 u_2^3 u_3^4 u_5 u_{1,5} + u_2^3 u_3^3 \\
& u_4^2 u_{1,3} - u_2^3 u_3^3 u_4^2 u_{1,5} - 2 u_2^3 u_3^3 u_4 u_5 u_{1,3} + 4 u_2^3 u_3^3 u_4 u_5 u_{1,4} - 2 u_2^3 u_3^3 u_4 u_5 u_{1,5} + u_2^3 \\
& u_3^3 u_5^2 u_{1,3} - 4 u_2^3 u_3^3 u_5^2 u_{1,4} + 3 u_2^3 u_3^3 u_5^2 u_{1,5} + u_2^3 u_3^2 u_4^3 u_{1,3} - u_2^3 u_3^2 u_4^3 u_{1,5} - 2 u_2^3 u_3^2 \\
& u_4^2 u_5 u_{1,3} + 2 u_2^3 u_3^2 u_4^2 u_5 u_{1,5} + u_2^3 u_3^2 u_4 u_5^2 u_{1,3} - u_2^3 u_3^2 u_4 u_5^2 u_{1,5} - 2 u_2^3 u_3 u_4^3 u_5 u_{1,3} \\
& + 2 u_2^3 u_3 u_4^3 u_5 u_{1,5} + u_2^3 u_3 u_4^2 u_5^2 u_{1,3} - u_2^3 u_3 u_4^2 u_5^2 u_{1,5} + 4 u_2^3 u_3 u_4 u_5^3 u_{1,3} - 4 u_2^3 u_3 u_4 \\
& u_5^3 u_{1,4} - 3 u_2^3 u_3 u_5^4 u_{1,3} + 4 u_2^3 u_3 u_5^4 u_{1,4} - u_2^3 u_3 u_5^4 u_{1,5} + u_2^3 u_3^3 u_5^2 u_{1,3} - u_2^3 u_3^3 u_5^2 u_{1,5} \\
& - 3 u_2^3 u_4 u_5^4 u_{1,3} + 2 u_2^3 u_4 u_5^4 u_{1,4} + u_2^3 u_4 u_5^4 u_{1,5} + 2 u_2^3 u_5^5 u_{1,3} - 2 u_2^3 u_5^5 u_{1,4} + u_2^2 \\
& u_3^5 u_4 u_{1,4} - u_2^2 u_3^5 u_4 u_{1,5} - u_2^2 u_3^5 u_5 u_{1,4} + u_2^2 u_3^5 u_5 u_{1,5} + u_2^2 u_3^4 u_4^2 u_{1,2} - u_2^2 u_3^4 u_4^2 u_{1,5} \\
& - 2 u_2^2 u_3^4 u_4 u_5 u_{1,2} + u_2^2 u_3^4 u_4 u_5 u_{1,4} + u_2^2 u_3^4 u_4 u_5 u_{1,5} + u_2^2 u_3^4 u_5^2 u_{1,2} - u_2^2 u_3^4 u_5^2 u_{1,4} - \\
& u_2^2 u_3^3 u_4^3 u_{1,2} - u_2^2 u_3^3 u_4^3 u_{1,3} + 2 u_2^2 u_3^3 u_4^3 u_{1,5} - u_2^2 u_3^3 u_4^2 u_5 u_{1,2} + u_2^2 u_3^3 u_4^2 u_5 u_{1,5} + 5 u_2^2 \\
& u_3^3 u_4 u_5^2 u_{1,2} + 3 u_2^2 u_3^3 u_4 u_5^2 u_{1,3} - 8 u_2^2 u_3^3 u_4 u_5^2 u_{1,4} - 3 u_2^2 u_3^3 u_5^3 u_{1,2} - 2 u_2^2 u_3^3 u_5^3 u_{1,3} \\
& + 8 u_2^2 u_3^3 u_5^3 u_{1,4} - 3 u_2^2 u_3^3 u_5^3 u_{1,5} + 3 u_2^2 u_3^2 u_4^3 u_5 u_{1,2} - 3 u_2^2 u_3^2 u_4^3 u_5 u_{1,5} - 3 u_2^2 u_3^2 u_4^2 \\
& u_5^2 u_{1,2} + 3 u_2^2 u_3^2 u_4^2 u_5^2 u_{1,3} - 3 u_2^2 u_3^2 u_4^2 u_5^2 u_{1,5} - 6 u_2^2 u_3^2 u_4 u_5^3 u_{1,2} + 8 u_2^2 u_3^2 u_4 u_5^3 u_{1,4} + \\
& u_2^2 u_3^2 u_4 u_5^3 u_{1,5} + 3 u_2^2 u_3^2 u_5^4 u_{1,2} + 3 u_2^2 u_3^2 u_5^4 u_{1,3} - 8 u_2^2 u_3^2 u_5^4 u_{1,4} + 2 u_2^2 u_3^2 u_5^4 u_{1,5} - 3 \\
& u_2^2 u_3 u_4^3 u_5^2 u_{1,2} + 3 u_2^2 u_3 u_4^3 u_5^2 u_{1,3} + 5 u_2^2 u_3 u_4^2 u_5^3 u_{1,2} - 6 u_2^2 u_3 u_4^2 u_5^3 u_{1,3} + u_2^2 u_3 u_4^2 \\
& u_5^3 u_{1,5} - u_2^2 u_3 u_4 u_5^4 u_{1,2} + 3 u_2^2 u_3 u_4 u_5^4 u_{1,3} - u_2^2 u_3 u_4 u_5^4 u_{1,4} - u_2^2 u_3 u_4 u_5^4 u_{1,5} - u_2^2 u_3 \\
& u_5^5 u_{1,2} + u_2^2 u_3 u_5^5 u_{1,4} + u_2^2 u_3^3 u_5^3 u_{1,2} - 2 u_2^2 u_3^3 u_5^3 u_{1,3} + u_2^2 u_3^3 u_5^3 u_{1,5} - 2 u_2^2 u_4^2 u_5^4 u_{1,2} \\
& + 3 u_2^2 u_4^2 u_5^4 u_{1,3} - u_2^2 u_4^2 u_5^4 u_{1,5} + u_2^2 u_4 u_5^5 u_{1,2} - u_2^2 u_4 u_5^5 u_{1,4} - u_2^2 u_5^6 u_{1,3} + u_2^2 u_5^6 u_{1,4} \\
& - u_2 u_3^5 u_4^2 u_{1,2} + u_2 u_3^5 u_4^2 u_{1,5} + 2 u_2 u_3^5 u_4 u_5 u_{1,2} - 2 u_2 u_3^5 u_4 u_5 u_{1,4} - u_2 u_3^5 u_5^2 u_{1,2} \\
& + 2 u_2 u_3^5 u_5^2 u_{1,4} - u_2 u_3^5 u_5^2 u_{1,5} + u_2 u_3^4 u_4^3 u_{1,2} - u_2 u_3^4 u_4^3 u_{1,5} - 3 u_2 u_3^4 u_4 u_5^2 u_{1,2} \\
& + 4 u_2 u_3^4 u_4 u_5^2 u_{1,4} - u_2 u_3^4 u_4 u_5^2 u_{1,5} + 2 u_2 u_3^4 u_5^3 u_{1,2} - 4 u_2 u_3^4 u_5^3 u_{1,4} + 2 u_2 u_3^4 u_5^3 u_{1,5} \\
& - 2 u_2 u_3^3 u_4^3 u_5 u_{1,2} + 2 u_2 u_3^3 u_4^3 u_5 u_{1,3} + 4 u_2 u_3^3 u_4^2 u_5^2 u_{1,2} - 3 u_2 u_3^3 u_4^2 u_5^2 u_{1,3} - u_2 u_3^3 \\
& u_4^2 u_5^2 u_{1,5} - 2 u_2 u_3^3 u_4 u_5^3 u_{1,2} + 2 u_2 u_3^3 u_4 u_5^3 u_{1,5} + u_2 u_3^3 u_5^4 u_{1,3} - u_2 u_3^3 u_5^4 u_{1,5} - 3 u_2 \\
& u_3^2 u_4^3 u_5^2 u_{1,3} + 3 u_2 u_3^2 u_4^3 u_5^2 u_{1,5} - 2 u_2 u_3^2 u_4^2 u_5^3 u_{1,2} + 4 u_2 u_3^2 u_4^2 u_5^3 u_{1,3} - 2 u_2 u_3^2 u_4^2 \\
& u_5^3 u_{1,5} + 4 u_2 u_3^2 u_4 u_5^4 u_{1,2} + u_2 u_3^2 u_4 u_5^4 u_{1,3} - 4 u_2 u_3^2 u_4 u_5^4 u_{1,4} - u_2 u_3^2 u_4 u_5^4 u_{1,5} \\
& - 2 u_2 u_3^2 u_5^5 u_{1,2} - 2 u_2 u_3^2 u_5^5 u_{1,3} + 4 u_2 u_3^2 u_5^5 u_{1,4} + 2 u_2 u_3 u_4^3 u_5^3 u_{1,2} - 2 u_2 u_3 u_4^3
\end{aligned}$$

$$\begin{aligned}
& u_5^3 u_{1,5} - 3 u_2 u_3 u_4^2 u_5^4 u_{1,2} + u_2 u_3 u_4^2 u_5^4 u_{1,3} + 2 u_2 u_3 u_4^2 u_5^4 u_{1,5} - 2 u_2 u_3 u_4 u_5^5 u_{1,3} \\
& + 2 u_2 u_3 u_4 u_5^5 u_{1,4} + u_2 u_3 u_5^6 u_{1,2} + u_2 u_3 u_5^6 u_{1,3} - 2 u_2 u_3 u_5^6 u_{1,4} - u_2 u_4^3 u_5^4 u_{1,2} + u_2 \\
& u_4^3 u_5^4 u_{1,3} + 2 u_2 u_4^2 u_5^5 u_{1,2} - 2 u_2 u_4^2 u_5^5 u_{1,3} - u_2 u_4 u_5^6 u_{1,2} + u_2 u_4 u_5^6 u_{1,3} + u_3^5 u_4^2 u_5 u_{1,2} \\
& - u_3^5 u_4^2 u_5 u_{1,5} - 2 u_3^5 u_4 u_5^2 u_{1,2} + u_3^5 u_4 u_5^2 u_{1,4} + u_3^5 u_4 u_5^2 u_{1,5} + u_3^5 u_5^3 u_{1,2} - u_3^5 u_5^3 u_{1,4} \\
& - u_3^4 u_4^3 u_5 u_{1,2} + u_3^4 u_4^3 u_5 u_{1,5} - u_3^4 u_4^2 u_5^2 u_{1,2} + u_3^4 u_4^2 u_5^2 u_{1,5} + 5 u_3^4 u_4 u_5^3 u_{1,2} - 3 u_3^4 u_4 \\
& u_5^3 u_{1,4} - 2 u_3^4 u_4 u_5^3 u_{1,5} - 3 u_3^4 u_5^4 u_{1,2} + 3 u_3^4 u_5^4 u_{1,4} + 3 u_3^3 u_4^3 u_5^2 u_{1,2} - u_3^3 u_4^3 u_5^2 u_{1,3} \\
& - 2 u_3^3 u_4^3 u_5^2 u_{1,5} - 3 u_3^3 u_4^2 u_5^3 u_{1,2} + 2 u_3^3 u_4^2 u_5^3 u_{1,3} + u_3^3 u_4^2 u_5^3 u_{1,5} - 3 u_3^3 u_4 u_5^4 u_{1,2} - \\
& u_3^3 u_4 u_5^4 u_{1,3} + 3 u_3^3 u_4 u_5^4 u_{1,4} + u_3^3 u_4 u_5^4 u_{1,5} + 3 u_3^3 u_5^5 u_{1,2} - 3 u_3^3 u_5^5 u_{1,4} - 3 u_3^2 u_4^3 \\
& u_5^3 u_{1,2} + 2 u_3^2 u_4^3 u_5^3 u_{1,3} + u_3^2 u_4^3 u_5^3 u_{1,5} + 5 u_3^2 u_4^2 u_5^4 u_{1,2} - 4 u_3^2 u_4^2 u_5^4 u_{1,3} - u_3^2 u_4^2 u_5^4 u_{1,5} \\
& - u_3^2 u_4 u_5^5 u_{1,2} + 2 u_3^2 u_4 u_5^5 u_{1,3} - u_3^2 u_4 u_5^5 u_{1,4} - u_3^2 u_5^6 u_{1,2} + u_3^2 u_5^6 u_{1,4} + u_3 u_4^3 u_5^4 u_{1,2} \\
& - u_3 u_4^3 u_5^4 u_{1,3} - 2 u_3 u_4^2 u_5^5 u_{1,2} + 2 u_3 u_4^2 u_5^5 u_{1,3} + u_3 u_4 u_5^6 u_{1,2} - u_3 u_4 u_5^6 u_{1,3} \}]
\end{aligned}$$

Thus GL2-connection having α -surfaces totally geodesic does not exist in 5D (for A-Sh) ...

Let us start looking for a connection by dropping the totally geodesic condition, but concentrating on GL2-property and Ricci tensor.

```

> RNE := { } :for i to 5 do for j to 6 do RNE := { op(RNE),
      coeffs( expand( numer( simplify( subs( S2, subs( Qsyst,
      ToJet( ContractIndices( DirectionalCovariantDerivative(vM[i], Quadu[j], C), ω &t ω,
      [[1, 1], [2, 2]]), {u(var)}, notation = jetnumbers) ) ) ) ), {q[_]} ) } :od:od: nops(RNE)
210
(7.17)

```

```

> Sconn := solve(RNE, ChrCoe) :
> FreePar := { } :for i to nops(Sconn) do if lhs(Sconn[i]) = rhs(Sconn[i]) then FreePar
      := { op(FreePar), lhs(Sconn[i]) } fi:od: FreePar, nops(%)
{a334, a435, a443, a445, a511, a512, a514, a515, a521, a522, a524, a525, a531, a535,
a541, a545, a551, a552, a554, a555}, 20
(7.18)

```

```

> ChrCoe1 := ChrCoe\FreePar
ChrCoe1 := {a111, a112, a113, a114, a115, a121, a122, a123, a124, a125, a131, a132,
a133, a134, a135, a141, a142, a143, a144, a145, a151, a152, a153, a154, a155, a211,
a212, a213, a214, a215, a221, a222, a223, a224, a225, a231, a232, a233, a234, a235,
a241, a242, a243, a244, a245, a251, a252, a253, a254, a255, a311, a312, a313, a314,
a315, a321, a322, a323, a324, a325, a331, a332, a333, a335, a341, a342, a343, a344,
a345, a351, a352, a353, a354, a355, a411, a412, a413, a414, a415, a421, a422, a423,
a424, a425, a431, a432, a433, a434, a441, a442, a444, a451, a452, a453, a454, a455,
a513, a523, a532, a533, a534, a542, a543, a544, a553}
(7.19)

```

```

> [ nops(ChrCoe), nops(FreePar), nops(ChrCoe1) ]
[125, 20, 105]
(7.20)

```

```

> FreePar := {a334, a435, a443, a445, a511, a512, a514, a515, a521, a522, a524, a525,
a531, a535, a541, a545, a551, a552, a554, a555} :

```

```
> ChrCoe1 := {a111, a112, a113, a114, a115, a121, a122, a123, a124, a125, a131, a132,
a133, a134, a135, a141, a142, a143, a144, a145, a151, a152, a153, a154, a155, a211,
a212, a213, a214, a215, a221, a222, a223, a224, a225, a231, a232, a233, a234, a235,
a241, a242, a243, a244, a245, a251, a252, a253, a254, a255, a311, a312, a313, a314,
a315, a321, a322, a323, a324, a325, a331, a332, a333, a335, a341, a342, a343, a344,
a345, a351, a352, a353, a354, a355, a411, a412, a413, a414, a415, a421, a422, a423,
a424, a425, a431, a432, a433, a434, a441, a442, a444, a451, a452, a453, a454, a455,
a513, a523, a532, a533, a534, a542, a543, a544, a553} :
```

```
> Sconn := solve(RNE, ChrCoe1) :
```

```
> CC := DGsimplify(subs(Sconn, C)) :
```

```
> TorC := TorsionTensor(CC) :
```

```
REN := {seq(seq(coeffs(expand(numer(subs(S2,
ToJet(Hook(convert(ContractIndices(DD[i] & DD[j], TorC, [[1, 2], [2, 3]]),
'DGvector'), ω), {u(var), q(var)}, notation = jetnumbers))))), {q[i]}), j = i + 1 .. 4), i
= 1 .. 4) : nops(REN)
```

42

(7.21)

```
> tmp := factor(eliminate(REN union DGinfo(ContractIndices(TorsionTensor(CC), [[1,
2]]), "CoefficientSet"), FreePar)) :
```

```
> tmp[2]
```

{ }

(7.22)

```
> tmnum := 0 :for i to nops(tmp[1]) do if lhs(tmp[1][i]) = rhs(tmp[1][i]) then tmnum
:= tmnum + 1 :fi:od: tmnum
```

0

(7.23)

```
> Sconn2 := tmp[1] : CC := DGsimplify(subs(Sconn2, CC)) :
```

```
> {seq(seq(seq(coeffs(expand(numer(simplify(subs(S2, subs(Qsyst,
ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CC), ω
& ω, [[1, 1], [2, 2]]), {u(var)}, notation = jetnumbers))))), {q[i]}), i = 1 .. 5), j = 1
.. 6) union DGinfo(ContractIndices(TorsionTensor(CC), [[1, 2]]), "CoefficientSet")
{0}
```

(7.24)

```
> RicCC := RicciTensor(FromJet(CC, {u(var)})) : RicCc := DGsimplify(subs(S3,
ToJet(RicCC, {u(var)}, notation = jetnumbers))) :
```

```
> RicCcSkew := SymmetrizeIndices(RicCc, [1, 2], "SkewSymmetric") :
```

```
RicCcSymm := SymmetrizeIndices(RicCc, [1, 2], "Symmetric") :
```

```
> ω
```

$$dx1 + \frac{1}{q[] - u_2} dx2 + \frac{1}{q[] - u_3} dx3 + \frac{1}{q[] - u_4} dx4 + \frac{1}{q[] - u_5} dx5$$

(7.25)

```
> ω0 := evalDG((q[] - u_2) (q[] - u_3) (q[] - u_4) (q[] - u_5) · ω) : ω0b
:= DGsimplify(evalDG(subs(q[] = 0, ω0))) :
```

```
ω1a := evalDG(1/ε · (subs(q[] = q[] + ε, ω0) - ω0)) : ω1 := evalDG(subs(ε = 0,
ω1a)) : ω1b := DGsimplify(evalDG(subs(q[] = 0, ω1))) :
```

```
ω2a := evalDG(1/ε · (subs(q[] = q[] + ε, ω1) - ω1)) : ω2 := evalDG(subs(ε = 0,
```

$$\begin{aligned}
& \omega 2a)) : \omega 2b := DGsimplify \left(evalDG \left(\frac{1}{2} \cdot subs(q_{[]} = 0, \omega 2) \right) \right) : \\
& \omega 3a := evalDG \left(\frac{1}{\varepsilon} \cdot (subs(q_{[]} = q_{[]} + \varepsilon, \omega 2) - \omega 2) \right) : \omega 3 := evalDG(subs(\varepsilon = 0, \\
& \omega 3a)) : \omega 3b := DGsimplify \left(evalDG \left(\frac{1}{6} \cdot subs(q_{[]} = 0, \omega 3) \right) \right) : \\
& \omega 4a := evalDG \left(\frac{1}{\varepsilon} \cdot (subs(q_{[]} = q_{[]} + \varepsilon, \omega 3) - \omega 3) \right) : \omega 4 := evalDG(subs(\varepsilon = 0, \\
& \omega 4a)) : \omega 4b := DGsimplify \left(evalDG \left(\frac{1}{24} \cdot subs(q_{[]} = 0, \omega 4) \right) \right) : \\
& > evalDG(\omega 0 - \omega 0b - \omega 1b \cdot q_{[]} - \omega 2b \cdot q_{[]}^2 - \omega 3b \cdot q_{[]}^3 - \omega 4b \cdot q_{[]}^4) \\
& \quad \quad \quad 0 \, dx1 \tag{7.26}
\end{aligned}$$

$$\begin{aligned}
& > \omega 3b \\
& \quad \quad \quad - ((u_2 + u_3 + u_4 + u_5) \, dx1) + dx2 + dx3 + dx4 + dx5 \tag{7.27}
\end{aligned}$$

$$\begin{aligned}
& > Qua\omega := evalDG([\omega 0b \&s \omega 0b, \omega 0b \&s \omega 1b, 2 \, \omega 0b \&s \omega 2b + \omega 1b \&s \omega 1b, \omega 0b \&s \omega 3b \\
& \quad + \omega 1b \&s \omega 2b, 2 \, \omega 0b \&s \omega 4b + 2 \, \omega 1b \&s \omega 3b + \omega 2b \&s \omega 2b, \omega 1b \&s \omega 4b + \omega 2b \\
& \quad \&s \omega 3b, 2 \, \omega 2b \&s \omega 4b + \omega 3b \&s \omega 3b, \omega 3b \&s \omega 4b, \omega 4b \&s \omega 4b]) : tmp\Omega := \{ \} : \textbf{for } i \\
& \quad \textbf{to nops}(Qua\omega) \textbf{ do for } j \textbf{ to } 6 \textbf{ do } tmp\Omega := \{ op(tmp\Omega), ContractIndices(Quad||j, \\
& \quad Qua\omega[i], [[1, 1], [2, 2]]) \} : \textbf{od:od: } tmp\Omega \\
& \quad \quad \quad \{0\} \tag{7.28}
\end{aligned}$$

$$\begin{aligned}
& > Vt0 := Annihilator([\omega 0, \omega 1, \omega 2, \omega 3]) [1] : Vt1 := Annihilator([\omega 0, \omega 1, \omega 2]) [2] : Vt2 \\
& \quad := Annihilator([\omega 0, \omega 1]) [3] : Vt3 := Annihilator([\omega 0]) [4] : \\
& \quad Vt01 := ContractIndices(TorsionTensor(CC), evalDG(Vt0 &t Vt1), [[2, 1], [3, 2]]) : \\
& > factor(Vt0) \\
& \quad - \left(\frac{(u_3 - u_5)(u_4 - u_5)(u_2 - u_5)}{(q_{[]} - u_5)^4} D_{x1} \right) - \left(\frac{(q_{[]} - u_2)^4 (u_4 - u_5)(u_3 - u_5)}{(q_{[]} - u_5)^4 (u_2 - u_4)(u_2 - u_3)} D_{x2} \right) \\
& \quad + \frac{(q_{[]} - u_3)^4 (u_4 - u_5)(u_2 - u_5)}{(u_2 - u_3)(q_{[]} - u_5)^4 (u_3 - u_4)} D_{x3} \\
& \quad - \left(\frac{(u_3 - u_5)(u_2 - u_5)(q_{[]} - u_4)^4}{(q_{[]} - u_5)^4 (u_3 - u_4)(u_2 - u_4)} D_{x4} \right) + D_{x5} \tag{7.29}
\end{aligned}$$

Thus there exist GL2-connections, with respect to which the null Lax surfaces are totally geodesic. The 4 parameters can be fixed uniquely by the condition $\text{Trace}(\text{Torsion})=0$.

$$\begin{aligned}
& > seq(factor(ContractIndices(Qua\omega[i], evalDG(Vt0 &t Vt0)), [[1, 1], [2, 2]])), i = 1 \\
& \quad \dots nops(Qua\omega) \\
& \quad \frac{q_{[]}^8 (u_4 - u_5)^2 (u_3 - u_5)^2 (u_2 - u_5)^2}{(q_{[]} - u_5)^8}, - \frac{4 \, q_{[]}^7 (u_4 - u_5)^2 (u_3 - u_5)^2 (u_2 - u_5)^2}{(q_{[]} - u_5)^8}, \\
& \quad \frac{28 \, q_{[]}^6 (u_4 - u_5)^2 (u_3 - u_5)^2 (u_2 - u_5)^2}{(q_{[]} - u_5)^8}, \tag{7.30}
\end{aligned}$$

$$\begin{aligned}
& - \frac{28 q_{[]}^5 (u_4 - u_5)^2 (u_3 - u_5)^2 (u_2 - u_5)^2}{(q_{[]} - u_5)^8}, \\
& \frac{70 q_{[]}^4 (u_4 - u_5)^2 (u_3 - u_5)^2 (u_2 - u_5)^2}{(q_{[]} - u_5)^8}, \\
& - \frac{28 q_{[]}^3 (u_4 - u_5)^2 (u_3 - u_5)^2 (u_2 - u_5)^2}{(q_{[]} - u_5)^8}, \\
& \frac{28 q_{[]}^2 (u_4 - u_5)^2 (u_3 - u_5)^2 (u_2 - u_5)^2}{(q_{[]} - u_5)^8}, - \frac{4 (u_4 - u_5)^2 (u_3 - u_5)^2 (u_2 - u_5)^2 q_{[]}}{(q_{[]} - u_5)^8}, \\
& \frac{(u_4 - u_5)^2 (u_3 - u_5)^2 (u_2 - u_5)^2}{(q_{[]} - u_5)^8}
\end{aligned}$$

> $QuT := evalDG(add(add(xi[i, j] \cdot dM[i] \&s dM[j], j = i..5), i = 1..5)) :$
 $Sqt := solve(\{coeffs(expand(numer(ContractIndices(QuT, Vt0 \&t Vt0, [[1, 1], [2, 2]]))),$
 $\{q_{[]}\}\}, \{seq(seq(xi[i, j], j = i..5), i = 1..5)\} : Sqt[10..15]$
 $\{\xi_{3,3} = \xi_{3,3}, \xi_{3,4} = \xi_{3,4}, \xi_{3,5} = \xi_{3,5}, \xi_{4,4} = \xi_{4,4}, \xi_{4,5} = \xi_{4,5}, \xi_{5,5} = \xi_{5,5}\}$ (7.31)

> $Vqt := DGsimplify(evalDG(subs(\{xi[3, 3] = \Lambda_1, xi[3, 4] = \Lambda_2, xi[3, 5] = \Lambda_3, xi[4, 4] = \Lambda_4,$
 $xi[4, 5] = \Lambda_5, xi[5, 5] = \Lambda_6\}, subs(Sqt, QuT)))) :$

> $EW := eliminate(DGinfo(DGsimplify(evalDG(subs(S3, Vqt - RicCcSymm)))) ,$
 $"CoefficientSet"), \{\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4, \Lambda_5, \Lambda_6\}) :$

This proves that symmetrized Ricci tensor of the canonical GL2-connection is a section of the bundle of quadrics annihilating the dual (tangent) to the canonical (cotangent) rational normal curve!

Here is the connection:

> CC (7.32)

>
>
>

II. Let us check if there exists a unique symmetric connection preserving GL2-structure

> $ChrSeq := \{seq(seq(seq(((a \parallel i) \parallel j) \parallel k), k = 1..5), j = i..5), i = 1..5)\} : nops(ChrSeq);$
 $CsC := Connection(subs(\{seq(seq(seq(((a \parallel i) \parallel j) \parallel k) = (((a \parallel j) \parallel i) \parallel k), k = 1..5), j = 1$
 $..(i - 1)), i = 1..5\}, add(add(add(((a \parallel i) \parallel j) \parallel k) \cdot vM[k] \&t dM[i] \&t dM[j], k = 1$
 $..5), j = 1..5), i = 1..5))) : TorsionTensor(CsC)$

75

$((0 D_x) dx) dx$ (7.33)

> $RNE := \{ \} : \text{for } i \text{ to } 5 \text{ do for } j \text{ to } 6 \text{ do } RNE := \{op(RNE),$
 $coeffs(expand(numer(simplify(subs(S2, subs(Qsyst,$
 $ToJet(ContractIndices(DirectionalCovariantDerivative(vM[i], Quadu[j], CsC), \omega$
 $\&t \omega, [[1, 1], [2, 2]]), \{u(x, y, t, z, s)\}, notation = jetnumbers))))), \{q_{[]}\}) : \text{od:od:}$

> nops(RNE)

227

(7.34)

> elRN := eliminate(RNE, ChrSeq) :

> elRN[2]

$$\{u_{1,1}, u_{1,2}, u_1 u_{1,1}, u_1 u_{1,2}, u_1 (3 u_1 u_{1,2} - 2 u_2 u_{1,1}), u_1 (3 u_1 u_{1,2} + u_2 u_{1,1}), u_1 (6 u_1 u_{1,2} - u_2 u_{1,1}), u_2 u_{1,1}, u_2 u_{1,2}, u_3 u_{1,1}, u_{1,1} (4 u_1^2 + u_3), u_1 u_2 u_{1,1}, u_1 u_3 u_{1,1}, u_1 u_{1,1} (3 u_1^2 + u_3), -2 u_1 u_3 u_{1,1}, 2 u_1 u_3 u_{1,1}, 3 u_1 u_{1,2} + u_2 u_{1,1}, u_1^2 u_{1,1} - 3 u_2 u_{1,2}, u_1^2 u_{1,1} + 3 u_2 u_{1,2}, u_1^3 u_{1,1} + 3 u_1 u_2 u_{1,2} - u_2^2 u_{1,1}, 12 u_1^2 u_{1,2} - u_1 u_2 u_{1,1} + 6 u_3 u_{1,2}, 6 u_1^3 u_{1,2} - u_1^2 u_2 u_{1,1} + u_2 u_3 u_{1,1}, 12 u_1^2 u_2 u_{1,2} - 3 u_1^2 u_3 u_{1,1} - 2 u_1 u_2^2 u_{1,1} - u_3^2 u_{1,1}, 6 u_1^3 u_{1,2} + 5 u_1^2 u_2 u_{1,1} + 6 u_2^2 u_{1,2} + u_2 u_3 u_{1,1}, 3 u_1^3 u_{1,2} - u_1^2 u_2 u_{1,1} - 3 u_1 u_3 u_{1,2} + u_2 u_3 u_{1,1}\} \quad (7.35)$$

$$\begin{aligned} &> \text{simplify}(\text{subs}(\{u_{1,1}=0\}, \{u_{1,1}, u_{1,2}, u_1 u_{1,1}, u_1 u_{1,2}, u_1 (3 u_1 u_{1,2} - 2 u_2 u_{1,1}), u_1 (3 u_1 u_{1,2} \\ &\quad + u_2 u_{1,1}), u_1 (6 u_1 u_{1,2} - u_2 u_{1,1}), u_2 u_{1,1}, u_2 u_{1,2}, u_3 u_{1,1}, u_{1,1} (4 u_1^2 + u_3), u_1 u_2 u_{1,1}, \\ &\quad u_1 u_3 u_{1,1}, u_1 u_{1,1} (3 u_1^2 + u_3), -2 u_1 u_3 u_{1,1}, 2 u_1 u_3 u_{1,1}, 3 u_1 u_{1,2} + u_2 u_{1,1}, u_1^2 u_{1,1} \\ &\quad - 3 u_2 u_{1,2}, u_1^2 u_{1,1} + 3 u_2 u_{1,2}, u_1^3 u_{1,1} + 3 u_1 u_2 u_{1,2} - u_2^2 u_{1,1}, 12 u_1^2 u_{1,2} - u_1 u_2 u_{1,1} \\ &\quad + 6 u_3 u_{1,2}, 6 u_1^3 u_{1,2} - u_1^2 u_2 u_{1,1} + u_2 u_3 u_{1,1}, 12 u_1^2 u_2 u_{1,2} - 3 u_1^2 u_3 u_{1,1} - 2 u_1 u_2^2 u_{1,1} \\ &\quad - u_3^2 u_{1,1}, 6 u_1^3 u_{1,2} + 5 u_1^2 u_2 u_{1,1} + 6 u_2^2 u_{1,2} + u_2 u_3 u_{1,1}, 3 u_1^3 u_{1,2} - u_1^2 u_2 u_{1,1} \\ &\quad - 3 u_1 u_3 u_{1,2} + u_2 u_3 u_{1,1}\})) \\ &\{0, u_{1,2}, u_1 u_{1,2}, u_2 u_{1,2}, 3 u_1 u_{1,2}, 3 u_1^2 u_{1,2}, 6 u_1^2 u_{1,2}, 6 u_1^3 u_{1,2}, -3 u_2 u_{1,2}, 3 u_2 u_{1,2}, \\ &\quad 3 u_1 u_2 u_{1,2}, 12 u_1^2 u_2 u_{1,2}, 6 u_1^3 u_{1,2} + 6 u_2^2 u_{1,2}, 12 u_1^2 u_{1,2} + 6 u_3 u_{1,2}, 3 u_1^3 u_{1,2} \\ &\quad - 3 u_1 u_3 u_{1,2}\} \end{aligned} \quad (7.36)$$

$$\begin{aligned} &> \text{simplify}(\text{subs}(\{u_{1,1}=0\}, \{u_{1,1}, u_1 u_{1,1}, u_2 u_{1,1}, (u_1^2 + u_3) u_{1,1}, u_1 u_2 u_{1,1}, u_1 (u_1^2 + u_3) u_{1,1}, \\ &\quad u_2 (u_1^2 + u_3) u_{1,1}, -8 u_1 u_{1,1}, -7 u_1 u_{1,1}, -6 u_1 u_{1,1}, -5 u_1 u_{1,1}, -3 u_1 u_{1,1}, -u_1 u_{1,1}, -5 \\ &\quad u_1^2 u_{1,1}, -u_1^2 u_{1,1}, -5 u_1 u_2 u_{1,1}, -u_1 u_{1,1} (6 u_1^2 + u_3), -5 u_1^2 u_2 u_{1,1}, -u_2 u_{1,1} (6 u_1^2 + u_3), \\ &\quad -16 u_1^2 u_{1,1} - u_3 u_{1,1}, -15 u_1^2 u_{1,1} - u_3 u_{1,1}, -12 u_1^2 u_{1,1} - u_3 u_{1,1}, -9 u_1^2 u_{1,1} - u_3 u_{1,1}, \\ &\quad -6 u_1^2 u_{1,1} - u_3 u_{1,1}, -5 u_1^3 u_{1,1} - 5 u_1 u_3 u_{1,1}, -6 u_1^4 u_{1,1} - 7 u_1^2 u_3 u_{1,1} - u_3^2 u_{1,1}, -21 \\ &\quad u_1^3 u_{1,1} - 6 u_1 u_3 u_{1,1} + 4 u_2^2 u_{1,1}, -6 u_1^3 u_{1,1} - u_1 u_3 u_{1,1} + u_2^2 u_{1,1}, -u_1^3 u_{1,1} - u_1 u_3 u_{1,1} \\ &\quad + 2 u_2^2 u_{1,1}\})) \\ &\{0\} \end{aligned} \quad (7.37)$$

> Sss1 := solve(RNE, ChrSeq)

Sss1 :=

(7.38)

So it does not exist in 5D (for dKP) as expected ...

>

III. Finally let us compute the canonical projective totally geodesic connection

> CsC := Connection(subs({seq(seq(seq(((a||i)||j)||k)=(((a||j)||i)||k), k=1..5), j=1..(i-1)), i=1..5}), add(add(add(((a||i)||j)||k)·vM[k]&t dM[i]&t dM[j], k=1

..5), j = 1 ..5), i = 1 ..5))) :

$$\begin{aligned} & \text{> } DD := \{YY, ZZ, TT, SS\} : \text{ChrCoe} := \{seq(seq(seq(((a \parallel i) \parallel j) \parallel k), k = 1 ..5), j = i ..5), i \\ & \quad = 1 ..5)\} : \\ & TGE := \{ \} : \text{for } i \text{ to } 4 \text{ do for } j \text{ to } 4 \text{ do } TGE := \{op(TGE), coeffs(expand(subs(S2, \\ & \quad subs(Qsyst, ToJet(Hook(convert(DirectionalCovariantDerivative(DD[i], DD[j], CsC), \\ & \quad 'DGvector'), \omega\omega), \{u(x, y, t, z, s), q(x, y, t, z, s)\}, notation = jetnumbers))\}), \{q_{[], q_1\}) \\ & \quad \} : \text{od:od:} \\ & [nops(ChrCoe), nops(TGE)] \end{aligned} \quad [75, 84] \quad (7.39)$$

$$\text{> } Sss2 := solve(TGE, ChrCoe \setminus \{a155, a255, a355, a455, a555\}) : nops(Sss2) \quad 70 \quad (7.40)$$

$$\begin{aligned} & \text{> } tmp := \{ \} : \text{for } i \text{ to } nops(Ss2) \text{ do if } lhs(Sss2[i]) = rhs(Sss2[i]) \text{ then } tmp := \{op(tmp), \\ & \quad lhs(Sss2[i])\} : \text{fi:od: } tmp \end{aligned} \quad \{ \} \quad (7.41)$$

$$\begin{aligned} & \text{> } Sss2 \\ & \left\{ \begin{aligned} & a111 = 2 a155, a112 = 0, a113 = 0, a114 = 0, a115 = 0, a121 = a255, a122 = a155, a123 \end{aligned} \right. \quad (7.42) \end{aligned}$$

$$\begin{aligned} & = 0, a124 = 0, a125 = 0, a131 = a355 - \frac{1}{2} u_{1,1}, a132 = 0, a133 = a155, a134 = 0, a135 \\ & = 0, a141 = a455 - u_{1,2}, a142 = -\frac{1}{2} u_{1,1}, a143 = 0, a144 = a155, a145 = 0, a151 = \\ & -u_1 u_{1,1} + \frac{1}{2} a555 - \frac{3}{2} u_{2,2}, a152 = -u_{1,2}, a153 = -\frac{1}{2} u_{1,1}, a154 = 0, a221 = u_{1,1}, \\ & a222 = 2 a255, a223 = 0, a224 = 0, a225 = 0, a231 = u_{1,2}, a232 = \frac{1}{2} u_{1,1} + a355, a233 \\ & = a255, a234 = 0, a235 = 0, a241 = \frac{3}{2} u_1 u_{1,1} + u_{2,2}, a242 = a455, a243 = \frac{1}{2} u_{1,1}, \\ & a244 = a255, a245 = 0, a251 = 2 u_1 u_{1,2} + \frac{3}{2} u_2 u_{1,1} + u_{2,3}, a252 = \frac{1}{2} u_1 u_{1,1} \\ & + \frac{1}{2} a555 - \frac{1}{2} u_{2,2}, a253 = 0, a254 = \frac{1}{2} u_{1,1}, a331 = -u_1 u_{1,1} + u_{2,2}, a332 = 2 u_{1,2}, \\ & a333 = 2 a355, a334 = 0, a335 = 0, a341 = -u_1 u_{1,2} - \frac{1}{2} u_2 u_{1,1} + u_{2,3}, a342 \\ & = \frac{1}{2} u_1 u_{1,1} + 2 u_{2,2}, a343 = a455 + u_{1,2}, a344 = a355, a345 = 0, a351 = -\frac{5}{2} u_1^2 u_{1,1} \\ & - u_1 u_{2,2} + u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + u_{3,3}, a352 = u_1 u_{1,2} + u_2 u_{1,1} + 2 u_{2,3}, a353 = \\ & -\frac{1}{2} u_1 u_{1,1} + \frac{1}{2} a555 + \frac{1}{2} u_{2,2}, a354 = u_{1,2}, a441 = u_1^2 u_{1,1} - u_1 u_{2,2} - 2 u_2 u_{1,2} \\ & + u_{3,3}, a442 = -2 u_1 u_{1,2} - u_2 u_{1,1} + 2 u_{2,3}, a443 = 2 u_1 u_{1,1} + 3 u_{2,2}, a444 = 2 a455, \\ & a445 = 0, a451 = u_1^2 u_{1,2} - u_1 u_{2,3} - 2 u_2 u_{2,2} - u_3 u_{1,2} + u_{3,4}, a452 = -\frac{3}{2} u_1^2 u_{1,1} \end{aligned}$$

$$\begin{aligned}
& -2 u_1 u_{2,2} - u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + 2 u_{3,3}, a453 = \frac{1}{2} u_2 u_{1,1} + 3 u_{2,3}, a454 = u_1 u_{1,1} \\
& + \frac{1}{2} a555 + \frac{3}{2} u_{2,2}, a551 = -u_1^3 u_{1,1} + u_1^2 u_{2,2} + 4 u_1 u_2 u_{1,2} - 2 u_1 u_3 u_{1,1} + u_2^2 u_{1,1} \\
& - u_1 u_{3,3} - 2 u_2 u_{2,3} - 3 u_3 u_{2,2} + u_{4,4}, a552 = 2 u_1^2 u_{1,2} - 2 u_1 u_{2,3} - 4 u_2 u_{2,2} \\
& - 2 u_3 u_{1,2} + 2 u_{3,4}, a553 = -4 u_1^2 u_{1,1} - 3 u_1 u_{2,2} - u_3 u_{1,1} + 3 u_{3,3}, a554 = 2 u_1 u_{1,2} \\
& + 2 u_2 u_{1,1} + 4 u_{2,3} \}
\end{aligned}$$

$$\Rightarrow pt := \{a155=0, a255=0, a355=0, a455=0, a555=0\} :$$

$$\Rightarrow Sss20 := \{op(subs(pt, Sss2)), op(pt)\}$$

$$Sss20 := \{a111=0, a112=0, a113=0, a114=0, a115=0, a121=0, a122=0, a123=0, \quad (7.43)$$

$$\begin{aligned}
& a124=0, a125=0, a131 = -\frac{1}{2} u_{1,1}, a132=0, a133=0, a134=0, a135=0, a141 = \\
& -u_{1,2}, a142 = -\frac{1}{2} u_{1,1}, a143=0, a144=0, a145=0, a151 = -u_1 u_{1,1} - \frac{3}{2} u_{2,2}, a152 \\
& = -u_{1,2}, a153 = -\frac{1}{2} u_{1,1}, a154=0, a155=0, a221 = u_{1,1}, a222=0, a223=0, a224 \\
& =0, a225=0, a231 = u_{1,2}, a232 = \frac{1}{2} u_{1,1}, a233=0, a234=0, a235=0, a241 \\
& = \frac{3}{2} u_1 u_{1,1} + u_{2,2}, a242=0, a243 = \frac{1}{2} u_{1,1}, a244=0, a245=0, a251 = 2 u_1 u_{1,2} \\
& + \frac{3}{2} u_2 u_{1,1} + u_{2,3}, a252 = \frac{1}{2} u_1 u_{1,1} - \frac{1}{2} u_{2,2}, a253=0, a254 = \frac{1}{2} u_{1,1}, a255=0, \\
& a331 = -u_1 u_{1,1} + u_{2,2}, a332 = 2 u_{1,2}, a333=0, a334=0, a335=0, a341 = -u_1 u_{1,2} \\
& - \frac{1}{2} u_2 u_{1,1} + u_{2,3}, a342 = \frac{1}{2} u_1 u_{1,1} + 2 u_{2,2}, a343 = u_{1,2}, a344=0, a345=0, a351 \\
& = -\frac{5}{2} u_1^2 u_{1,1} - u_1 u_{2,2} + u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + u_{3,3}, a352 = u_1 u_{1,2} + u_2 u_{1,1} \\
& + 2 u_{2,3}, a353 = -\frac{1}{2} u_1 u_{1,1} + \frac{1}{2} u_{2,2}, a354 = u_{1,2}, a355=0, a441 = u_1^2 u_{1,1} - u_1 u_{2,2} \\
& - 2 u_2 u_{1,2} + u_{3,3}, a442 = -2 u_1 u_{1,2} - u_2 u_{1,1} + 2 u_{2,3}, a443 = 2 u_1 u_{1,1} + 3 u_{2,2}, a444 \\
& =0, a445=0, a451 = u_1^2 u_{1,2} - u_1 u_{2,3} - 2 u_2 u_{2,2} - u_3 u_{1,2} + u_{3,4}, a452 = -\frac{3}{2} u_1^2 u_{1,1} \\
& - 2 u_1 u_{2,2} - u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + 2 u_{3,3}, a453 = \frac{1}{2} u_2 u_{1,1} + 3 u_{2,3}, a454 = u_1 u_{1,1} \\
& + \frac{3}{2} u_{2,2}, a455=0, a551 = -u_1^3 u_{1,1} + u_1^2 u_{2,2} + 4 u_1 u_2 u_{1,2} - 2 u_1 u_3 u_{1,1} + u_2^2 u_{1,1} \\
& - u_1 u_{3,3} - 2 u_2 u_{2,3} - 3 u_3 u_{2,2} + u_{4,4}, a552 = 2 u_1^2 u_{1,2} - 2 u_1 u_{2,3} - 4 u_2 u_{2,2} \\
& - 2 u_3 u_{1,2} + 2 u_{3,4}, a553 = -4 u_1^2 u_{1,1} - 3 u_1 u_{2,2} - u_3 u_{1,1} + 3 u_{3,3}, a554 = 2 u_1 u_{1,2}
\end{aligned}$$

$$+ 2 u_2 u_{1,1} + 4 u_{2,3}, a555 = 0 \}$$

> CsC0 := subs(Sss20, CsC) :

> DGsimplify(CsC0)

(7.44)

> $\Gamma_{131} = -\frac{1}{2} u_{1,1},$

$$\Gamma_{141} = -u_{1,2}, \Gamma_{142} = -\frac{1}{2} u_{1,1},$$

$$\Gamma_{151} = -u_1 u_{1,1} - \frac{3}{2} u_{2,2}, \Gamma_{152} = -u_{1,2}, \Gamma_{153} = -\frac{1}{2} u_{1,1},$$

$$\Gamma_{221} = u_{1,1},$$

$$\Gamma_{231} = u_{1,2}, \Gamma_{232} = \frac{1}{2} u_{1,1},$$

$$\Gamma_{241} = \frac{3}{2} u_1 u_{1,1} + u_{2,2}, \Gamma_{243} = \frac{1}{2} u_{1,1},$$

$$\Gamma_{251} = 2 u_1 u_{1,2} + \frac{3}{2} u_2 u_{1,1} + u_{2,3}, \Gamma_{252} = \frac{1}{2} u_1 u_{1,1} - \frac{1}{2} u_{2,2}, \Gamma_{254} = \frac{1}{2} u_{1,1},$$

$$\Gamma_{331} = -u_1 u_{1,1} + u_{2,2}, \Gamma_{332} = 2 u_{1,2},$$

$$\Gamma_{341} = -u_1 u_{1,2} - \frac{1}{2} u_2 u_{1,1} + u_{2,3}, \Gamma_{342} = \frac{1}{2} u_1 u_{1,1} + 2 u_{2,2}, \Gamma_{343} = u_{1,2},$$

$$\Gamma_{351} = -\frac{5}{2} u_1^2 u_{1,1} - u_1 u_{2,2} + u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + u_{3,3}, \Gamma_{352} = u_1 u_{1,2} + u_2 u_{1,1}$$

$$+ 2 u_{2,3}, \Gamma_{353} = -\frac{1}{2} u_1 u_{1,1} + \frac{1}{2} u_{2,2}, \Gamma_{354} = u_{1,2},$$

$$\Gamma_{441} = u_1^2 u_{1,1} - u_1 u_{2,2} - 2 u_2 u_{1,2} + u_{3,3}, \Gamma_{442} = -2 u_1 u_{1,2} - u_2 u_{1,1} + 2 u_{2,3}, \Gamma_{443} = 2 u_1 u_{1,1} + 3 u_{2,2},$$

$$\Gamma_{451} = u_1^2 u_{1,2} - u_1 u_{2,3} - 2 u_2 u_{2,2} - u_3 u_{1,2} + u_{3,4}, \Gamma_{452} = -\frac{3}{2} u_1^2 u_{1,1} - 2 u_1 u_{2,2} - u_2 u_{1,2} - \frac{1}{2} u_3 u_{1,1} + 2 u_{3,3}, \Gamma_{453} = \frac{1}{2} u_2 u_{1,1} + 3 u_{2,3}, \Gamma_{454} = u_1 u_{1,1} + \frac{3}{2} u_{2,2},$$

$$\Gamma_{551} = -u_1^3 u_{1,1} + u_1^2 u_{2,2} + 4 u_1 u_2 u_{1,2} - 2 u_1 u_3 u_{1,1} + u_2^2 u_{1,1} - u_1 u_{3,3} - 2 u_2 u_{2,3}$$

$$- 3 u_3 u_{2,2} + u_{4,4}, \Gamma_{552} = 2 u_1^2 u_{1,2} - 2 u_1 u_{2,3} - 4 u_2 u_{2,2} - 2 u_3 u_{1,2} + 2 u_{3,4}, \Gamma_{553} = -4 u_1^2 u_{1,1} - 3 u_1 u_{2,2} - u_3 u_{1,1} + 3 u_{3,3}, \Gamma_{554} = 2 u_1 u_{1,2} + 2 u_2 u_{1,1} + 4 u_{2,3}$$